Testing hypotheses of habitat use and temporal activity in relation to body plan in a Mediterranean lizard community

Journal:	<i>Canadian Journal of Zoology</i>
Manuscript ID	cjz-2021-0083.R2
Manuscript Type:	Article
Date Submitted by the Author:	20-May-2021
Complete List of Authors:	Rugiero, Lorenzo; Institute for Development Ecology Conservation and Cooperation Capula, Massimo; Museo Civico di Zoologia Dendi, Daniele; Institute for Development Ecology Conservation and Cooperation; Rivers State University of Science and Technology, Department of Applied and Environmental Biology; University of Lomé, Department of Zoology Petrozzi, Fabio; Ecolobby Fa, Julia; Manchester Metropolitan University, Department of Natural Sciences; Center for International Forestry Research (CIFOR) Funk, Stephan M.; NatureEritage Burke, Russell; Hofstra University, Department of Biology Luiselli, Luca; Institute for Development Ecology Conservation and Cooperation; Rivers State University of Science and Technology, Department of Applied and Environmental Biology; University of Lomé, Department of Zoology
Is your manuscript invited for consideration in a Special Issue?:	Not applicable (regular submission)
Keyword:	Sauria, Mediterranean, resource partitioning, bauplan, autoecological constraints, eco-physiological attributes

SCHOLARONE™
Manuscripts

1 **Testing hypotheses of habitat use and temporal activity in relation to body plan**
2 **in a Mediterranean lizard community**

3

4 Lorenzo Rugiero¹, Massimo Capula², Daniele Dendi^{1,3,4}, Fabio Petrozzi⁵, Julia E. Fa^{6,7}, Stephan M.
5 Funk⁸, Russell L. Burke⁹, Luca Luiselli^{1,3,4}

6

7 1 Institute for Development, Ecology, Conservation and Cooperation, via G. Tomasi di
8 Lampedusa 33, I-00144 Rome, Italy; lore.rho@gmail.com; d.dendi@ideccngo.org;
9 l.luiselli@ideccngo.org

10 2 Museo Civico di Zoologia, Rome, Italy; m.capula@comune.roma.it

11 3 Department of Applied and Environmental Biology, Rivers State University of Science and
12 Technology, P.M.B. 5080, Port Harcourt, Nigeria

13 4 Department of Zoology, University of Lomé, Lomé, Togo; lucamaria.luiselli@uniroma3.it
14 5 Ecolobby, Rome, Italy; fapetrozzi@gmail.com

15 6 Department of Natural Sciences, School of Science and the Environment, Manchester
16 Metropolitan University, Manchester, M1 5GD, UK; jfa949@gmail.com

17 7 Center for International Forestry Research (CIFOR), CIFOR Headquarters, Bogor, 16115,
18 Indonesia.

19 8 Nature Heritage, Jersey, Channel Islands, UK; smf@natureheritage.org

20 9 Department of Biology, Hofstra University, New York, USA; Russell.l.burke@hofstra.edu

21

22

23

24 Abstract

25 A body plan (= bauplan) is a suite of morphological characters shared by phylogenetically related
26 animals at some point during their development. Despite its value, the bauplan concept is still rarely
27 employed to characterize functional groups in community ecology. Here, we examine habitat use
28 and spatio-temporal activity correlates of an entire seven-species community of lizards with
29 different bauplans. The study was carried out in three locations in central Italy, encompassing a
30 complex landscape with a patchy mosaic of a wide variety of habitats and microclimates. We tested
31 four hypotheses regarding niche breadth, habitat use and activity patterns. The first hypothesis,
32 niche complementarity, in which species with similar body shapes should non-randomly partition
33 available habitats, was not supported. By contrast, the hypotheses that larger-bodied species should
34 have a wider niche breadth, that slower species should inhabit habitat types of higher cover, and
35 species inhabiting open sunny habitats should exhibit more seasonally variable activity patterns,
36 were all supported by the data. Sympatric lizard communities in our study area were clearly
37 organized by autecological constraints and eco-physiological attributes.

38
39 **Key words:** Sauria; *Lacerta bilineata*; *Podarcis muralis*; *Podarcis sicula*; *Hemidactylus turcicus*;
40 *Tarentola mauritanica*; *Anguis veronensis*; *Chalcides chalcides*; Mediterranean; resource
41 partitioning; bauplan; autecological constraints; eco-physiological attributes

43 Introduction

44 In evolutionary biology, a suite of characters shared by a group of phylogenetically related
45 animals at some point during their development is defined as body plan or “bauplan” (Woodger
46 1945; Rieppel 2006; Willmore 2012). The concept has represented an important element in
47 evolutionary developmental biology, evo-devo (Tsessarsky 2020), and the evolutionary ecology of
48 organisms (e.g., Hall 1999; Willmore 2012). However, bauplan has not been explicitly used in
49 community ecology studies, despite this being a central concept for distinguishing the various

guilds within truly “functional groups”. Most often, community ecology studies define the various guilds either in terms of phylogenetic units (for instance, varanids versus agamids, etc.) or “natural history-characterized” units (for instance, terrestrial versus arboreal, etc.). Thus, the concept of bauplan may be the ideal synthesis between phylogenetic and natural history characteristics (Stankowich and Stensrud 2019) and, therefore, particularly useful for determining assembly rules of biotic communities. To our knowledge, bauplan has not been explicitly used in community ecology studies.

Lizards are traditionally the most popular model organisms in community ecology studies (e.g., Pianka 1986; Flesch et al. 2017; Grundler et al. 2017; Jiménez-Robles and De la Riva 2019). This group of terrestrial vertebrates are particularly suited for this kind of study because they are easily observed, are found in a large range of temperate and tropical ecosystems, and often exhibiting a remarkable array of species diversity (Helmer et al. 1989; Maura et al. 2011; Zakkak et al. 2015; Simbula et al. 2019; Vacheva et al. 2020). Although sympatric lizard species often display patterns of niche partitioning along the trophic axis (Luiselli 2008; Sheu et al. 2020), the spatial dimension of the niche is instead the usual primary partitioning axis (Toft 1985; Arnold 1987; Gonçalves-Sousa et al. 2019; Sillero et al. 2020). Given this, the study of habitat use patterns by sympatric lizards can greatly contribute to ecological community theory (Arnold 1987; Gonçalves-Sousa et al. 2019; Sillero et al. 2020). For instance, previous data demonstrated remarkable discrepancies in the habitat selection patterns even within the same lizard clades, with some studies documenting a clear ontogenetic shift in habitat use (e.g., Jenssen et al. 1998) whereas others failed in finding any ontogenetic effect on structural niche use (e.g., Powell and Russell 1992).

Like Caribbean *Anolis* Daudin, 1802 lizards (Losos 2011; Pringle et al. 2019) and Australian desert lizards (Pianka 1986), European lizards offer an interesting and logistically convenient system for testing predictions of community ecology theory (e.g., Sillero et al. 2020). These animals

are often abundant, conspicuous, and approachable, and therefore it is often easier to observe and record data on their spatial ecologies than for less abundant and more elusive animals.

Microhabitat features useful in studying the ecology of European lizards can be divided into structural and climatic (Arnold 1987). Structural features include whether the microhabitat is flat or elevated, whether its surfaces are more or less continuous or broken, the nature of the substrate (e.g., rock, stones, vegetation) and the types of refuges used. These are responsible for much of the apparent differences in species' spatial distribution, for instance, species that climb high have elevated scores for occurrence on rock or its functional equivalents, and for using crevices as refuges (Arnold 1987). Climatic features include humidity, temperature and shadiness of the habitats and are often inter-correlated with structural features. These are more important in large scale studies of niche ecology than in smaller scale studies (Escoriza and Amat 2021).

In the present study we examined habitat use and temporal activity correlates of an entire community of lizards in three locations in central Italy (Tolfa Mountains, province of Rome). These study sites consist of a complex landscape composed of habitat patches that include a variety of environmental typologies typical of the area. Habitats range from open dry Mediterranean maquis to habitat types with damp vegetation, tall trees with cooler climate and higher humidity. Using this wide range of habitat features potentially available to lizards, we explored the spatio-temporal patterns of lizard community structure. This community is composed of species belonging to three main types of bauplan: type (i) was a typical "lacertid bauplan" with long legs, long tail and slender body, allowing them to run quickly on the ground but also climb efficiently on vertical substrates (three species: the larger sized *Lacerta bilineata* Daudin, 1802 and the smaller-sized *Podarcis muralis* Laurenti, 1768 and *Podarcis sicula* Rafinesque, 1810); type (ii) was a typical "gecko bauplan" with adhesive toe pads allowing them to climb well even on vertical surfaces (two species: *Hemidactylus turcicus* (Linnaeus, 1758) and *Tarentola mauritanica* (Linnaeus, 1758)); type (iii)

was an elongated, slender “snake-like” bauplan, allowing them to only move on ground and burrow underground (two species: *Anguis veronensis* Pollini, 1818, *Chalcides chalcides* Linnaeus, 1758).

Using this suite of sympatric species with different bauplans, we analysed four explicit hypotheses. The “niche complementarity” theory (Schoener 1974) which suggests that bauplans that use similar structural habitats tend to differ in climatic requirements and vice versa, to minimize potential interspecific competition. In lizards, species with similar bauplans tend to select structurally similar habitats (e.g., Arnold 1987) and hence potentially compete when resources are limited (Toft 1985). Therefore, we hypothesized that species with similar bauplans should non-randomly partition available habitats, showing less overlap in habitat niche overlap than for species belonging to different bauplans. For instance, the different species belonging to type (i) should differ in shadiness/vegetation cover of the habitats they occupy, with some species being linked to open sunny habitats and others to closed wet habitats. Again, the same habitat partitioning pattern should occur between geckos (type (ii)) or between the “elongated lizards” (type (iii)). For instance, in Salamanca (Spain) the lizard community was distributed in clusters, with species of the same genus (same bauplan according to our criteria presented herein) segregated almost totally by inhabiting different habitats, whereas species of different genera presented partial segregation, sharing some habitats (Sillero and Gomes 2016). In the latter study, ground-dwelling species showed partial spatial segregation whereas the saxicolous species presented a high degree of spatial segregation (Sillero and Gomes 2016). If the niche complementarity theory is supported, we predicted that (a) the habitat niche overlap between species belonging to a same bauplan group should be significantly lower than that observed between species of different bauplan groups, and (b) an evidence of a community structure compatible with non-random niche partitioning should be detected.

125 As a second hypothesis, we tested whether the species with largest body size exhibited a
126 wider habitat niche breadth in the patchy mosaic landscape than smaller sized species. This
127 hypothesis is based on the fact that, in lizards, home range sizes are generally positively correlated
128 with individual's body size (e.g., Turner et al. 1969; Christian and Waldschmidt 1984) and therefore
129 we predicted that the space requirements of larger bodied species exceeded the space available in
130 habitat types within the relatively narrow mosaics of woods, maquis and pastures in the study area.
131 Thus, in our study case, we predicted that the largest bodied species, *Lacerta bilineata*, would
132 exhibit a wider habitat niche breadth than the other, smaller-sized species, as it has much larger
133 individual home range size than the other species (Saint Girons and Bradshaw 1989).

134

135 Lizards represent important prey for a variety of predators in Mediterranean ecosystems
136 (carnivores, birds of prey and snakes; e.g., Rugiero et al. 1995; Capizzi and Luiselli 1996), and
137 thereby rely on running speed and/or cryptic colourations/behaviours to avoid predation
138 (Vanhooydonck and Van Damme 2003). As a third hypothesis, we predicted that lizard species with
139 slower movements and cryptic colouration would inhabit more closed habitats than those that are
140 able to run more quickly for antipredator reasons. In our study case, *Anguis veronensis* is much
141 slower in movements than any other lizard species in the assemblage, and therefore we predicted
142 that it should inhabit habitats that are much denser in vegetation than any other species.

143

144 In Mediterranean environments, the high ambient temperatures (> 35°C) typical during
145 summers may represent a constraint for reptile above-ground activities (Carretero and Llorente
146 1995; Rouag et al. 2007; Zamora-Camacho et al. 2013; Bouam et al. 2016). As a fourth hypothesis
147 we predicted that those species inhabiting open sunny habitats would exhibit more seasonally
148 variable activity patterns, with peaks during the early spring months, and scarce above-ground
149 activity in the summer (Burke and Ner 2005). By contrast, species that are very linked to closed and
150 cooler habitats would show a more constant above-ground activity pattern throughout the year.

151 Thus, the number of observed individuals by month should be seasonally more variable in the
152 species from open-sunny habitats than those in more vegetated and cooler habitats.

153
154 Using a suite of statistical procedures including univariate, multivariate and null model
155 (Monte Carlo methods) analyses, we tested each of the four hypotheses presented above to describe
156 the main “functional” characteristics of the studied lizard community.

157

158 **Materials and Methods**

159 *Study area*

160 Field data were collected in a woodland area surrounding the villages of Manziana, Oriolo Romano
161 and Canale Monterano (approximate coordinates: E 12° 05' N 42° 06'), about 50 km northwest of
162 Rome and just outside west of the regional natural park Bracciano-Martignano (Latium, Central
163 Italy). The study area was a mixed oak woodland with *Quercus cerris* Linnaeus, 1753 and *Q.*
164 *frainetto* Ten, 1819 as dominant species, and with open grasslands surrounding the wooded patches.
165 In the drier parts of the forest, the trees were mainly *Quercus ilex* Linnaeus, 1753; *Acer campestre*
166 Linnaeus, 1753 and *Tilia* sp. were also common inside the main forest patches. *Rubus ulmifolius*
167 Schott, 1818, *Rubus Cesium* Linnaeus, 1753, *Cytisus scoparius* Linnaeus and Link, 1822, *Smilax*
168 *aspera* Linnaeus, 1753, *Pteridium aquilinum* L. (Kuhn, 1753), *Prunus spinosa* Linnaeus, 1753,
169 *Hedera helix* Linnaeus, 1753, *Crataegus monogyna* Jacq, 1775, *Rosa canina* Linnaeus, 1753 and
170 *Sorbus domestica* Linnaeus, 1753 were the most common undergrowth species.

171

172 *Monitoring*

173 Details of the monitoring protocol were described by Rugiero et al. (2021) for *L. bilineata*. In
174 the present paper, we also include the dataset presented in Rugiero et al. (2021). Monitoring was
175 conducted during April-October 1991, 1992, 1993, 1996, and 1997. On each monitoring day, two or
176 three researchers walked independently along haphazard transects without revisiting the same spots

177 to avoid multiple sightings of the same individuals and, thus, statistical pseudoreplication (Hurlbert
178 1984). No fixed distance occurred between transects and there was no fixed time or fixed number of
179 walked transects daily. Data collection was restricted to sunny days between 0900 to 1600 hrs to
180 minimize the impact of inter-daily weather conditions. We observed lizards without any
181 interference such as capture. We considered only those individuals that were observed active above-
182 ground, i.e., not under stones, tree branches or other ground objects. We identified observed lizards
183 by species, sex based on external features for those species that can be distinguished by them (not
184 for the two gecko species and *C. chalcides*) and age class (adult or juvenile), based on both body
185 size and, for some species (*L. bilineata*, *P. muralis*), dorsal coloration. For each sighted lizard we
186 recorded a habitat category defined by the dominance of a specific bushy plant taxon. Six habitat
187 types were distinguished (Figure 1):

- 188 (1) *Rubus* spp., that was the most wet habitat available to lizards at the study area and had in
189 May a vegetation cover (established at 300 random 1 x 1 m spots by eye) = $72.3 \pm 21.4\%$;
- 190 (2) *Cytisus scoparius*, with a vegetation cover = $42.1 \pm 33.1\%$;
- 191 (3) *Spartium junceum* Linnaeus, 1753, with a vegetation cover = $38.7 \pm 11.1\%$;
- 192 (4) open grassland with no bushy species, with a vegetation cover = $12.1 \pm 3.3\%$;
- 193 (5) stony wall (locally called “muretti a secco”), with a vegetation cover = $52.1 \pm 43.2\%$. This
194 habitat was the only human-made feature available to lizards at the study areas;
- 195 (6) *Smilax aspera*, with a vegetation cover = $59.4 \pm 27.1\%$.

196 The vegetation cover calculations at each site were always undertaken by the same person;
197 cover percentages were assessed at the moment of sighting of individual lizards. The relative
198 availability of the various habitat types at the study areas is summarized in Table 1.

199 Mean body mass (g) of the various species was obtained from measurements made by one of
200 us (MC) during previous studies on the genetics of these species (e.g., Capula and Ceccarelli 2003)
201 and from the literature (e.g., Ferrandino et al. 2001 for *Chalcides chalcides*). Also, no individual
202 animals were killed for the genetic studies that preceded the current research.

203

204

205

206 *Statistical procedures*

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

Niche breadths of habitats used by species and, for those species in which it was possible to discriminate sexes, by sex, were assessed by Simpson's (1949) index (B_S) and by a standardized equation transformed from Levins' (1968) formula (B_L) with its values ranging from 0 (maximum specialisation) to 1 (maximum opportunism) (Luiselli 1992).

To verify whether one sex had, independently for each species, a higher habitat niche breadth than the other (hypothesis that can be anticipated because home ranges in reptiles are generally much larger in males than in females; see Rocha 1999), we pooled the various species and examined their average habitat niche breadth. We also examined the habitat breadth hypothesis by season, using spring (April, May and June), summer (July, August) and autumn (September and October) as season categories.

A non-metric multidimensional scaling (nMDS) analysis with the cosine distance, using PAST 4.3 software (<https://www.nhm.uio.no/english/research/infrastructure/past/>) was carried out to cluster lizard individuals by sex, age class and species in relation to habitat availability, with the centroids being reported in the output of the analyses for the various species, individual categories and habitat availability.

For the simulations study, datasets were inspected to determine the nonrandom structure in the studied lizard community by contrasting the actual data matrix as given in the original literature source (or as in the original data) with random 'pseudo-communities' generated by Monte Carlo simulations (Gotelli and Graves 1996). We used the EcoSim software (Aquired Intelligence Corp., Kesey-Bear, VT, USA) to calculate habitat niche overlap indices between sympatric species and to generate Monte Carlo simulations. For each lizard sighting, we parameterized resource (habitat)

items data as presence versus absences. As too many zeroes in the matrices can distort error levels while too often rejecting structure, we fixed zeroes prior to any analyses (Pianka 1986). Pianka's (1986) overlap formula was calculated for all lizard's group combinations; then, the original species utilization matrices were randomized by shuffling the original values among the resource states (habitats). We used three randomization algorithms RA2, RA3 and RA4 of Lawlor (1980) as they have been shown to be particularly robust for niche overlap studies (Gotelli and Graves 1996). RA2 explores the assemblage structure in the generalist–specialist nature of the resource utilization matrix by conserving guild structure, but destroying observed niche breadth (Gotelli and Graves 1996). RA3 explores the guild structure by conserving niche breadth for each species, but destroying guild structure manifested by the zero structure of the resource utilization matrix (Gotelli and Graves 1996). RA4 retains both the niche breadth of the lizard species and the zero states in the resource utilization matrix, so among the lizards only the original non-zero electivities were randomly reassigned among the set of resource states originally used by that consumer (Lawlor 1980). For each pair of species, 30,000 random Monte Carlo permutations were generated. This number of permutations is enough to avoid biases in the results in calculations (Lehsten and Harmand 2006). Niche overlap values were calculated for each randomly generated matrices, and species-pair and community-summary statistics were computed (Friggens and Brown 2005). Actual overlap values were then compared to the distributions of the expected values, with the nonrandom structure being assumed when $P(\text{obs} < \text{exp}) = 0.05$ or less either with RA2, RA3 or RA4 (Gotelli and Graves 1996). In all cases, we define the resource use based on its availability (%) in the field.

For the seasonal analyses, we processed the data independently for spring (April, May and June), summer (July, August) and autumn (September and October). Contingency table analysis by χ^2 tests was carried out to analyze the frequency differences of lizard individuals observed by season and by species. Pearson's correlation coefficient was used to analyse the relationship between lizard

body mass and habitat niche breadth, and between the observed sample sizes per species and the habitat niche breadth. In the text, means are presented ± 1 Standard Deviation, with alpha set at 5%.

Results

General considerations

A total of 7257 lizard sightings, belonging to seven distinct species, were observed in the study area (Table 1). The commonest species was *P. muralis* (n = 4708 sightings), followed by *P. sicula* (n = 1706), *L. bilineata* (n = 397) and *C. chalcides* (n = 394). The other species were remarkably less common in the study area: *T. mauritanica* was seen only 31 times, *H. turcicus* 11 times and *A. veronensis* 10 times. Lacertidae species accounted for 93.8% of the total number of observed lizard individuals. A nMDS plot arranged the various groups of lizard individuals in clearly distinct sectors of the multidimensional space by sex, age class and species (Figure 2). The coordinate 1 explained 87.3% of the overall variance, and coordinate 2 explained 11.9%. *Chalcides chalcides*, *P. sicula* and *L. bilineata* juveniles were joined in the same cluster; *P. muralis* and *A. veronensis* were separate from each other; *L. bilineata* females clustered very close with the overall habitat availability; *T. mauritanica* and *H. turcicus* were separated from the other clusters but did not cluster jointly in the multidimensional space.

Hypothesis 1: the “niche complementarity” theory

Overall, there were no differences between the two groups (same bauplan, mean niche overlap = 0.57 ± 0.37 ; different bauplan, mean niche overlap = 0.51 ± 0.28 ; $t = 0.433$, $P = 0.669$). The three random reorganization algorithms indicated inconsistent patterns of variation in mean habitat overlaps from the overlaps actually observed for the seven species along the six habitat variables: in RA3 the observed overlap was significantly higher than the mean simulated value, whereas in RA4 the observed overlap was significantly lower, and in RA2 it was random (Table 2).

280 Thus, no evidence of a community structure compatible with overall interspecific competition was
281 detected from our analyses. Therefore, the hypothesis that species with the same bauplan should
282 non-randomly partition the available habitats and show less overlap in habitat niche overlap than
283 species with different bauplans was not confirmed by our data.

284

285 In *L. bilineata*, the habitat niche overlap was very high between males and females ($O =$
286 0.929), low between males and juveniles ($O = 0.343$), and intermediate between females and
287 juveniles ($O = 0.554$). In *P. sicula*, all the individual categories exhibited extremely high habitat
288 niche overlaps: $O = 0.998$ between males and females, $O = 0.981$ between males and juveniles and
289 $O = 0.989$ between females and juveniles. In *P. muralis*, the trend was very similar as that of *P.*
290 *sicula*, with very high overlap between males and females ($O = 0.994$), males and juveniles ($O =$
291 0.970) or females and juveniles ($O = 0.939$). For the other sympatric species it was impossible to
292 calculate these intraspecific overlap values because we were unable to sex them in the field.

293

294 *Hypothesis 2: larger species have wider niche breadth*

295 The largest species in the assemblage (*L. bilineata*) showed the widest habitat niche breadth
296 ($B_L = 0.38$), followed by *P. muralis* ($B_L = 0.28$) and *P. sicula* ($B_L = 0.22$). The other four species had
297 a considerably narrower niche breadth: *A. veronensis* ($B_L = 0.16$), *H. turcicus* ($B_L = 0.09$), *C.*
298 *chalcides* ($B_L = 0.08$), and *T. mauritanica* ($B_L = 0$). These niche breadth values were not associated
299 with respective sample sizes for each species ($r = 0.462$, $n = 7$, $P = 0.297$), but were significantly
300 dependent on the relative body size of each species ($r = 0.857$, $r^2 = 0.734$, $n = 7$, $P = 0.0018$). The
301 niche breadth values, by category of individuals within each species, are given in Table 3. Pooling
302 the various species, males had on average niche breadth values similar to females ($x = 0.31 \pm 0.08$
303 versus 0.29 ± 0.25), so it may concluded that species body size, and not the sex, significantly
304 influenced the habitat niche breadth of these lizards. Overall, the null hypothesis 2 of larger species
305 have no wider niche breath was rejected, thus, hypothesis 2 was supported by our field data. The

niche breadth values did not vary by seasons (in all cases, $P > 0.05$), and are, therefore, not further presented here.

Hypothesis 3: the slower species inhabit the most covered habitat types

At our study areas, the frequency of occurrence of the various species in the two habitats with densest vegetation (*Rubus* and *Smilax*) differed significantly from equality (observed-versus-expected $\chi^2 = 136.8$, $df = 6$, $P < 0.0001$), with the slowest/most cryptically coloured species (*A. veronensis*) being observed in these two habitats much more frequently (50% of the cases) than any other species (in descending order: *P. muralis* 38%, *L. bilineata* 28%, *P. sicula* 10.4%, *C. chalcides* 1.5%, *T. mauritanica* and *H. turcicus* 0%). Overall, our data rejected null hypothesis 3 of no impact of species' speed, thus, supporting the hypothesis 3.

Hypothesis 4: the species inhabiting open sunny habitats exhibit more seasonally variable activity patterns

The percentage of individuals observed by season, for each of the study species, are given in Figure 3. Contingency table analysis revealed that there were significant differences among species ($\chi^2 = 1262$, $df = 24$, $P < 0.0001$), with *L. bilineata* and *C. chalcides* being mostly observed in the springtime and the two *Podarcis* species more regularly throughout the year. If we consider the species that occurred more frequently in the most open-sunny habitat (open grasslands) versus those that occurred more frequently in the most covered habitats (i.e., *C. chalcides* versus *A. veronensis*), there were significant differences ($\chi^2 = 135.7$, $df = 2$, $P < 0.0001$), with the former species showing a strong seasonality (peak of activity during springtime; sequential $\chi^2 = 274.8$, $df = 2$, $P < 0.0001$) and the latter showing no significant seasonality in its activity patterns ($\chi^2 = 4.6$, $df = 2$, $P = 0.148$).

331 (Figure 3). Thus, the null hypothesis of no impact of sunnyness was rejected, thus supporting our
332 hypothesis 4.

333

334 **Discussion**

335 *General considerations*

336 The studied lizard assemblages were characterized by a clear prevalence (over 90%) of
337 Lacertidae species in terms of frequency of observed individuals: the two species of *Podarcis* were
338 the most common, followed by *L. bilineata*, which is also the largest bodied of the seven saurians
339 species present. Among the non-Lacertidae species, only *C. chalcides* were recorded frequently,
340 while the remaining three species were clearly more rarely observed. These general frequency
341 differences among lizard families mirror previous data from Mediterranean communities (e.g.,
342 Rugiero 2004; Santos and Poquet 2010), that are often inhabited by a remarkable variety of
343 sympatric Lacertidae (Pérez Mellado 1982; Zakkak et al. 2015). The nMDS analysis showed that
344 two clear groups of species were recognizable: one group of "open" areas, consisting of *C.*
345 *chalcides*, *P. sicula* and *L. bilineata* juveniles, and one group of "close vegetation" consisting of *P.*
346 *muralis* and *A. veronensis*, with the other species being not clearly assignable to any of these two
347 well-defined groups of taxa. This result appears to mirror the known habitat characteristics that
348 have been described for these species (e.g., Corti et al. 2011), however this is not sufficient per se to
349 define whether the various species co-occurrences in the various habitats were determined
350 essentially by ecological constraints ("bauplan" characteristics mediated by climatic and structural
351 factors sensu Arnold 1987) or by interspecific relationships (niche partitioning, e.g., Luiselli 2008).
352 However, the four hypotheses tested in the present paper allow us to make some inferences on the
353 reasons behind the observed habitat use patterns by species at the study areas in Mediterranean
354 central Italy.

355

356 *Testing the four hypotheses*

357 In our first hypothesis, we predicted that habitat niche overlap between species with the
358 same bauplans should be significantly lower than the overlaps between species of different bauplan
359 groups. Thus, we hypothesized that species with a same bauplan would have non-randomly
360 partitioned the habitat types. We found no consistent support for either of these predictions, indeed,
361 patterns of habitat use overlap were not associated with bauplan. In *Anolis* lizards, species that use
362 similar habitats tend to evolve similar body plans (i.e., ecomorphological convergence, Williams
363 1983; Losos et al. 1998). In cases where species with similar body plans overlap in habitat use, they
364 may partition resources in other ways, such as temporally or along other resource axes (Luiselli
365 2008). Thus, our observations of a central Italian lizard community are not similar to those of
366 *Anolis* lizards, perhaps because many tropical communities have higher species richness and
367 competition can be quite intense (e.g., on small islands, Calsbeek and Cox 2010).

368
369 Concerning hypothesis 2, the wider habitat niche breadth of the largest bodied species (*L.*
370 *bilineata*) is interesting because in this species we detected a significant diversification of habitat
371 use in relation to age. The strong propensity of juveniles to use open environments, which led them
372 to be classified in the group also consisting of *P. sicula* and *C. chalcides*, changes ontogenetically,
373 with adults being significantly more generalized (Rugiero et al. 2021). This is a further confirmation
374 that it is the body size, and not other ecological characteristics, that is strongly associated with the
375 wider habitat niche breadth of this species. Furthermore, habitat niche breadth was nearly identical
376 in 1-2 years-old juvenile *L. bilineata* population cohorts in comparison to similarly-sized lizards of
377 other species. Intraspecific competition avoidance and cannibalism appear to be the main reasons
378 pushing juveniles to minimize habitat overlap with adults in the large-sized *L. bilineata* at these
379 study areas (Rugiero et al. 2021). The same pattern was also recorded from *Anolis* lizards, with
380 adult male density having significant effects on juvenile perch height, perch width, and substrate
381 use, thus suggesting that strong age-class competition may contribute to the ontogenetic differences
382 in habitat choice (Delaney and Warner 2017a, 2017b).

383

384 Our third hypothesis, that lizard species with slower movements and cryptic colouration
385 would inhabit more closed habitats, was supported in that *Anguis veronensis* was observed in the
386 two habitat types with the densest vegetations significantly more frequently than was any other
387 species. However, the patterns of dense habitat use for other lizard species did not indicate
388 decreasing movement speed and crypticity. It is possible that this relationship is non-linear, and
389 the benefits of dense vegetation are outweighed by other factors for lizards of a threshold movement
390 speed and/or crypticity.

391

392 Our fourth hypothesis was also related to vegetation density; we predicted that lizard species
393 that use habitats with denser vegetation would show more temporally consistent, unseasonal
394 activity, whereas those that use more open habitats would exhibit more seasonally variable activity.
395 We found that lizard species that occurred primarily in open habitat had distinct seasonal activity
396 peaks, while those that occurred primarily in densely vegetated habitats had unseasonal activity
397 patterns. This interesting result should be examined more fully with detailed studies of energy
398 budgets and metabolic rates because these dramatic differences in behaviour associated with
399 adjacent habitats could have important life history implications.

400

401 Our data supported three of the four hypotheses of our study, with the niche
402 complementarity hypothesis being clearly rejected by our study. The niche complementarity
403 hypothesis was the only one that involved interspecific relationships as a causal factor for its
404 acceptance, given that, under its assumptions, the habitat differences between species with a same
405 bauplan should be due to the need to minimize interspecific competition (Schoener 1974). On the
406 other hand, the other three hypotheses are based on size-related autecological exigencies
407 (hypothesis 2), morphological constraints and physical performance (hypothesis 3), and eco-
408 physiological constraints (hypothesis 4). Therefore, we think that these lizard communities are more

clearly organized by autecological constraints and eco-physiological attributes, whereas the interspecific interactions produced apparent patterns of habitat niche divergences that are unlikely due to interspecific competition and the ghost-of-the-competition-past (Connell 1980; Alatalo and Lundberg 1983; Cressman and Krivan 2013; Bottin et al. 2016). This general conclusion was also confirmed by our null model analyses, that did not provide any evidence for a competitively-structured assemblage of species under any of the randomization algorithms (Lawlor 1980) that we used and that have been shown elsewhere to be powerful tools for detecting structure signs of nonrandom niche partitioning in biotic communities (e.g., Gotelli and Graves 1996; Solida et al. 2011; Vignoli and Luiselli 2012). Nonetheless, our data cannot give firm conclusions on this issue, as we would need to examine habitat use in sites with and without competitors to examine variation in habitat use in absence of potential competition.

Acknowledgements. We are indebted to N. Sillero and an anonymous reviewer for helpful comments on the submitted draft. Since we did not capture lizard individuals for this study, no special authorization was needed from the competent governmental authorities.

References

- Alatalo, R.V., and Lundberg, A. 1983. Laboratory Experiments on Habitat Separation and Foraging Efficiency in Marsh and Willow Tits. *Ornis Scandinavica* **14**(2): 115–122.
- Arnold, E.N. 1987. Resource partition among lacertid lizards in southern Europe. *Journal of Zoology* **1**(4): 739–782.
- Bottin, M., Soininen, J., Alard, D., and Rosebery, J. 2016. Diatom cooccurrence shows less segregation than predicted from niche modeling. *PloS one* **11**(4): e0154581.
- Bouam, I., Necer, A., Saoudi, M., Tahar-Chaouch, L., and Khelfaoui, F. 2016. Diet and daily activity patterns of the lacertid lizard *Psammodromus algirus* (Sauria: Lacertidae) in a semi-arid Mediterranean region. *Zoology and Ecology* **26**(3): 244–252.
- Burke, R.L., and Ner, S.E. 2005. Seasonal and diel activity patterns of Italian wall lizards, *Podarcis sicula campestris*, in New York. *Northeastern Naturalist* **12**(3): 349–360.
- Calsbeek, R., and Cox, R. M. 2010. Experimentally assessing the relative importance of predation and competition as agents of selection. *Nature*, **465**(7298): 613–616.
- Capizzi, D., and Luiselli, L. 1996. Feeding relationships and competitive interactions between phylogenetically unrelated predators (owls and snakes). *Acta Oecologica* **17**(4): 265–284.

- Capula, M., and Ceccarelli, A. 2003. Distribution of genetic variation and taxonomy of insular and mainland populations of the Italian wall lizard, *Podarcis sicula*. *Amphibia-Reptilia* **24**(4): 483–495.
- Carretero, M.A., and Llorente, G.A. 1995. Thermal and temporal patterns of two Mediterranean Lacertidae. Papers submitted from 7th O.G.M. of Societas Europaea Herpetologica, Barcelona, September 15-19, 1993. Edited by G.A Llorente, A. Montori, X. Santos, and M.A. Carretero. *Scientia herpetologica*: 213–223. Asociación Herpetológica Española, Barcelona.
- Christian, K.A., and Waldschmidt, S. 1984. The relationship between lizard home range and body size: a reanalysis of the data. *Herpetologica* **40**(1): 68–75. JSTOR.
- Connell, J.H. 1980. Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos* **35**: 131–138.
- Corti, C., Capula, M., Luiselli, L., Razzetti, E., and Sindaco, R. 2011. *Reptilia. Fauna d'Italia*. Calderini editore, Bologna.
- Cressman, R., and Křivan, V. 2013. Two-patch population models with adaptive dispersal: the effects of varying dispersal speeds. *Journal of Mathematical Biology* **67**(2): 329–358. Springer.
- Delaney, D.M., and Warner, D.A. 2017a. Adult male density influences juvenile microhabitat use in a territorial lizard. *Ethology* **123**(2): 157–167.
- Delaney, D.M., and Warner, D.A. 2017b. Effects of age-and sex-specific density on behaviour and survival in a territorial lizard (*Anolis sagrei*). *Animal Behaviour* **129**: 31–41.
- Escoriza, D., and Amat, F. 2021. Habitat Partitioning and Overlap by Large Lacertid Lizards in Southern Europe. *Diversity* **13**(4): 155.
- Ferrandino, I., Viscardi, G., and Grimaldi, M.C. 2001. An immunohistochemical study of adenohypophyseal cells in the viviparous reptile *Chalcides chalcides*. *The Histochemical Journal* **33**(1): 1–8.
- Flesch, A.D., Rosen, P.C., and Holm, P. 2017. Long-term changes in abundances of Sonoran Desert lizards reveal complex responses to climatic variation. *Global Change Biology* **23**(12): 5492–5508.
- Friggens, M.M., and Brown, J.H. 2005. Niche partitioning in the cestode communities of two elasmobranchs. *Oikos* **108**(1): 76–84.
- Gonçalves-Sousa, J.G., Mesquita, D.O., and Ávila, R.W. 2019. Structure of a lizard assemblage in a semiarid habitat of the Brazilian Caatinga. *Herpetologica* **75**(4): 301–314.
- Gotelli, N.J., and Graves, G.R. 1996. *Null models in ecology* Smithsonian Institution Press. Washington, DC, USA.
- Grundler, M.R., Pianka, E.R., Pelegrin, N., Cowan, M.A., and Rabosky, D.L. 2017. Stable isotope ecology of a hyper-diverse community of scincid lizards from arid Australia. *PloS one* **12**(2): e0172879.
- Hall, B.K. 1999. *Evolutionary Developmental Biology*. In 2nd edition. Kluwer, Dordrecht, The Netherlands.
- Helmer, W., Strijbosch, H., and Scholte, P.T. 1989. Distribution and ecology of lizards in the Greek province of Evros. *Amphibia-reptilia* **10**(2): 151–174.
- Hurlbert, S.H. 1984. Pseudoreplication and the design of ecological field experiments. *Ecological Monographs* **54**(2): 187–211.
- Jenssen, T.A., Hovde, K.A., and Taney, K.G. 1998. Size-related habitat use by nonbreeding *Anolis carolinensis* lizards. *Copeia* **1998**(3): 774–779.
- Jiménez-Robles, O., and De la Riva, I. 2019. Lizards in the mist: Thermal niches constrained by habitat and microclimates in the Andes of southern Bolivia. *Journal of Biogeography* **46**(8): 1676–1686.
- Lawlor, L.R. 1980. Structure and stability in natural and randomly constructed competitive communities. *The American Naturalist* **116**(3): 394–408.

- Lehsten, V., and Harmand, P. 2006. Null models for species co-occurrence patterns: assessing bias and minimum iteration number for the sequential swap. *Ecography* **29**(5): 786–792.
- Levins, R. 1968. *Evolution in changing environments: some theoretical explorations*. Princeton University Press.
- Losos, J.B. 2011. *Lizards in an evolutionary tree: ecology and adaptive radiation of anoles*. Univ of California Press.
- Losos, J.B., Jackman, T.R., Larson, R., de Queiroz, K., and Rodriguez-Schettino, L. 1998. Contingency and determinism in replicated adaptive radiations of island lizards. *Science* **279**: 2115–2118.
- Luiselli, L. 1992. The diet of the Slow worm, *Anguis f. fragilis* Linnaeus, 1758, in the Tarvisio Forest (Carnic Alps, NE Italy)(Squamata: Sauria: Anguidae). *Herpetozoa* **5**(3/4): 91–94.
- Luiselli, L. 2008. Do lizard communities partition the trophic niche? A worldwide meta-analysis using null models. *Oikos* **117**(3): 321–330.
- Maura, M., Vignoli, L., Bologna, M.A., Rugiero, L., and Luiselli, L. 2011. Population density of syntopic, differently sized lizards in three fragmented woodlands from Mediterranean Central Italy. *Community Ecology* **12**(2): 249–258.
- Pérez Mellado, V. 1982. Estructura en una taxocenosis de Lacertidae (Sauria, Reptilia) del sistema central. Mediterránea. Serie de Estudios Biológicos, N. 6 (diciembre 1982); pp. 39-64. Universidad de Alicante. Departamento de Biología.
- Pianka, E.R. 1986. *Ecology and Natural History of Desert Lizards* Princeton University Press. Princeton, New Jersey.
- Powell, G.L., and Russell, A.P. 1992. Locomotor correlates of ecomorph designation in *Anolis*: an examination of three sympatric species from Jamaica. *Canadian Journal of Zoology* **70**(4): 725–739.
- Pringle, R.M., Kartzinel, T.R., Palmer, T.M., Thurman, T.J., Fox-Dobbs, K., Xu, C.C., et al. 2019. Predator-induced collapse of niche structure and species coexistence. *Nature* **570**(7759): 58–64.
- Rieppel, O. 2006. ‘Type’ in morphology and phylogeny. *Journal of Morphology* **267**(5): 528–535.
- Rocha, C.F.D. 1999. Home range of the tropidurid lizard *Liolaemus lutzae*: sexual and body size differences. *Revista Brasileira de Biologia* **59**(1): 125–130.
- Rouag, R., Djilali, H., Gueraiche, H., and Luiselli, L. 2007. Resource partitioning patterns between two sympatric lizard species from Algeria. *Journal of Arid Environments* **69**(1): 158–168.
- Rugiero, L. 2004. Composition of the reptile communities in five urban protected areas of different isolation degrees. *Herpetozoa* **16**(3/4): 151–155.
- Rugiero, L., Capula, M., Di Vittorio, M., Dendi, D., Meek, R., and Luiselli, L. 2021. Ontogenetic habitat use and density of the green lizard (*Lacerta bilineata*) in contrasted landscapes in France and Italy. *Conservation* **1**(1): 1–16.
- Rugiero, L., Capula, M., Filippi, L., and L. Luiselli. 1995. Food habits of Mediterranean populations of the Smooth snake (*Coronella austriaca*). *Herpetological Journal* **5**: 316–318.
- Saint Girons, H., and Bradshaw, S.D. 1989. Sédentarité, déplacements et répartition des individus dans une population de *Lacerta viridis* (Laurenti, 1768)(Lacertilia, Lacertidae). *Bijdragen tot de Dierkunde* **59**(2): 63–70.
- Santos, X., and Poquet, J.M. 2010. Ecological succession and habitat attributes affect the postfire response of a Mediterranean reptile community. *European Journal of Wildlife Research* **56**(6): 895–905.
- Schoener, T.W. 1974. Resource partitioning in ecological communities. *Science* **185**(4145): 27–39.
- Sheu, Y., Zurano, J.P., Ribeiro-Junior, M.A., Ávila-Pires, T.C., Rodrigues, M.T., Colli, G.R., et al. 2020. The combined role of dispersal and niche evolution in the diversification of Neotropical lizards. *Ecology and evolution* **10**(5): 2608–2625.

- Sillero, N., Argaña, E., Matos, C., Franch, M., Kaliontzopoulou, A., and Carretero, M.A. 2020. Local segregation of realised niches in lizards. *ISPRS International Journal of Geo-Information* **9**(12): 764.
- Sillero, N., and Gomes, V. 2016. Living in clusters: the local spatial segregation of a lizard community. *Basic and Applied Herpetology* **30**: 61–75.
- Simbula, G., Luiselli, L., and Vignoli, L. 2019. Lizards and the city: A community study of Lacertidae and Gekkonidae from an archaeological park in Rome. *Zoologischer Anzeiger* **283**: 20–26.
- Simpson, E.H. 1949. Measurement of Diversity. *Nature* **163**(4148): 688–688.
- Solida, L., Celant, A., Luiselli, L., Grasso, D.A., Mori, A., and Fanfani, A. 2011. Competition for foraging resources and coexistence of two syntopic species of Messor harvester ants in Mediterranean grassland. *Ecological Entomology* **36**(4): 409–416.
- Stankowich, T., and Stensrud, C. 2019. Small but spiny: the evolution of antipredator defenses in Madagascar tenrecs. *Journal of Mammalogy* **100**(1): 13–20.
- Toft, C.A. 1985. Resource partitioning in amphibians and reptiles. *Copeia* 1985 (1): 1–21.
- Tsessarsky, A.A. 2020. Role of paedomorphosis in the emergence of the skull bauplan in Acipenseriformes (Actinopterygii). *Biology Bulletin Reviews* **10**(5): 427–440.
- Turner, F.B., Jennrich, R.I., and Weintraub, J.D. 1969. Home ranges and body size of lizards. *Ecology* **50**(6): 1076–1081.
- Vacheva, E.D., Naumov, B.Y., and Tzankov, N.D. 2020. Diversity and habitat preferences in lizard assemblages (Reptilia: Sauria) from model territories in western Bulgaria. *Acta Zool. Bulg* **72**(3): 385–396.
- Vanhooydonck, B., and Van Damme, R. 2003. Relationships between locomotor performance, microhabitat use and antipredator behaviour in lacertid lizards. *Functional Ecology* **17**(2): 160–169.
- Vignoli, L., and Luiselli, L. 2012. Dietary relationships among coexisting anuran amphibians: a worldwide quantitative review. *Oecologia* **169**(2): 499–509.
- Williams, E.E. 1983. Ecomorphs, Faunas, Island Size, and Diverse End Points in Island Radiations of Anolis. *In* Lizard ecology: studies of a model organism. *Edited by* R.B. Huey, E.R. Pianka, and T.W. Schoener. Harvard University Press, Cambridge, Mass. pp. 326–370.
- Willmore, K.E. 2012. The body plan concept and its centrality in evo-devo. *Evolution: Education and Outreach* **5**(2): 219–230.
- Woodger, J.H. 1945. On biological transformations. *In* Essays on Growth and Form, presented to D'Arcy Wentworth Thompson. *Edited by* W.E. Le Gros Clark and P.B. Medawar. Oxford University Press, Oxford. pp. 95–120.
- Zakkak, S., Halley, J.M., Akriotis, T., and Kati, V. 2015. Lizards along an agricultural land abandonment gradient in Pindos Mountains, Greece. *Amphibia-Reptilia* **36**(3): 253–264. Brill.
- Zamora-Camacho, F.J., Reguera, S., Moreno-Rueda, G., and Pleguezuelos, J.M. 2013. Patterns of seasonal activity in a Mediterranean lizard along a 2200 m altitudinal gradient. *Journal of Thermal Biology* **38**(2): 64–69.

583

584 **Table 1.** Synthesis of the number of lizard species observed at the study area by habitat type and by individual category (male, female or juvenile).
585 Data from all seasons are pooled.

586

Genus	Species	Category	Symbol	<i>Rubus</i>	<i>Cytisus</i>	<i>Spartium</i>	grassland	stony walls	<i>Smilax aspera</i>	TOTAL
<i>Podarcis</i>	<i>sicula</i>	males	PsM	51	77	13	369	114	41	665
<i>Podarcis</i>	<i>sicula</i>	females	PsF	39	73	22	401	133	28	696
<i>Podarcis</i>	<i>sicula</i>	juveniles	PsJ	7	17	8	211	91	11	345
<i>Podarcis</i>	<i>muralis</i>	males	PmM	598	233	189	31	916	176	2143
<i>Podarcis</i>	<i>muralis</i>	females	PmF	668	321	273	9	894	203	2165
<i>Podarcis</i>	<i>muralis</i>	juveniles	PmJ	133	16	4	5	231	11	400
<i>Chalcides</i>	<i>chalcides</i>		Cc	6	41	33	311	3	0	394
<i>Anguis</i>	<i>veronensis</i>	males	AvM	1	0	0	0	2	1	4
<i>Anguis</i>	<i>veronensis</i>	females	AvF	1	0	0	0	0	0	1
<i>Anguis</i>	<i>veronensis</i>	juveniles	AvJ	2	1	0	0	2	0	5
<i>Tarentola</i>	<i>mauritanica</i>		Tm	0	0	0	0	31	0	31
<i>Hemidactylus</i>	<i>turcicus</i>		Ht	0	0	0	3	8	0	11
<i>Lacerta</i>	<i>bilineata</i>	males	LbM	51	18	10	10	22	7	118

22

<i>Lacerta</i>	<i>bilineata</i>	females	LbF	34	22	14	20	16	11	117
<i>Lacerta</i>	<i>bilineata</i>	juveniles	LbJ	7	9	12	111	22	1	162
HABITAT TYPE AVAILABILITY (%)			Availability	20.69	16.73	15.41	16.87	17.54	12.75	

Draft

588 **Table 2.** Testing the “niche complementarity” hypothesis using habitat niche overlap simulation
589 analysis for lizards at the study area in central Italy. The used algorithms were Lawlor’s (1980)
590 RA2, RA3 and RA4 algorithms, after 30,000 Monte Carlo simulations. These simulations were
591 done from data of all seasons being pooled.
592

	RA2	RA3	RA4
Observed mean Overlap	0.53870	0.53870	0.53870
Mean simulated Overlap	0.57851	0.42382	0.76684
Variance simulated Overlap	0.00095	0.00053	0.00092
Standardized Effect Size	-1.29004	5.01184	-7.50843
Random seed	-1645502148	-1800479482	-1525427312
p(observed $\leq$ expected)	0.10400	0.99999	0.00001
p(observed $\geq$ expected)	0.89600	0.00001	0.99999
Observed variance	0.09309	0.09309	0.09309
Mean of simulated variances	0.06089	0.07930	0.01587
Variance of simulated variances	0.00006	0.00004	0.00002
p(observed $\leq$ expected)	1.00000	0.97000	0.99999
p(observed $\geq$ expected)	0.00000	0.03000	0.00001

593

594

595 **Table 3.** Testing the hypothesis that larger species should have wider niche breadth using the values
 596 of habitat niche breadth, calculated for each individual category of the various species, at the study
 597 area in central Italy. Data from all seasons are pooled. B_S = Simpson's formula; B_L = Levins'
 598 formula.
 599

Genus	Species	Category	B_S	B_L
<i>Podarcis</i>	<i>sicula</i>	males	2.77	0.25
<i>Podarcis</i>	<i>sicula</i>	females	2.60	0.23
<i>Podarcis</i>	<i>sicula</i>	juveniles	2.23	0.18
<i>Podarcis</i>	<i>muralis</i>	males	3.48	0.35
<i>Podarcis</i>	<i>muralis</i>	females	3.20	0.31
<i>Podarcis</i>	<i>muralis</i>	juveniles	2.24	0.18
<i>Chalcides</i>	<i>chalcides</i>		1.56	0.08
<i>Anguis</i>	<i>veronensis</i>	males	2.67	0.24
<i>Anguis</i>	<i>veronensis</i>	females	1.00	0.00
<i>Anguis</i>	<i>veronensis</i>	juveniles	2.78	0.25
<i>Tarentola</i>	<i>mauritanica</i>		1.00	0.00
<i>Hemidactylus</i>	<i>turcicus</i>		1.66	0.09
<i>Lacerta</i>	<i>bilineata</i>	males	3.81	0.40
<i>Lacerta</i>	<i>bilineata</i>	females	5.24	0.61
<i>Lacerta</i>	<i>bilineata</i>	juveniles	2.01	0.14

600

601

602 **Figure 1.** Some of the habitat types at the study area, during the early spring: (a) *Rubus*; (b)
603 grassland; (c) and (d) two types of stony walls

604

605 **Figure 2.** Scatter plot of a non-metric multidimensional scaling (nMDS), using the cosine distance,
606 graphically representing the resemblance matrix between lizard individuals and in relation to the
607 habitat availability. Shown are the centroids of clusters of lizard individuals by sex, age class and
608 species and of habitat availability. Data from all seasons are pooled. Symbols: Cc = *Chalcides*
609 *chalcides*; Av = *Anguis veronensis*; Ht = *Hemidactylus turcicus*; Lb = *Lacerta bilineata*; Pm =
610 *Podarcis muralis*; Ps = *Podarcis sicula*; Tm = *Tarentola mauritanica*; M = males; F = females; J =
611 juveniles; Availability = habitat availability.

612

613 **Figure 3.** Testing the hypothesis that species inhabiting open sunny habitats should exhibit more
614 seasonally variable activity patterns using the percent of lizard individuals observed by season at the
615 study area in central Italy. Symbols: Cc = *Chalcides chalcides* (n = 394); Av = *Anguis veronensis* (n
616 = 10); Ht = *Hemidactylus turcicus* (n = 11); LB = *Lacerta bilineata* (n = 397); Pm = *Podarcis*
617 *muralis* (n = 4708); Ps = *Podarcis sicula* (n = 1706); Tm = *Tarentola mauritanica* (n = 28); M =
618 males; F = females; J = juveniles

619

620 Figure 1

621

622 Figure 2

623

624

625 Figure 3

626

Draft