**Transcriptome Analysis Reveals Key Genes and Pathways
Associated with Mummify Piglets**

Journal:	<i>Genome</i>
Manuscript ID	gen-2021-0026.R1
Manuscript Type:	Article
Date Submitted by the Author:	01-Jun-2021
Complete List of Authors:	Wang, Shujie; Sichuan Agricultural University Wu, Pingxian; Sichuan Agricultural University Wang, Kai; Sichuan Agricultural University Ji, Xiang; Sichuan Agricultural University Chen, Dong; Sichuan Agricultural University Jiang, Anan; Sichuan Agricultural University - Chengdu Campus Liu, Yihui; Sichuan Animal Husbandry Station Xiao, Weihang ; Sichuan Agricultural University - Chengdu Campus Jiang, Yanzhi; College of Life Science, Sichuan Agricultural University Zhu, Li; Sichuan Agricultural University - Chengdu Campus Xu, Xu; Sichuan Provincial Animal Husbandry and Food Bureau Li, Mingzhou; Sichuan Agricultural University Li, Xuwei ; Sichuan Agricultural University - Chengdu Campus Tang, Guoqing ; Sichuan Agricultural University - Chengdu Campus
Keyword:	Transcriptome, Pregnancy, Mummify Piglets, Blood
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

1 Transcriptome Analysis Reveals Key Genes and Pathways
2 Associated with Mummify Piglets
3 Shujie Wang¹, Pingxian Wu¹, Kai Wang¹, Xiang Ji¹, Dong Chen¹, Anan Jiang¹, Yihui Liu¹ Weihang
4 Xiao¹, Yanzhi Jiang², Li Zhu¹, Xu Xu³, Mingzhou Li¹, Xuewei Li¹, Guoqing Tang^{1,*}
5 ¹ Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province,
6 Sichuan Agricultural University, Chengdu 611130, Sichuan, China
7 ² College of Life Science, Sichuan Agricultural University, Yaan 625014, Sichuan, China
8 ³ Sichuan Animal Husbandry Station, Chengdu, 610041, Sichuan, China
9 ***Corresponding author:** Guoqing Tang, tyq003@163.com
10 **Co-first authors:** Shujie Wang and Pingxian Wu contributed equally.
11 **Authors' Email:** Shujie Wang, Email: 670186296@qq.com.
12 Pingxian Wu, Email: wupingxianxian@163.com.
13 Kai Wang, Email: 614351818@qq.com.
14 Xiang Ji, Email: 1084481695@qq.com.
15 Dong Chen, Email: 1123278154@qq.com.
16 Yihui Liu, Email: YihuiSky@163.com.
17 Anan Jiang, Email: 30769245@qq.com.
18 Weihang Xiao, Email: weihang.xiao@sicau.edu.cn.
19 Yanzhi Jiang, Email: jiangyz04@163.com.
20 Li Zhu, Email: zhuli7508@163.com.
21 Xu Xu, Email: 908517582@qq.com.
22 Mingzhou Li, Email: mingzhou.li@163.com.
23 Xuewei Li, Email: lixuewei9125@126.com.
24 Guoqing Tang, Email: tyq003@163.com.
25
26 **Abstract:** China is the country with the largest pork consumption in the world. However, the
27 incidence of high mummify piglets (3-5%) is one of the important factors that cause the slow

improvement of pig reproductive capacity, and the genetic mechanism is still unclear. This study aimed to identify candidate genes related to high mummify piglets. RNA-seq technology was used to comparative transcriptome profiling of blood from high piglets mummified and healthy sow at different stages of pregnancy (35d, 56d, 77d and 98d). A total of 137 to 420 DEGs were detected in each stage. Seven differentially expressed genes were significantly differentially expressed at various stages. *IL-9R*, *TLR8*, *ABLIM3*, *FSH- α* , *ASCC1*, *PRKCZ*, and *GCK* may play an important role in course of mummify piglets. The differential genes we identified between the groups were mainly enriched in immune and inflammation regulation, and others were mainly enriched in reproduction. Considering the function of candidate genes, *IL-9R* and *TLR8* were suggested as the most promising candidate genes involved in mummify piglet traits. We speculate that during pregnancy, it may be the combined effects of the above-mentioned inflammation, immune response, and reproduction-related signal pathways that affect the occurrence of mummifying piglets, and further affect pig reproduction.

Keywords: Transcriptome, Pregnancy, Mummify Piglets, Blood

1 Introduction

Reproductive traits are one of the most important economic traits that affect the economic benefits of pig production. The living piglets determine the greatest potential for the economic benefits of the pig farm. Generally, the number of stillbirths accounts for 3%~8% of the total number born (TNB), which has an important influence on the number born alive. With the development of intensive and large-scale pig breeding, mummify pigs have caused more and more economic losses.

50 Blood is a connective tissue, which not only ensures nutrient and oxygen supply of
51 all organs of the body, but also the communication between them. Blood plays an
52 important role at all levels of the life system, bringing together many transcriptional
53 variations in the process of life activities, and specific gene expression patterns in the
54 blood can directly reflect the physiological state information of specific tissues.
55 Peripheral blood is a highly dynamic tissue type with a high rate of cell turnover,
56 resulting in high diversity of RNA content over time(OGAWA 1993). Peripheral blood
57 interacts with every organ and tissue of the body, thus presenting a gene expression
58 profile that can reflect the subject's physiological and pathological conditions, behavior,
59 growth stage and lifestyle(Huang and Zeng 2020; Koutroumpakis et al. 2020; Shidid et
60 al. 2020; Zhou et al. 2021). During pregnancy, blood, as the direct source of fetal
61 nutrients, is the key to maintaining normal pregnancy and ensuring the normal growth
62 and development of the fetus, as well as the key to ensuring the functional
63 communication between the mother and the fetus. At present, peripheral blood
64 transcriptome is widely used in immune response, premature fetal delivery, and
65 abortion(Mach et al. 2013; Mallikarjun et al. 2017; Shen et al. 2014). Based on blood
66 transcriptome research, it is widely used in human diseases and pregnancy-related
67 detection(You et al. 2019). The analysis of pregnant maternal blood transcriptome can
68 be used to predict spontaneous fetal preterm delivery and may be used in the detection
69 of fetal malformations(Ling et al. 2017; Ronzoni et al. 2019). At present, transcription
70 studies related to porcine reproductive traits are mainly focused on the differences in
71 transcription of specific tissues to reveal the overall gene expression changes in the

tissues directly involved in pregnancy, such as placenta, ovary, endometrial tissue, etc. (Holly and LaVoie 2017; Kaczynski et al. 2019; Zhang et al. 2019). However, many studies related to pig pregnancy are carried out after slaughter, which will cause research limitations and waste of sow. Studies have shown that peripheral blood can be used as a detection means for detecting pregnancy status(Kamat et al. 2016). Therefore, based on the perspective of the whole blood transcriptome, it is helpful to efficiently explore the transcriptional regulation mechanism related to reproductive traits.

High-throughput RNA-Seq has become a method of choice to achieve inexpensive and routine comprehensive analysis of animal transcriptomes or genomes(do Nascimento et al. 2018; Seo et al. 2016). This technology is mainly used to quantitatively determine transcripts or genes (known and unknown), expression patterns of mRNA splice variants, and analysis of SNPs, which can be used as potential biomarkers for specific production traits(Cánovas et al. 2010; Giraldez et al. 2019; Maatz et al. 2014). Research has been conducted using RNA-Seq in agricultural animals to explore global gene expression patterns in organizations related to economically important features such as milk production, feed efficacies, carcass growth, and so on(Bai et al. 2016; Horodyska et al. 2019; Ramayo-Caldas et al. 2018; Zhuo et al. 2015).

Sow reproductive performance is an important factor for the profitability of pig production. In the past two decades, the total number of piglets born per litter was increased through genetic selection. At the same time, as the number of litters increases, the incidence of stillbirth often reduces the sow's reproductive efficiency(Wang et al.

2019). As far as we know, almost no research has been conducted to study the differences in gene expression of blood transcriptome in mummified piglets. Using RNA-Seq comparison of sow blood between healthy and high mummified piglets in four different stages of gestation (35d, 56d, 77d and 98d) found DEG, which may help explain the cause of stillbirth mummies and better reproductive performance to improve livestock breeding.

2 Methods

2.1 Animals

The purebred large white sows selected for this experiment were all from a company in Zhejiang, China. A total of 27778 parities were collected for this trial (from 2011 to 2017). In this experiment, the third parity sows (with two reproductive records) were selected, and the non-mummified reproductive record was defined as the no mummified piglets (NM) group, and the mummified record higher than 2 piglets per litter was defined as high mummified piglets (HM). In the early stage of the experiment, the sows were tested for sow reproductive and respiratory syndrome (PRRSV), pseudorabies virus (PRV), swine fever virus (CRFV), circovirus (PSV) and parvovirus (PPV). The test results were negative. The semen of the same boar was used for mating for pregnancy test. During the test, feeds with consistent nutritional levels were uniformly used. Each sow was raised in a single pen. Usually, 35d was the start time of the mummy pig. With a three-week interval, three subsequent time nodes were selected (35d, 56d, 77d and 98d). Blood samples from four pregnancy stages were taken, and Trizol lysate was added to the lysate, shaken quickly and evenly, and then placed in liquid nitrogen for

freezing. After the test, the farrowing situation of the sows was recorded, and the screening was performed again according to the actual farrowing situation during the test period, and samples of 8 sows without mummify piglets and 8 sows with high mummify piglets were selected for RNA-seq. The reproduction information of 16 pigs used for transcription sequencing is shown in Table 1.

2.2 Blood collection and total RNA extraction

Total RNA was isolated from the porcine vein blood using the Trizol Kit (Thermo Fisher Scientific, USA) according to the manufacturer's recommendations. All samples were treated with DNase I (Thermo Fisher Scientific, USA). Sample concentrations and RNA integrity number (RIN) quality were measured using the Nano Drop 2000 (Thermo Fisher Scientific, USA); the RIN values indicated high-quality total RNA in all the samples ($RIN \geq 7$).

2.3 Library construction and sequencing

The library construction was performed according to the instructions of the Illumina mRNA-seq library kit. First, the total RNA was separated and purified, and then fragmented. Using a short fragment of mRNA as a template, reverse transcription was performed to synthesize cDNA, and the obtained cDNA was purified. The double-stranded cDNA was repaired at the end, the poly-A tail and the sequencing adapter index were added. Then select the appropriate fragments for PCR amplification to obtain the final cDNA library (PrimeScript™ RT reagent Kit (Perfect Real Time), Taraka, Japan). Sample concentrations and RIN quality were measured using the Agilent 2100 (Agilent Technologies, CA, USA). Finally, the qualified library used

Illumina HiSeq 2000 PE150 (Illumina, USA) for pair-end sequencing, with a sequencing depth of 5G/sample.

2.4 Sequencing data quality control

Sampling FastQC software to perform quality control on the raw data of the obtained raw sequence data, removed low-quality reads, sequencing adapter sequences, and contain more than 10% of N information to obtain high-quality data. Clean data was used for subsequent analysis.

2.5 Sequence Alignment

Took the Sus_scrofa.Sscrofa11.1dna_sm.toplevel (Release-94) genome data of the Ensemble database as the reference genome. HISAT2 was used to compare the clean data to the reference genome. StringTie was used maximum flow algorithm to assemble the reads on the comparison and expressive assessment.

2.6 Screening of differentially expressed genes

EdgeR was used to screen differentially expressed genes in different stages of pregnancy. The screening criteria were $|FC| \geq 1.5$ and $p\text{-value} \leq 0.05$, of which $FC \geq 1.5$ was upregulated differentially expressed genes, and $FC \leq -1.5$ was downregulated.

2.7 Differential gene GO and KEGG analysis

DAVID software was used to analyze the differentially expressed genes by GO analysis and KEGG pathway analysis. The significantly enriched pathways ($P\text{-value} < 0.05$) were screened by the hypergeometric test method, and pathways related to the target traits and gene functions were further analyzed.

2.8 Differential gene PCR verification

In order to further verify the accuracy of the differentially expressed genes obtained by sequencing, 12 differentially expressed genes were randomly selected in the results, *β-actin* was used as the internal reference gene, and RT-qPCR (Quantitative Real-time PCR) was used to quantitatively verify the differentially expressed genes, using $2^{-\Delta\Delta Ct}$ the method calculates the expression amount of the quantitative results. The primer sequence of each gene is shown in Table 2.

3 Results

3.1 Transcriptome Profiles

In this study, a total of 64 libraries were established from the blood of NW and HW group in 35d, 56d, 77d, and 98d. The gene expression of NW and HW group's blood transcriptome at different stages of pregnancy was systematically analyzed by high-throughput RNA sequencing. After quality control, clean reads of samples ranging from 20,828,466 to 34,601,836 (28,343,637 on average), GC contents of the samples were between 56.55% and 59.67%, and $\geq Q30$ (%) were 93.51% to 95.70% (Table S1). In total reads of the samples, over 41,656,932 high-quality reads per sample were mapped to the SSC genome and used for gene expression analysis, where "Uniq Mapped Reads" ranging from 28,807,331 (45.26%) to 51,638,131 (75.45%) and "Multiple Map Reads" ranging from 8,366,761 (16.72%) to 31,989,604 (50.26%). Besides, the number of Reads aligned to the positive strand in the SSC genome were 18,564,928 to 32,012,639 and the number of reads aligned to the negative strand in the SSC genome were between 18,820,558 and 32,415,364 (Table 3).

3.2 DEGs and Functional Annotations between NM and HM group

The relative expression of gene was normalized as fragments per kilobase of exon model per million mapped reads (FPKM), which was proportional to the number of cDNA fragments originated by gene transcription. Statistical analysis of DEGs in blood of NM and HM group at different stages of pregnancy were shown in Figure 2. In the blood genome, differential gene expression analysis of NM_35d vs. HM_35d showed that 378 genes were significantly differentially expressed (Fold change ≥ 1.5 and $p < 0.05$, the same below), including 148 upregulated and 230 downregulated genes. Moreover, there were 338 significantly expressed genes from NM_56d vs. HM_56d, among which 191 were upregulated genes and 147 were downregulated genes. The number of 137 DEGs were detected from NM_77d vs. HM_77d, the number of upregulated genes were 68 and the downregulated genes were 69. There were 420 DEGs from NM_98d vs. HM_98d, including 220 upregulated genes and 200 downregulated genes (Figure 1). Normalized counts and logarithmic values of fold change ($\log_2FC$) were visualized in correlation, MA, and Volcano plots (Figure 2).

3.2.1 Analysis of Differential Gene Expression Function between HM_35d vs. NM_35d

To further determine the biological functions of the DGEs identified by HM and NM transcription at 35d gestation, we analyzed the identified DGEs by GO and KEGG. GO analysis showed that a total of 62 significant GO items were enriched (*Supplement S1*), including nuclear inflammatory response, positive regulation of tumor necrosis factor production, glycosomal structure composition, innate immune response, ribosome, Toll-like receptor signaling pathway, and ATPase Binding, apoptosis signal

pathway, neuronal apoptosis process, negative regulation of cysteine endopeptidase activity in apoptosis signal pathway and positive regulation of interferon- β production (Figure 3A). Through KEGG enrichment analysis, it was found that the differential genes between NM_35d vs. HM_35d were mainly enriched in 10 pathways, including ribosome, diabetes, fluid shear stress and atherosclerosis, Toll-like receptor signaling pathway, and prolactin signaling Pathways and amino acid metabolism. Ovarian steroids play a key role in the reproduction process of animals. Our results also enriched some pathways related to ovarian steroid production (Figure 4A).

3.2.2 Analysis of Differential Gene Expression Function between HM_56d vs. NM_56d

The enrichment analysis of differentially expressed genes in the HM_56d vs. NM_56d group identified 55 GO term (*Supplement S1*) and 14 KEGG pathways, respectively. Among them, GO term was mainly related to innate immune response, Toll-like receptor signaling pathway, calcium ion binding, positive regulation of interferon- β production, positive regulation of angiogenesis, cartilage development, and regulation of intracellular pH (Figure 3B). The KEGG pathway was mainly related to diseases (such as bladder disease and malaria), apoptosis, Toll-like receptor signaling pathway, and nicotinate and niacinamide metabolism pathways (Figure 4B).

3.2.3 Analysis of Differential Gene Expression Function between HM_77d vs. NM_77d

HM_77d vs. NM_77d difference gene GO and KEGG enrichment analysis enriched 77 GO entries (*Supplement S1*) and 6 KEGG pathways. GO term was mainly

related to development process, apoptosis process, placental vascular development, Toll-like receptor 7 signaling pathway, negative regulation of innate immune response, cysteine-type endopeptidase activity involved in apoptosis signaling pathway, ATPase activity and amino acids Transshipment and other related (Figure 3C). KEGG enrichment was mainly related to diabetes, glucagon signaling pathway, antifolate, amino acid metabolism, FoxO signaling pathway, etc. (Figure 4C).

3.2.4 Analysis of Differential Gene Expression Function between HM_98d vs. NM_98d

HM_98d vs. NM_98d difference gene GO and KEGG enrichment analysis enriched 90 GO entries (*Supplement S1*) and 11 KEGG pathways. GO entries are mainly related to dephosphorylation, oxygen transport, apoptosis signaling pathway, hematopoiesis, sodium and potassium ion transport, ATPase binding, transmembrane receptor protein tyrosine kinase signaling pathway, and ATPase-coupled intramembrane lipid transporter activity and the process of heme biosynthesis are related (Figure 3D). KEGG enrichment analysis mainly includes porphyrin and chlorophyll metabolism, hematopoietic cell lineage and sphingolipid signaling pathway, amino acid metabolism and diseases. In addition, we also found that these differential genes are related to inflammation, FoxO signaling pathway, Toll-like receptor signaling pathway, and prolactin signaling pathway (Figure 4D).

3.3 RT-qPCR verification of transcriptome sequencing

To verify the accuracy of the transcriptome sequencing data, we randomly selected 12 differentially expressed genes for RT-qPCR verification, and further compared the

quantitative results with the sequencing results. The results showed that the quantitative results of the 12 differentially expressed genes we screened were basically consistent with the results of transcription sequencing, indicating that the transcription sequencing data was accurate and reliable (Table 5).

4 Discussion

Sow reproductive performance provides direct economic benefits to the livestock industry. The incidence of high mummify piglets (3-5%) is one of the important factors that cause the slow improvement of pig reproductive capacity. The previous research on mummify piglets mainly focuses on nutrition, disease, management, and behavior (Anil et al. 2009; Links et al. 2010; N et al. 2015; Segura-Correa and Solorio-Rivera 2013; Sharma and Saikumar 2010). However, the genetic mechanism affecting the mummify piglets is still unclear, and it has always been one of the difficult problems in pig reproductive traits research. Transcriptome analysis is an efficient and fast tool that has been widely used in animal husbandry (Jäger et al. 2011; Kim et al. 2010; Lin et al. 2019). The purpose of this study was to identify candidate genes associated with the blood of mummification. Annotation of the sequence data using the pig genome as a reference revealed expression of 41,656,932 to 69,203,672 reads in the sow blood transcriptome, and an average of 56,595,204 high-quality reads per sample was mapped to the SSC genome.

The differential expression of blood-related genes is one of the causes of genetic variation in sows during pregnancy, indicating that the regulatory mechanism of pregnancy has changed. In this research, we determined that 1,062 accurately annotated

270 unique genes were differentially expressed. From these genes, 416 genes were
271 upregulated, and 646 genes were downregulated. These genes may be an important
272 factor in inducing mummify piglets. Therefore, it is necessary to use these DEGs for
273 further functional studies to determine the key genes that affect sow reproduction. The
274 comparative analysis of the NM and HM group showed that the differential genes were
275 mainly enriched in some immune and inflammation-related pathways, such as Toll-like
276 receptor signaling pathway, apoptosis signaling pathways, positive regulation of tumor
277 necrosis factor production, innate immune response, placental vascular development,
278 interferon regulation, NF-kappaB signaling, diabetes and FoxO signaling pathways,
279 etc.; on the other hand, it was mainly enriched in some pathways related to reproduction.
280 Such as calcium ion binding, prolactin signaling pathway, development process,
281 ovarian steroid production, and so on.

282 Mummify piglet is a pro-inflammatory state with excessive activation of
283 inflammatory cells and cytokines that may promote placental insufficiency. The
284 functions related to *Interleukin (IL)-9* are mediated by its specific receptor, *IL-9R*. In
285 fact, it is known that in vitro lymphocytes, natural killer cells, and mast cells produce
286 *IL-9*, particularly following exposure to *TGF- β* and *IL-4*(Donninelli et al. 2020; Schmitt
287 et al. 1994). *IL-9* could reduce and regulate infiltration and phagocytic activity of
288 macrophages(Donninelli et al. 2020). *Toll-like receptors (TLR)* are the family of pattern
289 recognition receptors (PRRs). They recognize pathogen associated molecular patterns
290 (PAMPs) and initiate protective immune response(Ao et al. 2019). *TLR8* is one of the
291 main components of the Toll-like receptor family, which participates in the innate

immune process and resists inflammation. *TLR8* is known to be primarily expressed in monocytes/macrophages and myeloid dendritic cells (DCs)(Hornung et al. 2002). DCs are professional antigen-presenting cells (APCs) that play a vital role in shaping adaptive immunity. DC maturation begins when endogenous or exogenous danger molecules are recognized by Toll-like receptors, triggering upregulation of costimulatory molecules and production of immune-polarizing cytokines(Dowling et al. 2008). Study have found that the expression of *TLR8* in endometrial and placental tissues may help prevent pathogens from invading the uterine cavity during the pig estrus cycle and pregnancy(Inkyu et al. 2018). During pregnancy, inflammation may contribute to various disorders so further studies are needed to reveal how exactly the regulation of mentioned gene affects the fetus development leading to mummify piglets.

Embryonic development is a precisely regulated and complex process. *Progesterone receptor (PGR)* play an important role in this. The expression of *PGR* was correlated with *Actin Binding LIM Protein Family Member 3(ABLIM3)*(A et al.). The protein encoded by *ABLIM3* interacts with actin filaments and may be part of adhesion junctions in several cell types. *ABLIM3* was up regulated at 35d and 98d compared with NW group, which suggests that PGR may be one of the causes of mummifying piglets. We also found *Follicle Stimulating Hormone (FSH) α* was downregulated at 35d.*FSH* is a hormone secreted by basophils in the anterior pituitary gland. *FSH* plays a key role in the development and regulation of male and female reproductive systems by acting on *FSHR*, which is mainly expressed in granulosa cells and supporting cells(Simoni et al. 1997). In females, *FSH* induces follicle growth and

maturation and promotes ovulation and luteinization triggered by Luteinizing hormone(Howles 2000). *FSH* treatment of porcine preantral follicles improves the quality of oocytes by promoting growth and higher frequency of embryonic developmental competence(Wu et al. 2007). In the early pregnancy of sows, the expression of *FSH* can be changed to alter the pattern of estradiol, especially progesterone, thereby improving the conditions for embryo survival(Br et al. 2011). The downregulation of *FSH-α* may cause the growth of porcine embryos to be blocked, which caused the occurrence of mummifying piglets.

The formation of mummy piglets may be caused by a variety of diseases. *PRKCZ*, *GCK*, and *LOC100511937* were enriched in diabetes-related pathways in this study. Diabetes not only causes various complications, such as retinopathy, nephropathy, ischaemic heart disease, and neuropathy, but also lead to congenital anomalies, stillbirth, and excessive fetal growth(Schaefer-Graf et al. 2018). Nuclear factor kappa-B (NF-κB) single pathway is mainly related to diabetes(Zhang et al. 2019). Differentially expressed genes *Activating Signal Cointegrator 1 Complex Subunit 1 (ASCC1)* and *TLR8* were related to the NF-κB pathway(Jung et al. 2002; Knierim et al. 2016; Stein et al. 2018). *ASCC1* also caused a profound disturbance of neuromotor unit development and result in prenatal onset spinal muscular atrophy with bone fractures(Knierim et al. 2016). We speculated that differentially expressed genes related to diabetes may interrelate with mummifying piglets.

5 Conclusions

In this study, transcriptome data was generated by RNA-Seq technology, which

will help to further understand the molecular sequences and functions of blood reproduction related genes of pregnant sow at different stages. There were differences in the expression of genes at different pregnancy stages. *IL-9R*, *TLR8*, *ABLIM3*, *FSH- α* , *ASCC1*, *PRKCZ*, and *GCK* may play an important role in course of mummifying piglets. These findings will provide valuable resources for explain the cause of stillbirth mummified piglets and better reproductive performance to improve livestock breeding.

Declarations

Ethics approval and consent to participate

All experimental procedures were performed in accordance with the Institutional Review Board (IRB14044) and the Institutional Animal Care and Use Committee of the Sichuan Agricultural University under permit number DKY-B20140302.

Consent for publication

Not applicable.

Conflicts of interest

The authors declare no conflict of interest.

Funding

This study was supported by grants from the Sichuan Science and Technology Program (2020YFN0024), the Sichuan Innovation Team of Pig (sccxtd-2021-08), the National key R&D Program of China #2018YFD0501204, the National Natural Science Foundation of China (31530073, C170102) and the Earmarked fund for the China Agriculture Research System (No. CARS-35-01A).

Authors' contributions

SW, PW, and KW performed experiments; SW, PW, PW, XJ and DC analyzed data and prepared figures and tables; GT and PW edited and revised manuscript; GT, PW, YL, AJ, WX, YJ, LZ, YZ, XX, XQ, ML and XL conceived, designed research, and wrote this paper. PW, DC, KW, SW, XY, YL, AJ, WX, YJ, LZ, YZ, XX, XQ, ML, XL and GT approved

final version of this manuscript.

Acknowledgements

Not applicable.

References

- A, J.R.W., Yang, Z.B., C, X.P.Z., and Xue, Q.D. Estrogen receptor 1 and progesterone receptor are distinct biomarkers and prognostic factors in estrogen receptor-positive breast cancer: Evidence from a bioinformatic analysis. *Biomedicine & Pharmacotherapy* **121**.
- Anil, S.S., Anil, L., and Deen, J. 2009. Effect of lameness on sow longevity. *Journal of the American Veterinary Medical Association* **235**(6): 734-738.
- Ao, D., Xia, P., Jiang, S., Chen, N., and Zhu, J. 2019. Comparative transcriptome analysis of TLR8 signaling cells revealed the porcine TLR8 specific differentially expressed genes. *Developmental & Comparative Immunology* **98**.
- Bai, X., Zheng, Z., Liu, B., Ji, X., Bai, Y., and Zhang, W. 2016. Whole blood transcriptional profiling comparison between different milk yield of Chinese Holstein cows using RNA-seq data. *BMC Genomics* **17 Suppl 7**(Suppl 7): 512. doi:10.1186/s12864-016-2901-1.
- Br, Uuml, Ssow, K.-P., Schneider, F., Wollenhaupt, K., and Tuchscherer, A. 2011. Endocrine Effects of GnRH Agonist Application to Early Pregnant Gilts. *Journal of Reproduction and Development* **57**(2): 242-248. doi:10.1262/jrd.10-0210.
- Cánovas, A., Rincon, G., Islas-Trejo, A., Wickramasinghe, S., and Medrano, J.F. 2010. SNP discovery in the bovine milk transcriptome using RNA-Seq technology. *Mamm Genome* **21**(11-12): 592-598. doi:10.1007/s00335-010-9297-z.
- do Nascimento, N.C., Guimaraes, A.M.S., Dos Santos, A.P., Chu, Y., Marques, L.M., and

- 386 Messick, J.B. 2018. RNA-Seq based transcriptome of whole blood from
387 immunocompetent pigs (*Sus scrofa*) experimentally infected with *Mycoplasma suis*
388 strain Illinois. *Vet Res* **49**(1): 49. doi:10.1186/s13567-018-0546-6.
- 389 Donninelli, G., Saraf-Sinik, I., Mazziotti, V., Capone, A., and Volpe, E. 2020. Interleukin-9
390 regulates macrophage activation in the progressive multiple sclerosis brain. *Journal of*
391 *Neuroinflammation* **17**(1).
- 392 Dowling, D., Hamilton, C.M., and O'Neill, S.M. 2008. A comparative analysis of cytokine
393 responses, cell surface marker expression and MAPKs in DCs matured with LPS
394 compared with a panel of TLR ligands. *Cytokine* **41**(3): 254-262.
- 395 Giraldez, M.D., Spengler, R.M., Etheridge, A., Goicochea, A.J., Tuck, M., Choi, S.W., Galas,
396 D.J., and Tewari, M. 2019. Phospho-RNA-seq: a modified small RNA-seq method that
397 reveals circulating mRNA and lncRNA fragments as potential biomarkers in human
398 plasma. *Embo j* **38**(11). doi:10.15252/emboj.2019101695.
- 399 Holly, A., and LaVoie. 2017. Transcriptional control of genes mediating ovarian follicular growth,
400 differentiation, and steroidogenesis in pigs. *Molecular Reproduction & Development*.
- 401 Hornung, V., Rothenfusser, S., Britsch, S., Krug, A., Jahrsdorfer, B., Giese, T., Endres, S., and
402 Hartmann, G. 2002. Quantitative Expression of Toll-Like Receptor 1–10 mRNA in
403 Cellular Subsets of Human Peripheral Blood Mononuclear Cells and Sensitivity to CpG
404 Oligodeoxynucleotides. *Journal of Immunology* **168**(9): 4531-4537.
- 405 Horodyska, J., Hamill, R.M., Reyer, H., Trakooljul, N., Lawlor, P.G., McCormack, U.M., and
406 Wimmers, K. 2019. RNA-Seq of Liver From Pigs Divergent in Feed Efficiency
407 Highlights Shifts in Macronutrient Metabolism, Hepatic Growth and Immune Response.

- 408 Front Genet **10**: 117. doi:10.3389/fgene.2019.00117.
- 409 Howles, C.M. 2000. Role of LH and FSH in ovarian function. Molecular & Cellular Endocrinology
- 410 **161**(1-2): 25-30.
- 411 Huang, C., and Zeng, Y. 2020. Association of peripheral blood dendritic cells with recurrent
- 412 pregnancy loss. American Journal of Reproductive Immunology.
- 413 Inkyu, Yoo, Jisoo, Han, Soohyung, Lee, Wonchul, Jung, Ji, and Hoon. 2018. Analysis of stage-
- 414 specific expression of the toll-like receptor family in the porcine endometrium
- 415 throughout the estrous cycle and pregnancy. Theriogenology.
- 416 Jäger, M., Ott, C.E., Grünhagen, J., Hecht, J., and Lienau, J. 2011. Composite transcriptome
- 417 assembly of RNA-seq data in a sheep model for delayed bone healing. BMC Genomics
- 418 **12**(1): 158.
- 419 Jung, D.-J., Sung, H.-S., Goo, Y.-W., Lee, H.M., Park, O.K., Jung, S.-Y., Lim, J., Kim, H.-J.,
- 420 Lee, S.-K., Kim, T.S., Lee, J.W., and Lee, Y.C. 2002. Novel transcription coactivator
- 421 complex containing activating signal cointegrator 1. Mol Cell Biol **22**(14): 5203-5211.
- 422 doi:10.1128/mcb.22.14.5203-5211.2002.
- 423 Kaczynski, P., Goryszewska, E., Baryla, M., and Wacławik, A. 2019. Prostaglandin F2 α
- 424 stimulates angiogenesis at the embryo-maternal interface during early pregnancy in
- 425 the pig. Theriogenology.
- 426 Kamat, M.M., Vasudevan, S., Maalouf, S.A., Townson, D.H., Pate, J.L., and Ott, T.L. 2016.
- 427 Changes in Myeloid Lineage Cells in the Uterus and Peripheral Blood of Dairy Heifers
- 428 During Early Pregnancy. Biology of Reproduction.
- 429 Kim, N.K., Park, H.R., Lee, H.C., Yoon, D., Son, E.S., Kim, Y.S., Kim, S.R., Kim, O.H., and Lee,

- 430 C.S. 2010. Comparative studies of skeletal muscle proteome and transcriptome
431 profilings between pig breeds. *Mammalian Genome* **21**(s5-6): 307-319.
- 432 Knierim, E., Hirata, H., Wolf, N.I., Morales-Gonzalez, S., Schottmann, G., Tanaka, Y., Rudnik-
433 Schöneborn, S., Orgeur, M., Zerres, K., Vogt, S., van Riesen, A., Gill, E., Seifert, F.,
434 Zwirner, A., Kirschner, J., Goebel, H.H., Hübner, C., Stricker, S., Meierhofer, D.,
435 Stenzel, W., and Schuelke, M. 2016. Mutations in Subunits of the Activating Signal
436 Cointegrator 1 Complex Are Associated with Prenatal Spinal Muscular Atrophy and
437 Congenital Bone Fractures. *Am J Hum Genet* **98**(3): 473-489.
438 doi:10.1016/j.ajhg.2016.01.006.
- 439 Koutroumpakis, F., Ahsan, M., Rivers, C.R., Schwartz, M., and Binion, D. 2020. S0808 The
440 Biomarker Peripheral Blood Eosinophilia Predicts Failure to Achieve Mucosal Healing
441 in Inflammatory Bowel Disease. *The American Journal of Gastroenterology* **115**(1):
442 S414-S415.
- 443 Lin, Y., Lv, H., Jiang, M., Zhou, J., Song, S., and Hou, X. 2019. Functional analysis of the dairy
444 cow mammary transcriptome between early lactation and mid-dry period. *Journal of*
445 *Dairy Research* **86**(1): 63-67.
- 446 Ling, Wei, Pengyun, Qiao, Ying, Shi, Yan, Ruan, Jie, and Yin. 2017. Triclosan/triclocarban
447 levels in maternal and umbilical blood samples and their association with fetal
448 malformation - ScienceDirect. *Clinica Chimica Acta* **466**: 133-137.
- 449 Links, I.J., Whittington, R.J., Kennedy, D.J., Grewal, A., and Sharrock, A.J. 2010. An
450 association between encephalomyocarditis virus infection and reproductive failure in
451 pigs. *Australian Veterinary Journal* **63**(5): 150-152.

- 452 Maatz, H., Jens, M., Liss, M., Schafer, S., Heinig, M., Kirchner, M., Adami, E., Rintisch, C.,
453 Dauksaite, V., Radke, M.H., Selbach, M., Barton, P.J., Cook, S.A., Rajewsky, N.,
454 Gotthardt, M., Landthaler, M., and Hubner, N. 2014. RNA-binding protein RBM20
455 represses splicing to orchestrate cardiac pre-mRNA processing. *J Clin Invest* **124**(8):
456 3419-3430. doi:10.1172/jci74523.
- 457 Mach, N., Gao, Y., Lemonnier, G., Lecardonnel, J., and Rogel-Gaillard, C. 2013. The peripheral
458 blood transcriptome reflects variations in immunity traits in swine: towards the
459 identification of biomarkers. *Bmc Genomics* **14**.
- 460 Mallikarjun, Bidarimath, Chandrakant, and Tayade. 2017. Pregnancy and spontaneous fetal
461 loss: A pig perspective. *Molecular Reproduction & Development*.
- 462 N, M., Smit, D, J., Spencer, L, J., Patterson, K, M., Dyck, T, W., and Dixon. 2015. Effects of
463 dietary enrichment with a marine oil-based n-3 LCPUFA supplement in sows with
464 predicted birth weight phenotypes on birth litter quality and growth performance to
465 weaning. *Animal An International Journal of Animal Bioscience*.
- 466 OGAWA, M. 1993. Differentiation and proliferation of hematopoietic stem cells. *Blood* **81**.
- 467 Ramayo-Caldas, Y., Ballester, M., Sánchez, J.P., González-Rodríguez, O., Revilla, M., Reyer,
468 H., Wimmers, K., Torrallardona, D., and Quintanilla, R. 2018. Integrative approach
469 using liver and duodenum RNA-Seq data identifies candidate genes and pathways
470 associated with feed efficiency in pigs. *Sci Rep* **8**(1): 558. doi:10.1038/s41598-017-
471 19072-5.
- 472 Ronzoni, S., Steckle, V., D'Souza, R., Murphy, K.E., Lye, S., and Shynlova, O. 2019. Cytokine
473 Changes in Maternal Peripheral Blood Correlate With Time-to-Delivery in Pregnancies

- 474 Complicated by Premature Prelabor Rupture of the Membranes.
- 475 Schaefer-Graf, U., Napoli, A., Nolan, C.J., and Diabetic Pregnancy Study, G. 2018. Diabetes
476 in pregnancy: a new decade of challenges ahead. *Diabetologia* **61**(5): 1012-1021.
477 doi:10.1007/s00125-018-4545-y.
- 478 Schmitt, E., Germann, T., Goedert, S., Hoehn, P., and Rude, E. 1994. IL-9 production of naive
479 CD4+ T cells depends on IL-2, is synergistically enhanced by a combination of TGF-
480 beta and IL-4, and is inhibited by IFN-gamma. *Journal of Immunology* **153**(9): 3989-
481 3996.
- 482 Segura-Correa, J.C., and Solorio-Rivera, J.L. 2013. Risk factors for stillborn pigs and
483 mummified fetuses in two swine farms in southeastern Mexico. *Livestock Research for*
484 *Rural Development* **25**(10).
- 485 Seo, M., Kim, K., Yoon, J., Jeong, J.Y., Lee, H.J., Cho, S., and Kim, H. 2016. RNA-seq analysis
486 for detecting quantitative trait-associated genes. *Sci Rep* **6**: 24375.
487 doi:10.1038/srep24375.
- 488 Sharma, R., and Saikumar, G. 2010. Porcine parvovirus- and porcine circovirus 2-associated
489 reproductive failure and neonatal mortality in crossbred Indian pigs. *Tropical Animal*
490 *Health & Production* **42**(3): 515-522.
- 491 Shen, J., Zhou, C., Zhu, S., Shi, W., Hu, M., Fu, X., Wang, C., Wang, Y., Zhang, Q., and Yu,
492 Y. 2014. Comparative Transcriptome Analysis Reveals Early Pregnancy-Specific
493 Genes Expressed in Peripheral Blood of Pregnant Sows. *Plos One* **9**.
- 494 Shidid, S., Kohlhoff, S., Norowitz, Y., Hammerschlag, M., and Smith-Norowitz, T. 2020.
495 Chlamydia pneumoniae decreases CD4+ and CD8+ T effector memory IL-2 responses

- 496 in stimulated peripheral blood mononuclear cells in non-asthmatic subjects. *Journal of*
497 *Allergy & Clinical Immunology* **145**(2): AB163.
- 498 Simoni, M., Gromoll, J., and Nieschlag, E. 1997. The Follicle-Stimulating Hormone Receptor:
499 *Biochemistry, Molecular Biology, Physiology, and Pathophysiology* ¹.
500 *Endocrine Reviews* **18**(6): 739-773.
- 501 Stein, T., Wollschlegel, A., Te, H., Weiss, J., Joshi, K., Kinzel, B., Billich, A., Guntermann, C.,
502 and Lehmann, J.C.U. 2018. Interferon regulatory factor 5 and nuclear factor kappa-B
503 exhibit cooperating but also divergent roles in the regulation of pro-inflammatory
504 cytokines important for the development of TH1 and TH17 responses. *Febs j* **285**(16):
505 3097-3113. doi:10.1111/febs.14600.
- 506 Wang, H., Hu, C., Cheng, C., Cui, J., Ji, Y., Hao, X., Li, Q., Ren, W., Deng, B., Yin, Y., Deng,
507 J., and Tan, C. 2019. Unraveling the association of fecal microbiota and oxidative
508 stress with stillbirth rate of sows. *Theriogenology* **136**: 131-137.
509 doi:10.1016/j.theriogenology.2019.06.028.
- 510 Wu, J., Xu, B., and Wang, W. 2007. Effects of luteinizing hormone and follicle stimulating
511 hormone on the developmental competence of porcine preantral follicle oocytes grown
512 in vitro. *Journal of Assisted Reproduction & Genetics* **24**(9): 419-424.
- 513 You, Y.A., Yoo, J.Y., Kwon, E.J., and Kim, Y.J. 2019. Blood Microbial Communities During
514 Pregnancy Are Associated With Preterm Birth. *Other* **10**.
- 515 Zhang, Y.Y., Tan, R.Z., Zhang, X.Q., Yu, Y., and Yu, C. 2019. Calycosin Ameliorates Diabetes-
516 Induced Renal Inflammation via the NF-κB Pathway In Vitro and In Vivo. *Med Sci Monit*
517 **25**: 1671-1678. doi:10.12659/msm.915242.

518 Zhou, Y., Jiang, D., Chu, X., Cheng, W., and Dai, Y. 2021. Prognostic value of peripheral blood
519 T lymphocyte subsets in clear cell renal cell carcinoma. *Translational Andrology &*
520 *Urology* **10**(1): 326-335.

521 Zhuo, Z., Lamont, S.J., Lee, W.R., and Abasht, B. 2015. RNA-Seq Analysis of Abdominal Fat
522 Reveals Differences between Modern Commercial Broiler Chickens with High and Low
523 Feed Efficiencies. *PLoS One* **10**(8): e0135810. doi:10.1371/journal.pone.0135810.

524

Draft

Figure 1. Results of differentially expressed genes.

Figure 2. Correlation, MA, and Volcano plots. (A), (B), (C) and (D) for HM_35d vs. NM_35d, HM_56d vs. NM_56d, HM_77d vs. NM_77d, HM_98d vs. NM_98d, respectively.

Figure 3. Top of GO term enrichment results between HM and NM group. (A), (B), (C) and (D) for HM_35d vs. NM_35d, HM_56d vs. NM_56d, HM_77d vs. NM_77d, HM_98d vs. NM_98d, respectively.

Figure 4. Top20 of KEGG pathway enrichment results between HM and NM group. (A), (B), (C) and (D) for HM_35d vs. NM_35d, HM_56d vs. NM_56d, HM_77d vs. NM_77d, HM_98d vs. NM_98d, respectively.

Table 1 Descriptive statistics of sample sows

Group	Sample ID	Parity	TNB (mean ± SD)	mummify piglets (mean ± SD)
HM	HM1	3	10.67±1.70	2.00±0.82
	HM2	3	10.67±2.05	2.33±0.94
	HM3	3	11.33±1.70	3.33±0.47
	HM4	3	13.67±0.94	2.33±0.47
	HM5	3	15.00±2.94	2.67±0.94
	HM6	3	11.00±2.16	2.33±0.47
	HM7	3	13.33±1.25	2.33±0.47
	HM8	3	12.00±1.63	2.33±0.94
NM	NM1	3	11.67±0.47	\
	NM2	3	11.33±0.47	\
	NM3	3	10.33±1.25	\
	NM4	3	12.00±0.82	\
	NM5	3	10.67±2.05	\
	NM6	3	12.67±0.94	\
	NM7	3	12.67±0.47	\
	NM8	3	11.00±0.82	\

Mean and SD concentrations were given as the mean ± SD.

Table 2 DEG verifying primer sequence

Gene	Sequence 5`-3`	Temperature (°C)
<i>β-actin</i>	F: AAGGACCTCTACGCCAACAC	60
	R: CTGGCTGATCCACATCTGCT	
<i>DENND5A</i>	F: GAACATCGGGGAGATCAAGA	60
	R: AGGAAGGCATAGCGCTTGTA	
<i>GLB1L</i>	F: CAGGACCGTGGCTTTATGTT	60
	R: CCCTTTTCCCCATCAAAAAT	

<i>GAREM2</i>	F: AGCGAGACATCCTGCTCATT R: AAGATACGGTCGGGGAAGAC	60
<i>TLR8</i>	F: TGTCATTGCAGAGTGCACAA R: GAGAAACGCCCCATCTGTAA	60
<i>TAF10</i>	F: GGAGGATTACACGCCTACGA R: GTGTACTTGCGGTCCTTGCT	60
<i>RAB5C</i>	F: CAAGGGACAATTCCACGAGT R: GGCTGGCCTGTCTCTGTAAC	60
<i>UCP2</i>	F: GGACGTGGTCAAGACGAGAT R: CAAACATCACCACGTTCCAG	60
<i>CASQ1</i>	F: CTGGCACTGCTCTACCATGA R: GCGTCCTTCTCAGAGTCCAC	60
<i>APBB1</i>	F: ACCACTCGAAACTGGTGGAC R: GGGTTTAGCAACAGGCACAT	60
<i>ZNF76</i>	F: CACCGCATAATTACCAAG R: CTAAGCCACCTCTGTCTGC	60
<i>SDHD</i>	F: GGAGGCTCAGTGTTCTTTGC R: CTGGGTGACAGGTGAATGTG	60

Table 3 Summary of RNA-seq data

Sample ID	Total Reads	Q30 (%)	GC (%)	Mapped Reads	Mapped Rate (%)
HM1_35d	62487618	0.9527	0.5798	59,427,456	95.10%
HM2_35d	51966574	0.9569	0.5856	49,209,432	94.69%
HM3_35d	58295166	0.9541	0.5936	55,514,468	95.23%
HM4_35d	58043220	0.9519	0.5717	54,250,938	93.47%
HM5_35d	68443314	0.9486	0.5655	64,195,352	93.79%
HM6_35d	66772370	0.9522	0.5732	63,314,238	94.82%
HM7_35d	53646658	0.9568	0.5883	51,359,576	95.74%
HM8_35d	53525350	0.9475	0.5812	50,751,552	94.82%
HM1_56d	57225494	0.9376	0.587	52,833,491	92.33%
HM2_56d	64458452	0.9469	0.5789	60,867,068	94.43%

HM3_56d	53301044	0.9455	0.5807	49,352,210	92.59%
HM4_56d	61513794	0.9523	0.5824	57,769,587	93.91%
HM5_56d	54895964	0.9515	0.5751	52,079,253	94.87%
HM6_56d	56536532	0.9518	0.5762	52,946,444	93.65%
HM7_56d	69203672	0.955	0.5908	66,212,991	95.68%
HM8_56d	48029722	0.9354	0.5792	45,182,924	94.07%
HM1_77d	54168224	0.9468	0.5888	50,653,336	93.51%
HM2_77d	65511674	0.9474	0.5831	61,411,916	93.74%
HM3_77d	56785464	0.9536	0.5798	53,869,048	94.86%
HM4_77d	62523506	0.95	0.5908	59,389,094	94.99%
HM5_77d	47961798	0.951	0.5848	45,348,361	94.55%
HM6_77d	54940596	0.9505	0.5938	52,196,592	95.01%
HM7_77d	66831048	0.9451	0.5962	62,457,604	93.46%
HM8_77d	52014634	0.9456	0.5788	49,515,092	95.19%
HM1_98d	56399724	0.9514	0.5856	53,692,958	95.20%
HM2_98d	59162000	0.9488	0.5852	56,082,792	94.80%
HM3_98d	50040926	0.9472	0.569	46,449,843	92.82%
HM4_98d	54032580	0.9518	0.5813	51,193,623	94.75%
HM5_98d	59433650	0.9495	0.5787	56,776,080	95.53%
HM6_98d	60600680	0.9501	0.5712	57,408,991	94.73%
HM7_98d	63334688	0.9514	0.583	60,108,091	94.91%
HM8_98d	53100274	0.9497	0.5838	50,152,050	94.45%
NM1_35d	67733656	0.9537	0.5821	64,066,674	94.59%
NM2_35d	56988580	0.9509	0.5843	54,420,235	95.49%
NM3_35d	54736262	0.9488	0.5916	52,039,877	95.07%
NM4_35d	54480012	0.9456	0.5779	51,603,294	94.72%
NM5_35d	44462950	0.9376	0.5727	41,921,124	94.28%
NM6_35d	56231816	0.9471	0.5899	53,389,888	94.95%
NM7_35d	62600302	0.957	0.5876	59,644,575	95.28%
NM8_35d	60311294	0.9503	0.5844	56,895,987	94.34%
NM1_56d	49060696	0.946	0.5758	45,826,270	93.41%
NM2_56d	48614586	0.9555	0.581	46,789,104	96.24%
NM3_56d	68034576	0.9523	0.5818	65,508,362	96.29%
NM4_56d	49025624	0.9351	0.5797	45,185,990	92.17%
NM5_56d	41656932	0.9408	0.5838	38,551,517	92.55%

NM6_56d	49501900	0.9376	0.5801	46,657,280	94.25%
NM7_56d	54737078	0.9465	0.5833	51,559,191	94.19%
NM8_56d	67506748	0.9545	0.5763	63,780,694	94.48%
NM1_77d	57242542	0.9543	0.5967	54,315,473	94.89%
NM2_77d	49863948	0.9415	0.5888	47,001,979	94.26%
NM3_77d	47847380	0.939	0.5834	44,649,962	93.32%
NM4_77d	49162716	0.9417	0.5801	46,141,262	93.85%
NM5_77d	55187380	0.9412	0.5945	51,921,259	94.08%
NM6_77d	57036950	0.9424	0.5932	53,566,878	93.92%
NM7_77d	52837370	0.9481	0.5891	49,817,624	94.28%
NM8_77d	59280262	0.9485	0.5882	55,923,949	94.34%
NM1_98d	55946166	0.9504	0.5714	51,054,754	91.26%
NM2_98d	56111920	0.9507	0.5858	53,544,943	95.43%
NM3_98d	56379812	0.9452	0.5894	53,054,585	94.10%
NM4_98d	60114526	0.952	0.5895	56,770,456	94.44%
NM5_98d	55500824	0.9475	0.5673	51,792,137	93.32%
NM6_98d	63648164	0.95	0.5919	60,796,935	95.52%
NM7_98d	51596890	0.9503	0.5856	48,510,687	94.02%
NM8_98d	59359224	0.9508	0.5828	56,312,913	94.87%

Table 4 Detail information on the top 10 most differentially expressed genes of HM vs. NM group

Group	Gene name	log2FC	P-value
HM_35d vs. NM_35d	<i>ATP6</i>	-4.55	2.39E-05
	<i>LOC100158003</i>	4.44	1.33E-05
	<i>GLB1L</i>	-3.25	1.80E-05
	<i>GAREM2</i>	-2.95	3.96E-05
	<i>DENND5A</i>	2.44	2.51E-03
	<i>TMOD4</i>	-2.13	1.31E-04
	<i>FBLN2</i>	1.74	1.34E-04
	<i>ARL4A</i>	-1.71	6.42E-04
	<i>BAMBI</i>	-1.69	1.41E-03
	<i>ENSSSCG00000024263</i>	1.67	7.81E-03
HM_56d vs. NM_56d	<i>TLR8</i>	5.61	9.52E-04
	<i>TAF10</i>	-5.01	1.73E-06
	<i>GAREM2</i>	-4.24	2.55E-08

HM_77d vs. NM_77d	<i>APBB1</i>	-3.30	2.54E-06
	<i>ENSSSCG00000038670</i>	2.63	6.77E-04
	<i>CCDC62</i>	2.59	1.70E-03
	<i>INO80B</i>	2.04	6.43E-03
	<i>ENSSSCG00000037879</i>	-1.94	3.64E-03
	<i>INO80E</i>	1.88	4.11E-04
	<i>CIPC</i>	1.74	9.27E-04
	<i>TLR8</i>	4.06	8.46E-03
	<i>GPX3</i>	3.43	1.21E-04
	<i>UCP2</i>	-2.99	3.04E-04
HM_98d vs. NM_98d	<i>RAB5C</i>	-2.31	1.21E-03
	<i>GLB1L</i>	-1.81	1.88E-03
	<i>GAREM2</i>	-1.75	4.42E-03
	<i>ENSSSCG00000032137</i>	1.67	1.38E-03
	<i>FARP2</i>	1.39	9.79E-03
	<i>SH3BP4</i>	1.37	9.72E-04
	<i>COL6A2</i>	-1.33	2.98E-03
	<i>CCNB1IP1</i>	4.68	7.94E-13
	<i>GAREM2</i>	-3.59	2.94E-06
	<i>CASQ1</i>	3.31	5.57E-07
HM_98d vs. NM_98d	<i>SDHD</i>	3.17	6.56E-06
	<i>ZNF76</i>	-3.01	2.39E-06
	<i>IL9R</i>	2.38	1.58E-08
	<i>LGALS7</i>	-2.36	2.79E-03
	<i>APBB1</i>	-2.24	1.74E-04
	<i>ENSSSCG00000038322</i>	-2.21	1.02E-03
	<i>CCDC107</i>	2.11	6.77E-08

Table5 Summary of qPCR and RNA-Seq results

Gene name	Stage	RNA-Seq (log2FC)	qPCR validation (log2FC)
<i>DENND5A</i>	35d	2.44	1.54
<i>GAREM2</i>	35d	-2.95	-2.77
<i>GLB1L</i>	35d	-3.25	-1.97
<i>APBB1</i>	56d	-3.30	-2.38

<i>TAF10</i>	56d	-5.01	-3.57
<i>TLR8</i>	56d	5.61	4.1
<i>GLB1L</i>	77d	-1.81	-0.98
<i>RAB5C</i>	77d	-2.31	-1.23
<i>UCP2</i>	77d	-2.99	-2.12
<i>CASQ1</i>	98d	3.31	3.01
<i>SDHD</i>	98d	3.17	1.77
<i>ZNF76</i>	98d	-3.01	-2.38

Draft