Frozen out: unanswered questions about winter biology

Journal:	<i>Environmental Reviews</i>
Manuscript ID	er-2020-0127.R1
Manuscript Type:	Perspective
Date Submitted by the Author:	30-Apr-2021
Complete List of Authors:	<p>Sutton, Alex; Kansas State University; University of Guelph, Department of Integrative Biology Studd, Emily; University of Alberta, Department of Biological Sciences Fernandes, Timothy; University of Toronto Mississauga, Department of Biology Bates, Amanda; Memorial University of Newfoundland, Department of Ocean Sciences Bramburger, Andrew; Watershed Hydrology and Ecology Research Division Environment and Climate Change Canada Cooke, Steven; Carleton University Department of Biology, Biology Hayden, Brian; University of New Brunswick, Biology Department Henry, Hugh A. L.; University of Western Ontario, Department of Biology Humphries, Murray; McGill University, Department of Natural Resource Sciences Martin, Rosemary; University of Toronto Mississauga, Department of Biology McMeans, Bailey; University of Toronto Mississauga, Department of Biology Moise, Eric; Canadian Forest Service O'Sullivan, Antoin M.; University of New Brunswick, Biology Department Sharma, Sapna; York University, Department of Biology Templer, Pamela H; Boston University, Department of Biology</p>
Is this manuscript invited for consideration in a Special Issue? :	Not applicable (regular submission)
Keyword:	Climate change, Communities, Ecosystems, Individuals, Populations

Frozen out: unanswered questions about winter biology

Alex O. Sutton^{1,2,*}, Emily Studd³, Timothy Fernandes⁴, Amanda E. Bates⁵, Andrew Bramburger⁶, Steven J. Cooke⁷, Brian Hayden⁸, Hugh A. L. Henry⁹, Murray Humphries¹⁰, Rosemary Martin⁴, Bailey McMeans⁴, Eric Moise¹¹, Antin M. O’Sullivan⁸, Sapna Sharma¹², Pamela H. Templer¹³

¹ Division of Biology, Kansas State University, Manhattan, Kansas, United States of America

² Department of Integrative Biology, University of Guelph, Guelph, ON, Canada

³ Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

⁴ Department of Biology, University of Toronto Mississauga, Mississauga, ON, Canada

⁵ Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, NL, Canada

⁶ Watershed Hydrology and Ecology Research Division, Environment and Climate Change Canada, Burlington, Ontario, Canada

⁷ Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, Ottawa, ON, Canada

⁸ Canadian Rivers Institute, Biology Department, University of New Brunswick, Fredericton, NB, Canada

⁹ Department of Biology, University of Western Ontario, London, ON, Canada

¹⁰ Department of Natural Resource Sciences, Macdonald Campus, McGill University, Ste-Anne-de-Bellevue, QC, Canada

¹¹ Natural Resources Canada - Canadian Forest Service, 26 University Drive, Corner Brook, NL

¹² Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, Canada

¹³ Department of Biology, Boston University, 5 Cummington Mall, Boston, United States of America

Running title: Unanswered questions about winter biology

Statement of authorship: All authors contributed substantially to the conception, writing and revisions of this manuscript.

Data accessibility statement: No data was used in this manuscript and, as a result, there is no data to deposit in a public repository.

Key words: Climate change, Communities, Ecosystems, Individuals, Populations

Number of figures, tables: 3 figure, 0 tables

* *corresponding author:* Alex O. Sutton, alexosutto@gmail.com

Draft

Abstract

Winter conditions impose dramatic constraints on temperate, boreal, and polar ecosystems, and shape the abiotic and biotic interactions underpinning these systems. At high latitudes, winter can last longer than the growing season and may have a disproportionately large impact on organisms and ecosystems. Even so, our understanding of the ecological implications of winter is often lacking. Indeed, even what exactly defines winter is currently unclear and boundaries that delineate this season are blurred across marine, freshwater, and terrestrial realms and fields of biology. Here, we discuss the complexity of defining winter and highlight the importance of maintaining the capacity to test hypotheses across seasons, realms, and domains of life. We then outline questions drawn from diverse fields of research that address current gaps in our understanding of winter ecology and how winter influences multiple levels of biological organization, from individuals to ecosystems. Finally, we highlight potential consequences of changes to both the length and severity of winter due to climate change and discuss the role winter may play in mediating ecosystem function in the future.

Introduction

Ecologists have traditionally regarded winter as a dormant time of year; a time when most organisms actively or passively depress metabolism or seek out warmer climates until spring arrives. This view, in combination with the arrangement of academic calendars and logistical difficulties associated with field work during the coldest months of the year, has restricted the scope of research conducted during winter and an underrepresentation of the ecological importance of winter (Marchand 2013; Studd et al. 2021). Despite the many challenges, a growing number of ecologists who brave winter conditions are striving to reveal its many snow-covered and ice-encased secrets. In direct contrast to historical attitudes around winter ecology, recent studies have provided strong evidence that winter is far from dormant (Berge et al. 2015; Hampton et al. 2017; McMeans et al. 2020). In some systems, winter can be a period of unexpectedly high productivity (Darnis et al. 2012), while in others, specialized organisms remain active year-round, maintaining interactions and energy flow between species (Humphries et al. 2017), and even underpinning species coexistence (McMeans et al., 2020). Regardless of an organism's overwintering strategy, winter conditions can shape survival and reproduction (Berteaux et al. 2017), population dynamics (Kreyling 2010), and food web structure (Tremblay et al. 2012; McMeans et al. 2015). Thus, winter is a critical period for which we need a better understanding.

Winter imposes unique challenges on survival for many taxa; however, it can also introduce ecological opportunities, such as reduced competition and access to otherwise rare resources that are critical for many species at high latitudes (Bokhorst et al. 2009; Farmer et al. 2015; Forrest 2016). However, at high latitudes, winter is changing more rapidly than any other season (IPCC 2013). With the continued acceleration of global climate change, boreal and arctic

systems have experienced some of the most dramatic warming on the planet, with annual average temperatures rising 2.7°C in the past 30 years (IPCC, 2013; Box et al. 2019), much higher than rates of change in tropical ($0.26 \pm 0.05^{\circ}\text{C}$; Malhi and Wright 2004) and temperate systems (0.9°C ; Zhang et al. 2000). Water bodies that historically froze for months every year now go full winters without freezing over (Sharma et al. 2019). In less extreme situations, freeze-thaw events are becoming more common due to the ever-increasing frequency of winter whiplash – drastic and rapid oscillations in temperature and precipitation (Cohen 2016; Swain et al. 2018; Casson et al. 2019). For organisms and systems that are adapted to cold, dark, and snowy periods, a warming climate that drives winter temperatures beyond critical thresholds (e.g., 0°C , the freezing point of water) will alter outcomes at all levels of biological organization and introduce a cascade of biological consequences across systems.

Recent climate change research has led to the realization that winter warming and winter weirding (i.e. deviations from traditional norms of winter) may have far greater impacts on biodiversity than previously appreciated, generating increased motivation for studying winter and the effects of winter conditions on the ecology of species and ecosystems. As more researchers begin to focus on this understudied season, there is a need to pause and identify the major knowledge gaps. Here, we bring together a group of scientists with diverse expertise across ecosystems, taxa, and biological scales to pose questions we believe are most crucial to advancing our understanding of how winter influences individuals, populations, communities, and ecosystems, and how climate change impacts these relationships. We start by discussing how researchers have attempted to define what winter is, highlighting the range of perceived definitions and the complexity that exists in defining this season across systems and organisms. We then underscore the importance of using a definition of winter that allows for explicit

hypothesis testing about the role of winter in shaping biological phenomena across time, taxa, and systems. Next, we focus on fundamental questions that will advance our understanding of natural systems, explicitly considering organism-, population-, community-, and ecosystem-scale impacts. We conclude by considering the management implications of continuing to ignore the impacts of winter on wildlife and highlight current knowledge gaps surrounding winter in the context of continued climate change.

What is winter?

Images of temperate, boreal and Arctic winters commonly depict snow, ice, and cold weather; however, defining winter for these latitudes can be difficult. In the broadest terms, winter is the coldest season of the year caused by the shifting of the Earth's rotational axis during its migration around the sun which results in spatially heterogeneous fluxes of incoming solar radiation. While each part of the planet goes through 'winter', the seasonal amplitude of temperature is most severe in the northern hemisphere (Serreze 2015; Turner 2015), driven primarily by the interactions of land mass and oceanic heat fluxes. Land masses increase average summer temperature and decrease winter average temperatures, whereas water masses reduce variation in average annual temperature (Kraus and Morrison 1966; Dickinson 2015). In the northern hemisphere at north-temperate and Arctic latitudes, winter is characterized by long, cold, dry periods, where temperatures can reach -49 °C, and daylight can be almost entirely absent (Serreze 2015). These fluxes of energy, heat, and light from solar radiation underlie variation in what winter is, but are not necessarily accounted for when authors define winter itself.

Defining winter remains a contentious issue and definitions are often specific to a focal system or geographic area. Meteorological definitions confine winter to the three coldest months of the year (Huschke 1959), while more recent works suggest delineations of winter that depend on the satisfaction of specific environmental criteria, such as maximum daily temperatures (Mayes et al. 2015) or lake and river ice cover (Shuter et al. 2012). However, these definitions of winter can vary dramatically between biomes, geographical locations, terrestrial and aquatic ecosystems, and even freshwater and marine systems (e.g., Kielland et al. 2006; Contosta et al. 2019). Such definitions may limit the range of testable ecological hypotheses, as they outline a ‘bounding box’ for winter that depends on focal variables (e.g., ice cover, temperature). As such, the focus is shifted towards the temporal axis on which winter’s bounding box sits (i.e. the length and timing of when winter-defining characteristics are achieved), as opposed to the focal variables themselves. Compartmentalized definitions that constrain “winter” to system-specific conditions hinder our ability to adopt a general framework for describing winter, and likely limit the translation of existing knowledge between levels of biological organization and across geographic locations (Figure 1).

In light of ongoing climate change, it is imperative to forecast how continuing changes in climate will impact organisms. Although characterising winter at a local or system-specific scale has value, generalizing more broadly will also allow for an understanding of how winter is changing across space. This could, in turn, allow researchers to identify ecosystems that are experiencing the most rapid changes, or even quantify how variation in winter conditions influence individuals, populations, communities, and ecosystems. Without a generalizable definition outlining the bounds of winter, or how to characterize winter outside of single focal

systems, it will be difficult, if not impossible, to develop broad characterizations of how winter is changing and how these ongoing changes influence organisms.

Numerous studies have proposed unifying definitions of winter to facilitate broader comparisons across geographic areas and between ecosystems (e.g. Ladwig et al. 2016; Contosta et al. 2020). These proposed definitions attempt to define winter based on select characteristics and typically rely on the fact that winter is considered to be a ‘frigid’ season (Contosta et al. 2020). For example, Ladwig et al. (2016) defined the timing and duration of winter based on the 90% freeze/frost probability for each site in their analysis. Similarly, Contosta et al. (2020) define the start and end of winter based on the occurrence of four days of below freezing air temperatures and snow cover within a seven-day period. In contrast to these meteorological definitions of winter that rely on temperature or snow cover data, we propose a two-step approach that can be used to test hypotheses across temporal (e.g. inter-annual) and spatial (e.g. between systems and geographically) scales. First, we propose that photoperiod should be used as the overarching characteristic to bound winter to facilitate temporal comparisons (‘meteorological’ definition of winter), as photoperiod represents a causal mechanism driving seasonal conditions. Though the magnitude of photoperiodic variation can be extensive and varies widely based on latitude, the phenology of photoperiodic cycles changes little from year to year and is relatively consistent across biomes and geographic regions. By delineating boundaries that are relatively static in time, the magnitude and phenology of focal variable fluctuations within that window can be used to quantify system-specific winter conditions, while also facilitating comparisons across time.

Once photoperiod has been used to bound the length of winter, a condition-dependent definition using maxims (a maxim is a fundamentally held truth or principle) can be leveraged to

outline the characteristics of contemporary winters that facilitate and motivate hypothesis generation in winter ecology (Studd et al. 2021). Studd et al. (2021) proposed nine maxims of winter (1. Cold, 2. Frozen, 3. Snowy, 4. Dark, 5. Less productive, 6. Variable, 7. Deadly, 8. Understudied, and 9. Changing), that provide a clear way to both define and compare winter between systems, and spatial and temporal scales (Figure 2). For example, maxims can highlight similar constraints between systems imposed by a suite of environmental factors. Predictable changes in photoperiod lead to dark winters and phenomena like ‘polar night’ (when night lasts for longer than 24 hours) at high latitudes, while snow and ice cover can cause winter to be dark in aquatic systems. Additionally, while these maxims clarify ‘what winter is’, they also allow for broad comparisons to be made across systems and can be used to assess the relative impact that predictable changes in abiotic conditions (i.e. shorter photoperiods and colder temperatures) have on organisms, communities, and ecosystems. Maxim seven (‘winter is deadly’) illustrates this point because in terrestrial ecosystems winter may be most deadly due to snowy conditions or cold temperatures. In contrast, in limnic and marine environments winter may be deadly not because temperatures fluctuate greatly, but because of ice cover that potentially limits oxygen concentrations, or the growth of photosynthetic organisms that many species rely on for food (Studd et al. 2021). Establishing maxims that exist for various regions that experience winter can help to guide ecological research and provide a way to organize comparisons between systems that otherwise seem disparate.

We have outlined a series of questions within the context of these maxims of winter, arranged according to levels of biological organization (Organisms, Populations, Communities, Ecosystems). The final sections review potential challenges to natural resource management and climate change research driven by winter processes (Management Concerns, Climate Change).

All questions are preceded by a brief review of known influences of winter on each level of organization and key areas of uncertainty, where questions are referred to explicitly in-text to help direct the reader. All questions focus on the direct and indirect effects of winter at each level of biological organization to highlight key uncertainties and expand our understanding of the importance of winter (Figure 3). Many of these questions can be applied across systems and taxa to understand the broad consequences of winter. However, some questions are specific to a particular system (e.g. terrestrial, limnic or marine) and we state alongside these questions which system or taxa they are directed towards in these cases.

Organisms (general and physiology/behaviour)

Environmental stressors are experienced, and responded to, at the individual-level, and thus a comprehensive understanding of organismal biology is central to determining the effects of winter. The reduction of temperature-dependent biochemical reaction rates coupled with the threat of chill injury and thermal stress has given rise to many unique survival strategies (Questions 1-3; Sinclair 1999; Shuter et al. 2012; Convey 2013; Bartlett et al. 2020). Despite an ability to (partially) mitigate stress and mortality, performance nonetheless varies across a myriad of dimensions (e.g. taxa, thermal variability; Williams et al. 2012 a,b), and this context-dependency will necessitate the consideration of many intrinsic and extrinsic factors (Questions 4-5). Colder temperatures can also induce oxidative stress, and the efficacy of physiological strategies to cope with radical production varies extensively (Questions 6-7; Lalouette et al. 2011; Stier et al. 2014; see Blagojevic et al. 2011). How these stressors influence organism life history and overwinter survival outcomes remains very poorly understood (Questions 8-9). Moreover, while ectotherms can benefit from reduced energy depletion, costs increase for

endotherms as temperatures fall below their thermal-neutral range (Romano et al. 2019). These unique winter maintenance and activity costs likely influence other aspects of organismal physiology, including immunity and the microbiome (Question 10); however, very little has been done to directly quantify these influences.

Cold exposure is also necessary for the initiation of many life history processes. Many insects and plants have minimum chilling requirements prior to the resumption of development (Sung & Amasino 2004; Forrest 2016). Some fish species also require low temperatures during winter to optimize reproductive success (Farmer et al. 2015). However, cues (e.g. photothermal regimes) that emerge during winter and their resultant influences have yet to be fully unpacked. Extensive snow and ice cover limits oxygen availability in terrestrial and aquatic systems (though oxygen saturation through convective mixing is still possible in large lakes: Yang et al. 2017), introduces thermal refugia for endotherms (Pauli et al. 2013), influences mobility (Parker et al. 1984; Richard et al. 2014), and attenuates light (Blanchfield et al. 2009). The consequences of these environmental conditions on organism life history are not well studied, though they likely mediate activity, survival, and reproductive outcomes across systems.

Consequences of winter stress are further modulated by variability in abiotic conditions (Question 11; e.g., extreme events: Bokhorst et al. 2009). Both mid-winter warm spells and cold snaps impact energetics (Question 12; ectotherms: Hahn and Denlinger 2011; endotherms: Humphries and Umbanhowar 2007), and can induce injury or death across both terrestrial and aquatic systems (Salt 1961; Donaldson et al. 2008; Lukatkin et al. 2012). Often, these temperature extremes occur in quick succession, resulting in weather whiplash and freeze-thaw cycles (Question 13). Snowmelt, for example, exposes organisms to refreeze, with consequences ranging from cold injury and ice encasement to substantial energy savings (Marshall and Sinclair

2012; Gudleifsson 1993; Hoback and Stanley 2001). Likewise, the exposure of soils to cold temperatures can result in frost heave and plant root damage (Question 14; Groffman et al. 2001; Bergsten et al. 2011). Though generally more stable, convective upwelling and wind fetch in marine systems can drive large daily temperature swings upwards of 18°C (Bates et al. 2018). Our understanding of how these winter processes influence organism-level outcomes remains rudimentary in many systems, despite their potentially far-reaching implications.

Beyond temperature fluxes, winter also presents a unique suite of anthropogenic disturbance regimes. Artificial snowscapes such as managed ski trails can alter habitat quality for arthropods (Negro et al. 2010). Similarly, ice manipulation along shipping lanes facilitates the penetration of light and oxygen into the water column. Even for a given disturbance that can be experienced across multiple seasons (e.g. catch and release angling), the combination of intrinsic and extrinsic conditions specific to the winter period might influence organism responses. Our ability to predict how anthropogenic disturbances will affect overwintering organisms is limited by the lack of current consideration for such relationships (Question 15). Quantitative analyses across disturbance gradients are needed to better understand and manage systems that are impacted by anthropogenic activity during the winter.

Winter effects are often considered in a short-term context (e.g. acute metabolic changes, mortality), but can also be expressed over time as individuals exhibit chronic and sub-lethal energetic costs of winters past (Questions 16 - 17). Abnormally warm winters lead to, for example, delayed invertebrate emergence (Stålhandske et al. 2015). Winter experiences can also result in stress “memories”, encoded in the form of epigenetic markers with transgenerational consequences (Questions 18 - 19; Bruce et al. 2007). In addition to winter experiences dictating performance in other seasons (i.e. through carry-over effects), the reverse is also true as

processes such as resource accumulation, behavioural modification, and physiological preparation during the active season can all play essential roles in subsequent overwintering success (Questions 19 - 20; Sinclair 2015; Gilbert et al. 2010; Fernandes & McMeans 2019). The number of seasonal cycles experienced may also shape these seasonal strategies and their resultant outcomes (Question 21). However, the identification of these relationships has only recently begun. Developing a better understanding how winter conditions directly or indirectly shape individual physiology and ecology will be central to successful management moving forward.

Questions:

1. How entrained are (aspects of) winter strategies? How much do the strategies utilized depend on individual experience, energetic condition, and environmental context?
2. What winter conditions drive the breakdown of seasonal buffering mechanisms? What are the sublethal implications of such a breakdown?
3. To what extent is behaviour constrained by physiological state during winter?
4. What is the relative importance of intrinsic vs. extrinsic cues and triggers that influence decisions (e.g., when to stir) during winter?
5. How do light, temperature, and oxygen (aquatic only) regimes experienced during winter interact to shape life history outcomes and decisions (e.g., reproduction, growth, migration) in both the short and long term?
6. How do winter conditions influence the relative risk and impact of oxidative damage?
7. What role does the production and accumulation of reactive oxygen species play in mediating physiological responses to winter and winter-time decision making?
8. What are the critical winter oxygen levels for survival and activity in aquatic organisms?

- 295 9. To what extent does aerobic scope constrain the activity of ectotherms at low temperatures?
- 296 10. How does winter shape the microbiome of wild organisms?
- 297 11. To what extent do the frequency, duration, intensity, and timing of winter stressors
- 298 affect performance?
- 299 12. How do the consequences of weather whiplash events differ amongst ectothermic and
- 300 endothermic organisms?
- 301 13. How much more costly are weather whiplash events for terrestrial species relative to
- 302 temperature fluctuations that remain below the important biophysical and physiological threshold
- 303 of 0°C?
- 304 14. What is the relative importance of physical effects driven by winter processes (e.g., frost
- 305 heave, frazil ice, wave regimes/wind fetch) for determining stress/physiological risk in
- 306 organisms?
- 307 15. What is the impact of winter disturbance (e.g., anthropogenic noise, capture and handling,
- 308 storms, forest harvest) on individual performance relative to disturbances experienced in other
- 309 seasons? How does this cost vary when the disturbance is the same (e.g., catch-and-release
- 310 angling) vs. season-specific (e.g., winter ice storms vs. summer heat wave)?
- 311 16i. How do levels of stress imposed by winter (e.g. freezing, hypoxia) interact or shape the
- 312 ability of organisms to cope with stress during the summer (e.g. heat, drought)?
- 313 16ii. Do traits that maximize survival during harsh winters trade-off with those that maximize
- 314 survival during harsh summers?
- 315 17. What is the influence of winter stressors relative to other factors (e.g. natural enemies, host
- 316 quality) in the establishment of alien organisms?
- 317 18. What is the relative importance of carry-over effects that emerge in winter vs. from winter?

19. To what extent are winter carry-over effects responsible for shaping major life history events (i.e. migration, spawning cycles, post-winter emergence phenology)?

20. What is the role of nutritional quality (e.g., micro- and macronutrients) in preparing for or sustaining life in winter? Are there specific micronutrients that are most limiting?

21. What is the relative energetic cost of seasonal acclimatization?

22. To what extent do winter stressors drive life history characteristics in short vs. long-lived organisms?

Populations

Though environmental conditions during winter can directly shape ecological and physiological consequences at the individual level, emergent properties that manifest only at higher levels of biological organization necessitate broad-scale questions and investigations (Question 23).

Studies of a variety of taxa from diverse biomes have found that winter is an important portion of the annual cycle with consequences on population vital rates and population growth (Altwegg et al. 2005; Altwegg et al. 2006; Gamelon et al. 2017; Woodworth et al. 2017; Bargmann et al. 2020). However, despite the growing recognition of the importance of winter in understanding population growth, there remains a wide array of questions to be answered about the influence of winter on populations.

Leveraging the growing number of published studies investigating how winter influences populations can provide insight into broader trends of the mechanisms by which events in winter influence populations and how interspecies variability in life history traits may influence their susceptibility to the effects of winter (Question 24; Altwegg et al. 2005). Abiotic winter conditions are a potentially important selective filter, but variation in the behavioural and

physiological responses of individuals to these harsh abiotic conditions can also buffer or magnify the effects of these conditions on populations (Leirs et al. 1997; Coulson et al. 2002; Uelman Jr et al. 2016; Froy et al. 2019; Masoero et al. 2020). Broader life-history characteristics, such as life span (Sæther and Bakke 2000), overwintering strategy (e.g. hibernation, migration or remaining active) and behaviour during winter (Sutton et al. 2016; Sutton et al. 2021), could also mediate or enhance the effects of winter on population vital rates (Questions 25 - 27). Additionally, if animals aggregate in smaller patches of suitable habitat it will also be important to understand how density-dependent processes are likely to interact with winter to shape population dynamics (Questions 28 - 29).

Understanding broader patterns that describe the varying impacts of winter can predict population-level responses to winter for data-deficient species and future population trends from across a species' range, rather than restricting inference to a single population. Evidence suggests that populations at certain range extents may be more sensitive to winter severity and changes to environmental conditions (Ringsby et al. 1999; Veteli et al. 2005; Waite and Strickland 2006). These populations could be used to predict future responses to winter conditions as climate change continues to cause shifts in environmental conditions. Furthermore, comparative analyses of multiple populations across a species' range, may provide insight into the generalizability of responses to winter and allow us to better predict how species, not just focal populations, respond to a range of winter conditions (Question 30). Synthesis across systems (e.g. aquatic vs. terrestrial) will also help to identify possibly divergent mechanisms by which winter can influence population dynamics (Question 31).

Questions:

23. To what extent do carry-over effects on individuals contribute to population dynamics?

- 364 24. Does winter have a disproportionate effect on certain population vital rates (e.g. survival,
365 fecundity)?
- 366 25. Relative to other environmental processes, what is the role of winter as a selective filter on
367 populations?
- 368 26i. To what extent do winter conditions currently serve as environmental filters against the
369 persistence of species in mid/high latitude/elevation regions?
- 370 26ii. What are the most important filters in terrestrial vs. aquatic systems?
- 371 27. Are certain types of species (e.g. r vs. k selected species) more susceptible to winter as a
372 selective filter?
- 373 28. How does winter interact with density-dependent processes to influence population growth?
- 374 29. How do aggregations of wildlife in the winter interact with environmental conditions to
375 modulate density-dependent processes?
- 376 30i. How flexible and variable are responses to winter within species or populations?
- 377 30ii. How generalizable are certain winter 'strategies' across individuals, populations, or
378 species?
- 379 31. Does winter have similar effects on both aquatic and terrestrial population dynamics?
380

381 **Communities**

382 The mechanisms that regulate the composition, structure, and dynamics of biological
383 communities remain poorly understood. Community composition is broadly influenced by
384 ecological filtering, dispersal capabilities, and environmental tolerances of the component
385 species, while structure (relative abundance and distribution of taxa) and dynamics are shaped by
386 species' environmental preferences (i.e. optima) and by the nature and intensity of their

interactions with one another and their environment. As with other subdisciplines of ecology, most of our understanding of community processes has been derived from tropical systems or examinations of temperate areas during the warmer months (Hampton et al. 2015). At high latitudes, winter represents a relatively un-investigated period that is likely to yield novel insight into the manner by which species coexist under seasonally-fluctuating environmental conditions and community interaction regimes (Questions 32 - 33).

Outside of the tropics, seasonal fluctuations in solar irradiance are associated with often dramatic changes in basic habitat parameters including ambient temperature, photoperiod, and the availability of liquid water. At the base of the food web, this often leads to decreased primary productivity (e.g. Bramburger et al. 2020) and associated reductions in food resource availability in many ecosystems. Sub-zero temperatures, oxygen limitations, and resource scarcity impose physiological constraints upon many taxa and can be sufficiently harsh to induce mortality or dormancy in non-hardy taxa, driving migration and hibernation in others that are able change their behavior to avoid the most severe winter conditions. Further, snow and ice cover often constrains habitat connectivity and limits dispersal of propagules among suitable patches. Consequently, winter communities are effectively less diverse than their summer counterparts. The duration and severity of winter conditions are often potent structuring agents of regional biodiversity, and taxa that thrive in areas characterized by harsh winters are often specifically adapted to winter conditions and able to capitalize upon the hardship of others (Question 36).

The relative lack of biological activity during the winter presents various challenges and opportunities for overwintering species, resulting in shifting community dynamics. For taxa that are active during the frozen season, particularly consumers, the relative inaccessibility of food resources can cause intensification of interspecific competition and shift the food web towards

410 top down pressure on the few resources that remain (Humpries et al. 2017). In contrast, with
411 fewer organisms active under harsh conditions, those species that are tolerant may enjoy a
412 release from competitive interactions, predation, or grazing pressure (Question 36; McMeans et
413 al., 2020). Scarce food resources and changes in consumer abundance or pressure can also de-
414 couple trophic linkage characteristics of warmer seasons and could restructure winter food webs
415 (Questions 34 - 35). A poignant example of this decoupling in both aquatic and terrestrial food
416 webs is migration out of a system or metabolic arrest (e.g. hibernation, diapause), which
417 effectively cuts off species interactions (McMeans et al., 2015; Humpries et al., 2017). Re-
418 structuring can occur when consumers shift their diet towards different resources during winter.
419 For example, in aquatic and terrestrial food webs, allochthonous, detrital based pathways may
420 become more important (McMeans et al., 2015). Several reports also suggest that aquatic
421 consumers shift towards benthic or littoral prey during winter, which may be less susceptible to
422 winter declines in abundance than their pelagic, phytoplankton-supported counterparts (Question
423 35; Caldwell et al. 2020).

424 Presence of winter-active organisms may indirectly affect other members of the
425 community that are only active in warmer months, by modifying or consuming a shared resource
426 (Question 36). For example, a winter active predator may shift its diet to feed on a small
427 herbivore which is abundant during the winter but otherwise ignored when other prey species are
428 available. Decreases in the number of those herbivores as a result of winter predation could lead
429 to food shortages in spring for other predator species that hibernated through the winter. In
430 contrast, winter active organisms could facilitate spring or summer active organisms. For
431 example, winter active detritivores could break down leaf litter deposited in autumn, making
432 nutrients more readily available for spring-active detritivores lower down in the detrital

processing chain (Heard 1994). This is particularly relevant in lotic systems where terrestrial derived allochthonous material (predominantly leaf litter) may be decomposed and mineralised during winter resulting in the availability of nutrients to fuel primary production in the spring (Question 37) Furthermore, differential responses to environmental conditions throughout winter could also cause shifts in species interactions, further modulating the strength of species interactions (Question 38).

Questions:

32. Does the occurrence of winter stabilize or destabilize populations, food webs and community dynamics?

33. Are taxa being differentially influenced by changing environmental conditions throughout winter, and what physiological (e.g. thermal preference, energetic status) and ecological traits (e.g. functional group) explain these differences?

34. Are winter food webs supported by fewer resources, characterized by fewer trophic linkages and dampened top down regulation?

35. If primary production declines in the winter, do winter food webs shift towards reliance on detrital-based pathways?

36. How important is winter in providing a refuge from negative species interactions (e.g. by weakening competition)?

37. How does the seasonal input of terrestrial derived material in fall and spring influence allochthonous-autochthonous reliance in stream and river communities?

38. How do changing environmental conditions throughout winter influence the strength of trophic interactions (e.g. competition, predation)?

456 **Ecosystems**

457 Studies of ecosystems in the context of winter have focused primarily on how the movement,
458 storage, and flow of energy (i.e. the carbon cycle) and nutrients (primarily nitrogen) vary in
459 response to fluctuating winter conditions at a single site or among sites along climatic gradients.
460 While biological activity, including productivity, decomposition, and nutrient mineralization,
461 typically slows over winter, the contributions of processes in winter to annual cycles can be
462 substantial in regions where winter is relatively protracted (Question 39; Hobbie et al. 2002). The
463 transition from winter to spring (snow melt and soil thaw) can be a time of accelerated transport
464 and loss of nutrients (Hobbie and Chapin 1996), and soil freezing and other stressors associated
465 with winter can have a substantial effect on primary production over the subsequent growing
466 season (Kreyling 2010; Reinmann et al. 2019). Many ecosystem-scale experiments have
467 demonstrated the effects of current and projected changes in winter climate, through snow-
468 removal, snow-addition, and snowmelt manipulation experiments (Campbell et al. 2014; Henry
469 et al. 2018).

470 Despite recent progress in the study of how ecosystems function in winter and how these
471 processes are likely to be altered by climate change (described in Sanders-DeMott and Templer
472 2017), ecosystem studies can be limited by the spatial and temporal scales of experiments and
473 observations, and by an inability to capture the complexity of the interactions
474 involved. Specifically, ecosystem dynamics reflect an integration of biological processes at
475 multiple scales of organization (i.e. organismal, population, community, and ecosystem scales)
476 and are subject to the inherent shortcomings in quantifying these processes (Question 40).
477 Ecosystem-scale experiments are further complicated by the influence of abiotic components
478 (e.g. the geology, topography, climate, and hydrology of the system). When winter conditions

are manipulated experimentally at the plot level using shoveling, shelters, heaters, etc..., experimental artifacts often arise. For example, contrary to what would occur at the landscape level, plots subject to snow removal can remain artificially wet in spring because of runoff or groundwater recharge from the surrounding area. Likewise, regarding biotic factors, while plot level ecosystem experiments can address the responses of sessile organisms such as plants and soil microorganisms to variability in winter conditions, potentially important interactions with animals such as herbivores or detritivores can be underestimated (or when examined not completely understood) because these mobile organisms can move into and out of experimental plots (Question 41; Moise and Henry 2010; Templer et al. 2012). Regarding temporal scale, the manipulation of winter conditions in the short term can be useful for examining the direct effects of climate or weather on ecosystem processes, but important indirect effects that can occur via long term shifts in species composition (Komatsu et al. 2019) or legacy effects (Question 42) are often not captured.

A further characteristic of winter ecosystem experiments is that they have been conducted disproportionately at a select number of sites (Question 43). Gradient analyses across multiple sites have imparted greater generality to our understanding of how variability in winter conditions can affect ecosystem processes (Question 44), but they are nevertheless vulnerable to the influence of confounding factors. Combining both approaches by coordinating ecosystem experiments that manipulate winter conditions at many sites across natural climate gradients may be an ideal solution, but it requires greater logistical and financial support (Fraser et al. 2013).

Ecosystem-level responses to variability in winter conditions are particularly important to understand in the context of global climate change, given the key role of ecosystems in driving climate change feedback (Questions 45 - 46; e.g. changes in the net flux of carbon in response to

warming alter atmospheric greenhouse gas concentrations). Answering the following questions will help address the substantial knowledge gaps regarding the spatial and temporal scales of these responses, and the extent to which biotic complexity must be integrated into our understanding of these responses.

Questions:

39. How does winter influence energy and nutrient transfer through communities and ecosystems?

40. How effectively can we scale up ecosystem processes measured in experiments conducted in specific sites to the regional scale, utilizing our understanding of geology, topography, and hydrologic connectivity?

41. Given that animals may operate at much larger spatial scales than experimental plots, how can interactions between animals and primary producers be better addressed in ecosystem-level winter experiments?

42. What are the legacy effects of acute frost damage in soils or to tree canopies on terrestrial ecosystems over the longer term (e.g. years to decades), and what mechanisms underlie these legacy effects?

43. What novel insights can be gained by coordinating winter ecosystem experiments across many sites along climatic gradients; what should be manipulated and measured?

44. Do ecosystem-level processes respond more to average winter conditions or to extreme winter events?

45. How does winter constrain or influence important plant functional groups such as vegetation with nitrogen fixers, pests, or invasive species, that can have disproportionate influences on ecosystem processes?

46. What are the contributions of biological processes in winter to annual carbon and nutrient cycles and to processes in other seasons, including the growing season, as well as shoulder seasons such as spring and fall?

Management Concerns

Winter presents challenges to wild organisms but also does so for those tasked with managing and conserving wildlife and plant populations because of knowledge gaps in winter ecology and the difficulty in accessing field sites in the winter (Question 47). Winter climate regimes are shifting, increasing the need for adaptive natural resource management and consideration of winter processes (e.g., winter forest management; Rittenhouse and Rissman 2015). Indeed, failure to explicitly incorporate winter into management strategies can influence populations and management during other seasons (Questions 48 - 49). Predictive tools are therefore required to incorporate both exposure and species' sensitivities to winter conditions, in particular at higher latitudes where life persists for more than half the year in winter conditions.

The capacity of both terrestrial and aquatic animals to be buffered from winter conditions, escape predation during the winter, and emerge from winter with adequate energy reserves have strong management implications. For instance, quantifying and identifying seasonal variability as a key climate-change driver of biological responses provides insights with significance for population biology, the trajectories of species invasions and geographic range expansions (Rahel and Olden 2008; Pecl et al. 2017), and the outcomes of species interactions, such as disease dynamics (Question 50; Begon et al., 2009) and coexistence (Question 51; McMeans et al., 2020). Characterizing winter habitats of wildlife, especially for organisms that aggregate or have specific hibernation requirements, is essential for habitat management

including both protection (e.g., limiting development activities; McKinnon and Hnytka 1988) and restoration (e.g., increasing prevalence of habitat types that enable overwinter survival; Elphick and Oring 1998; Taft and Haig 2003). This includes the availability of microhabitats for many species to survive winter during hibernation or torpor, such as (e.g., animals that burrow within leaf litter and soil, or use rocky crevices as overwinter refugia). Moreover, understanding the vulnerability of important wildlife and ecosystems to different stressors such as human disturbance (e.g., wildlife responses to winter recreation; Cassirer et al. 1992) are also necessary to devise plans that reduce conflict between humans and wildlife (Question 52; Braunisch et al. 2011). Habitat used in the winter is rarely classified as “critical,” which suggests valuable winter habitat is more susceptible to loss or disturbance, potentially limiting population growth or recovery (Question 53; Cunjak 1996). Refining natural resource management policies and practices to consider events throughout the annual cycle has the potential to greatly improve management outcomes (Question 54; e.g., Zeug et al. 2012; Osnas et al. 2016), recognizing that carry-over effects are often manifested across seasons, especially in winter (e.g., Midwood et al. 2015).

Questions:

47. What type of training is needed to prepare natural resource practitioners to consider winter in management decisions?
48. How do land use decisions, management/fishing regulations, and habitat restoration influence wild organisms during winter?
49. To what extent do winter threats contribute to the imperilment of wildlife?
50. How does winter contribute to disease dynamics (transmission, pathogenicity)?

51. How will invasive and range shifting species alter community structure and function during winter?

52. Are “so-called” winter construction windows beneficial or detrimental to wildlife? How can they be altered to be more effective?

53. What specific types of policies and regulations are needed to protect critical winter habitats?

54. What are the management implications for rapid/unpredictable responses to changes in winter conditions/seasonal transitions (e.g. hunting/angling seasons and regulations, flow regulation in reservoirs + rivers during spawning runs and migrations, etc.)?

Climate Change

Climate-change is a dynamic process, involving changes to many environmental variables with distinct signatures in different seasons and many possible patterns of change. During summer months, shifts in air and ocean circulation cause increases in the frequency of extreme events, including heat waves and wildfires. However, climate change arguably has a greater impact in winter than summer, for example rates of warming in Northern Europe in winter outpace those seen in summer (Question 55; Elliott & Elliott 2010). The rapidity of such changes to high latitude winters presents a unique challenge to researchers studying winter processes, as scientific interest in this season increases, so too do the forces driving its attenuation. In addition, winter conditions across the globe are changing in potentially very different ways and it will be important to characterize how winter is changing and what factors may influence these changes across space and time (Questions 56 - 57). In both freshwater and marine systems, the ice-covered period has been steadily decreasing over the last century (Sharma et al. 2016). This has reciprocal effects for biotic communities as spring melt occurs earlier in the year causing

disequilibria with the phenology of natural (Shuter et al. 2012) and managed events (Horstkotte et al. 2017). In terrestrial systems, a feedback loop whereby a reduced albedo effect increases winter warming is speeding up snowpack loss in northern regions (Kim et al. 2018) with implications for snow associated biotic communities (Berteaux et al. 2016) and with potentially more devastating effects than aquatic ecosystems which may be buffered from climate change (Question 58). However, in some regions, instead of winter warming, harsher winters with longer cold spells are expected, including changes in lake and sea ice, and storm regimes (Question 59). This presents ecologists with the variation necessary to investigate the consequences of different rates of climate change on natural systems and the potential to investigate the long-term consequences of these winter climate change relative to climate change during other seasons (Question 60).

In some cases, strong biological responses to these environmental changes have emerged. For instance, species geographic ranges are extending polewards across multiple biomes as species are able to survive winter in regions that were once inhospitable (Parmesan 2006; Poloczanska et al. 2013; Pecl et al. 2015). Similar trends have been observed across a diversity of taxa, including plant and algal communities, marine and freshwater fishes, plankton, mammals, and insects (IPCC 2014). While this pattern of warming winters and changes in resident biotic communities is well documented, the mechanisms through which changing winter conditions drive biospheric changes are less well understood.

At the individual level, it is important to understand how climate change may directly impact strategies used to cope with harsh winter conditions (Question 61) or the potential benefits of these strategies (Question 62). Tracking these shifts in coping strategies and the benefits of these shifts could also illuminate broader patterns of rapid evolution (Question 63)

and the timescales over which changes in traits occur (Question 64). It could also shed light on the potential downstream consequences of changing environmental conditions on individuals (Question 65) or broader scales (Questions 66 - 68).

As individuals are directly impacted by winter climate change, it is likely that interactions between species within a community will experience rapid change. Future research should investigate how differential impacts of climate change on species may cause shifts in biotic interactions (Question 69). Additionally, it will also be important to characterize how changes in winter characteristics (e.g. length of winter, magnitude of temperature changes and precipitation) may simultaneously act to shape biotic interactions (Questions 70 - 71).

Finally, effort should also be made to refine predictive models, with an emphasis on winter precipitation (Question 72). This could allow for targeted research on areas experiencing the greatest changes and will allow researchers to better predict future climate change to forecast how species may continue to be shaped by changing abiotic conditions.

Questions:

55. How is winter changing and to what degree does this ongoing change vary geographically?

56. Are the effects of winter dependent on spatial scale (e.g. differ with size of the water body)?

57i. To what extent does vulnerability to ice loss vary spatially?

57ii. Are regions that are most susceptible to ice loss also experiencing the greatest loss?

58. Will terrestrial organisms be affected by climate change more quickly than aquatic/semi-aquatic organisms because aquatic environments are more thermally buffered (both to temperature extremes and variation in temperature) than terrestrial environments?

59. Can we define and predict the spatially-explicit impacts of changing jet streams on winters conditions?

- 639 60. How do legacy effects of winter climate change compare in magnitude to other aspects of
640 climate change such as summer warming, drought and elevated CO₂?
- 641 61. Are changing winters influencing behavioural/physiological strategies used to cope with
642 harsh environmental conditions?
- 643 62. Under what conditions are different winter coping strategies best suited, and which strategies
644 are likely to be most advantageous under climate change?
- 645 63. What species, or traits, will exhibit rapid adaptation to contemporary changes in winter and
646 spring phenology?
- 647 64. Over what timescales will organisms respond to changes in winter and spring phenology and
648 how are these timescales associated with generation time, intrinsic variation in
649 physiological/behavioural/ecological responses to environmental cues?
- 650 65. How will rapid misalignment between photoperiod and climatic cycles influence anticipatory
651 physiological responses to the initiation and conclusion of “winter” and what are the downstream
652 impacts of misaligned abiotic cues?
- 653 66. Which features of winter have the greatest influence on species and, consequently,
654 ecosystem responses to climate change?
- 655 67. How does the increased prevalence of freeze-thaw/rain on snow events through winter
656 impact terrestrial ecology at different scales (e.g., habitat quality/availability, mortality,
657 metabolism)?
- 658 68. How do changing winters impact light transmission and winter primary productivity in
659 aquatic ecosystems?
- 660 69. Given that species will be affected differently by changing winters, how will biotic
661 interactions and energy transfer through food webs shift under winter climate change?

70. How will changing winters influence the timing and magnitude of resource pulse dynamics in the coming years?

71. With the weakening of winter environmental filters as a result of climate change, what response do we expect from species formerly limited by them, and how will it impact local communities?

72. How can current climate change models be modified to better predict winter precipitation?

Conclusions

Although historically understudied, winter has recently been shown to be an important driver of ecological processes with consequences on all levels of biological organization from individuals to ecosystems. We have endeavored to highlight some key questions that will advance our understanding of the importance of winter and continue to fill in knowledge gaps associated with winter ecology. In light of ongoing climate change it is important to understand not only the importance of winter as a driver of ecological processes, but how its impacts may be changing over time. We hope that these questions stimulate future work on understanding the important, but understudied season of winter.

Acknowledgements

We would like to thank Brent Sinclair for providing suggestions on an early draft of the manuscript.

References

Altwegg, R., Roulin, A., Kestenholz, M., and Jenni, L. 2006. Demographic effects of extreme winter weather in the barn owl. *Oecologia* **149**(1): 44-51.

- Altwegg, R., Dummermuth, S., Anholt, B. R., and Flatt, T., 2005. Winter weather affects asp viper *Vipera aspis* population dynamics through susceptible juveniles. *Oikos* **110**(1): 55-66.
- Bargmann, T., Wheatcroft, E., Imperio, S., Vetaas, O. R. 2020. Effects of winter weather and hunting on wild reindeer population dynamics in Hardangervidda National park. *Pop Ecol* **62**(1): 91-104.
- Bartlett, J. C., Convey, P., Hayward, S. A. 2020. Surviving the Antarctic winter-life stage cold tolerance and ice entrapment survival in the invasive Chironomid Midge *Eretmoptera murphyi*. *Insects* **11**(3): 147-162.
- Bates, A.E., Helmuth, B., Burrows, M.T., Duncan, M.I., Garrabou, J., Guy-Haim, T., Lima, F., Queiros, A.M., Seabra, R., Marsh, R. and Belmaker, J., 2018. Biologists ignore ocean weather at their peril. *Nature* **560**: 299-301.
- Begon, M., Telfer, S., Smith, M. J., Burthe, S., Paterson, S., Lambin, X. 2009. Seasonal host dynamics drive the timing of recurrent epidemics in a wildlife population. *Proc R Soc B* **276**(1662): 1603-1610.
- Berge, J., Daase, M., Renaud, P. E., Ambrose, W. G., Darnis, G., Last, K. S., *et al.* 2015. Unexpected Levels of Biological Activity during the Polar Night Offer New Perspectives on a Warming Arctic. *Curr Biol* **25**(19): 2555–2561.
- Bergsten, U., Goulet, F., Lundmark, T., Löfvenius, M. O. 2011. Frost heaving in a boreal soil in relation to soil scarification and snow cover. *Can J Forest Res* **31**(6): 1084-1092.
- Berteaux, D., Gauthier, G., Domine, F., Ims, R. A., Lamoureux, S. F., Lévesque, E. *et al.* 2017. Effects of changing permafrost and snow conditions on tundra wildlife: critical places and times 1. *Arctic Science* **3**(2): 65–90.
- Blagojevic, D. P., Grubor-Lajsic, G. N., Spasic, M. B. 2011. Cold defence responses: the role of oxidative stress. *Front Biosci* **3**: 416-427.
- Blanchfield, P. J., Tate, L. S., Plumb, J. M., Acolas, M. L., Beaty, K. G. 2009. Seasonal habitat selection by lake trout (*Salvelinus namaycush*) in a small Canadian shield lake: constraints imposed by winter conditions. *Aquat Ecol* **43**(3): 777-787.
- Bokhorst, S.F., Bjerke, J.W., Tømmervik, H., Callaghan, T.V., Phoenix, G.K. 2009. Winter warming events damage sub-Arctic vegetation: Consistent evidence from an experimental manipulation and a natural event. *J Ecol* **97**(6): 1408-1415.
- Box, J. E., Colgan, W. T., Christensen, T. R., Schmidt, N. M., Lund, M., Parmentier, F.-J. *et al.* 2019. Key indicators of Arctic climate change. 1971–2017. *Environ Res Lett* **14**(4): 045010.

- Bramburger, A. J., Reavie, E. D., Sgro, G. V., Estepp, L. R., Chraibi, V. L. S., Pillsbury, R. W. 2020. Size matters: diatom establishment and extirpation timing in the Laurentian Great Lakes has been influenced by cell size. *J Plankton Res* **42**(3): 255-264.
- Braunisch, V., Patthey, P., Arlettaz, R. 2011. Spatially explicit modeling of conflict zones between wildlife and snow sports: prioritizing areas for winter refuges. *Ecol Appl* **21**(3): 955-967.
- Bruce, T. J. A., Matthes, M. C., Napier, J. A., Pickett, J. A. 2007. Stressful "memories" of plants: Evidence and possible mechanisms. *Plant Sci* **173**(6): 603-608.
- Casson, N. J., Contosta, A. R., Burakowski, E. A., Campbell, J. L., Crandall, M. S., Creed, I. F *et al.* 2019. Winter weather whiplash: Impacts of meteorological events misaligned with natural and human systems in seasonally snow-covered regions. *Earth's Future* **7**(12): 1434–1450.
- Caldwell, T. J., Chandra, S., Feher, K., Simmons, J. B., Hogan, Z. 2020. Ecosystem response to earlier ice break-up date: Climate-driven changes to water temperature, lake-habitat-specific production, and trout habitat and resource use. *Glob Change Biol* **26**(10): 5475-5491.
- Campbell, J. L., Soggi, A. M., Templer, P. H. 2014. Increased nitrogen leaching following soil freezing is due to decreased root uptake in a northern hardwood forest. *Glob Change Biol* **20**(8): 2663-2673.
- Cassirer, E. F., Freddy, D. J., Ables, E. D. 1992. Elk responses to disturbance by cross-country skiers in Yellowstone National Park. *Wild Soc Bull* **20**(4): 375-381.
- Cohen, J. 2016. An observational analysis: Tropical relative to Arctic influence on midlatitude weather in the era of Arctic amplification. *Geophys Res Lett* **43**(10): 5287–5294.
- Contosta, A. R., Casson, N. J., Nelson, S. J., Garlick, S. 2020. Defining frigid winter illustrates its loss across seasonally snow-covered area of eastern North America. *Environ Res Lett* **15**(3): 034020.
- Contosta, A. R., Casson, N. J., Garlick, S., Nelson, S. J., Ayres, M. P., Burakowski, E. A., *et al.* 2019. Northern forest winters have lost cold, snowy conditions that are important for ecosystems and human communities. *Ecol Appl* **29**(7): e01974.
- Convey, P. 2013. Overwintering strategies of terrestrial invertebrates in Antarctica-the significance of flexibility in extremely seasonal environments. *EJE* **93**(3): 489-505.
- Coulson, T., Catchpole, E. A., Albon, S. D., Morgan, J. T. B., Pemberton, J. M., Clutton-Brock, T. H. *et al.* 2002. Age, sex, density, winter weather and population crashes in Soay sheep. *Science* **292**(5521): 1528-1531.
- Cunjak, R. A. 1996. Winter habitat of selected stream fishes and potential impacts from land-use activity. *C J Fish Aquat Sci* **53**(S1): 267-282.

- 778
779 Darnis, G., Robert, D., Pomerleau, C., Link, H., Archambault, P., Nelson, R. J. 2012. Current
780 state and trends in Canadian Arctic marine ecosystems: II. Heterotrophic food web, pelagic-
781 benthic coupling, and biodiversity. *Climatic Change* **115**(1): 179–205.
782
- 783 Dickinson, R. E. 2015. Land-Atmosphere Interactions: Overview. In *Encyclopedia of*
784 *Atmospheric Sciences: Second Edition*, 240–243. Elsevier Inc.
785
- 786 Donaldson, M. R., Cooke, S. J., Patterson, D. A., Macdonald, J. S. 2008. Cold shock and fish. *J*
787 *Fish Biol* **73**(7): 1491-1530.
788
- 789 Elliott, J. M., Elliott, J.A. 2010. Temperature requirements of Atlantic salmon *Salmo salar*,
790 brown trout *Salmo trutta* and Arctic charr *Salvelinus alpinus*: predicting the effects of climate
791 change. *J Fish Biol* **77**(8): 1793-1817.
792
- 793 Elphick, C. S., Oring, L. W. 1998. Winter management of Californian rice fields for waterbirds. *J*
794 *App Ecol* **35**(1): 95-108.
795
- 796 Farmer, T.M., Marschall, E.A., Dabrowski, K., Ludsins, S.A. 2015. Short winters threaten
797 temperate fish populations. *Nat Comm* **6**(1): 1-10.
798
- 799 Fernandes, T., McMeans, B. C. 2019. Coping with the cold: energy storage strategies for
800 surviving winter in freshwater fish. *Ecography* **42**(12): 2037-2052.
801
- 802 Forrest, J.R. 2016. Complex responses of insect phenology to climate change. *Curr Opin Insect*
803 *Sci* **17**: 49-54.
804
- 805 Fraser, L. H., Henry, H. A. L., Carlyle, C. N., White, S. R., Beierkuhnlein, C., Cahill, J. F. *et al.*
806 2013. Coordinated distributed experiments: an emerging tool for testing global hypotheses in
807 ecology and environmental science. *Front Ecol Environ* **11**(3): 147-155.
808
- 809 Froy, H., Sparks, A. M., Watt, K., Sinclair, R., Bach, F., Pilkington, J. G., *et al.* 2019.
810 Senescence in immunity against helminth parasites predicts adult mortality in a wild mammal.
811 *Science* **365**(6459): 1296-1298.
812
- 813 Gamelon, M., Grøtan, V., Nilsson, A.L., Engen, S., Hurrell, J.W., Jerstad, K., *et al.* 2017.
814 Interactions between demography and environmental effects are important determinants of
815 population dynamics. *Sci Advances* **3**(2): e1602298.
816
- 817 Gilbert, C., McCafferty, D., Le Maho, Y., Martrette, J. M., Giroud, S., Blanc, S., *et al.* 2010. One
818 for all and all for one: The energetic benefits of huddling in endotherms. *Biol Rev* **85**(3): 545-
819 569.
820
- 821 Groffman, P. M., Driscoll, C. T., Fahey, T. J., Hardy, J. P., Fitzhugh, R. D., Tierney, G. L. 2001.
822 Colder soils in a warmer world: A snow manipulation study in a northern hardwood forest
823 ecosystem. *Biogeochemistry* **56**(2): 135-150.

- Gudleifsson, B. E. 1993. Metabolic and cellular impact of ice encasement on herbage plants. In: Jackson, M.B., Black, C.R. (ed.): *Interacting Stress on Plants in a Changing Climate*. 407-422. Springer-Verlag, Berlin.
- Hahn, D. A., Denlinger, D. L. 2011. Energetics of insect diapause. *Annu Rev Entomol* **56**: 103-121.
- Hampton, S. E., Galloway, A. W., Powers, S. M., Ozersky, T., Woo, K. H., Batt, R. D., *et al.* 2017. Ecology under lake ice. *Ecol Lett* **20**(1): 98-111.
- Hampton, S. E., Moore, M. V., Ozersky, T., Stanley, E. H., Polashenski, C. M., Galloway, A. W. 2015. Heating up a cold subject: prospects for under-ice plankton research in lakes. *J Plankton Res* **37**(2): 277-284.
- Heard, S. B. 1994. Processing chain ecology: resource condition and interspecific interactions. *J Anim Ecol*: 451-464.
- Henry, H. A., Abedi, M., Alados, C. L., Beard, K. H., Fraser, L. H., Jentsch, A., *et al.* 2018. Increased soil frost versus summer drought as drivers of plant biomass responses to reduced precipitation: results from a globally coordinated field experiment. *Ecosystems* **21**(7): 1432-1444.
- Hoback, W. W., Stanley, D. W. 2001. Insects in hypoxia. *J Insect Phys* **47**(6): 533-542.
- Hobbie, S. E., Schimel, J. P., Trumbore, S. E., Randerson, J. R. 2002. Controls over carbon storage and turnover in high-latitude soils. *Glob Change Biol* **6**(S1): 196-210.
- Hobbie, S. E., Chapin, F. S. 1996. Winter regulation of tundra litter carbon and nitrogen dynamics. *Biogeochemistry* **35**(2): 327-338.
- Horstkotte, T., Utsi, T. A., Larsson-Blind, Å., Burgess, P., Johansen, B., Käyhkö, J., *et al.* 2017. Human-animal agency in reindeer management: Sámi herders' perspectives on vegetation dynamics under climate change. *Ecosphere* **8**(9): e01931.
- Humphries, M. M., Studd, E. K., Menzies, A. K., Boutin, S. 2017. To everything there is a season: Summer-to-winter food webs and the functional traits of keystone species. *Integr Comp Biol* **57**(5): 961-976.
- Humphries, M. M., Umbanhowar, J. 2007. Filtering environmental variability: activity optimization, thermal refuges, and the energetic responses of endotherms to temperature. In *The impact of environmental variability on ecological systems*, 61-87. Springer, Dordrecht.
- Huschke, R. E. 1959. *Glossary of Meteorology*. American Meteorology Society, Boston, Massachusetts, USA.

- 870 IPCC. 2013. IPCC Climate Change 2013: The Physical Science Basis. (T. F. Stocker, Ed.).
871 Cambridge University Press.
872
- 873 IPCC. 2014. Climate Change 2014: impacts, adaptation and vulnerability. Cambridge University
874 Press.
875
- 876 Kielland, K., Olson, K., Ruess, R. W., Boone, R. D. 2006. Contribution of winter processes to
877 soil nitrogen flux in taiga forest ecosystems. *Biogeochemistry* **81**(3): 349-360.
878
- 879 Kim, S. Y., Noguera, J. C., Velando, A. 2019. Carry-over effects of early thermal conditions on
880 somatic and germline oxidative damages are mediated by compensatory growth in sticklebacks. *J*
881 *Anim Ecol* **88**(3): 473-483.
882
- 883 Komatsu, K. J., Avolio, M. L., Lemoine, N. P., Isbell, F., Grman, E., Houseman, G. R. *et al.*
884 2019. Global change effects on plant communities are magnified by time and the number of
885 global change factors imposed. *P Natl Acad Sci USA* **116**(36): 17867-17873.
886
- 887 Kraus, E. B., Morrison, R. E. 1966. Local interactions between the sea and the air at monthly and
888 annual time scales. *Q J Roy Meteor Soc* **92**(391): 114–127.
889
- 890 Kreyling, J. 2010. Winter climate change: a critical factor for temperate vegetation performance.
891 *Ecology* **91**(7): 1939–1948.
892
- 893 Ladwig, L. M., Ratajczak, Z. R., Ocheltree, T. W., Hafich, K. A., Churchill, A. C., Frey, S. J., *et*
894 *al.* 2016. Beyond arctic and alpine: the influence of winter climate on temperate ecosystems.
895 *Ecology* **97**(2): 372-382.
896
- 897 Lalouette, L., Williams, C. M., Hervant, F., Sinclair, B. J., Renault, D. 2011. Metabolic rate and
898 oxidative stress in insects exposed to low temperature thermal fluctuations. *Comp Biochem Phys*
899 *A* **158**(2): 229-234.
900
- 901 Leirs, H., Stenseth, N. C., Nichols, J. D., Hines, J. E., Verhagen, R., Verheyen, W. 1997.
902 Stochastic seasonality and nonlinear density-dependent factors regulate population size in an
903 African rodent. *Nature* **389**(6647): 176-180.
904
- 905 Lukatkin, A., Brazaityte, A., Bobinas, C. and Duchovskis, P. 2012. Chilling injury in chilling-
906 sensitive plants: a review. *Agriculture* **99**(2): 111-124.
907
- 908 Malhi, Y., Wright, J. 2004. Spatial patterns and recent trends in the climate of tropical rainforest
909 regions. *Phil Trans R Soc B: Biol Sci*, **359**(1443): 311-329.
910
- 911 Marchand, P. J. 2013. *Life in the Cold: An Introduction to Winter Ecology* (4th ed.). Lebanon,
912 NH: University Press of New England.
913
- 914 Marshall, K. E., Sinclair, B. J. 2012. Threshold temperatures mediate the impact of reduced snow
915 cover on overwintering freeze-tolerant caterpillars. *Naturwissenschaften* **99**(1): 33-41.

- Masoero, G., Laaksonen, T., Morosinotto, C., Korpimäki, E. 2020. Climate change and perishable food hoards of an avian predator: Is the freezer still working? *Glob Change Biol* **26**(10): 5414-5430.
- Mayes Boustead, B. E., Hilberg, S. D., Shulski, M. D., Hubbard, K. G. 2015. The accumulated winter season severity index (AWSSI). *J Appl Meteor Clim* **54**(8): 1693-1712.
- McKinnon, G. A., Hnytka, F. N. 1988. The effect of winter pipeline construction on the fishes and fish habitat of Hodgson Creek, NWT, No. 1598. Fisheries and Oceans, Canada.
- McMeans, B. C., McCann, K. S., Guzzo, M. M., Bartley, T. J., Bieg, C., Blanchfield, P. J., *et al.* 2020. Winter in water: Differential responses and the maintenance of biodiversity. *Ecol Lett* **23**(6): 922-938.
- McMeans, B. C., McCann, K. S., Humphries, M., Rooney, N., Fisk, A. T. 2015. Food web structure in temporally-forced ecosystems. *Trends Ecol Evol* **30**(11): 662-672.
- Midwood, J. D., Larsen, M. H., Boel, M., Aarestrup, K., Cooke, S. J. 2015. An experimental field evaluation of winter carryover effects in semi-anadromous brown trout (*Salmo trutta*). *J Exp Zool Part A* **323**(9): 645-654.
- Moise, E. R. D., Henry, H. A. L. 2010. Like moths to a street lamp: exaggerated animal densities in plot-level global change field experiments. *Oikos* **119**(5): 791-795.
- Negro, M., Isaia, M., Palestini, C., Schoenhofer, A., Rolando, A. 2010. The impact of high-altitude ski pistes on ground-dwelling arthropods in the Alps. *Biodivers Conserv* **19**(7): 1853-1870.
- Osnas, E. E., Zhao, Q., Runge, M. C., Boomer, G. S. 2016. Cross-seasonal effects on waterfowl productivity: Implications under climate change. *J Wild Manage* **80**(7): 1227-1241.
- Parker, K. L., Robbins, C. T., Hanley, T. A. 1984. Energy expenditures for locomotion by mule deer and elk. *J Wild Manage*: 474-488.
- Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Syst* **37**: 637-669.
- Pauli, J. N., Zuckerberg, B., Whiteman, J. P., Porter, W. 2013. The subnivium: a deteriorating seasonal refugium. *Front Ecol Environ* **11**(5): 260-267.
- Pecl, G. T., Araujo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I-C. 2017. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**: 6332.

- Pecl, G.T., Frusher, S., Hobday, A.J., Jennings, S.M., Ogier, E.M., Sullivan, A., *et al.* 2015. Developing adaptation pathways for climate-impacted and at risk fisheries in south-east Australia. In 3rd ICES-PICES-IOC International Symposium on the Effect of Climate Change on the World's Oceans, 118.
- Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J., *et al.* 2013. Global imprint of climate change on marine life. *Nat Clim Change* **3**(10): 919-925.
- Rahel, F. J., Olden, J. D. 2008. Assessing the effects of climate change on aquatic invasive species. *Conserv Biol* **22**(3): 521-533.
- Reinmann, A.B., Susser, J.R., Demaria, E.M., Templer, P.H. 2019. Declines in northern forest tree growth following snowpack decline and soil freezing. *Glob Change Biol* **25**(2): 420-430.
- Richard, J. H., Wilmshurst, J., Côté, S. D. 2014. The effect of snow on space use of an alpine ungulate: recently fallen snow tells more than cumulative snow depth. *Can J Zoo* **92**(12): 1067-1074.
- Ringsby, T. H., Sæther, B. E., Altwegg, R., Solberg, E. J. 1999. Temporal and spatial variation in survival rates of a house sparrow, *Passer domesticus*, metapopulation. *Oikos*: 419-425.
- Rittenhouse, C. D., Rissman, A. R. 2015. Changes in winter conditions impact forest management in north temperate forests. *J Environ Manage* **149**: 157-167.
- Romano, A. B., Hunt, A., Welbergen, J. A., Turbill, C. 2019. Nocturnal torpor by superb fairy-wrens: a key mechanism for reducing winter daily energy expenditure. *Biol Lett* **15**(6): 20190211.
- Salt, R. 1961. Principles of insect cold-hardiness. *Annu Rev Ento* **6**(1): 55-74.
- Sanders-DeMott, R., Templer, P. H. 2017. What about winter? Integrating the missing season into climate change experiments in seasonally snow covered ecosystems. *Methods Ecol Evol* **8**(10): 1183–1191.
- Sæther, B. E., Bakke, Ø. 2000. Avian life history variation and contribution of demographic traits to the population growth rate. *Ecology* **81**(3): 642-653.
- Serreze, M. C. 2015. Arctic and Antarctic: Arctic Climate. In *Encyclopedia of Atmospheric Sciences: Second Edition*, 107–115. Elsevier Inc.
- Shuter, B. J., Finstad, A. G., Helland, I. P., Zweimüller, I., Hölker, F. 2012. The role of winter phenology in shaping the ecology of freshwater fish and their sensitivities to climate change. *Aquat Sci* **74**(4): 637-657.

- Sharma, S., Blagrove, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D. 2019. Widespread loss of lake ice around the Northern Hemisphere in a warming world. *Nat Clim Change* **9**(3): 227–231.
- Sharma, S., Magnuson, J. J., Batt, R. D., Winslow, L. A., Korhonen, J., Aono, Y. 2016. Direct observations of ice seasonality reveal changes in climate over the past 320–570 years. *Sci Reports* **6**: 25061.
- Sinclair, B. J. 1999. Insect cold tolerance: how many kinds of frozen? *Eur J Entomol* **96**: 157–164.
- Sinclair, B. J. 2015. Linking energetics and overwintering in temperate insects. *J Therm Biol* **54**: 5–11.
- Stålhandske, S., Lehmann, P., Pruisscher, P., Leimar, O. 2015. Effect of winter cold duration on spring phenology of the orange tip butterfly, *Anthocharis cardamines*. *Ecol Evol* **5**(23): 5509–5520.
- Stier, A., Massemin, S., Criscuolo, F. 2014. Chronic mitochondrial uncoupling treatment prevents acute cold-induced oxidative stress in birds. *J Comp Phys B* **184**(8): 1021–1029.
- Studd, E. K., Bates, A. E., Bramburger, A. J., Fernandes, T., Hayden, B., Henry, H. A. L., Humphries, M. M., Martin, R., McMeans, B. C., Moise, E. R. D., O'Sullivan, A. M., Sharma, S., Sinclair, B. J., Sutton, A. O., Templer, P. H., Cooke, S. J. 2021. Nine maxims for the ecology of cold-climate winters. *BioScience*: biab032.
- Sung, S., Amasino, R. M. 2004. Vernalization and epigenetics: how plants remember winter. *Curr Opin Plant Biol* **7**(1): 4–10.
- Sutton, A. O., Strickland, D., Norris, D. R. 2016. Food storage in a changing world: implications of climate change for food-caching species. *Clim Change Res* **3**: 12.
- Sutton, A. O., Strickland, D., Freeman, N. E., Norris, D. R. 2021. Climate-driven carry-over effects negatively influence population growth rate in a food-caching boreal passerine. *Glob Change Biol* **27**(5): 983–992.
- Swain, D. L., Langenbrunner, B., Neelin, J. D., Hall, A. 2018. Increasing precipitation volatility in twenty-first-century California. *Nat Clim Change* **8**(5): 427–433.
- Templer, P. H., Mack, M. C., III, F. C., Christenson, L. M., Compton, J. E., Crook, H. D., *et al.* 2012. Sinks for nitrogen inputs in terrestrial ecosystems: a meta-analysis of ¹⁵N tracer field studies. *Ecology* **93**(8): 1816–1829.
- Taft, O. W., Haig, S. M. 2003. Historical wetlands in Oregon's Willamette Valley: implications for restoration of winter waterbird habitat. *Wetlands* **23**(1): 51–64.

- 1051 Tremblay, J.-É., Robert, D., Varela, D. E., Lovejoy, C., Darnis, G., Nelson, R. J. 2012. Current
1052 state and trends in Canadian Arctic marine ecosystems: I. Primary production. *Climatic Change*
1053 **115**(1): 161–178.
- 1054
1055 Turner, J. 2015. Arctic and Antarctic: Antarctic Climate. In *Encyclopedia of Atmospheric*
1056 *Sciences: Second Edition*, 98–106. Elsevier Inc.
- 1057
1058 Uelmen Jr, J. A., Lindroth, R. L., Tobin, P. C., Reich, P. B., Schwartzberg, E. G., Raffa, K. F.
1059 2016. Effects of winter temperatures, spring degree-day accumulation, and insect population
1060 source on phenological synchrony between forest tent caterpillar and host trees. *Forest Ecol*
1061 *Manage* **362**: 241-250.
- 1062
1063 Veteli, T. O., Lahtinen, A., Repo, T., Niemelä, P., Varama, M. 2005. Geographic variation in
1064 winter freezing susceptibility in the eggs of the European pine sawfly (*Neodiprion sertifer*). *Agri*
1065 *Forest Ento* **7**(2): 115-120.
- 1066
1067 Waite, T.A., Strickland, D. 2006. Climate change and the demographic demise of a hoarding bird
1068 living on the edge. *Proc R Soc B* **273**(1603): 2809-2813.
- 1069
1070 Williams, C. M., Hellmann, J. J., Sinclair, B. J. 2012a. Lepidopteran species differ in
1071 susceptibility to winter warming. *Climate Res* **53**(2): 119-130.
- 1072
1073 Williams, C. M., Marshall, K. E., MacMillan, H. A., Dzurisin, J. D .K., Hellmann, J .J., Sinclair,
1074 B. J. 2012b. Thermal variability increases the impact of autumnal warming and drives depression
1075 in an overwintering butterfly. *PLoS ONE* **7**(3): e34470.
- 1076
1077 Woodworth, B. K., Wheelwright, N. T., Newman, A. E., Schaub, M., Norris, D. R. 2017. Winter
1078 temperatures limit population growth rate of a migratory songbird. *Nat Comm* **8**(1): 1-9.
- 1079
1080 Yang, B., Young, J., Brown, L., Wells, M. 2017. High-frequency observations of temperature
1081 and dissolved oxygen reveal under-ice convection in a large lake. *Geophys Res Lett* **44**(24): 12-
1082 18.
- 1083
1084 Zeug, S. C., Bergman, P. S., Cavallo, B. J., Jones, K. S. 2012. Application of a life cycle
1085 simulation model to evaluate impacts of water management and conservation actions on an
1086 endangered population of Chinook salmon. *Environ Monit Assess* **17**(5): 455-467.
- 1087 Zhang, X., Vincent, L.A., Hogg, W.D., Niitsoo, A. 2000. Temperature and precipitation trends in
1088 Canada during the 20th century. *Atmosphere-ocean*, **38**(3): 395-429.
- 1089

Figure 1 Conceptualization of how different definitions outlining boundaries for winter imposed divergent interpretations of winter across years. In instances where winter is defined by specific focal conditions, the period defined as ‘winter’ is mobile and adjusts to changes in these conditions across years. Thus, the length of winter is the predominant characteristic that varies. However, when winter is defined by a fixed, photoperiod window, conditions within that window are likely to change. In this way, winter processes can be more easily compared longitudinally and geographically. Light blue lines represent seasonal snow depth, with the light blue box representing winter as defined by minimum snow depth. Black lines represent seasonal water temperature, with the black box representing winter as defined by temperature. The yellow box represents winter defined by photoperiod.

Figure 2 Winter can vary across space and between systems (e.g. panels A – C). Additionally, winter can vary over time (panels D and E) with potentially high inter-annual variability in environmental conditions. Researchers interested in understanding the consequences of this variability can experimentally manipulate winter conditions (panel F) and monitor the effects on individuals, populations, communities, and ecosystems. As climate change continues to cause winter to change, it is imperative to understand how winter influences a variety of ecological processes at different biological scales. Photo credit: A) Antóin M. O’Sullivan, B) Brian Hayden, C) Ocean Networks Canada, D) and E) Alex O Sutton, F) Pamela H. Templer.

Figure 3 Conceptual diagram outlining direct and indirect effects of winter on various levels of biological organization with unanswered questions highlighting where key uncertainties remain in winter biology. This diagram demonstrates that uncertainties exist across all levels of

1113 biological organization and is a non-exhaustive list of the possible interactions within or between
1114 levels of biological organization. Furthermore, it displays the range of impacts, both direct and
1115 indirect, that winter can have. Climate change also has the potential to influence all proposed
1116 interactions and is likely to influence the magnitude of the effects of winter on all levels of
1117 biological organization. Questions that appear throughout the text have been mapped onto this
1118 diagram to identify what effects of winter these questions address. Black arrows represent direct
1119 effects, gray arrows represent indirect effects and the arrowhead represent the direction of the
1120 proposed effect. Single headed arrows indicate a unidirectional relationship, while two-headed
1121 arrows represent bi-directional relationships.

1122

Draft

1123 Figure 1

1124

1125 **Figure 2**

1126

Figure 3