Canadian Journal of Physics

Qualitative aspects of Rastall gravity with barotropic fluid

Journal:	<i>Canadian Journal of Physics</i>
Manuscript ID	cjp-2020-0061.R2
Manuscript Type:	Article
Date Submitted by the Author:	17-May-2021
Complete List of Authors:	Singh, Dr. Ashutosh ; Jabalpur Engineering College, Department of Applied Mathematics Raushan, Rakesh; Banaras Hindu University Faculty of Science, DST-CIMS Chaubey, R.; Banaras Hindu University, DST - CIMS
Keyword:	Rastall gravity, equation of state, dynamical system, cyclic universe, thermodynamics
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Qualitative aspects of Rastall gravity with barotropic fluid

Ashutosh Singh^{1*}, Rakesh Raushan^{2†}, R. Chaubey^{2‡}

¹Department of Applied Mathematics,
Jabalpur Engineering College, Jabalpur, India

²CIMS, Institute of Science,
Banaras Hindu University, Varanasi, India

Abstract

We investigate the dynamical evolution of homogeneous and isotropic flat universe filled with a barotropic fluid satisfying linear equation of state in Rastall gravity. Using dynamical system approach, we find the fixed points of system and study their stability. We further explore the thermodynamic aspects at apparent horizon by investigating the validity of generalized second law of thermodynamics with equilibrium description.

Keywords: Rastall gravity, dynamical system, cyclic universe, thermodynamics

PACS: 98.80-Jk, 98.80-k, 95.30.Tg

1 Introduction

Cosmological modelling requires a theory of gravity and description of matter content in the universe. In Rastall's theory of gravity [1–3], the matter source is described by energy-momentum tensor and metric of external space. The usual conservation law ($T_{;j}^{ij} = 0$) of Einstein's gravity is modified in this theory and, covariant divergence of energy-momentum tensor is related to covariant divergence of Ricci scalar. In empty space, this theory coincides with Einstein's gravity [1]. Rastall's gravity lacks Lagrangian formulation at least in context of Riemannian geometry, however in a phenomenological way, it produces the violation of classical conservation laws in an explicit covariant form (a distinguishing property which is termed as gravitational anomaly) [4]. Rastall's gravity may be considered as a phenomenological way to implement some concepts that are typical of quantum fields in curved space-time. The particle creation mechanism in cosmology also leads to violation of classical conservation laws and in this sense, Rastall's gravity may be considered as a classical formulation of that quantum phenomena, as covariant divergence of energy momentum tensor is directly related with covariant divergence of Ricci scalar [5]. For resolving the issue of Lagrangian formulation of Rastall gravity, Moraes and Santos have done investigation with gravity theories where geometry is non-minimally coupled with matter fields [6]. Shabani and Ziaie have derived Lagrangian formulation for Rastall theory by assuming perfect

*theprabhu.09@gmail.com

†rakesh.raushan4@bhu.ac.in

‡rchaubey@bhu.ac.in

fluid matter content with barotropic equation of state and claimed that Rastall gravity can be identified from $f(R, T)$ gravity Lagrangian [7]. Rastall's gravity has extensively been studied in different cosmological contexts. [8–27]. A qualitative study of Rastall gravity by using phase plane analysis may give insight on class of solutions available in this theory. We aim to study Rastall gravity by using dynamical system analysis for exploring possibilities of non-singular solutions. The stability analysis of Rastall model with quadratic equation of state have been investigated by Silva et al. [11]. Dynamical system analysis of cosmological models has been an active and interesting area of theoretical research [28–40].

The connection between gravity and thermodynamics has been an interesting topic since development of black hole thermodynamics [41–43]. Jacobson [44] was first to demonstrate the derivation of Einstein's field equations from Clausius relation at cosmological horizon collectively with equivalence principle. Eling et al [45] demonstrated that by using terms other than Einstein-Hilbert action, one may produce entropy due to the non-equilibrium thermodynamical aspects of generalized $f(R)$ theory [46]. By using field equations and applying thermodynamical laws to cosmological horizon, one can find rate of total horizon entropy in different gravitational setups [12, 15, 19, 23, 47–57]. The behaviour of expanding universe have also been studied extensively from the thermodynamic point of view. The generalized second law of thermodynamics (GSLT) states that the entropy of horizon plus entropy of matter inside horizon is an increasing function of time [42]. Moradpour [12] had investigated GSLT in flat-FRW Rastall gravity framework. A modified expansion law for the emergent cosmic space in Rastall gravity have been used to recover Friedmann-like equations [15]. The exploration of thermodynamic properties at apparent horizon in Rastall gravity with entropy correction have been done by Bamba et al [19]. Das et al [20] have discussed cosmological consequences and universal thermodynamics in Rastall gravity by illustrating a complete cosmological history starting from inflation to late-time acceleration. Cruz et al [23] have presented a thermodynamic discussion of Rastall gravity by introducing chemical potential.

Our aim in this paper is to study the role of barotropic fluid satisfying the linear equation of state in Rastall's gravity framework by employing dynamical system analysis. We also discuss the thermodynamics of homogeneous and isotropic universe in resulting gravitational setup. Since entropy of a physical system should be a positive quantity [58], we analyze the conditions on Rastall and other parameters by applying this condition to entropy of cosmological horizon.

The paper is organized as follows: in section 2, we write the cosmological equations and study the dynamics of isotropic models by finding the critical points and their behaviour. In section 3, we discuss validity of generalized second law of thermodynamics in resulting cosmology with equilibrium description. In section 4, we give our conclusion.

2 Dynamics of the model

In this section, we briefly introduce the basic cosmological equations and study phase plane in Rastall's gravity framework.

The line element of FRW universe can be written as

$$ds^2 = dt^2 - a^2(t) \left(\frac{dr^2}{1 - \kappa r^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right) \quad (1)$$

where $a(t)$ denotes scale factor and $\kappa = 0, +1, -1$ for flat, closed and open spatial sections respectively. In Rastall theory of gravity, the energy-momentum tensor is not conserved, and therefore $T_{;i}^i = \lambda R$, where R is Ricci scalar and λ is Rastall parameter [1]. Rastall parameter

may be interpreted as a measure of tendency of matter field to couple with geometry or vice-versa. The equivalent versions of gravitational field equations for metric tensor g_{ij} can be written as [1, 59]

$$\begin{aligned} R_{ij} - \frac{1}{2}(1 - 2k\lambda)Rg_{ij} &= kT_{ij} \\ G_{ij} &= k \left(T_{ij} - \frac{k\lambda}{4k\lambda - 1} Tg_{ij} \right) \\ R_{ij} &= k \left(T_{ij} - \frac{1}{2} \frac{2k\lambda - 1}{4k\lambda - 1} Tg_{ij} \right) \end{aligned} \quad (2)$$

where $R_{ij}, G_{ij}, R, g_{ij}, T_{ij}$ are Ricci tensor, Einstein's tensor, Ricci scalar, metric tensor, energy-momentum tensor of fluid respectively. For $\lambda = 0$ and $k = 8\pi$ in units $G = c = 1$, the Einstein field equations of General relativity can be re-obtained whenever $T_{ij}^j = 0$ [1]. We may also take $\frac{6k\lambda-1}{4k\lambda-1} = \frac{8\pi}{k}$ as considered in references [19, 47]. Note that k is the gravitational coupling and may be considered as free parameter for modelling requirements with care to the satisfaction of equations. However, $k\lambda = \frac{1}{4}, \frac{1}{6}$ should be avoided in the Rastall gravity [19, 47]. For a perfect fluid source $T_i^j = \text{diag}(\rho, -p, -p, -p)$, the corresponding field equations in Rastall gravity [15] are as follows:

$$3(1 - 4k\lambda)H^2 - 6k\lambda\dot{H} + 3(1 - 2k\lambda)\frac{\kappa}{a^2} = k\rho \quad (3)$$

$$3(1 - 4k\lambda)H^2 + 2(1 - 3k\lambda)\dot{H} + (1 - 6k\lambda)\frac{\kappa}{a^2} = -kp \quad (4)$$

where $H = \frac{\dot{a}}{a}$ is the Hubble parameter and ρ, p are energy density and pressure of energy-momentum source. In this paper, we take $\kappa = 0$. The Bianchi identity ($G_{ij}^j = 0$) leads to the equation of continuity [59] as follows:

$$(3k\lambda - 1)\dot{\rho} + 3k\lambda\dot{p} + (4k\lambda - 1)3H(\rho + p) = 0 \quad (5)$$

A functional dependence of one variable on the other may be postulated from motivation on the physical grounds. In this paper, we analyse the perfect fluid model with a general linear equation of state $p = \alpha\rho - \rho_0$ where α and ρ_0 are free parameters. This equation may be taken as generalization of a homogeneous linear equation of state $p = \alpha\rho$. The advantage of using the general linear equation of state is in the possibility of describing the dark energy with positive squared sound speed (c_s^2) because with equation of state $p = \alpha\rho$, the dark energy is hydrodynamically unstable, (since $c_s^2 = \frac{\partial p}{\partial \rho} = \alpha < 0$) [60]. This non-homogeneous linear equation of state provides the description of both hydrodynamically stable ($\alpha > 0$) and unstable ($\alpha < 0$) fluids. If α and ρ are small enough as compared to ρ_0 in any scenario, then we may get negative pressure in the model.

In this paper, we take perfect fluid with following barotropic equation of state which have also been investigated in different cosmological set-ups as [60–63]

$$p = \alpha\rho - \rho_0 \quad (6)$$

where α and ρ_0 are constants. 'Barotropic fluid' means some matter or field with equation of state $p = f(\rho)$. Barotropic fluids have long been studied in the context of dark energy [28, 60–66]. For flat-FRW metric, from equations (3) and (4), we have

$$k(\rho - p) = 6(1 - 4k\lambda)H^2 + 2(1 - 6k\lambda)\dot{H} \quad (7)$$

$$k(\rho + p) = -2\dot{H} \quad (8)$$

$$k(\rho + 3p) = 6(4k\lambda - 1)H^2 + 6(2k\lambda - 1)\dot{H} \quad (9)$$

We are using criterion for classical stability from equation $p = \alpha\rho - \rho_0$. The equation of state parameter γ is defined as $\gamma = \frac{p}{\rho}$ and $\alpha = \frac{\partial p}{\partial \rho}$ may be interpreted as sound speed squared (c_s^2). For classical stability of the model, we should have $0 \leq \alpha \leq 1$. From equation (6 and 8), we have $\rho + p = (1 + \alpha)\rho - \rho_0$. Null energy condition (NEC) is valid when $\rho \geq \frac{\rho_0}{1+\alpha}$. In the phantom region, the fluid violates NEC. It means that the energy density is increasing (decreasing) in the future for an expanding (contracting) universe. The trace of energy-momentum tensor T is given by $\rho - 3p$ as

$$k(\rho - 3p) = -2(2H^2 + \dot{H})(12k\lambda - 3) \quad (10)$$

From equations (3,5,6,8), we have

$$H^2 = \frac{k[3k\lambda(1 + \alpha) - 1]}{3(4k\lambda - 1)}\rho - \frac{k^2\lambda\rho_0}{4k\lambda - 1} \quad (11)$$

$$\rho = \frac{1}{1 + \alpha} \left(\frac{a_0}{a^3} \right)^{\frac{1}{K}} + \frac{\rho_0}{1 + \alpha} \quad (12)$$

where $K = \frac{3k\lambda(1+\alpha)-1}{(4k\lambda-1)(1+\alpha)}$ for $\alpha \neq -1$. Note that K depends on α, k, λ , therefore dynamics of model and its fate also depend on the values of α, k, λ . If we consider the case of $0 \leq \alpha \leq 1$, then for $\frac{1}{K} > 0$, we should have either $k\lambda > \max\left(\frac{1}{4}, \frac{1}{3(1+\alpha)}\right)$ or $k\lambda < \min\left(\frac{1}{4}, \frac{1}{3(1+\alpha)}\right)$. With $\frac{1}{K} > 0$, the expanding cosmological model in Rastall gravity will recover de Sitter expansion scenario (since with $a \rightarrow \infty, \rho \rightarrow \text{constant}$ and therefore $p \rightarrow \text{constant}$) and with $\frac{1}{K} < 0$, the universe may end-up with rip singularity (since with $a \rightarrow \infty, \rho \rightarrow \infty$ and therefore $p \rightarrow \infty$). For $\alpha = -1$, we have

$$\rho = \log \left(\frac{a_0}{a^{3(4k\lambda-1)\rho_0}} \right) \quad (13)$$

where a_0 is integration constant. We will concentrate on the broader class of models where $\alpha \neq -1$. From equations (3) and (5) with $3k\lambda(1 + \alpha) \neq 1$, we have

$$\dot{H} = -\frac{1}{6\lambda}\rho - \frac{(4k\lambda - 1)}{2k\lambda}H^2 \quad (14)$$

$$\dot{\rho} = \frac{1 - 4k\lambda}{[3k\lambda(1 + \alpha) - 1]}3H[(1 + \alpha)\rho - \rho_0] \quad (15)$$

The stability nature of the fixed points can be found by carrying out a linear stability analysis of a dynamical system. In brief (see [29–31] for details), this involves analysing the behaviour of linear perturbations around the fixed points, which obey the equations $y'_i = My_j$. The matrix M is the Jacobian matrix of the dynamical system. The eigenvalues λ_i of the Jacobian matrix evaluated at fixed points tell us about the linear stability character of fixed points. If real part of eigenvalues is non-zero, the fixed point is said to be hyperbolic point. The point is a repeller if all the real parts of eigenvalues are positive. Any small deviations from this point will cause the system to move away from this state. If all real parts of eigenvalues are negative, point is said to be an attractor. In this case, if system is perturbed away from this state then, it will return to the equilibrium state. If some of the eigenvalues are positive, while others are negative, then point is a saddle point [64]. Furthermore, phase planes are useful in visualizing the behaviour of oscillatory systems where the trajectories can ‘spiral in’ towards zero, ‘spiral out’ towards infinity, or reach neutrally stable situations called centers. For centers, the eigenvalues

of fixed points are purely imaginary [28]. Linear stability analysis of anisotropic cosmological models in different gravitational settings can be found here [32–36]. Dynamical system approach have also been used to study bouncing and cyclic universes in different cosmological set-ups in literature [11, 28, 37–40].

We now study the structure of dynamical system via phase plane analysis by using variables H and ρ . In our analysis, we shall consider critical points at finite distance only. The fixed / critical points are characterized by simultaneous conditions $\dot{H} = 0$ and $\dot{\rho} = 0$. Equation (11) may serve as constraint equation for the system. Solving equations (14,15) for $\dot{H} = 0$ and $\dot{\rho} = 0$, we get

1. $P_1: H = 0, \rho = 0$
2. $P_2: H = \frac{\sqrt{k\rho_0}}{\sqrt{3(1+\alpha)(1-4k\lambda)}}, \rho = \frac{\rho_0}{1+\alpha}$
3. $P_3: H = -\frac{\sqrt{k\rho_0}}{\sqrt{3(1+\alpha)(1-4k\lambda)}}, \rho = \frac{\rho_0}{1+\alpha}$

The critical points P_2, P_3 will exist under the conditions $\rho_0 > 0, k > 0, \alpha > -1, \lambda > 0$ with $0 < k\lambda < \frac{1}{4}$. The condition $\rho_0 > 0, \alpha > -1$ ensures $\rho \geq 0$ in the (ρ, H) phase-space. If we allow $\rho < 0$ in the cosmological model then one should have $\rho_0 < 0$ as well.

From equation (6), for $\rho_0 = 0$, the equation of state takes the form $p = \alpha\rho$. For $\rho_0 \neq 0$ case, we have

- At P_1 , the eigenvalues are $\lambda_{1,2} = \pm \frac{i\sqrt{\lambda\rho_0}\sqrt{4k\lambda-1}}{\sqrt{2}\sqrt{(1+\alpha)3k\lambda-1}}$. This point represents an empty flat (Minkowski) model. The point will be center for either

$$\left(\lambda\rho_0 > 0, k\lambda > \max\left\{\frac{1}{4}, \frac{1}{1+\alpha}\right\} \right) \text{ or } \left(\lambda\rho_0 > 0, k\lambda < \min\left\{\frac{1}{4}, \frac{1}{1+\alpha}\right\} \right)$$

This point will act as saddle point if any of the following condition holds:

1. $(\lambda\rho_0 < 0, k\lambda > \max\{\frac{1}{4}, \frac{1}{1+\alpha}\})$
 2. $(\lambda\rho_0 < 0, k\lambda < \min\{\frac{1}{4}, \frac{1}{1+\alpha}\})$
 3. $(\lambda\rho_0 > 0, \frac{1}{4} < k\lambda < \frac{1}{1+\alpha})$
 4. $(\lambda\rho_0 > 0, \frac{1}{1+\alpha} < k\lambda < \frac{1}{4})$
- At P_2 , the eigenvalues are as follows:

$$\lambda_1 = \frac{\sqrt{3k\rho_0}\sqrt{(1+\alpha)(1-4k\lambda)}}{(1+\alpha)k\lambda-1}, \lambda_2 = \frac{\sqrt{k\rho_0}\sqrt{1-4k\lambda}}{\sqrt{1+\alpha}\sqrt{3k\lambda}}$$

The critical point P_2 will exist under the conditions $\rho_0 > 0, k > 0, \alpha > -1, \lambda > 0$ with $0 < k\lambda < \frac{1}{4}$. With condition $k\lambda > \frac{1}{1+\alpha}$, the point P_2 will be an unstable point (repeller) and with $k\lambda < \frac{1}{1+\alpha}$, it will be a saddle point. With $\rho_0 > 0, \alpha > -1$, we have $\rho \geq 0$ in (ρ, H) phase-space. This point represents expanding spatially flat de-Sitter scenario.

- At P_3 , the eigenvalues are as follows:

$$\lambda_1 = -\frac{\sqrt{3k\rho_0}\sqrt{(1+\alpha)(1-4k\lambda)}}{(1+\alpha)k\lambda-1}, \lambda_2 = -\frac{\sqrt{k\rho_0}\sqrt{1-4k\lambda}}{\sqrt{1+\alpha}\sqrt{3k\lambda}}$$

The critical point P_3 will exist under the conditions $\rho_0 > 0, k > 0, \alpha > -1, \lambda > 0$ with $0 < k\lambda < \frac{1}{4}$. With condition $k\lambda > \frac{1}{1+\alpha}$, the point P_3 will be a stable point (attractor) and with $k\lambda < \frac{1}{1+\alpha}$, it is a saddle point. With $\rho_0 > 0, \alpha > -1$, we have $\rho \geq 0$ in (ρ, H) phase-space. This point represents contracting spatially flat de-Sitter scenario.

For $\rho_0 = 0$ case, only one fixed point $P_1: H = 0, \rho = 0$ will exist for system and the eigenvalues are $(\lambda_1, \lambda_2) = (0, 0)$ for this point. We assume that ρ as non-negative, the case $\rho < 0$ may occur dynamically in phase-space, but these cases may regarded as un-physical.

Now, we plot few figures depending on arbitrary values of α, ρ_0, k and λ to display the linear stability nature of critical points P_1, P_2, P_3 existing for $p = \alpha\rho - \rho_0$ case in contrast with critical point P_1 of $p = \alpha\rho$ case. One may also check for different set of values of parameters for stability nature of these points.

For $k = 0.25, \lambda = 0.75, \alpha = 0.25, \rho_0 = 0.5$ phase plane diagram is given in figure (1). In this

Figure 1: ρ vs H for $k = 0.25$, $\lambda = 0.75, \alpha = 0.25, \rho_0 = 0.5$

Figure 2: ρ vs H for $k = 0.25$, $\lambda = 0.75, \alpha = 0.25, \rho_0 = 0$

case, all the 3 critical points exists and point P_1 is center, points P_2, P_3 are saddle points. The critical points P_2 and P_3 are located on the phantom divide line $\rho = \frac{\rho_0}{1+\alpha}$. In figure (2), $\rho - H$ phase plane diagram for $k = 0.25, \lambda = 0.75, \alpha = 0.25, \rho_0 = 0$ has been given. Here, only one fixed point P_1 exists and it is attractor in one direction and repeller in another direction. Therefore, this point is saddle in nature. In expanding models of these cases, since $\frac{3}{K} = 3.15$ (from equation (12)), $\rho \rightarrow \frac{\rho_0}{1+\alpha}$ when $a \rightarrow \infty$.

For $k = 0.125, \lambda = 0.125, \alpha = 1.5, \rho_0 = 0.5$ phase plane diagram is given in figure (3). The critical point P_1 is center, points P_2, P_3 are saddle points. In figure (4), for $k = 0.125, \lambda = 0.125, \alpha = 1.5, \rho_0 = 0$ phase plane diagram has existence of only one critical point P_1 and it is a saddle point. In expanding models of these cases, since $\frac{3}{K} = 7.96$ (from equation (12)), $\rho \rightarrow \frac{\rho_0}{1+\alpha}$ when $a \rightarrow \infty$.

In figure (5), $\rho - H$ phase plane diagram for $k = 0.125, \lambda = 0.125, \alpha = -1.5, \rho_0 = -0.5$ has 3 critical points. Point P_1 is saddle point because eigenvalues are of opposite signs. Point P_2 is repeller and point P_3 is attractor because eigenvalues are negative. The critical points P_2 and P_3 are located on the phantom line $\rho = \frac{\rho_0}{1+\alpha}$. In figure (6), for $k = 0.125, \lambda = 0.125, \alpha = -1.5, \rho_0 = 0$ phase plane diagram has only one critical point P_1 which is a saddle point. In expanding models of these cases, since $\frac{3}{K} = -1.374$ (from equation (12)), so $\rho \rightarrow \infty$ when $a \rightarrow \infty$. Since $\rho \rightarrow \infty$, due to $p = \alpha\rho - \rho_0$ we may have $p \rightarrow \infty$.

Figure 3: ρ vs H for $k = 0.125$,
 $\lambda = 0.125$, $\alpha = 1.5$, $\rho_0 = 0.5$

Figure 4: ρ vs H for $k = 0.125$,
 $\lambda = 0.125$, $\alpha = 1.5$, $\rho_0 = 0$

Figure 5: ρ vs H for $k = 0.125$,
 $\lambda = 0.125$, $\alpha = -1.5$, $\rho_0 = -0.5$

Figure 6: ρ vs H for $k = 0.125$,
 $\lambda = 0.125$, $\alpha = -1.5$, $\rho_0 = 0$

Figure 7: ρ vs H for $k = -0.25$,
 $\lambda = -0.75$, $\alpha = 1.25$, $\rho_0 = 0.5$

Figure 8: ρ vs H for $k = -0.25$,
 $\lambda = -0.75$, $\alpha = 1.25$, $\rho_0 = 0$

Figure 9: ρ vs H for $k = -0.25$, $\lambda = -0.75$, $\alpha = -1.25$, $\rho_0 = -0.5$

Figure 10: ρ vs H for $k = -0.25$, $\lambda = -0.75$, $\alpha = -1.25$, $\rho_0 = 0$

With $k = -0.25$, $\lambda = -0.75$, $\alpha = 1.25$, $\rho_0 = 0.5$ phase plane diagram (given in figure (7)) has only one critical point P_1 and this point is center because eigenvalues are purely imaginary. In figure (8), for $k = -0.25$, $\lambda = -0.75$, $\alpha = 1.25$, $\rho_0 = 0$ only one critical point P_1 is a saddle point. In expanding models of these cases, since $\frac{3}{K} = -6.3528$ (from equation (12)), we have $\rho \rightarrow \infty$ when $a \rightarrow \infty$.

For $k = -0.25$, $\lambda = -0.75$, $\alpha = -1.25$, $\rho_0 = -0.5$ phase plane diagram has been given in figure (9). It has only one critical point P_1 and it is a center. In figure (10), for $k = -0.25$, $\lambda = -0.75$, $\alpha = -1.25$, $\rho_0 = 0$, the model has only one critical point P_1 and it is a saddle point. In expanding models of these cases, since $\frac{3}{K} = -0.1641$ (from equation (12)), we have $\rho \rightarrow \infty$ when $a \rightarrow \infty$.

We note that in Rastall gravity model with linear equation of state, the $\rho - H$ phase space has critical point P_1 which may act as center depending on the values of α , ρ_0 , k and λ . In the case of centers, curves are having closed trajectories around this point in phase space and eigenvalues are purely imaginary and conjugated. The real parts of eigenvalues are zero for cyclic (or non-singular) solutions, as corresponding critical points are centers and thus marginally stable. Due to the properties of center critical point, the nearby trajectories do not converge to the point but rotate about it, realizing oscillating solutions. This is in contrast with $p = \alpha\rho$ case, where only critical point is having saddle nature. Oscillating universes avoids the singularity and replace it with a cyclical evolution from a previous incarnation of universe. During oscillation, the universe undergoes a series of turnaround and bounce where they achieve maxima and minima of radius respectively [67–71]. At the extrema $\dot{a}(t_i) = 0$, where $i = ta, b$ denotes turnaround and bounce respectively. In the contraction and expansion phases of universe, $\dot{a} < 0$ and $\dot{a} > 0$ respectively. The minimal condition for bounce and turnaround in FRW space-time may be stated as [71]

- $\dot{a}(t_b) = 0$ & $\ddot{a}(t) > 0$ for $t \in (t_b - \epsilon, t_b) \cup (t_b, t_b + \epsilon)$ with small $\epsilon > 0$.
- $\dot{a}(t_{ta}) = 0$ & $\ddot{a}(t) < 0$ for $t \in (t_{ta} - \epsilon, t_{ta}) \cup (t_{ta}, t_{ta} + \epsilon)$ with small $\epsilon > 0$.

Equivalently, $H = 0$ at bounce or turnaround point. In small neighbourhood of bounce and turnaround, $\dot{H} > 0$ and $\dot{H} < 0$ respectively. We now discuss the possibility of conditions that lead to non-singular solutions surrounding the critical points by considering the energy conditions or restrictions on the energy-momentum tensor T_{ij} . We now demonstrate connection between a bounce or turnaround with energy conditions in the presence of barotropic equation of state $p = \alpha\rho - \rho_0$.

The behaviour of point-wise energy conditions in a oscillating universe scenario may be of special interest. For perfect fluid source with $T_{ij} = (\rho, -p, -p, -p)$, the expression for energy conditions may be stated as

- Null energy condition (NEC) $\Leftrightarrow \rho + p \geq 0$
- Weak energy condition (WEC) $\Leftrightarrow \rho \geq 0, \rho + p \geq 0$
- Dominant energy condition (DEC) $\Leftrightarrow \rho \geq 0, \rho \pm p \geq 0$
- Strong energy condition (SEC) $\Leftrightarrow \rho + 3p \geq 0, \rho + p \geq 0$

For flat-FRW metric, from equations (3) and (4), at and near the turnaround and bounce, we have

- $\rho - p = \frac{2(1-6k\lambda)}{k} \dot{H}$
- $\rho + p = \frac{-2}{k} \dot{H}$
- $\rho + 3p = \frac{6(2k\lambda-1)}{k} \dot{H}$

The conditions for satisfaction of energy conditions in terms of parameters $k\lambda, \rho_0, \alpha$ are given in Table 1. At the bounce or turnaround, from equations (5) and (6), we have

Energy conditions	Bounce	Turnaround
WEC	$k < 0, \rho > \max\left(0, \frac{\rho_0}{1+\alpha}\right)$	$k > 0, \rho > \max\left(0, \frac{\rho_0}{1+\alpha}\right)$
NEC	$k < 0$	$k > 0$
SEC	$k < 0$ with $k\lambda < \frac{1}{2}$ $\rho > \max\left(\frac{\rho_0}{1+\alpha}, \frac{3\rho_0}{1+3\alpha}\right)$	$k > 0$ with $k\lambda < \frac{1}{2}$ $\rho > \max\left(\frac{\rho_0}{1+\alpha}, \frac{3\rho_0}{1+3\alpha}\right)$

Table 1: Constraints for satisfaction of energy conditions near bounce and turnaround point

$$\ddot{\rho} = 3P(\rho + p)\dot{H}$$

where, $P = \frac{1-4k\lambda}{3k\lambda(1+\alpha)-1}$. For $P > 0$, we have $\min\left(\frac{1}{3(1+\alpha)}, \frac{1}{4}\right) < k\lambda < \max\left(\frac{1}{3(1+\alpha)}, \frac{1}{4}\right)$. And for $P < 0$, $k\lambda < \min\left(\frac{1}{3(1+\alpha)}, \frac{1}{4}\right)$ or, $k\lambda > \max\left(\frac{1}{3(1+\alpha)}, \frac{1}{4}\right)$.

At the bounce $\dot{H} > 0$, the energy density will be

- ρ_{min} if NEC is violated with $P > 0$ or, NEC is satisfied with $P < 0$.
- ρ_{max} if NEC is violated with $P < 0$ or, NEC is satisfied with $P > 0$.

At the turnaround $\dot{H} < 0$, the energy density will be

- ρ_{min} if NEC is violated with $P < 0$ or, NEC is satisfied with $P > 0$.
- ρ_{max} if NEC is violated with $P > 0$ or, NEC is satisfied with $P < 0$.

The violation or satisfaction of energy conditions near bounce or turnaround depend on $\alpha, \rho_0, k\lambda$. So, one may identify the evolution of non-singular universes on the basis of existence of center fixed point along-with violations of energy conditions. Violation of NEC is also affecting the energy density value at bounce and turnaround respectively. By employing barotropic fluid satisfying quadratic equation of state $p = \alpha\rho + w\rho^2$, the field equation in the Rastall gravity framework has been solved analytically [11]. With the low energy regime equation of state [60] of form $p = \alpha\rho - \rho_0$, the present model has peculiar feature of having empty flat (Minkowskian) fixed point (which may act as center in few cases) and thus may exhibit oscillating behaviour of universe in the model. In literature, these kind of cyclic solutions are illustrated using dynamical system technique in General relativity framework with Chaplygin gas [28].

3 Horizon thermodynamics

In the present section, we investigate the thermodynamical behaviour of cosmological model by considering the universe as a thermodynamic system. For FRW geometry using apparent horizon, one can extract the Friedmann equations by applying first law of thermodynamics [46, 52]. The apparent horizon coincides with Hubble horizon for flat FRW model, but these two are different for non-flat FRW model. In case of dynamical FRW space-time, the apparent horizon has been argued to be associated with surface gravity and the gravitational entropy, and thus may be treated as causal horizon [72]. Therefore, one may consistently define thermodynamic laws with apparent horizon, since it is the appropriate one among various choices of cosmological horizons [72]. The accelerating universe enveloped by apparent horizon satisfies GSLT, whereas accelerating universe inside the event horizon does not [73]. From thermodynamics point of view in cosmological scenario, the enveloping surface should be apparent horizon and not the event horizon [74]. Apparent horizon as universe's boundary have been used in literature for examining the validity of GSLT in different cosmological set-ups [19, 23, 25, 52, 55, 57, 61, 73, 74].

In flat-FRW universe, radius of apparent horizon is related with Hubble parameter as $R_a = \frac{1}{H}$. Here, we consider the universe as a thermodynamic system with it's boundary being the apparent horizon surface. In case of FRW space-time, the entropy measured by a comoving observer is given by $S = S_a + S_i$, where the first term is entropy of apparent horizon and second term S_i correspond to the entropy of fluid content enclosed by horizon. GSLT states that the time variation of the entropy of horizon (S_a) plus fluid entropy (S_i) inside it should be positive definite [25]. First derivative of R_a with respect to cosmic time t is given by $\dot{R}_a = -\frac{\dot{H}}{H^2}$. The modified Bekestien entropy on the apparent horizon in Rastall's gravity is given as [47]

$$S_a = \frac{\tilde{A}}{4} \quad (16)$$

where $\tilde{A} = \left(1 + \frac{2\gamma}{4\gamma-1}\right) A$ with $A = 4\pi R_a^2$ and $\gamma = k\lambda$ being the dimensionless parameter. In terms of γ , we have $k = \frac{4\gamma-1}{6\gamma-1}8\pi$ and $\lambda = \frac{\gamma(6\gamma-1)}{(4\gamma-1)8\pi}$. It is worthwhile to mention that λ diverges at $\gamma = \frac{1}{4}$ and k diverges at $\gamma = \frac{1}{6}$. So, $\gamma = \frac{1}{4}, \frac{1}{6}$ cases are excluded in our study. In this paper, the units of $c = \hbar = G = 1$ has been considered. The Hawking temperature at apparent horizon is given as [48]

$$T_a = \frac{1}{2\pi R_a} \quad (17)$$

The differential amount of energy crossing the apparent horizon is given as [49]

$$-dE_a = \frac{1}{2} R_a^3 (\rho + p) H dt \quad (18)$$

The derivative of surface entropy can be written as $\dot{S}_a = - \left(1 + \frac{2\gamma}{4\gamma-1}\right) 2\pi \frac{\dot{H}}{H^3}$. Therefore, we have

$$T_a \dot{S}_a = - \left(1 + \frac{2\gamma}{4\gamma-1}\right) \frac{\dot{H}}{H^2} \quad (19)$$

With help of Clausius relation ($-dE_a = T_a dS_a$), the first law of thermodynamics is given as [19]

$$\Omega dt = T_a dS_a + dE_a \quad (20)$$

The first law of thermodynamics holds when $\Omega \rightarrow 0$, which lead to the constraint $k\lambda = \frac{k-1}{2(3k-2)}$. We assume that horizon and barotropic fluid confined by it are in thermal equilibrium with each other. For generalized second law of thermodynamics to hold, we must have $d(S_i + S_a) \geq 0$. The Gibb's equation is of form:

$$T_i dS_i = dE_i + p dV \quad (21)$$

where T_i is temperature inside the horizon, $E_i = \rho V$ is internal energy, S_i is the entropy inside horizon and $V = \frac{4}{3}\pi R_a^3$ and therefore

$$T_i dS_i = \frac{8\pi}{k} \frac{\dot{H}^2}{H^4} + \frac{8\pi}{k} \frac{4\gamma-1}{(1+\alpha)3\gamma-1} \frac{\dot{H}}{H^2} \quad (22)$$

Therefore, $T\dot{S} = T_a \dot{S}_a + T_i \dot{S}_i$ can be written as

$$T\dot{S} = \frac{8\pi}{k} \left(A \frac{\dot{H}}{H^2} + \frac{\dot{H}^2}{H^4} \right) \quad (23)$$

where $A = \frac{\gamma(1-3\alpha)}{3\gamma(1+\alpha)-1}$. Cosmological data indicates that $\dot{H} \leq 0$ [75]. For generalized second law of thermodynamics to be valid in the Rastall's gravity model with linear equation of state, we only need $A \leq 0$ for the expanding cosmologies. The fluid satisfying equation (6) will be hydrodynamically stable for $0 \leq \alpha \leq 1$ [60]. With the condition $0 \leq \alpha \leq 1$, we plot A where any point (α, γ) lying on the surface satisfies the constraint $A \leq 0$ (see figure 11a). If we lift the condition of classical stability of the model, the behaviour of A for different (α, γ) with the condition $A \leq 0$ has been shown in the figure 11b.

We know that $q+1 = -\frac{\dot{H}}{H^2}$, where q denotes the deceleration parameter which is an indicator of the rate at which the expansion of universe is slowing down. The cosmological model is decelerating for $q > 0$. The negative values of q gives information about the general dynamical behaviour of the universe: de Sitter expansion happens at $q = -1$, accelerating power-law expansion can be achieved for $-1 < q \leq 0$ and a super-exponential expansion happens for $q < -1$. In terms of deceleration parameter q , equation (23) may be written as

$$T\dot{S} = \frac{8\pi}{k} [(q+1)^2 - A(q+1)] \quad (24)$$

The above equation can also be used to discuss the validity of GSLT, especially when one is discussing the transition of universe from the decelerated era to the accelerated era. In the following discussion, we assume $k > 0$. In the framework of Rastall gravity contracting cosmologies, with $A \leq 0$ the accelerated contracting universe (the case when $-1 < q < 0$), the GSLT will be valid. In the decelerated contracting universe, the validity of GSLT demands $A \leq 0$. Here, one can observe that in de Sitter expansion era, $T\dot{S} = 0$. In the phase-space analysis, we have observed that points P_2 and P_3 represents expanding and contracting spatially flat de Sitter model. One can conclude that at these critical points, the modified Bekestien entropy has critical value as well.

Figure 11: The constraint $A \leq 0$ with admissible values of (α, γ) .

4 Conclusion

In this paper, we have studied phase-space of Rastall gravity model with barotropic fluid satisfying linear equation of state. We obtain the critical points and presented the conditions for their existence and stability. We have discussed the behaviour of critical points with corresponding figures. For the critical point P_1 (that is, $(H, \rho) = (0, 0)$) eigenvalues have non-zero imaginary part which means that oscillating solutions may exists in Rastall model with linear equation of state. However, other two points P_2 and P_3 may be attractor or repeller or of saddle nature for different values of parameters. Point P_2 represents expanding de Sitter scenario and point P_3 represents contracting de Sitter scenario. Stability of critical points does not mean that model is stable. In order to say that model is stable, we must investigate classical stability and quantum stability. We discuss the classical stability of the model (classical fluctuations may be regarded to be stable when the speed of sound satisfies $0 \leq c_s^2 \leq 1$). This criterion in present model has been observed by constraint $0 \leq \alpha \leq 1$, which is consistent from literature [60]. Quantum stability is beyond the scope of present work.

Equation of state $p = \alpha\rho - \rho_0$ takes the form $p = \alpha\rho$ for $\rho_0 = 0$. We also have discussed the stability of critical points with $\rho_0 = 0$ case. The Minkowskian critical point has always stability of saddle nature.

We have explored the validity of generalized second law of thermodynamics in resulting expanding and contracting cosmologies with equilibrium description. We may conclude that GSLT will not be valid, at least in the super-exponential expansion with $A < 0$ and in the neighbourhood of bouncing and turnaround point during periodic evolution of the universe.

Acknowledgement

We are very thankful to the anonymous learned reviewers for insightful and interesting comments, which have been helpful in improvement of the manuscript.

References

- [1] P. Rastall, Phys. Rev. D **6**, 3357 (1972)
- [2] P. Rastall, Can. J. Phys. **54**, 66 (1976)
- [3] C.M. Will, Theory and Experiments in Gravitational Physics, Cambridge University Press, UK (1993)
- [4] N.D. Birrell, P.C.W. Davies, Quantum fields in curved space, Cambridge University Press, Cambridge (1982)
- [5] C.E.M. Batista, M.H. Daouda, J.C. Fabris, O.F. Piattella and D.C. Rodrigues, Phys. Rev. D **85**, 084008 (2012)
- [6] W.A.G. De Moraes, A.F. Santos, Gen. Relativ. Grav. **51** 167 (2019)
- [7] H. Shabani and A.H. Ziaie, EPL **129** 20004 (2020)
- [8] J.C. Fabris, M.H. Daouda and O.F. Piattella, Phys. Lett. B **711**, 232 (2012)
- [9] M.H. Daouda, J.C. Fabris and O.F. Piattella, AIP Conf. Proc. **1471**, 57 (2012)
- [10] J.C. Fabris, O.F. Piattella, D.C. Rodrigues, C.E.M. Batista, and M.H. Daouda, Int. J. Mod. Phys. Conf. Ser. **18**, 67 (2012)
- [11] G.F. Silva, O.F. Piattella, J.C. Fabris, L. Casarini, T.O. Barbosa, Gravit. Cosmol. **19**, 156 (2013)
- [12] H. Moradpour, Phys. Lett. B **757**, 187 (2016)
- [13] H. Moradpour, N. Sadeghnezhad and S.H. Hendib, Can. J. Phys. **95**, 1257 (2017)
- [14] H. Moradpour, Y. Heydarzade, F. Darabi et al. Eur. Phys. J. C **77**, 259 (2017)
- [15] F-F. Yuan and P. Huang, Class. Quantum Grav. **34**, 077001 (2017)
- [16] F. Darabi, H. Moradpour, I. Licata, Y. Heydarzade, C. Corda, Eur. Phys. J. C **78**, 25 (2018)
- [17] M. Visser, Phys. Lett. B **782**, 83 (2018)
- [18] F. Darabi, K. Atazadeh and Y. Heydarzade, Eur. Phys. J. Plus **133**, 249 (2018)
- [19] K. Bamba, A. Jawad, S. Rafique and H. Moradpour, Eur. Phys. J. C **78**, 986 (2018)
- [20] D. Das, S. Dutta and S. Chakraborty, Eur. Phys. J. C **78**, 810 (2018)
- [21] S. Hansraj, A. Banerjee, P. Channuie, Annals of Physics **400**, 320 (2019)
- [22] W. Khylllep, J. Dutta, Phys. Lett. B **797**, 134796 (2019)
- [23] M. Cruz, S. Lepe, G. Morales-Navarrete, Class. Quantum Grav. **36** 225007 (2019)
- [24] Z.-Xi Yu, H. Wei, Nucl. Phys. B **960** 115179 (2020)
- [25] A. Singh and K.C. Mishra, Eur. Phys. J. Plus **135**, 752 (2020)

- [26] A.H. Ziaie, H. Moradpour and H. Shabani, Eur. Phys. J. Plus **135** 916 (2020)
- [27] S. Ghaffari, A.A. Menon, H. Moradpour, A.H. Ziaie, Mod. Phys. Lett. A **35** 2050276 (2020)
- [28] A. Salehi, Phys. Rev. D **94** 123519 (2016)
- [29] A.A. Coley, Dynamical Systems and Cosmology, Springer (2003)
- [30] G.F.R. Ellis and J. Wainwright, Dynamical systems in cosmology, Cambridge University Press, Cambridge (2005)
- [31] S. Bahamonde, C.G. Bohmer, S. Carloni, E.J. Copeland, W. Fang and N. Tamanini, Phys. Rep. **775 - 777**, 1-122 (2018)
- [32] R. Chaubey, R. Raushan, Astrophys Space Sci **361**, 215 (2016)
- [33] R. Raushan, R. Chaubey, Can. J. Phys. **95** 1049 (2017)
- [34] M. Sharif and S. Mumtaz, Gen. Relativ. Gravit. **50**, 63 (2018)
- [35] R. Raushan, R. Chaubey, Int. J. Geom. Methods Mod. Phys. **16**, 02, 1950023 (2019)
- [36] S. Angit, R. Raushan, R. Chaubey, Int. J. Geom. Methods Mod. Phys. **16**, 1950127 (2019)
- [37] J. De-Santiago, J.L. Cervantes-Cota and D Wands, Phys. Rev. D **87**, 023502 (2013)
- [38] T. Bandyopadhyay, U. Debnath, Can. J. Phys. **97**, 286 (2019)
- [39] J.D. Barrow and A. Paliathanasis, Eur. Phys. J. C **79**, 379 (2019)
- [40] R. Raushan, A. Singh, R. Chaubey and T. Singh, Int. J. Geom. Methods Mod. Phys. **17**, 2050064 (2020)
- [41] J.M. Bardeen, B. Carter and S.W. Hawking, Comm. Math. Phys. **31**, 161 (1973)
- [42] J.D. Bekenstien, Phys. Rev. D **7**, 2333 (1973)
- [43] S.W. Hawking, Comm. Math. Phys. **43**, 199 (1975)
- [44] T. Jacobson, Phys. Rev. Lett. **75**, 1260 (1995)
- [45] C. Eling, R. Guedens and T. Jacobson, Phys. Rev. Lett. **96**, 121301 (2006)
- [46] M. Akbar and R.G. Cai, Phys. Lett. B **648**, 243 (2007)
- [47] H. Moradpour and I.G. Salako, Adv. High Energy Phys. **2016** 3492796 (2016)
- [48] R.G. Cai, L.M. Cao and Y.P. Hu, Class. Quantum Grav. **26** 155018 (2009)
- [49] R. Bousso, Phys. Rev. D **71** 064024 (2005)
- [50] A. Sheykhi, M.H. Dehghani, R. Dehghani, Gen. Relativ. Gravit. **46**, 1679 (2014)
- [51] A. Sheykhi, B. Wang, R.G. Cai, Nucl. Phys. B **779**, 1-12 (2007)
- [52] R.G. Cai and L.M. Cao, Phys. Rev. D **75**, 064008 (2007)

- [53] T. Padmanabhan, *Phys. Rep.* **406**, 49 (2005)
- [54] A. Singh, A.K. Shukla, *Int. J. Mod. Phys. A* **35**, 2050054 (2020)
- [55] G.P. Singh, N. Hulke and A. Singh, *Can. J. Phys.* **96**, 992 (2018)
- [56] A. Singh, R. Raushan, R. Chaubey and T. Singh, *Int. J. Mod. Phys. A* **33**, 1850213 (2018)
- [57] A. Pourbagher & A. Amani, *Astrophys. Space Sci.* **364**, 140 (2019)
- [58] H.B. Callen, *Thermodynamics and introduction to thermostatics*, John Wiley & Sons, New York (1985)
- [59] M Capone et al., *J. Phys.: Conf. Ser.* **222**, 012012 (2010)
- [60] E. Babichev, V. Dokuchaev, Yu. Eroshenko, *Class. Quant. Grav.* **22**, 143 (2005)
- [61] G.P. Singh, N. Hulke and A. Singh, *Int. J. Geom. Methods Mod. Phys.* **15**, 1850129 (2018)
- [62] T. Singh and R. Chaubey, *Pramana* **71**, 447 (2008)
- [63] S.K. Tripathy, D. Behera, B. Mishra, *Eur. Phys. J. C* **75** 149 (2015)
- [64] K.N. Ananda, M. Bruni, *Phys. Rev. D* **74** 023523 (2006)
- [65] J. Kehayias and R.J. Scherrer, *JCAP* **12** 015 (2015)
- [66] S. Nojiri, S.D. Odintsov, S. Tsujikawa, *Phys. Rev. D.* **71** 063004 (2005)
- [67] J.D. Barrow, M.P. Dabrowski, *Mon. Not. R. Astron. Soc.* **275**, 850-862 (1995)
- [68] C. Molina-Paris, M. Visser, *Phys. Lett. B*, **455** 90–95 (1999)
- [69] T. Singh, R. Chaubey, A. Singh, *Eur. Phys. J. Plus*, **130** 31 (2015)
- [70] T. Singh, R. Chaubey, A. Singh, *Int. J. Mod. Phys. A*, **30** 1550073 (2015)
- [71] M. Novello, S.E.P. Bergliaffa, *Phys. Rep.* **463** 127–213 (2008)
- [72] D. Bak, S.J. Rey, *Class. Quantum Grav.* **17** L83 (2000)
- [73] J. Zhou, B. Wang, Y. Gong, E. Abdalla, *Phys. Lett. B* **652** 86 (2007)
- [74] K. Bamba, C.-Q. Geng, *JCAP* **11** 008 (2010)
- [75] A. Singh, *Astrophys. Space Sci.* **365** 54 (2020)