Canadian Journal of Fisheries and Aquatic Sciences

Where do you come from, where do you go: Early life stage drift and migrations of cod inferred from otolith microchemistry and genetic population assignment

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2020-0409.R2
Manuscript Type:	Article
Date Submitted by the Author:	20-May-2021
Complete List of Authors:	Hüssy, Karin; Technical University of Denmark, National Institute of Aquatic Sciences Albertsen, Christoffer; Technical University of Denmark, National Institute of Aquatic Sciences Hemmer-Hansen, Jakob; Technical University of Denmark, Vinther, Morten; Technical University of Denmark, National Institute of Aquatic Sciences Serre, Simon Hansen; Geological Survey of Denmark and Greenland Thomsen, Tonny Bernt; Technical University of Denmark Eero, Margit; Technical University of Denmark, National Institute of Aquatic Sciences
Keyword:	GENETICS < General, migrations, otolith microchemistry, stock mixing, spawning stock origin
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Where do you come from, where do you go: Early life stage drift and migrations of cod inferred from otolith microchemistry and genetic population assignment

Karin Hüsey^{1*}, Christoffer Moesgaard Albertsen¹, Jakob Hemmer-Hansen², Morten Vinther¹, Simon Hansen Serre³, Tonny Bernt Thomsen³, Margit Eero¹

Affiliations

¹ National Institute of Aquatic Resources, Technical University of Denmark, Kemitovet, building 201, Lyngby Campus, 2800 Kgs. Lyngby, Denmark

² National Institute of Aquatic Resources, Technical University of Denmark, Vejlsvvej 39, 8600 Silkeborg, Denmark

³ Geological Survey of Denmark and Greenland, Øster Voldgade 10, DK-1350 Copenhagen K., Denmark

* Corresponding author: kh@aqua.dtu.dk, Mobile +45 93511840

Competing interests: The authors declare there are no competing interests.

Abstract

This study investigates stock mixing of genetically distinct Atlantic cod (*Gadus morhua*) stocks in the Kattegat, an area geographically located between the North Sea and the Baltic Sea, by combining genetic population identification with habitat assignments from hatch to capture from otolith microchemistry. Cod captured in Kattegat were genetically assigned to either the North Sea or the endemic Kattegat population. Otolith chemical fingerprints differed significantly between populations during the larval and pelagic juvenile stage with higher strontium and lower barium and manganese concentrations in the North Sea population than the Kattegat population, indicating that North Sea cod are spawned in the North Sea or Skagerrak and drift into the Kattegat during the early life stages. Individual cod of both populations undertook frequent, but predominantly short term, migrations to other areas than the Kattegat, with < 25 % of individuals remaining resident within the Kattegat throughout their life. Across seasons and age classes, the two populations were both most frequently distributed in the Kattegat (67 %), with approximately 25 % of both population distributed in the western Baltic Sea and less than 10 % in the Skagerrak/North Sea. This study demonstrates the usefulness of this approach to infer population-specific connectivity and migration trajectories for individual fish and its potential applications in basic and applied fields of fish ecology and fisheries sciences.

Key words: genetics, migrations, otolith microchemistry, spawning stock origin, stock mixing

Introduction

Fish migrations and stock mixing are major issues for assessments of stock status and for fisheries management, which are usually constrained to geographical management areas that do not always correspond to the population's distribution boundaries (e.g. Kerr et al. 2017). Stock assessment approaches are being developed to account for spatial stock structure (Cadrin and Secor 2009; Goethel et al. 2011; Albertsen et al. 2018). However, an understanding of migration patterns and stock mixing must be gained before the potential of these assessment models can be fully realized. The Kattegat is a good case for testing different approaches to investigate migration and mixing, since it geographically is located between the North Sea and the Baltic Sea (Fig. 1). Atlantic cod (*Gadus morhua*) populations inhabit the entire area and area currently managed as four stocks: the North Sea/Skagerrak, the Kattegat, the western Baltic Sea/Sound and the eastern Baltic Sea, with distinct spawning areas and seasons (Brander 1994; Fox et al. 2008; Vitale et al. 2008; Munk et al. 2009; Hüsey, 2011; Börjesson et al. 2013). Cod from these areas are clearly distinct genetically (Nielsen et al. 2003; Berg et al. 2015; Barth et al. 2019), and we refer to these populations as the "North Sea", "Kattegat" and eastern and western "Baltic Sea" throughout this study. In the Kattegat, mixing of the local cod population (the Kattegat population) with adjacent populations (primarily the North Sea population) takes place (Knutsen et al. 2004; ICES 2015; André et al. 2016). However, the mechanisms and dynamics of the mixing and migrations patterns of cod in this area are currently not well understood. It is believed that connectivity between these populations exists through both drift of early life stages (Eero et al. 2016; Jonsson et al. 2016; André et al. 2016) as well as migration of adult individuals (Svedäng et al. 2007). There is no targeted cod fishery in Kattegat at present, and cod is mainly taken as bycatch in the trawl fishery for Norway lobster (*Nephrops norvegicus*) (ICES 2020). Lack of knowledge on population mixing hampers stock assessment

as well as limits the understanding of the recovery potential of the presently severely depleted Kattegat cod population (ICES 2019).

As an environmental transition zone between the North Sea and the Baltic Sea, Kattegat is well suited for the application of methods for inferring stock mixing. The bathymetry becomes successively shallower from > 120 m in the North Sea and Skagerrak, to 40 – 80 m in the northern Kattegat, to depths of 20 – 40 m in the southern Kattegat. The hydrography is characterized by deep-water inflow of saline water from the North Sea and outflow of freshwater from river-runoff in the surface. These topographic and hydrographic conditions lead to a salinity gradient ranging from fully marine saltwater in the North Sea to brackish water of very low salinity in the eastern Baltic Sea (Fig. S1). This salinity gradient provides an ideal setup for studying fish migrations based on chronological analyses of the chemical composition of their otoliths.

Otoliths consist of calcium carbonate (~98 %) and organic matrix (~2 %) and small quantities of trace elements. The biomineralization of the otolith is regulated by physiological processes resulting in both daily growth increments and distinct annual growth zones reflecting seasonally fluctuating conditions in temperature and food availability (Beckman and Wilson 1995; Weidman and Millner 2000; Høie and Folkvord 2006). Trace elements are absorbed primarily from the water across the gill surface and therefore provide a record of environmental conditions experienced by the fish (Watanabe et al. 1997; Campana 1999; Milton and Chenery 2001). The chemical composition of the water depends on the geo-chemistry of the surrounding catchment and therefore provides an area-specific chemical “fingerprint” which is reflected in the fish’s otoliths (Walther and Limburg 2012). The most prominent example of an element’s applicability for reconstruction of migration patterns is strontium (Sr), which since the mid 1990s has been considered a useful proxy for environmental salinity owing to its ability to track

fish's migrations across salinity gradients (Limburg 1995; Secor et al. 1995). Otolith chemistry has over the last three decades gained increasing attention as a tool for analysing fish stock dynamics, migration patterns, pollution exposure, connectivity between areas, and plays an increasingly important role as a fisheries management tool (Campana 1999; Campana and Thorrold 2001; Elsdon et al. 2008; Carlson et al. 2017). Statistically significant differences in element concentrations have previously been observed between cod stocks from the North Sea, eastern and western Baltic Sea, where concentrations of strontium (Sr) and magnesium (Mg) decreased continually from North Sea to Baltic Sea, while concentrations of barium (Ba), zirconium (Zr) and manganese (Mn) increased (Heidemann et al. 2012). The strongest differences occurred between the North Sea and the western Baltic (Mg, Mn) or between these two areas and the eastern Baltic (Sr, Ba, Zr). Within the eastern Baltic Sea, no strong differences in element compositions were observed between different spawning areas. Chemical fingerprints even suggested indications for small-scale stock-structuring and natal homing as important stock separating mechanisms over short distances of < 100 km within the Kattegat/Sound area (Svedäng et al. 2010).

We combined genetic population identification of cod originating from two different populations (North Sea and Kattegat), but caught together in the Kattegat, with chemical composition of otoliths analysed with a new state-of-the art regime-switching state-space migration model, to examine drift and migration patterns of each individual cod. In this study we demonstrate the applicability of this approach to provide new insights into the drift and migration patterns between geographically adjacent areas. More specifically, we first tested whether cod caught in the Kattegat consist of a genetic mixture of populations (North Sea and Kattegat). Then we tested whether North Sea cod are spawned in the same spawning areas as Kattegat cod by comparing the elemental composition of the otolith core, corresponding to the

chemical signature at the time of hatching, between the two populations. Subsequently, we tested if adult cod in the Kattegat perform return-migrations to the Skagerrak/North Sea or the Baltic Sea by assigning area occupation throughout the lifetime of each cod using the chronological record of otolith strontium concentration.

Materials and methods

Sample collection

Atlantic cod were sampled during research surveys in December 2016 in the Kattegat (Fig. 1). From each cod a dorsal muscle tissue sample was stored in an Eppendorph vial with 98 % ethanol for genotyping. Prior to the selection of otolith samples for this study, all cod were genetically assigned to either the North Sea or the Kattegat population (for details on genetic population assignment, see below) and their age determined by expert readers from the sectioned otoliths following routine procedures (Vitale et al. 2019). The bathymetry of the Kattegat is somewhat heterogeneous, where the area south of 57 degree north is shallower than the northern area. From the genotyped individuals, 306 otoliths were therefore selected to ensure adequate sample sizes aiming at a uniform geographic coverage (northern and southern Kattegat) within the size ranges < 35cm, 40 – 55 cm, > 60cm (corresponding age ranges: 0 – 1 years, 2 – 3 years, > 3 years), as well as of the two populations (North Sea and Kattegat). Equal sample distribution was not possible owing to limited catches in some of the groups. The spatial distribution and number of samples within groups are summarized in Fig. 1 and Table 1. Otoliths were extracted, cleaned of adhering tissue, air dried and stored in individually labelled paper bags.

Genetic assignment to population of origin

Genetic assignment to population of origin was conducted through the analyses of 192 genetic markers (single nucleotide polymorphism, SNPs) with high levels of population differentiation in comparisons involving North Sea, Kattegat and Eastern Baltic samples. The markers were selected from published studies (Heath et al. 2014, Barth et al. 2018; Nielsen et al. 2012) and re-analyzed in new baseline samples analyzed for the specific purpose in this study (586 fish in total baseline; see Supplementary information). DNA was extracted by Chelex resin (Estoup et al. 1996) and SNPs were genotyped on a Fluidigm Biomark HD system. After initial screening of the 192 markers, 5 markers were excluded because they did not provide reliable genotype information, resulting in a final data set composed on 187 SNPs. Since the markers originated from different studies, we re-assessed marker independence through analyses of linkage disequilibrium in new baseline samples from the North Sea, Kattegat and the eastern Baltic Sea, using the R package LDheatmap (Shin et al. 2006). These analyses found low levels of LD between the markers, confirming that they would provide independent information for population assignment (see Supplementary information, Fig. S2). Assignment to the most likely baseline sample (“North Sea”, “Kattegat”, “Eastern Baltic”) was conducted by calculating genotype likelihoods (following Rannala and Mountain (1997)) using the programme GeneClass2 (Piry et al. 2004). Population assignment to baseline samples was based on the highest assignment score (ranging between 0 and 100; a measure of the likelihood of the most likely population divided by the sum of all likelihoods) calculated for each fish. Re-analyses of the new panel of markers in baseline samples confirmed the high power for population assignment, as few baseline fish were mis-assigned and likelihood ratio distributions in pairwise comparisons were well separated. For details, see Supplementary information (Fig. S3).

Otolith preparation and chemical analysis

In the laboratory, otoliths were soaked in deionized water, cleaned for 10 minutes in an ultrasonic bath of deionized water, rinsed under deionized water and left to dry overnight under a laminar flow hood in acid-washed trays. Otoliths were embedded in Epoxy resin (Struers®)

and sectioned through the core using an Accutom-100 multi-cut sectioning machine to obtain a 1 cm wide block containing the rostral part of the otolith with the nucleus exposed at the sectioned surface. The surface of each section was polished with 3 μm abrasive paper mounted on rotating disks (Buehler®) to obtain a smooth surface and cleaned in the ultrasonic bath again as described above. Trace element analyses were carried out by Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA ICP-MS) at the Geological Survey of Denmark and Greenland (GEUS), employing a NWR213 frequency-quintupled Nd:YAG solid state laser system from Elemental Scientific Lasers (ESI) that was coupled to an ELEMENT 2 double-focusing, single-collector magnetic sector field ICP-MS from Thermo-Fisher Scientific. Each transect line analysis used a beam diameter of 40 μm and a laser fluence of $\sim 9.5 \text{ J/cm}^2$, a repetition rate of 10 Hz, and a travelling speed of 5 $\mu\text{m sec}^{-1}$. This study focused on the measurement of magnesium (^{25}Mg), calcium (^{43}Ca), manganese (^{55}Mn), copper (^{65}Cu), zinc (^{66}Zn), strontium (^{88}Sr) and barium (^{137}Ba), which all are elements known to have discriminatory power in cod from this area (Heidemann et al. 2012). Concentrations are reported in element:Ca ratios in ppm, using Ca as an internal standard element to account for any variable sample introduction parameters affecting the ablation yield such as variation in the amount of ablated material, laser energy, and ablation rate. Further details on operating conditions, data acquisition parameters, analytical protocols and data processing techniques are described in Serre et al. (2018) which describes the general procedure for analysing otoliths at Geological Survey of Denmark and Greenland. Additional details of the analytical settings are presented in the Supplementary information (Table S1), including analytical precision and accuracy of the LA ICP-MS data (Fig. S4). The otoliths were analysed along a transect from the nucleus to the dorsal edge of the otolith following the axis of maximum growth. The data thus represent elemental signatures spanning from hatch to death of each individual. Values >

4x standard deviations from the mean were treated as outliers and discarded (percentage of data discarded, Mg: 0.6 %, P: 0.3 %, Mn: 0.7, Cu: 1.9 %, Zn: 1.7 %, Sr: 0.4 %, Ba: 0.7 %). Otolith sections were digitized using a Leica DCF290 camera at a magnification of 380 μm pixel⁻¹ with a standard setup (8 bit/channel, 2048 x 1536 pixel frame). Otolith growth chronologies were obtained for each individual by measuring the widths of successive opaque and translucent growth bands along the laser track, from nucleus to edge using ImageJ (Rueden et al. 2017). Opaque zones were divided into three equally spaced sections in order to obtain a measure of sub-seasonal time of formation (spring, summer, fall). LA ICP-MS data were thereafter assigned to the corresponding zones of the otolith and element concentration values averaged by zone, thereby allowing to assess the data on a temporal scale representing different periods in the fish's life (Fig. 2).

Statistical analyses

Otolith size and growth patterns

Linear trends in the element – fish size relationship, if present, are usually removed by subtracting the common, within-group slope of the regression. Since otolith size and fish size are strongly correlated in cod (Li et al. 2008), we tested whether the otolith size at the end of the first year of life differed between year classes and populations using ANOVA, with subsequent Tukey's Honest Significant Difference test for post-hoc pairwise comparison. Since some of the elements analysed here are under physiological regulation, leading to the possibility of growth-related differences in element concentration (Hüssy et al. 2020), we tested whether there were differences in growth rate between the two populations. Growth patterns were compared by testing the linearity of the relationship between fish size on fish age using population as group covariate with an ANCOVA. All analyses were carried out in "R" ver. 3.4.2 (R Core Team 2020).

Spawning origin

Identification of spawning origin: Unfortunately, no larval otolith samples were available to be used as baseline, therefore an accurate assignment of the early life stages is not possible. The conclusions here are therefore based on the approach that if chemical fingerprints of the otolith core are not statistically different between populations, all individuals have been spawned in the same area (presumably the Kattegat). On the other hand, if the chemical fingerprints are significantly different between populations, individuals from the two populations have been spawned in different areas, implying that individuals of the North Sea genotype were spawned in either Skagerrak or the North Sea and had been advected into the Kattegat (Eero et al. 2016), while those of the Kattegat genotype were spawned in the Kattegat (Jonsson et al. 2016).

Year class effect on chemical fingerprint: The samples used in this study contain 8 different age classes, spanning the year classes 2008 - 2016 (Table 1). To rule out bias introduced by combining samples from different year classes, we tested whether there were differences in chemical fingerprints in between year classes using ANOVA, with subsequent Tukey's Honest Significant Difference test for post-hoc pairwise comparison in "R" ver. 3.4.2 (R Core Team 2020) prior to all subsequent analyses.

Chemical fingerprints: Analyses of variance are based on the assumptions that elemental concentrations are normally distributed and that variance is homogenous between groups, in the present case the two populations. Comparison of elemental concentrations between populations also requires that there is no size effect on element concentration. Element concentrations were tested for normality of distribution using Shapiro's test, and homogeneity of variance using Bartlett's test. Non-normally distributed data were (log+1)-transformed. Univariate (ANOVA) tests were used to test for differences in individual elements and multivariate tests (MANOVA) on all elements combined to test for overall differences between

the two populations using the packages “MVA” (Everitt and Hothorn 2011) and “MASS” (Venables and Ripley 2002) in “R” ver. 3.4.2 (R Core Team 2020). Stepwise Linear Discriminant Analysis (LDA) using centred and scaled coefficients was used to identify the elements that drive differences between populations and estimate their ability to correctly classify individuals into the correct group using the “lfa” package (Tang and Li 2016). Differences between populations were visualized using biplots of the first two discriminant functions against each other. In all three analyses, each specific period in the fish’s life was analysed separately. To rule out bias introduced by combining samples from different year classes, the elemental composition in the pelagic juvenile stage was compared between year classes within each population separately using ANOVA and MANOVA, prior to testing for differences in elemental concentration between populations.

Migration analyses

Because otolith Sr concentration is strongly correlated with ambient concentrations, with correlations generally > 0.95 found throughout the literature (review of studies in Hüseyin et al. (2020)) the analyses of migration patterns presented in this study rely on otolith Sr concentration exclusively. Some studies indicate a potential impact of seasonal temperature and growth on otolith Sr concentration (Sturrock et al. 2015; Walther et al. 2010), while others found that otolith Sr patterns were unrelated to water temperature and somatic growth (Clarke and Friedland 2004). To avoid introducing bias into the following analyses related to these potential seasonal effects of temperature and growth, otolith Sr concentrations in consecutive seasons and ages were tested using an ANOVA with subsequent Tukey’s Honest Significant Difference test for post-hoc pairwise comparison in “R” ver. 3.4.2 (R Core Team 2020). Migration patterns were analysed based on otolith strontium concentrations using a newly developed regime-switching state-space model. Specific details of this approach may be found

in Albertsen et al. (*accepted*). Here, only a brief summary of the approach is given. Throughout this paper, “habitat preference” is referring to geographic area (see Fig. 1) and “habitat assignment” means the geographic area each otolith Sr measurement along the otolith chronologies is assigned to.

Since LA ICP-MS transect data have inherent measurement variability, a regime-switching state-space model was developed to filter the signal from the noise and infer habitat preference and occupancy for each measurement along a transect. The model consisted of three layers: 1. A discrete, unobserved Markov chain to describe habitat preference, 2. Estimation of the true, unobserved element level in the otolith, related to habitat occupancy, given a specific habitat preference and temporal correlation in the element concentration in addition to habitat preference, and 3. Measurement variability.

Layer 1: A discrete, unobserved Markov chain is used to describe habitat preference at any given time of the fish’s life, similar to Fablet et al. (2007). For the Markov chain, a predetermined number of habitats must be given. For this application, the model included four habitats representing the four biological populations and management areas in the region. The only other restriction on habitat areas was that strontium levels increased from habitat 1 to 2 to 3 to 4 (mirroring geographic trends in salinity levels from the eastern Baltic Sea to the North Sea – see Fig. S1), to ensure areas to be identifiable.

Layer 2: Given the habitat preference at any given time of the fish’s life, the true, unobserved log-strontium concentration in the otolith was modelled by a first order autoregressive process where the mean depends on the current habitat preference. The true unobserved strontium concentration is interpreted as the average concentration that would be observed for a fish at the same geographical position throughout its life. Consequently, the habitat preference in layer 1 determines where an individual is moving towards while the true, unobserved strontium concentration in layer 2 reflects habitat occupancy. With this interpretation, measurement noise

may be observed in measured strontium concentrations, even for an immobile fish, masking the signal from the true unobserved concentration. Finally, the autoregressive process in layer 2 models temporal correlation in the element levels besides the habitat type to reflect that fish may move between habitats at time scales that are longer than the time between measurements. *Layer 3:* The signal-to-noise ratio will depend on inherent background spectral interferences and sample volume determined by the laser beam diameter (e.g. Lear et al. 2012). Further, variability at small time scales may occur from variability in unmodelled factors such as temperature and food rations. Therefore, observed log-strontium was modelled by a normal distribution where the mean was the true log-strontium level.

Model parameters were estimated simultaneously for all individuals by maximum likelihood using an approximate filter (Albertsen 2018) implemented in “R” ver. r75965 (R Core Team 2019) using the package “TMB” (version 1.7.15; Kristensen et al. 2016). For the estimated parameters, most likely habitat preference and true strontium level transects were obtained through an approximate smoother (Albertsen 2018) for each individual. Subsequently, most likely true strontium (Sr) levels at the edge of the otoliths, corresponding to the final observation, were combined with salinity (S) levels at capture locations to estimate the calibration curve (Fig. S5):

$$\log(Sr) = 6.241 (\pm 0.099 \text{ se}) + 0.442 (\pm 0.030 \text{ se}) \cdot \log(S) \quad (1)$$

This calibration curve was derived from cod otolith samples with associated water salinity levels from all four habitat areas, but are not presented in this study, and through comparison with experimentally derived values from the literature (Albertsen et al. *accepted*).

While the state-space migration model estimates four levels of habitat-specific salinity values, it needs to be informed of the threshold salinities that characterize the geographic boundaries from one habitat area to its neighbour. Salinity cut-off values between habitat areas of interest

(North Sea, Kattegat, western Baltic Sea and eastern Baltic Sea) were determined based on annual mean salinities for the years 2013-2016 from ICES Hydrographic database (covering the entire lifespan of all fish in the study) (Available at <https://ices.dk/data/data-portals/Pages/ocean.aspx>) and known depth distribution ranges of cod: 30 – 80 m in the Skagerrak and Kattegat (Casini et al. 2005), 10 – 40 m in the western Baltic Sea and 30 – 60 m in the eastern Baltic Sea (Oeberst 2008). Cut-off salinities selected at the boundaries between areas were 10 psu between eastern and western Baltic Sea, 17.5 psu between western Baltic Sea and Kattegat, and 34 psu between Kattegat and Skagerrak/North Sea. An overview over geographic areas with associated salinity levels and data selected is given in Fig. S1. Using the strontium – salinity calibration curve, the strontium levels corresponding to these salinities was calculated.

In summary, with this approach each Sr measurement along the individual chronological otolith transects is translated into a corresponding salinity value and be assigned to one of the four habitat preference areas (example in Fig. 2). In order to simplify the visualisation of habitat preferences, preferences were grouped by time interval (spring, summer, fall, and winter of each year of life) and the most frequently occurring assignment chosen to represent a single habitat preference per individual and time interval.

Habitat assignment precision: One of the key features of the regime-switching state-space model used for assigning habitat association is a strontium – salinity calibration curve, described in model (1). Here, we use the assignments of individuals to the Skagerrak/North Sea for demonstration, since this is the area where habitat assignments are most uncertain as the area-specific differences in salinity are less pronounced than between other areas and the logarithmic relationship of the calibration curve at the same time is flattening. The confidence intervals of the salinity – strontium calibration curve at the salinity cut-off between the Kattegat and the Skagerrak/North Sea (34 psu) were used to estimate the proportion of individuals

assigned to the Skagerrak/North Sea. Habitat assignments were carried out for salinities estimated from the mean parameter values of model (1) as well as from mean $\pm$ standard error of the mean.

Results

Characterisation of the two populations: Genetics, otolith size and growth

Genetic population identification: Among the 306 fish analysed for population assignment, 141 assigned to the Kattegat while 165 assigned to the North Sea (Table 1). No fish assigned to the Eastern Baltic Sea population. The assignment scores were generally very high (average score of 97.97, range: 61.35-100.00), indicating high support for population assignment for the majority of the individuals. A few individuals had scores below 75 (4 from the North Sea and 4 from the Kattegat).

Otolith size: Otolith size at the end of the first year of life was significantly different between populations but did not differ between year classes within populations (ANOVA, $df = 10$ and 191 , $p_{\text{population}} < 0.05$, $p_{\text{age}} > 0.05$). Pairwise comparison of all groups showed that the population-related difference was caused by North Sea individuals of the age groups 1 and 2 having larger otoliths (Tukey's HSD, all $p < 0.05$), while there were no differences among all other groups (Tukey's HSD, all $p > 0.05$) (Fig. 3). Since otolith size and fish size are strongly correlated in cod (Li et al. 2008), this suggests that there were only limited differences in fish size at first winter between year classes and populations. For the analyses of elemental composition during the first three life stages (larval, pelagic and demersal juvenile) it is therefore not necessary to remove any size-related trend as this is not present.

Growth patterns: Growth patterns were compared by testing the linearity of the relationship between fish size and age using population as group covariate. Fish size was linearly related to the age of the fish (Fig. 4) but with significant differences in the intercept and slopes between

populations with a smaller intercept and higher slope in the North Sea population (ANCOVA, $F = 2944$, $df = 4$ and 296 , $p < 0.05$, $r^2 = 0.98$). A higher slope in the North Sea population indicates faster growth than the Kattegat population, and needs to be considered in the discussion of growth-related differences in element concentration.

Spawning origin

Year class effect on chemical fingerprint: For both the North Sea and the Kattegat population, significant differences in chemical fingerprint during the pelagic juvenile stage were detected between the 8 year classes (North Sea: MANOVA, $df = 7$ on 292 , $p < 0.05$; Kattegat: MANOVA, $df = 7$ on 130 , $p < 0.05$). For both populations, these differences between year classes were attributable to Mg, Mn, Cu, and Zn (ANOVA, $df = 1$, $p < 0.05$ for all). No significant differences were found for P, Sr and Ba (ANOVA, $df = 1$, $p > 0.05$ for all). Even though there were significant differences between year classes within each population, these were restricted to elements that are under physiological control (Campana 1999; Hüseyin et al. 2020), and not in Sr and Ba, which are the element of key interest for inferring migration history. Year class effects were therefore not considered relevant in the subsequent analyses.

Pelagic juvenile stage: The chemical fingerprints in the pelagic juvenile stage differed significantly between the two populations (MANOVA, $df = 7$ on 292 , $p < 0.05$). Significant differences occurred in all elements except for P (ANOVA, $df = 1$, $p < 0.05$ for all except P). Of particular interest are Sr, Ba and Mn, where Sr was higher in the North Sea population, while Ba and Mn were higher in the Kattegat population (Table 2). Differences between year-classes within the two populations were not significant (MANOVA, $df = 7$ on 292 , $p > 0.05$). The LDA showed that 83 % of the North Sea population and 68 % of Kattegat population were correctly classified to their respective population with an overall classification success of 76 %

based on the first four discriminant functions. The biplot of the first two discriminant functions against each other with loadings (representing individual elements) showed that in particular Sr, Ba and Mn drive population-specific fingerprints in the pelagic juvenile stage (Fig. 5). In this plot, LD1 explains 41.4 % of the variation in chemical fingerprint between populations and LD2 21.1 %. A considerable part of the variation (37.5 %) is thus explained by dimensions that are orthogonal to the first two dimensions, and thereby difficult to represent visually. Together, the MANOVA and LDA results of the chemical fingerprint in the pelagic juvenile stage show that cod of different populations captured at the same locations in the Kattegat have different spawning origins.

Juvenile demersal life stages: In order to address during which life stage (pelagic juvenile, demersal juvenile, first winter, first, second etc. year of life) the North Sea cod enter Kattegat, the ANOVA, MANOVA and LDA analyses were repeated for each of the progressively older life stages. The results for the demersal juvenile stage mirrored those for the pelagic stage, with significant differences between populations (MANOVA, $df = 7$ on 292, $p < 0.05$), where differences were primarily driven by Sr, Ba and Mn (Fig. 5, Table S2). There were no statistical differences in concentration of any elements during the subsequent first winter (MANOVA, $df = 7$ on 292, $p > 0.05$) (Fig. 5, Table S3). The classification success decreased correspondingly from 83 % to 72 % and 50 % in the North Sea population and from 68 % to 62 % and 51 % in the Kattegat population (total classification success decreasing from 76 % to 67 % and 51 %), which is illustrated by the increasing overlap between groups in the LDA biplot (Fig. 5). From age 2 onward, differences between populations were no longer significant (MANOVA, $df = 7$ on 292, $p > 0.05$).

Adult migrations

The comparison of otolith Sr concentration between consecutive seasons and ages found significant differences overall (ANOVA, $df = 19$, $p < 0.05$), no significant differences were evident between any of the consecutive seasons and ages in the pairwise comparisons (Tukey HSD, all $p > 0.05$). This shows that there is no consistent seasonal effect on otolith Sr concentration. It was therefore assumed, that any change in otolith Sr was associated with a change in habitat occupation and not seasonally varying temperature or growth. From the regime-switching state-space migration model habitat area assignments (Fig. 1) were obtained for all time periods of each individual's life. Examination of individual cod's habitat area occupation patterns of the North Sea population (Fig. 6a) and the Kattegat population (Fig. 6b) showed that the majority of individuals from both populations showed migrations in and out of the Kattegat, with 25 % ($n = 21$) of the North Sea individuals and only 8 % ($n = 9$) of the Kattegat individuals remaining resident within the Kattegat from the first winter and until capture (Fig. 6, green). The majority of individuals from both populations showed evidence of migration to other areas at some point between the first winter and time of capture. By far the most frequent migration signature was alternating distribution in the Kattegat and western Baltic Sea (55 % ($n=47$) of North Sea cod; 68 % ($n = 78$) of Kattegat cod) (Fig. 6, blue green), while a proportion of fish from both populations also displayed signatures consistent with migration to the Skagerrak/North Sea and back to the Kattegat (31 % ($n=27$) of North Sea cod; 45 % ($n = 52$) of Kattegat cod) (Fig. 6, lime). Ten individuals (8.5 %) from the Kattegat population appeared to have visited the eastern Baltic Sea for shorter periods of time (Fig. 6, purple). In general, the migration patterns were not consistent between seasons, suggesting the absence of synchronized migrations between areas.

While the habitat area assignments of individuals (Fig. 6) provide insight into individual fish's movements between areas, they do not provide an easily interpretable picture of how much

time the two populations on average spend in the different areas. To get an estimate of how many individuals had been assigned to the four habitats at any given fish age and time of the year, percentages of habitat area assignments were calculated across all individuals for each age and season separately. The resulting percentages thus represent a population's distribution pattern in space (North Sea/Skagerrak, Kattegat, western and eastern Baltic Sea) and time (age and season). The most notable observation is that habitat distribution patterns are remarkably similar in the two populations (Fig. 7). Across all four seasons and all four age classes, the two populations were both most frequently distributed in the Kattegat, with 66.8% (range = 44 - 81 %) of North Sea population and 67.1 % (range = 53 - 79 %) of the Kattegat population (Fig. 8, green bars), making the Kattegat the primary habitat of distribution. Across seasons and age classes, approximately 25 % of both population was distributed in the western Baltic Sea (North Sea: mean = 24.9, range = 0 - 50 %; Kattegat: mean = 23.5, range = 8 - 37 %) (Fig. 7, blue green bars). Less than 10 % of both populations were distributed within the Skagerrak/North Sea (North Sea: mean = 8.3, range = 5 - 12 %; Kattegat: mean = 7.9, range = 3 - 13 %) across all seasons for the age classes 1 - 3 years (Fig. 7, lime bars). An apparently higher degree of migrations between the Skagerrak/North Sea and back to the Kattegat seemed to occur during age class 4 in the North Sea population, in that increasing percentages (mean = 26.6, range = 13 - 50 %) were distributed within the Skagerrak/North Sea. However, it is important to note that the sample sizes are relatively low for these age groups and inferred distribution patterns therefore somewhat uncertain. Surprisingly, in the Kattegat population 1.4 ± 1.3 % of individuals were assigned to the eastern Baltic Sea for restricted periods of time around fall and winter (Fig. 7, purple bars). Migration patterns of fish older than four years could not be analysed for the North Sea population, as no samples of this population were available.

Habitat assignment precision: Correct habitat area assignment relies heavily on the strontium – salinity calibration curve used as well as the salinity cut-off values used for defining the boundary between areas. The impact of salinity cut-off values selected for defining the boundary between areas on estimated habitat assignments was evaluated with a simple sensitivity analysis. Since this selection is more related to the study area, but is less relevant for the methodological approach as such, these result are not presented here, but may be found in the Supplementary information (Fig. S6) together with details on the hydrography in the study area (Fig. S1). The impact of uncertainty in the strontium – salinity calibration curve on the other hand is a generic issue. Here, we demonstrate the impact thereof on the number of individuals assigned to the Skagerrak/North Sea, because the environmental salinity at the boundary between this area and the Kattegat is at the upper end of the salinity range, where the calibration curve flattens. This exercise shows, that the uncertainty in the calibration curve on proportion of individuals assigned to the Skagerrak/North Sea results in an average deviation of 3.4% for both populations (North Sea: range 0 to 12.5%; Kattegat: range 0.9 to 6.5%) between the highest (mean - se) and lowest (mean + se) estimates (Fig. 8).

Discussion

Spawning origin: In this study, we found population-specific chemical fingerprints in the early life stages of cod in Kattegat. The elements driving this differentiation were Sr, Ba and Mn. These elements are under strong environmental control. Strontium has for over two decades been known to reflect ambient salinity, owing to the fact that the Sr content in marine habitats is fairly constant worldwide, and mixing with freshwater not only dilutes salinity but also the Sr concentration (Kraus and Secor 2004; Walther and Limburg 2012). Because otolith Sr concentration is strongly correlated with ambient concentrations, with correlations generally > 0.95 found throughout the literature (review of studies in Hüseyin et al. (2020)), otolith Sr is a

490 useful proxy for environmental salinity and is globally used to track movements of fish between
491 marine and freshwater and within estuaries (Bath et al. 2000; Elsdon and Gillanders 2003;
492 Miller 2011; Sturrock et al. 2012). While several studies report a significant impact of
493 temperature on otolith Sr, the results are largely conflicting, ranging from a positive influence
494 to a negative (Bath et al. 2000) – or no influence at all (Clarke and Friedland 2004; Walther et
495 al. 2010; Sturrock et al. 2015), explaining at best a few percent of the variation in otolith Sr
496 (Hüssy et al. 2020). Also food (Walther et al. 2010) and other physiological factors (Sturrock
497 et al. 2015) have been suggested to regulate otolith Sr concentrations. However, similar to other
498 studies on Atlantic cod from the North Sea to the Baltic Sea (Hüssy et al. 2016; Heimbrand et
499 al. 2020; Hüssy et al. 2021), the samples used in the present study showed no apparent
500 seasonality in otolith Sr patterns, suggesting that changes in otolith Sr are associated with a
501 change in habitat occupation rather than seasonally varying temperature or growth.

502 The higher Sr concentrations in the North Sea population are thus consistent with a spawning
503 area in more saline water than the Kattegat. Barium shows a nutrient-like distribution in the
504 aquatic environment that is strongly related to environmental salinity with depletion in surface
505 waters, with higher concentrations in freshwater and nearshore areas (Elsdon and Gillanders
506 2005; Walther and Limburg 2012). Highest Ba concentrations generally occur at salinities
507 between 5 and 20 psu (Walther and Limburg 2012). The Ba concentration in otoliths almost
508 exclusively reflects ambient concentrations (Bath et al. 2000; Elsdon and Gillanders 2003;
509 Hicks et al. 2010; Miller 2011; Reis-Santos et al. 2013). Consistent with the hypothesis that the
510 North Sea population is spawned in a more offshore environment, Ba concentrations were
511 higher in the Kattegat population. Manganese concentrations on the other hand are known to
512 increase in hypoxic areas owing to the reduction of manganese oxides from the sediment with
513 decreasing ambient oxygen content. Otolith Mn concentrations have proven useful for tracking
514 hypoxia exposure in e.g. Baltic cod (Limburg et al. 2011, 2015). Prolonged seasonal hypoxia

is known to occur in the Kattegat (Rosenberg et al. 1992, 1996), which is reflected in the much higher Mn concentration during the early life stages of the Kattegat population. Significant differences in particularly Mg, P, Cu and Zn were also detected between a few year classes within the two populations. These elements are under strong physiological control (Campana 1999; Sturrock et al. 2015; Limburg et al. 2018; Hüsey et al. 2020) and have been shown to reflect fish growth (Heimbrand et al. 2020; Hüsey et al. 2021). Otolith element concentrations of these elements were consistently higher in the North Sea populations. This corresponds with the larger size at age observed between populations and suggests a genetic component in growth regulation, and a mechanism promoting early settlement of North Sea cod. No differences between year classes in all other elements shows that the classification to population based on Sr, Ba and Mn is not influenced by inter-annual growth rate differences. These results provide consistent and biologically meaningful evidence for the hypothesis that the North Sea population was spawned in a more marine and offshore environment compared to the Kattegat population, presumably the Skagerrak or northern North Sea.

Early life stage drift: The rapidly decreasing classification success from the pelagic juvenile stage to the beginning of the first winter and the lack of significant differences in environmentally related elements (Sr, Ba, Mn) in subsequent years, suggests that the North Sea fish arrived in the Kattegat during the early life stages. Kattegat cod spawn within a restricted area of the Kattegat (Vitale et al. 2008; Börjesson et al. 2013), and early life stages of this population have in recent years largely been retained in the area (Jonsson et al. 2016). North Sea cod spawn along the southern and eastern edges of the Dogger Bank, in the German Bight, the Moray Firth and to the east of the Shetlands (Brander 1994; Fox et al. 2008; Munk et al. 2009). Drift simulations have shown that cod eggs, larvae and early juveniles spawned in the North Sea may be advected for more than 600 km during their drift period (Eero et al. 2016).

Particularly individuals spawned in the German Bight may frequently be transported into the Skagerrak (Eero et al. 2016). The early lifestages are entrained by the Jutland current running north along the west coast of Denmark into the eastern part of Skagerrak (Svansson 1975; Dyrssen 1993; Jakobsen 1997). During westerly winds, strong currents prevail from the Skagerrak towards the Baltic Sea along the coast of Sweden. An inflow of North Sea cod into the Kattegat during their early life stages is thus highly likely, where drift durations from the North Sea to the Kattegat may range from days to months (Eero et al. 2016). This inter-annual and inter-individual variability in drift duration explains the lack of a stronger separation of individuals in the pelagic juvenile stage and the gradually decreasing classification success between the two populations with age. Genetic studies of juvenile cod from the fjords from the Norwegian Skagerrak coast to the central Kattegat support this, in that a large proportion can be genetically similar to cod from offshore spawning areas in the eastern North Sea, with substantial temporal variation in advection (Knutsen et al. 2004; Stenseth et al. 2006; André et al. 2016). Owing to the low stock size of Kattegat cod and the influx of individuals from other areas, Jonsson et al. (2016) estimated that currently only approximately 34 % of the juveniles in Kattegat are from locally retained spawning areas compared to 83 % in the 1970s when the stock was larger.

Migrations between Kattegat and Skagerrak/North Sea: The present study showed that even though the majority of individuals undertook predominantly short-term migrations to other areas, they were most frequently distributed within the Kattegat, regardless of their population of origin. These results suggest the absence of strong synchronised migrations between areas. Only a few individuals (< 10 %) were distributed within the Skagerrak/North Sea with subsequent return to the Kattegat. The absence of North Sea cod individuals older than four years old suggests that the North Sea cod leave the Kattegat, presumably returning to the North

Sea once mature in support of natal homing as suggested by Svedäng et al. (2007). From our data it is not possible to deduce whether the migratory cod did in fact spawn during their stay in the Skagerrak/North Sea.

The habitat assignments largely depend on the salinity cut-off value selected as defining the boundary between areas and the accuracy of the strontium – salinity calibration curve. Since both salinity and strontium are subject to measurement errors, the calibration curve is subject to some uncertainty, in particular at the Kattegat/Skagerrak boundary, where area-specific differences in salinity are less pronounced than between other areas and the logarithmic relationship of the calibration curve at the same time is flattening. Although a rigorous cross-validation would be needed to estimate the power of our method, we did estimate effects from calibration curve uncertainty and different cut-off salinity values used for categorizing areas and found only minor variations in these proportions, suggesting the presented approach to be relatively robust. Consequently, our results suggest some migration between the Kattegat and Skagerrak/North Sea. We also found some evidence for inter-individual variability in migratory behavior with some fish displaying a more stationary signature while other fish seemed to display signatures in agreement with a pattern of recurrent migrations and longer time occupancy in the North Sea.

Information of juvenile and adult cod migrations is available from tagging experiments from the 1950s, 1990s and 2000s and have documented migration patterns similar to the ones inferred from the present study. These studies found that cod tagged as 1-year old juveniles in Kattegat only undertook limited migrations until the age of 2 years and 30 - 50 cm in length, when they moved offshore towards the south and west – towards the spawning grounds in the southern Kattegat and the eastern North Sea (Pihl and Ulmestrand 1993). Adult cod tagged at offshore and in coastal locations in the Skagerrak undertook long-distance migrations ranging from the southern North Sea to the central Kattegat (Danielssen 1969). Migrations of adult cod

tagged along the Swedish Kattegat coast were directed both south and north towards the Sound and the eastern North Sea, with some coherence between tagging location and migration direction (Righton et al. 2010; Svedäng et al. 2010). Of the cod tagged in the northern Kattegat, 12 % were recaptured in the Skagerrak or eastern North Sea, while cod tagged in the Sound and the southern/central Kattegat were primarily resident (Svedäng et al. 2010), suggesting migration patterns quite similar to the distribution patterns observed in the present study. Geolocation of individuals based on electronic tags (DST's) documented that the migrations of cod tagged in the northern Kattegat and Skagerrak are directional towards the North Sea and coincide with known spawning times (Svedäng et al. 2007). Subsequent returns to the areas where the cod were tagged also occurred, but quantification thereof was not possible (Svedäng et al. 2007). Linking genetics with tagging-based migration patterns, André et al. (2016) suggested a strong correspondence between population and philopatric migrations towards natal spawning grounds in the Skagerrak/Kattegat area. One may argue that these tagging studies represent migration patterns at a time when the cod stocks were not yet depleted and may therefore not apply to today's scenarios. However, given the similar conclusions from widely different analytical approaches and a large range of years, it seems that the migratory exchange between Kattegat and Skagerrak/North Sea is – and has always been - limited in immature cod, but extensive in mature cod and most likely linked to spawning migrations.

Migration between Kattegat and western Baltic Sea: This study also found that a large proportion (approximately 25 %) of individuals from both populations were distributed within the western Baltic Sea during all seasons and across all age classes. The salinity regimes in the western Baltic and Kattegat differ considerably from each other with mean bottom salinities of 10 – 15 psu and 30 - 33 psu respectively. This salinity gradient lies within the ideal range for discriminating habitat use (Kraus and Secor 2004; Walther and Limburg 2012), and the results

are therefore considered reliable. The migration patterns of tagged cod discussed above do not suggest a pronounced migration further south than the Sound. However, the dynamics in stock size have changed considerably over the last decades, with unknown impact on migratory behaviour. Considerable numbers of eggs and larvae spawned in the western Baltic Sea, more specifically the Great Belt and the Kiel Bay, may drift into the Kattegat and Skagerrak depending on wind conditions prevailing during the spawning season (Huwert et al. 2016). Provided that these individuals also show philopatric migration behaviour, one would expect to find seasons with high assignments to the western Baltic Sea. A clear seasonal pattern with higher proportions of western Baltic Sea signals in spring (the main spawning season of western Baltic cod) was, however, not evident. The occurrence of 2 % cod with a clear western Baltic/Sound otolith chemistry signal over most of their lives prior to being captured in the Kattegat further suggests considerable connectivity between these areas. Cod spawning in the Kattegat and western Baltic Sea likely belong to the same biological population or population-complex which has been found to display a hybrid zone genetic signature between the clearly differentiated North Sea and Baltic Sea populations (Nielsen et al. 2003). Consequently, some migration between areas is also well aligned with the overall patterns of genetic similarity observed in the geographical region in the populations. Further work will be needed to investigate potential patterns of minor genetic sub-structuring of spawning components within the Kattegat and western Baltic Sea. However, our results clearly indicate that current assumptions of closed units used for stock assessments of cod in the North Sea, Kattegat and the western Baltic Sea are likely violated by the dynamic patterns of exchange between areas.

Evaluation of the interdisciplinary approach: With this study we have demonstrated that an interdisciplinary approach combining information from genetics and otolith microchemistry coupled with advanced modelling tools may provide unprecedented new insights into complex

biological questions. Genotyping to identify population of origin and chemical fingerprinting of the otolith core to identify hatch area are well established methodologies. Combined, however, they provide a powerful tool to study connectivity in area use of early life stages. The most innovative aspect of this study is the demonstration of how time-series models of chronological microchemistry data from hatch to death can be combined with genetic information and used to infer migration patterns of individual fish as well as stock mixing proportions at different life stages and ages. While we used Atlantic cod in the geographic area between the North Sea and the Baltic Sea as a case study here, the approach is generally applicable where environmental gradients in water chemistry are present and sampling designs cover all areas/environments through which the fish can be expected to move. The approach outlined here therefore has wider applicability in both basic and applied research in fish and fisheries biology and could provide valuable data as direct input for conservation and management of fish stocks.

Acknowledgements

Thanks go to all technical staff of DTU Aqua and GEUS involved in the collection and processing of samples used in this study. Additional thanks to Kristian Ege Nielsen for sample preparation and testing of setup for the chemical analyses of samples. We thank Benjamin Heredia Dominguez, GEUS, for providing the figure on LA ICP-MS data accuracy and precision. We would also like to thank two anonymous reviewers for their valuable feedback and input on an earlier version of this paper.

Competing interests

The authors declare there are no competing interests.

Contributors' statement

KH: Conceptualization, Formal analysis, Methodology, Writing - Original Draft, Funding acquisition

CMA: Conceptualization, Formal analysis, Methodology, Writing - Review & Editing

JHH: Conceptualization, Methodology, Writing - Review & Editing, Investigation, Data curation, Funding acquisition, Project administration

MV: Conceptualization, Writing - Review & Editing

SHS: Methodology, Data curation, Writing - Review & Editing

TBT: Methodology, Data curation, Writing - Review & Editing

ME: Conceptualization, Writing - Review & Editing

Funding statement

The study was supported by the Danish Ministry for Environment and Food and the European Maritime Fisheries Fund through the projects *Sustainable management of Kattegat cod; better knowledge of stock components and migration* [grant number 33113-B-16-034] and *Management of mixed cod stocks in the transition zone between the North Sea and the Baltic Sea: How can this be achieved most efficiently?* [grant number 33113-B-19-140].

Data availability statement

Access to the data upon which this study is based may be obtained by contacting the corresponding author.

References

Albertsen, C.M. 2018. An approximate filter and smoother for general regime-switching state-space models *In* State-space modelling in marine science. PhD thesis, National Institute

- of Aquatic Resources, Technical University of Denmark, Kgs. Lyngby, Denmark.
- Albertsen, C.M, Hüsey, K., Serre, S.H., Hemmer Hansen, J., and Thomsen, T.B. (xxxx) Estimating migration patterns of fish from otolith chemical composition time-series, submitted to Can. J. Fish. Aquat. Sci. (*accepted*).
- Albertsen, C.M., Nielsen, A., and Thygesen, U.H. 2018. Connecting single-stock assessment models through correlated survival. ICES J. Mar. Sci. **75**(1): 235-244. doi: 10.1093/icesjms/fsx114.
- André, C., Svedäng, H., Knutsen, H., Dahle, G., Jonsson, P., Ring, A.-K., Sköld, M., and Jorde, P.E. 2016. Population structure in Atlantic cod in the eastern North Sea-Skagerrak-Kattegat: early life stage dispersal and adult migration. BMC Res. Notes **9**(1): 63. doi:10.1186/s13104-016-1878-9.
- Barth, J.M.I., Berg, P.R., Jonsson, P., Bonanomi, S., Corell, H., Hemmer-Hansen, J., Jakobsen, K.S., Johannesson, K., Jorde, P.E., Knutsen, H., Moksnes, P.O., Star, B., Stenseth, N.C., Svedäng, H., Jentoft, S., and André, C. 2018. Genome architecture enables local adaptation of Atlantic cod despite high connectivity. Mol. Ecol. **26**: 4452-4466. doi: 10.1111/mec.14207.
- Barth, J.M., Villegas-Ríos, D., Freitas, C., Mpland, E., Star, B., André, C., Knutsen, H., Bradbury, I., Dierking, J., Petereit, C., Righton, C., Metcalfe, J., Jakobsen, K.S., Olsen, E.M., and Jentoft, S. 2019. Disentangling structural genomic and behavioural barriers in a sea of connectivity. Mol. Ecol. **28**: 1394-1411. doi: 10.1111/mec.15010.
- Bath, G.E., Thorrold, S.R., Jones, C.M., Campana, S.E., McLaren, J.W., and Lam, J.W. 2000. Strontium and barium uptake in aragonitic otoliths of marine fish. Geochim. Cosmochim. Acta **64**(10): 1705–1714. doi:10.1016/S0016-7037(99)00419-6.
- Beckman, D., and Wilson, C.A. 1995. Seasonal timing of opaque zone formation in fish otoliths. In Recent Developments in Fish Otolith Research. Edited by D.H. Secor, J.M.

- 715 Dean, and S.E. Campana. University of South Carolina Press, Columbia, SC. pp. 27–44.
- 716 Berg, P.R., Jentoft, S., Star, B., Ring, K.H., Knutsen, H., Lien, S., Jakobsen, K.S., and André,
717 C. 2015. Adaptation to Low Salinity Promotes Genomic Divergence in Atlantic Cod
718 (*Gadus morhua* L.). *Gen. Biol. Evol.* **7**: 1644–1663. doi:10.1093/gbe/evv093.
- 719 Becker, R.A., and Wilks A.R. 1993. Maps in S, AT&T Bell Laboratories Statistics Research
720 Report [93.2]. Available at: <http://ect.bell-labs.com/sl/doc/93.2.ps>
- 721 Börjesson, P., Jonsson, P., Pacariz, S., Björk, G., Taylor, M.I., and Svedäng, H. 2013.
722 Spawning of Kattegat cod (*Gadus morhua*)—Mapping spatial distribution by egg surveys.
723 *Fish. Res.* **147**: 63–71. doi:10.1016/J.FISHRES.2013.04.012.
- 724 Brander, K.M. 1994. The location and timing of cod spawning around the British Isles. *ICES*
725 *J. Mar. Sci.* **51**: 71–89. doi: 10.1006/jmsc.1994.1007.
- 726 Cadrin, S.X., and Secor, D.H. 2009. Accounting for Spatial Population Structure in Stock
727 Assessment: Past, Present, and Future. *In* *The Future of Fisheries Science in North*
728 *America*. Springer Netherlands, Dordrecht. pp. 405–426. doi:10.1007/978-1-4020-9210-
729 7_22.
- 730 Campana, S.E. 1999. Chemistry and composition of fish otoliths: pathways, mechanisms and
731 applications. *Mar. Ecol. Prog. Ser.* **188**: 263–297. doi: 10.3354/meps188263.
- 732 Campana, S.E., and Thorrold, S.R. 2001. Otoliths, increments, and elements: keys to a
733 comprehensive understanding of fish populations? *Can. J. Fish. Aquat. Sci.* **58**(1): 30–38.
734 doi:10.1139/f00-177.
- 735 Carlson, A.K., Phelps, Q.E., and Graeb, B.D.S. 2017. Chemistry to conservation: using otoliths
736 to advance recreational and commercial fisheries management. *J. Fish Biol.* **90**(2): 505–
737 527. doi:10.1111/jfb.13155.
- 738 Casini, M., Cardinale, M., Hjelm, J., and Vitale, F. 2005. Trends in cpue and related changes in
739 spatial distribution of demersal fish species in the Kattegat and Skagerrak, eastern North

- 740 Sea, between 1981 and 2003. ICES J. Mar. Sci. **62**:671e682.
- 741 Clarke, L.M., and Friedland, K.D. 2004. Influence of growth and temperature on strontium
742 deposition in the otoliths of Atlantic salmon. J Fish Biol. **65**(3):744–759.
743 doi:10.1111/j.0022-1112.2004.00480.x.
- 744 Danielssen, D. 1969. On the migrations of the cod in the Skagerrak shown by tagging
745 experiments in the period 1954-1965. Fisk. Skr. Ser. Havundersøkelser **15**: 331–338.
746 Available from
747 [https://brage.bibsys.no/xmlui/bitstream/handle/11250/114468/sh_vol15_03\(21\)_1969.pdf](https://brage.bibsys.no/xmlui/bitstream/handle/11250/114468/sh_vol15_03(21)_1969.pdf?sequence=1)
748 [f?sequence=1](https://brage.bibsys.no/xmlui/bitstream/handle/11250/114468/sh_vol15_03(21)_1969.pdf?sequence=1) [accessed 25 March 2019].
- 749 Dyrssen, D. 1993. The Baltic-Kattegat-Skagerrak Estuarine System. Estuaries **16**(3): 446.
750 doi:10.2307/1352592.
- 751 Eero, M., Hemmer-Hansen, J., Hüsey, K., Huwer, B., Berg, C., Mariani, P., Mosegaard, H.,
752 Nielsen, A., Nielsen, E.E., Rindorf, A., Ulrich, C., Vinther, M., and Worsøe Clausen, C.
753 2016. Optimal bæredygtig udnyttelse af tilgængelige torskebestande for dansk fiskeri.
754 DTU Aqua scientific report (EMFF J.nr. 33010-13-k-0269). Charlottenlund, Denmark.
- 755 Elsdon, T., and Gillanders, B. 2003. Relationship between water and otolith elemental
756 concentrations in juvenile black bream *Acanthopagrus butcheri*. Mar. Ecol. Prog. Ser.
757 **260**: 263–272. doi:10.3354/meps260263.
- 758 Elsdon, T.S., and Gillanders, B.M. 2005. Alternative life-history patterns of estuarine fish:
759 barium in otoliths elucidates freshwater residency. Can. J. Fish. Aquat. Sci. **62**(5): 1143–
760 1152. doi:10.1139/f05-029.
- 761 Elsdon, T.S., Wells, B.K., Campana, S.E., Gillanders, B.M., Jones, C.M., Limburg, K.E.,
762 Secor, D.H., Thorrold, S.R., and Walther, B.D. 2008. Otolith chemistry to describe
763 movements and life-history parameters of fishes—hypotheses, assumptions, limitations
764 and inferences. In Oceanography and Marine Biology: An Annual Review. Edited by R.N.

- Gibson, A.R. J, and J.D.M. Gordon. CRC Press, Boca Raton, London, New York. pp. 297–330.
- Estoup, A., Largiadere, C.R., Perrot, E., and Chourrout, D. 1996. Rapid one tube DNA extraction for reliable PCR detection of fish polymorphic markers and transgenes. *Mol. Mar. Biol. Biotech.* **5**: 295–298.
- Everitt, B., and Hothorn, T. 2011. *An Introduction to Applied Multivariate Analysis with R*. Springer New York, New York, NY. doi:10.1007/978-1-4419-9650-3.
- Fablet, R., Daverat, F., and De Pontual, H. (2007). Unsupervised bayesian reconstruction of individual life histories from otolith signatures: case study of sr:ca transects of european eel (*anguilla anguilla*) otoliths. *Can. J. Fish. Aquat. Sci.* **64**(1): 152–165. doi: 10.1139/f06-173.
- Fox, C.J., Taylor, M., Dickey-Collas, M., Fossum, P., Kraus, G., Rohlf, N., Munk, P., van Damme, C.J.G., Bolle, L.J., Maxwell, D.L., and Wright, P.J. 2008. Mapping the spawning grounds of North Sea cod (*Gadus morhua*) by direct and indirect means. *Proc. R. Soc. B* **275**(1642): 1543–1548. doi:10.1098/rspb.2008.0201.
- Goethel, D.R., Quinn, T.J., and Cadrin, S.X. 2011. Incorporating Spatial Structure in Stock Assessment: Movement Modeling in Marine Fish Population Dynamics. *Rev. Fish. Sci.* **19**(2): 119–136. doi:10.1080/10641262.2011.557451.
- Heath, M.R., Culling, M.A., Crozier, W.W., Fox, C.J., Gurney, W.S.C., Hutchinson, W.F., Nielsen, E.E., O'Sullivan, M., Preedy, K.F., Righton, D.A., Speirs, D.C., Taylor, M.I., Wright, P.J., and Carvalho, G.R. 2014. Combination of genetics and spatial modelling highlights the sensitivity of cod (*Gadus morhua*) population diversity in the North Sea to distributions of fishing. *ICES J. Mar. Sci.* **71**: 794-807. doi: 10.1093/icesjms/fst185.
- Heidemann, F., Marohn, L., Hinrichsen, H.H., Huwer, B., Hüsey, K., Klügel, A., Böttcher, U., and Hanel, R. 2012. Suitability of otolith microchemistry for stock separation of Baltic

- 790 cod. Mar. Ecol. Prog. Ser. **465**: 217-226. doi:10.3354/meps09922.
- 791 Heimbrand, Y., Limburg, K., Hüsey, K., Casini, M., Sjöberg, R., Palmén Bratt, A.-M.,
792 Levinsky, S.-E., Karpushevskaya, A., Radtke, K., and Öhlund, J. 2020. Seeking the true
793 time: Exploring otolith chemistry as an age-determination tool. J. Fish Biol.
794 doi:10.1111/jfb.14422.
- 795 Hemmer-Hansen, J., Hüsey, K., Vinther, M., Albertsen, C. M., Storr-Paulsen, M., and Eero, M.
796 2020. Sustainable management of Kattegat cod; better knowledge of stock components
797 and migration. DTU Aqua Report no. 357-2020. National Institute of Aquatic Resources,
798 Technical University of Denmark. 42 pp. ISBN: 978-87-7481-281-4.
- 799 Hicks, A.S., Closs, G.P., and Swearer, S.E. 2010. Otolith microchemistry of two amphidromous
800 galaxiids across an experimental salinity gradient: A multi-element approach for tracking
801 diadromous migrations. J. Exp. Mar. Bio. Ecol. **394**(1): 86–97. doi:
802 10.1016/j.jembe.2010.07.018.
- 803 Høie, H., and Folkvord, A. 2006. Estimating the timing of growth rings in Atlantic cod otoliths
804 using stable oxygen isotopes. J. Fish Biol. **68**(3): 826–837. doi:10.1111/j.0022-
805 1112.2006.00957.x.
- 806 Huwer, B., Hinrichsen, H.-H., Hüsey, K., and Eero, M. 2016. Connectivity of larval cod in the
807 transition area between North Sea and Baltic Sea and potential implications for fisheries
808 management. ICES J. Mar. Sci. **73**(7): 1815-1824. doi:10.1093/icesjms/fsw043.
- 809 Hüsey, K. 2011. Review of western Baltic cod (*Gadus morhua*) recruitment dynamics. ICES
810 J. Mar. Sci., 68: 1459–1471. doi:10.1093/icesjms/fsr088.
- 811 Hüsey, K., Gröger, J., Heidemann, F., Hinrichsen, H.-H., & Marohn, L. (2016). Slave to the
812 rhythm: Seasonal signals in otolith microchemistry reveal age of eastern Baltic cod
813 (*Gadus morhua*). ICES J. Mar. Sci. **73**(4): 1019–1032. doi:10.1093/icesjms/fsv247.
- 814 Hüsey, K., Krüger-Johnsen, M., Thomsen, T.B., Heredia, B.D., Naeraa, T., Limburg, K.E.,

- 815 McQueen, K., Haase, S., Krumme, U., Casini, M., Mion, M., Radtker, K. (2021) It's
816 elemental, my dear Watson: Validating seasonal patterns in otolith chemical
817 chronologies. *Can. J. Fish. Aquat. Sci.* doi: 10.1139/cjfas-2020-0388.
- 818 Hüsey, Karin, Limburg, K.E., de Pontual, H., Thomas, O.R.B., Cook, P.K., Heimbrand, Y.,
819 Blass, M., and Sturrock, A.M. 2020. Trace Element Patterns in Otoliths: The Role of
820 Biomineralization. *Rev. Fish. Sci. Aquacult.* 1–33. doi:10.1080/23308249.2020.1760204.
- 821 ICES. 2015. Report of the Benchmark Workshop on Baltic Cod Stocks (WKBALTCOD).
822 ICES C. 2015/ACOM35 (2–6 March 2015, Rostock, Germany): pp 172.
- 823 ICES. 2019. Baltic Fisheries Assessment Working Group (WGBFAS). *ICES Sci. Rep.* **1:20**:
824 pp 653. doi:10.17895/ices.pub.5256.
- 825 ICES. 2020. Cod (*Gadus morhua*) in Subdivision 21 (Kattegat). *In* Report of the ICES
826 Advisory Committee, 2020. ICES Advice 2020, cod.27.21.
827 <https://doi.org/10.17895/ices.advice.5903>.
- 828 Jakobsen, F. 1997. Hydrographic investigation of the Northern Kattegat front. *Cont. Shelf Res.*
829 **17(5)**: 533–554. doi:10.1016/S0278-4343(96)00044-1.
- 830 Jonsson, P.R., Corell, H., André, C., Svedäng, H., and Moksnes, P.-O. 2016. Recent decline in
831 cod stocks in the North Sea-Skagerrak-Kattegat shifts the sources of larval supply. *Fish.*
832 *Oceanogr.* **25(3)**: 210–228. doi:10.1111/fog.12146.
- 833 Kerr, L., Hintzen, N., Cadrin, S., Worsøe Clausen, L., Dickey-Collas, M., Goethel, D.R.,
834 Hatfield, E.M.C., Kritzer, J., and Nash, R.D.M. 2017. Lessons learned from practical
835 approaches to reconcile mismatches between biological population structure and stock
836 units of marine fish. *ICES J. Mar. Sci.* **74(6)**: 1708–1722. doi:10.1093/icesjms/fsw188.
- 837 Knutsen, H., André, C., Jorde, P.E., Skogen, M.D., Thuróczy, E., and Stenseth, N.C. 2004.
838 Transport of North Sea cod larvae into the Skagerrak coastal populations. *Proc. R. Soc.*
839 *London. Ser. B Biol. Sci.* **271(1546)**: 1337–1344. doi:10.1098/rspb.2004.2721.

- 840 Kraus, R.T., and Secor, D.H. 2004. Incorporation of strontium into otoliths of an estuarine fish.
841 J. Exp. Mar. Bio. Ecol. **302**(1): 85–106. doi:10.1016/J.JEMBE.2003.10.004.
- 842 Kristensen, K., Nielsen, A., Berg, C. W., Skaug, H., Bell, B. M. (2016). TMB: Automatic
843 Differentiation and Laplace Approximation. Journal of Statistical Software, 70(5), 1-21.
844 doi:10.18637/jss.v070.i05.
- 845 Lear, J., Hare, D. J., Fryer, F., Adlard, P. A., Finkelstein, D. I., Doble, P. A. (2012). High-
846 Resolution Elemental Bioimaging of Ca, Mn, Fe, Co, Cu, and Zn Employing LA-ICP-MS
847 and Hydrogen Reaction Gas. Analytical Chemistry. **84**(15), 6707-6714.
848 doi:10.1021/ac301156f.
- 849 Li, L., Høie, H., Geffen, A. J., Heegaard, E., Skadal, J., Folkvord, A. (2008). Back-calculation
850 of previous fish size using individually tagged and marked Atlantic cod (*Gadus morhua*).
851 Can. J. Fish. Aquat. Sci. **65**, 2496–2508. doi:10.1139/F08-157.
- 852 Limburg, K. E. 1995. Otolith strontium traces environmental history of subyearling American
853 shad *Alosa sapidissima*. Mar. Ecol. Prog. Ser. **119**(1/3), 25-35.
854 <http://www.jstor.org/stable/24849794>.
- 855 Limburg, K.E., Olson, C., Walther, Y., Dale, D., Slomo, C.P., and Høie, H. 2011. Tracking
856 Baltic hypoxia and cod migration over millennia with natural tags. Proc. Natl. Acad. Sci.
857 U.S.A. **108**(22): E177–E182. doi: 10.1073/pnas.1100684108.
- 858 Limburg, K.E., Walther, B.D., Lu, Z., Jackman, G., Mohan, J., Walther, Y., Nissling, A.,
859 Weber, P.K., and Schmitt, A.K. 2015. In search of the dead zone: Use of otoliths for
860 tracking fish exposure to hypoxia. J. Mar. Syst. **141**: 167–178.
861 doi:10.1016/J.JMARSYS.2014.02.014.
- 862 Limburg, K.E., Wuenschel, M.J., Hüsey, K., Heimbrand, Y., and Samson, M. 2018. Making
863 the Otolith Magnesium Chemical Calendar-Clock Tick: Plausible Mechanism and
864 Empirical Evidence. Rev. Fish. Sci. Aquac. **26**(4): 479–493.

- doi:10.1080/23308249.2018.1458817.
- Miller, J.A. 2011. Effects of water temperature and barium concentration on otolith composition along a salinity gradient: implications for migratory reconstructions. *J. Exp. Mar. Bio. Ecol.* **405**(1): 42–52. doi:10.1016/j.jembe.2011.05.017.
- Milton, D.A., and Chenery, S.R. 2001. Sources and uptake of trace metals in otoliths of juvenile barramundi (*Lates calcarifer*). *J. Exp. Mar. Bio. Ecol.* **264**(1): 47–65. doi:10.1016/S0022-0981(01)00301-X.
- Munk, P., Fox, C.J., Bolle, L.J., van Damme, C.J.G., Fossum, P., and Kraus, G. 2009. Spawning of North Sea fishes linked to hydrographic features. *Fish. Oceanogr.* **18**(6): 458–469. doi:10.1111/j.1365-2419.2009.00525.x
- Nielsen, E.E., Cariani, A., Mac Aoidh, E., Maes, G.E., Milano, I., Ogden, R., Taylor, M., Hemmer-Hansen, J., Babbucci, M., Bargelloni, L., Bekkevold, D., Diopere, E., Grenfell, L., Helyar, S., Limborg, M.T., Martinsohn, J.T., McEwing, R., Panitz, F., Patarenello, T., Tinti, F., van Houndt, J.K.J., Volckaert, F.A.M., Waples, R.S., and Carvalho, G.R. 2012. Gene-associated markers provide tools for tackling illegal fishing and false eco-certification. *Nat. Com.* **3**: 851. doi: 10.1038/ncomms2975.
- Nielsen, E.E., Hansen, M.M., Ruzzante, D.E., Meldrup, D., and Grønkjær, P. 2003. Evidence of a hybrid-zone in Atlantic cod (*Gadus morhua*) in the Baltic and the Danish Belt Sea revealed by individual admixture analysis. *Mol. Ecol.* **12**: 1497–1508. doi: 10.1046/j.1365-294X.2003.01819.x.
- Oeberst, R. 2008. Distribution pattern of cod and flounder in the Baltic Sea based on international coordinated trawl surveys. ICES CM 2008/J:09.
- Pihl, L., and Ulmestrand, M. 1993. Migration pattern of juvenile cod (*Gadus morhua*) on the Swedish west coast. *ICES J. Mar. Sci.* **50**(1): 63–70. doi: 10.1006/jmsc.1993.1007.
- Piry, S., Alapetite, A., Cornuet, J.-M., Paetkau, D., Baudouin, L., and Estoup, A. 2004.

- 890 GeneClass2: A Software for Genetic Assignment and First-Generation Migrant Detection.
891 J. Hered. **95**: 536-539. doi:10.1093/jhered/esh074.
- 892 R Core Team (2020). R: A language and environment for statistical computing. R Foundation
893 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>. Venables,
894 W. N. & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth Edition. Springer,
895 New York. ISBN 0-387-95457-0.
- 896 Rannala, B., and Mountain, J.L. 1997. Detecting immigration by using multilocus genotypes.
897 Proc. Nat. Acad. Sci. USA **94**: 9197-9221. doi:10.1073/pnas.94.17.9197.
- 898 Reis-Santos, P., Tanner, S.E., Elsdon, T.S., Cabral, H.N., and Gillanders, B.M. 2013. Effects
899 of temperature, salinity and water composition on otolith elemental incorporation of
900 *Dicentrarchus labrax*. J. Exp. Mar. Bio. Ecol. **446**: 245–252.
901 doi:10.1016/J.JEMBE.2013.05.027.
- 902 Righton, D.A., Andersen, K.H., Neat, F., Thorsteinsson, V., Steingrund, P., Svedäng, H.,
903 Michalsen, K., Hinrichsen, H.H., Bendall, V., Neuenfeldt, S., Wright, P., Jonsson, P.,
904 Huse, G., Van Der Kooij, J., Mosegaard, H., Hüsey, K., and Metcalfe, J. 2010. Thermal
905 niche of Atlantic cod *Gadus morhua*: Limits, tolerance and optima. Mar. Ecol. Prog. Ser.
906 **420**: 1–13. doi:10.3354/meps08889.
- 907 Rosenberg, R., Cato, I., Förlin, L., Grip, K., and Rodhe, J. 1996. Marine environment quality
908 assessment of the Skagerrak - Kattegat. J. Sea Res. **35**(1–3): 1–8. doi:10.1016/S1385-
909 1101(96)90730-3.
- 910 Rosenberg, R., Loo, L.-O., and Möller, P. 1992. Hypoxia, salinity and temperature as
911 structuring factors for marine benthic communities in a eutrophic area. Netherlands J. Sea
912 Res. **30**: 121–129. doi:10.1016/0077-7579(92)90051-F.
- 913 Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and
914 Eliceiri, K.W. 2017. ImageJ2: ImageJ for the next generation of scientific image data.

- 915 BMC Bioinformatics **18**(1): 529. doi:10.1186/s12859-017-1934-z.
- 916 Secor, D. H., Henderson-Arzapalo, A., and Piccoli, P. M. 1995. Can otolith microchemistry
917 chart patterns of migration and habitat utilization in anadromous fishes? J. Exp. Mar. Biol.
918 Ecol. **192**: 15–33. Available from
919 <https://www.sciencedirect.com/science/article/pii/002209819500054U>.
- 920 Serre, S.H., Nielsen, K.E., Fink-Jensen, P., Thomsen, T., and Hüsey, K. 2018. Analysis of cod
921 otolith microchemistry by continuous line transects using la-icp-ms. Geological Survey
922 of Denmark and Greenland Bulletin, **41**:91–94. doi:1604-8156.
- 923 Shin, J.-H., Blay, S., McNeney, B., and Graham J. 2006. LDheatmap: An R Function for
924 graphical display of pairwise linkage disequilibria between single nucleotide
925 polymorphisms. J. Stat. Soft. **16**: Code Snippet 3. Available from
926 <https://www.jstatsoft.org/article/view/v016c03>.
- 927 Stenseth, N.C., Jorde, P.E., Chan, K.-S., Hansen, E., Knutsen, H., André, C., Skogen, M.D.,
928 and Lekve, K. 2006. Ecological and genetic impact of Atlantic cod larval drift in the
929 Skagerrak. Proc. R. Soc. B Biol. Sci. **273**(1590): 1085–1092.
930 doi:10.1098/rspb.2005.3290.
- 931 Sturrock, A.M., Hunter, E., Milton, J.A., EIMF, Johnson, R.C., Waring, C.P., and Trueman,
932 C.N. 2015. Quantifying physiological influences on otolith microchemistry. Methods
933 Ecol. Evol. **6**(7): 806–816. doi:10.1111/2041-210X.12381.
- 934 Sturrock, A.M., Trueman, C.N., Darnaude, A.M., and Hunter, E. 2012. Can otolith elemental
935 chemistry retrospectively track migrations in fully marine fishes? J. Fish Biol. **81**(2): 766–
936 795. doi:10.1111/j.1095-8649.2012.03372.x.
- 937 Svansson, A. 1975. Physical and chemical oceanography of the Skagerrak and the Kattegat.
938 Fishery Board of Sweden, Uddevalla. p. 93. Available from
939 https://gupea.ub.gu.se/bitstream/2077/49089/1/gupea_2077_49089_1.pdf.

- 940 Svedäng, H., André, C., Jonsson, P., Elfman, M., and Limburg, K.E. 2010. Migratory
941 behaviour and otolith chemistry suggest fine-scale sub-population structure within a
942 genetically homogenous Atlantic Cod population. *Environ. Biol. Fishes* **89**(3–4): 383–
943 397. doi:10.1007/s10641-010-9669-y.
- 944 Svedäng, H., Righton, D., and Jonsson, P. 2007. Migratory behaviour of Atlantic cod *Gadus*
945 *morhua*: natal homing is the prime stock-separating mechanism. *Mar. Ecol. Prog. Ser.*
946 **345**: 1–12. doi:10.3354/meps07140.
- 947 Tang, Y., and Li, W. 2016. lfda: An R Package for Local Fisher Discriminant Analysis and
948 Visualization. Available from <https://arxiv.org/abs/1612.09219>.
- 949 Venables, W.N., and Ripley, B.D. 2002. *Modern Applied Statistics with S*, Fourth edition.
950 Springer, New York. ISBN 0-387-95457-0, <https://www.stats.ox.ac.uk/pub/MASS4/>.
- 951 Vitale, F., Börjesson, P., Svedäng, H., and Casini, M. 2008. The spatial distribution of cod
952 (*Gadus morhua* L.) spawning grounds in the Kattegat, eastern North Sea. *Fish. Res.* **90**(1–
953 3): 36–44. doi:10.1016/j.fishres.2007.09.023.
- 954 Vitale, F., Worsøe Clausen, L., and Ní Chonchúir, G. (Eds.) 2019. *Handbook of fish age*
955 *estimation protocols and validation methods*. ICES Cooperative Research Report No. 346.
956 180 pp. doi:10.17895/ices.pub.5221.
- 957 Walther, B.D., Kingsford, M.J., O’Callaghan, M.D., and McCulloch, M.T. 2010. Interactive
958 effects of ontogeny, food ration and temperature on elemental incorporation in otoliths of
959 a coral reef fish. *Environ. Biol. Fish.* **89**(3–4):441–451. doi:10.1007/s10641-010-9661-6.
- 960 Walther, B.D., and Limburg, K.E. 2012. The use of otolith chemistry to characterize
961 diadromous migrations. *J. Fish Biol.* **81**(2): 796–825. doi:10.1111/j.1095-
962 8649.2012.03371.x.
- 963 Watanabe, T., Kiron, V., and Satoh, S. 1997. Trace minerals in fish nutrition. *Aquaculture*
964 **151**(1–4): 185–207. doi:10.1016/S0044-8486(96)01503-7.

965 Weidman, C.R., and Millner, R. 2000. High-resolution stable isotope records from North
966 Atlantic cod. *Fish. Res.* **46**(1–3): 327–342. doi:10.1016/S0165-7836(00)00157-0.
967

Draft

Figure captions

Figure 1. Map of the sampling area, where the management areas of the cod populations are 1. North Sea stock: North Sea and Skagerrak, 2. Kattegat stock: Kattegat, and 3. western Baltic Sea stock: Belt Sea, Sound and Arkona Sea. Pie charts indicate sampling locations, where the size of the symbols represents number of fish, and the colour represents populations North Sea (bright blue) and Kattegat (red). The colouring of the geographic areas indicate the habitat occupation areas used: North Sea/Skagerrak (lime), Kattegat (green), western Baltic Sea (blue green), and eastern Baltic Sea (purple). Map created using the packages “maps”, “mapdata” and “maptools” (Becker and Wilks 1993) in “R” (R Core Team 2020). The samples used in this study were collected during scientific surveys by the Technical University of Denmark in 2016.

Figure 2. Image of otolith cross-section, viewed under reflected light, with laser track indicated with a solid black line, and the corresponding otolith Sr profile where colours indicate the habitat area assignments (North Sea = lime, Kattegat = green, western Baltic = blue green). Translucent winter growth zones are outlined by vertical shaded bars linking the visual image with the corresponding sections of the Sr profile. This individual is 4 years old Kattegat population cod captured in the northern Kattegat had spent most of its life in the western Baltic/Kattegat and performed two migrations into the Skagerrak or North Sea during fall/winter of its second and third year of life.

Figure 3. Otolith size at the end of first year of life for the 6 age groups of cod captured in Kattegat, representing the year classes 2010 – 2015, where colours represent different populations: Kattegat (red) and North Sea (blue). Otolith sizes of age 0 individuals (year class 2016) are not included in this graph, because they were captured before the end of their first

winter. Horizontal lines indicate mean, box upper and lower limits the 25% and 75% percentiles, whiskers represent the highest and lowest values within 1.5 interquartile range and dots represent outliers.

Figure 4. Fish size in relation to age of cod captured in Kattegat with linear regression lines and confidence interval bands, where colours represent different populations: Kattegat (red) and North Sea (blue). The slope of the North Sea population is significantly higher than that of the Kattegat population, suggesting faster growth in North Sea cod.

Figure 5. Biplot of the first two discriminant functions of the LDA analysing otolith chemical fingerprints in the larval and pelagic juvenile stage (0 to ca. 4 months old), the demersal juvenile stage until the first winter zone (ca. 4 to 8 months old), and the first winter, with life stage indicated above each biplot. Colours represent the two populations: Kattegat (red) and North Sea (blue), and arrows the direction and strength of the loadings (= elements).

Figure 6. Area assignment for each individual fish and each time interval of its entire lifespan from the first winter to catch for cod from the Kattegat population and the North Sea population (population indicated above graph). The time from hatch to the first winter is not included in this plot, as the area assignment for this age group is based on a different habitat area levels and provides less precise assignments. Colours indicate the habitat area assignments: North Sea = lime, Kattegat = green, western Baltic = blue green, eastern Baltic = purple. Lifetime area use is shown for the two populations separately, where individuals are ordered according to latitude of their capture location in the Kattegat.

1017 **Figure 7.** Percentage distributions of habitat area assignments by age and season, representing
1018 a population's distribution pattern in space and time, for the Kattegat population and the North
1019 Sea population separately (population indicated above graph). Colours indicate percentage of
1020 individuals with an area use assigned to the North Sea (lime), the Kattegat (green), the western
1021 Baltic Sea (blue green), and the eastern Baltic Sea (purple).

1022

1023 **Figure 8.** Percentage of individuals assigned to the Skagerrak/North Sea (lime coloured area
1024 in Fig. 1) by age and season, for the Kattegat population and the North Sea population
1025 (population indicated above graph). Colour shading represents the percentage estimate based
1026 on the mean values of the otolith strontium – salinity calibration curve (dark lime), for mean -
1027 se values (no colour), and mean + se (transparent lime).

Image of otolith cross-section, viewed under reflected light, with laser track indicated with a solid black line, and the corresponding otolith Sr profile where colours indicate the habitat area assignments (North Sea = lime, Kattegat = green, western Baltic = blue green). Translucent winter growth zones are outlined by vertical shaded bars linking the visual image with the corresponding sections of the Sr profile. This individual is 4 years old Kattegat population cod captured in the northern Kattegat had spent most of its life in the western Baltic/ Kattegat and performed two migrations into the Skagerrak or North Sea during fall/winter of its second and third year of life.

290x179mm (150 x 150 DPI)

Otolith size at the end of first year of life for the 6 age groups of cod captured in Kattegat, representing the year classes 2010 – 2015, where colours represent different populations: Kattegat (red) and North Sea (blue). Otolith sizes of age 0 individuals (year class 2016) are not included in this graph, because they were captured before the end of their first winter. Horizontal lines indicate mean, box upper and lower limits the 25% and 75% percentiles, whiskers represent the highest and lowest values within 1.5 interquartile range and dots represent outliers.

85x101mm (300 x 300 DPI)

Fish size in relation to age of cod captured in Kattegat with linear regression lines and confidence interval bands, where colours represent different populations: Kattegat (red) and North Sea (blue). The slope of the North Sea population is significantly higher than that of the Kattegat population, suggesting faster growth in North Sea cod.

85x101mm (300 x 300 DPI)

Biplot of the first two discriminant functions of the LDA analysing otolith chemical fingerprints in the larval and pelagic juvenile stage (0 to ca. 4 months old), the demersal juvenile stage until the first winter zone (ca. 4 to 8 months old), and the first winter, with life stage indicated above each biplot. Colours represent the two populations: Kattegat (red) and North Sea (blue), and arrows the direction and strength of the loadings (= elements).

266x86mm (300 x 300 DPI)

Individual fish

Age 1

Age 2

Age 3

Age 4

Spring Summer Fall Winter Spring Summer Fall Winter Spring Summer Fall Winter Spring Summer Fall Winter

Season

The Author(s) or their Institution(s)

Table 1. Overview over samples used in this study by population and age.

	Population		
Age	Kattegat	North Sea	Total
0	23	79	102
1	7	28	35
2	32	42	74
3	14	8	22
4	52	7	59
5	5	1	6
6	7		7
8	1		1
Total	141	165	306

Table 2. Concentration of element/Ca ratios (ppm), in the pelagic juvenile stage by population (mean +/- standard deviation), together with ANOVA statistics indicating differences between populations (ns = not significant, *** < 0.001)

	Population				
Element	Kattegat	North Sea	df	F	<i>p</i>
Sr	1458.1 (128.9)	1604.4 (181.1)	1	12.34	***
Ba	24.3 (7.9)	17.1 (7.3)	1	7.56	***
Mn	12.4 (5.7)	8.2 (3.7)	1	101.18	***
Mg	61.1 (9.9)	73.0 (39.1)	1	13.38	***
P	276.9 (55.7)	312.7 (90.1)	1	0.001	ns
Cu	0.7 (0.1)	0.4 (0.2)	1	11.90	***
Zn	5.5 (2.6)	3.2 (1.5)	1	22.82	***