DIARRHEA DURATION AND PERFORMANCE OUTCOMES OF PRE-WEANED DAIRY CALVES SUPPLEMENTED WITH BACTERIOPHAGE

Journal:	<i>Canadian Journal of Animal Science</i>
Manuscript ID	CJAS-2021-0074.R1
Manuscript Type:	Article
Date Submitted by the Author:	17-Sep-2021
Complete List of Authors:	Schmoeller, Evandro; Universidade Federal de Pelotas, Núcleo de Pesquisa, Ensino e Extensão em Pecuária de Matos, Adriane; Federal University of Pelotas Rahal, Natalia; Federal University of Pelotas, Veterinary Feijó, Josiane; Federal University of Pelotas Brauner, Cassio; Universidade Federal de Pelotas, Zootecnia- FAEM; NUPEEC, Faculdade de Veterinária Del Pino, Francisco; Universidade Federal de Pelotas, Bioquímica; Núcleo de Pesquisa, Ensino e Extensão em Pecuária - UFPel, Faculdade de Veterinária Corrêa, Marcio; Núcleo de Pesquisa, Ensino e Extensão em Pecuária - UFPel, Faculdade de Veterinária; Universidade Federal de Pelotas, Clínicas Veterinária Rabassa, Viviane; Faculdade de Veterinaria da Universidade Federal de Pelotas, Departamento de Clinicas Veterinaria
Keywords:	phage therapy, growth, neonatal, bovine

SCHOLARONE™
Manuscripts

DIARRHEA DURATION AND PERFORMANCE OUTCOMES OF PRE-WEANED
DAIRY CALVES SUPPLEMENTED WITH BACTERIOPHAGE

Evandro Schmoeller^a, Adriane D. C. de Matos^a, Natalia M. Rahal^a, Josiane O. Feijo^a,
Cassio C. Brauner^{a,b}, Francisco Augusto B. Del Pino^{a,c}, Marcio N. Correa^{a,d}, Viviane R.
Rabassa^{a,d*}

^aNúcleo de Pesquisa, Ensino e Extensão em Pecuária – Livestock Research, Education and Extension Center, Universidade Federal de Pelotas, CEP 96010-900, Capão do Leão, RS, Brazil

^bDepartamento de Zootecnia, Faculdade de Agronomia Eliseu Maciel da Universidade Federal de Pelotas, CEP 96010-900, Capão do Leão, RS, Brazil

^cCentro de Ciências Químicas Farmacêuticas e de Alimentos, Universidade Federal de Pelotas, CEP 96010-900, Capão do Leão, RS, Brazil

^dDepartamento de Clínicas Veterinária, Faculdade de Veterinária da Universidade Federal de Pelotas, CEP 96010-900, Capão do Leão, RS, Brazil

*corresponding author: Viviane Rohrig Rabassa, Departamento de Clínicas Veterinária, Universidade Federal de Pelotas, Campus Universitário s/n, CEP 96160-000, Capão do Leão, RS, Brazil, +55 53 32757136, vivianerabassa@gmail.com

ABSTRACT

This study aimed to evaluate lytic bacteriophage supplementation in pre-weaned dairy calves over diseases occurrence, performance and biochemical parameters. Two hundred Holstein×Gyr crossbred female calves were divided into two groups: CON, no supplementation; and PHAGE, bacteriophage supplementation (1 g/day) from d 3 until d 70 of life. Calves were monitored daily for age at the first diarrheic episode and its duration. Fecal samples were cultured for isolation and PCR was performed to identify *E. coli* virulence genes and to confirm *Salmonella* spp. Performance outcomes were evaluated up to 80 d of age. Blood samples were collected to determine serum levels of total proteins, albumin, cholesterol, γ -glutamyl transferase and urea. PHAGE group had fewer days in diarrhea (PHAGE: 4.68 d, CON: 6.61 d; $P=0.03$). Fecal samples of three animals in PHAGE and nine in CON were positive for *E. coli* in PCR. Average daily gain mean of PHAGE was higher up to 80 d of life ($P<0.05$). PHAGE mean was lower for albumin and higher for urea ($P=0.004$ and $P<0.001$, respectively). Phage therapy during the pre-weaned period reduced the duration of neonatal diarrhea, providing greater weight gain for calves.

Key words: phage therapy, growth, neonatal, bovine.

INTRODUCTION

Appropriate calf management is fundamental for dairy cattle production, because the neonatal period presents several challenges to these animals. Metabolic instability, associated with the beginning of the immune response development makes these animals susceptible to diseases and increased mortality (Volpato et al., 2017). Among diseases of this period, neonatal calf diarrhea (NCD) is associated with significant economic losses due to high morbidity and mortality, cost of treatment and impaired development (Coura et al., 2015; Foster, 2009). NCD can be infectious (caused by viruses, bacteria and protozoa); or related to nutrition, influenced by temperature, quantity and quality of milk or milk replacer; and climate conditions (Smith, 2009). Regarding the primary cause, with imbalance of intestinal microbiota due to diarrhea, bacterial enteritis ends up occurring as an opportunistic infection in most clinical conditions (Cho et al., 2013).

Escherichia coli and *Salmonella* spp. are the main etiologic agents of bacterial NCD and are commonly treated with antibiotics (Dall Agnol et al., 2021). Antibiotic therapy has been questioned for stimulating factors of antimicrobial resistance and death of

microorganisms that act symbiotically to the intestine, participating in the development of intestinal tissue and the immune system (Macpherson et al., 2004). This imbalance can impair inflammatory response modulation (Lei et al., 2015), causing a competition between symbiotic agents and pathogenic microorganisms for nutrients and intestinal epithelial surface, prolonging or aggravating NCD (Backhed et al., 2005; Rakoff-Nahoum et al., 2004).

In order to decrease the indiscriminate use of antibiotics, phage therapy has emerged as an alternative in prophylactic control of bacterial diarrhea. Bacteriophages are mandatory intracellular viral particles that parasitize bacteria with high specificity (Santos et al., 2014). These phages can have different life cycles, known as lytic or lysogenic. In the lytic cycle, they become excellent tools in controlling pathogenic bacteria, as they use bacterial metabolism to replicate and produce endolysins, causing cell wall destruction and bacterial death (Harper and Enright, 2011). Studies have shown positive effects on maintaining ruminants' intestinal microbiota balance (Callaway et al., 2008; Sheng et al., 2006) and controlling septicemia in calves and birds (Barrow et al., 1998). Therefore, we aimed to evaluate lytic bacteriophage supplementation in pre-weaned dairy calves over NCD and respiratory diseases occurrence, performance and biochemical parameters of dairy calves.

MATERIAL AND METHODS

Animals and experimental conditions

The experiment was conducted in accordance with the Canadian Council on Animal Care in Science (2009) and approved by Ethics Committee on Animal Experimentation - CEEA of the Federal University of Pelotas (number 14807). The study was conducted on a commercial dairy farm in Passos, Minas Gerais, Brazil and sample size was established using G*Power software 3.9.1.2 version. Two hundred Holstein × Gyr crossbred female calves with different breed admixture ($1/2$ H + $1/2$ G; $3/4$ H + $1/4$ G, $5/8$ H + $3/8$ G) were maintained at an outdoor system known as “tropical housing”. Briefly, calves were neck-tied to a chain fixed to a wire stretched on the ground (tied to wood posts) and had access to shade provided by a shading screen. After birth, all calves were bottle-fed 10% BW of colostrum (i.e., 35 kg calf was fed with 3.5 L of colostrum), divided into two meals: the first one until the first 8 hours of life; and the second one until the first 24 hours. Colostrum quality was measured using Brix refractometry (K52-032, Kasvi, Brazil), considering good quality when values were $\geq 20\%$ (Quigley et al., 2013). Until 3 d of life calves were kept in individual pens and

were bucket or bottle fed 4 L of transition milk (milk from cows from days 2 to 5 post-partum), divided into two meals. From the third day, calves were bucket fed with a 3 L of milk replacer Nurture Prime, Nutron, Brazil (22% CP, 18% fat DM basis, 38% lactose DM basis), twice a day, up to 42 d of life. After d 42, 3 L of milk replacer was given only in the afternoon beginning the step-down weaning process. In addition, from the 3 d of life animals had free access to starter (12.59% CP and 15.2% starch DM basis) and water and at 60 d corn silage was offered (NRC, 2000). Weaning was performed at 70 d of life, with calves remaining in this housing system until 80 d.

To verify transfer of passive immunity (TPI), blood sample was collected within 48 h of life via jugular venipuncture using vacuum tube with EDTA (5mL). These were centrifuged at $2183 \times g$ to obtain plasma and total plasma protein (TPP) was analyzed in an optical refractometer (RFATC200, Benfer, Brazil). Failed TPI (FTPI) was determined in calves with TPP below 5.5 g/dL (Tyler et al., 1996), and calves with FTPI were homogeneously distributed between groups.

This study had a completely randomized design. Animals were homogeneously divided into two groups and blocked according to birth weight, breed admixture, birth order and FTPI. Bacteriophage group (PHAGE), $n=100$ ($50 \frac{1}{2}$, $48 \frac{3}{4}$, $2 \frac{5}{8}$ of Holstein), was fed 1g of a product containing bacteriophage (FAGOLAC, Bayer Animal Health, Brazil) added to milk replacer from d 3 d until weaning, and initial weight mean was 35.48 ± 0.75 kg. Control Group (CON), $n=100$ ($51 \frac{1}{2}$, $48 \frac{3}{4}$, $1 \frac{5}{8}$ of Holstein blood), was fed milk replacer without addition of the product, and initial weight mean was 35.31 ± 0.71 kg.

Calves were monitored daily only by a trained veterinary up to 80 d of life for occurrence of diarrhea and respiratory disease. For diarrhea, animals were considered sick when fecal score was ≥ 2 and for respiratory disease when greyish to yellowish nasal discharge was presented accompanied by fever and abnormal respiratory sounds. From these records, the following rates were determined: morbidity (number of sick calves/total number of calves enrolled in the experiment); mortality (number dead calves/ total number of calves enrolled in the experiment); case fatality rate (number of calves that died from diarrhea or respiratory disease/ total number of calves that presented diarrhea or respiratory disease); and relapse (number of animals that got sick twice or more).

Clinical diagnostics were based on evaluation of dehydration, respiratory and heart rates and rectal temperature (Smith, 2009). Sick calves were treated according to farm's protocol similar for both groups. For diarrhea, treatment with trimethoprim-sulfamethoxazole (20mg/kg B.W., once a day, intramuscular injection) was used for 3 d or more (according to clinical condition) associated with oral fluids. Penicillin (procaine benzylpenicillin 7,500UI/kg; benzathine benzylpenicillin 5,000UI/kg; dihydrostreptomycin 17.06mg/kg, once a day, intramuscular injection) associated with meloxicam (0.5 mg/kg B.W., once a day, intramuscular injection) were used for 3-5 days to treat respiratory disease.

Fecal score (FS) and duration of diarrhea

FS was determined on a scale of 0 to 4, where 0 = normal consistency, 1 = pasty, 2 = loose or watery feces, 3 = profuse diarrhea with liquefied stools, and 4 = profuse diarrhea with liquefied and bloody feces (adapted from Kertz and Chester-Jones, 2004). Onset of diarrheic episodes occurred when calves presented $FS \geq 2$ and ended when $FS \leq 1$. The total number of days calves remained with diarrheic feces ($FS \geq 2$), considering all diarrheic episodes of each calf, and duration of the first diarrheic episode (days) were determined throughout the first 30 d of life.

Fecal samples and bacterial characterization

For this analysis were included only samples from calves that did not receive farm's treatment for diarrhea at the time of sampling. Sixty-six fecal samples (PHAGE=31; CON=35) on the first day of the first diarrheic episode were collected directly from the rectum, packed in a sterile cup and frozen at -20°C until microbiological analysis. Subsequently, these samples were cultured in two types of medium, for *Salmonella* spp. and *E. coli* growth.

For *Salmonella* spp. identification, samples were initially grown in tetrathionate broth (Acumedia Manufacturers, Inc. Baltimore, Maryland). The broth was kept in a bacteriological oven (Olidex, Brazil) at 35-37°C for 72 hours. Every 24 hours of incubation, an aliquot was removed from the broth and added to MacConkey agar (Kasvi, Brazil), incubated again under the same conditions for 24 hours. Biochemical tests were performed with colonies with *Salmonella* spp. visual characteristics, following the technique described by Quinn et al. (1993).

For *E. coli*, lactose fermenting colonies found in MacConkey agar medium were submitted to biochemical characterization (Quinn et. Al., 1993). The PCR was performed for colonies characterized as *E. coli* and *Salmonella* spp.

Molecular methods

The genomic DNA extraction of *E. coli* and *Salmonella* spp. colonies was performed according to the protocol described by Green and Sambrook (2012). The DNA concentration was of samples was 5 ng/ μ L, with purity ratio ≥ 1.8 quantified by spectrophotometry using NanoVue Plus (GE Healthcare Life Sciences, Buckinghamshire, UK).

Amplification procedure followed the protocols described in the literature. For *E. coli*, the *USPA* gene was PCR amplified and sequenced to confirm genus and species designation, generating a fragment of 884 base pairs (bp) (Chen; Griffiths, 1998). After confirmation, the PCR amplification of the following virulence genes were attempted: (i) *EAE* (384 bp), intimin, present in the enteropathogenic *E. coli* (EPEC); (ii) *STXI* (180 bp), which encode for Shiga-toxin-producing *E. coli* (STEC); (iii) *HLYA* (534 bp), encoding for hemolysin, present in the enterohaemorrhagic *E. coli* (EHEC); and (iv) *ESTIA* (157 bp), encoding for thermostable toxin A. For genus confirmation of *Salmonella* spp., the *HILA* gene was PCR amplified and sequenced, generating a 413 bp fragment (Craciunas et al., 2012).

The conventional PCR was performed as follows: 12.5 μ L of Taq polymerase master mix 2x (Cellco), 8.5 μ L of water for PCR (Ludwig Biotec), 10 pmol (1 μ L) of each forward and reverse primers (Extend), and 10ng (2 μ L) of template DNA, for a final volume of 25 μ L. Amplification was performed on MJ Research® PTC 100 thermocycler (Bio-Rad Laboratories, Hercules, CA, USA) and PCR products were subjected to electrophoresis on an agarose gel 1.5% (for fragments larger than 200 bp) and 2% (for fragments smaller than 200 bp) with a standard molecular weight of 1kb (Invitrogen). Then, the amplified products were visualized on a transilluminator (Loccus, L-Pix Touch).

Performance outcomes

The assessment of calves' body weight, average daily gain (ADG), thoracic perimeter (chest circumference), withers height and rump width were performed at birth and weekly until 30 d of life, and subsequently evaluated at days 42, 60 and 80. The body weight was determined through the circumferential measurement of thoracic perimeter using a graduated tape in kg. Average daily gain was determined by dividing weight gain in the period for the

number of days in the same period. Withers' height (from the ground to scapular junction) and croup width (between the ischial tuberosities) were measured using a graduated (cm) tape. A total of 175 calves (PHAGE=83; CON=92) were included in these evaluations, calves that died before the end of experimental period were excluded.

Blood collection and biochemical analyzes

For biochemical analyses a sampling of both groups was used (PHAGE=43; CON=41). Blood samples were collected through jugular venipuncture at d 0 (up to 24 hours after birth) and at d 7, 14, 21, 30 and 60 using a vacuum system and a tube without anticoagulant. Samples were centrifuged at $2183 \times g$ for 15 min, serum was stored in 1.5 ml microtubes and frozen at -20°C until the time of analysis. The Labmax Plenno Automatic Biochemical Analyzer, standard model (Labtest, Brazil), was used and analyzes were performed following manufacturer's protocols and the reagent kits specifically (Labtest, Brazil) for each parameter: total proteins, albumin, cholesterol, γ -glutamyl transferase (GGT) and urea.

Statistical analysis

The T Test was used to compare birth weight and TPI between groups, as for performance and biochemical data PROC MIXED method was used, considering animal as random effect, and group and age, as well as their interactions, as a fixed effect. Each animal was considered an experimental unit. Normality was verified by PROC UNIVARIATE of residues. Comparisons between groups regarding number of diarrheic calves in the first 30 days of life were performed using Fisher's Exact Test. Diarrhea duration was analyzed using One-way ANOVA, while the categorical variables TPP, morbidity, mortality, case fatality rate, disease recurrence and positive PCR isolates for *E. coli* and *Salmonella* spp. were assessed by the Chi-square and Fischer's exact test. All analyses were performed on SAS University Edition (SAS Institute Inc., Cary, USA), and mean difference were considered significant when $P < 0.05$ and tendency when $0.05 < P > 0.1$.

RESULTS

There were no differences in TPI and birth weight between groups ($P=0.25$ and $P=0.35$, respectively). Morbidity, mortality, case fatality rate and disease relapse rates did not differ between groups (Tab. 2). First diarrheic episode duration and all days with diarrhea during the first month of life were shorter in PHAGE ($P < 0.001$ and $P=0.03$, respectively)

during the neonatal period (Fig. 1). On days 3, 5 to 13 and 25 of life, PHAGE had fewer diarrheic calves ($P \leq 0.05$; Tab.3; Fig. 2).

Regarding age at first diarrhea, there was no difference between the groups (PHAGE=6.15 and CON=5.57; $P=0.99$). Sixty-six feces' samples were tested in microbiological analysis and 27 of these had colonies with biochemical characteristics of *E. Coli* and 2 samples with colonies of *Salmonella* spp. Positive samples for *USPA* gene tended to be higher in CON group (CON:9; PHAGE: 3; $P=0.09$). In one *E. coli* positive sample from CON, *EAE* gene (intimin virulence factor) characteristic of EPEC was expressed. None of the samples was positive for *Salmonella* spp. in the molecular analysis.

The PHAGE group had higher ADG mean in the neonatal period (first 30 d of life; $P=0.004$), at the beginning of step-down weaning (up to 42 days of life; $P=0.004$) and until after weaning (80 d of life; $P=0.009$) (Fig. 3). The supplemented group had greater weight at the end of the experiment ($P<0.001$) and tended to have a greater thoracic perimeter ($P=0.06$), while withers height and croup width did not differ ($P>0.05$) between groups (Tab. 3).

Albumin and urea levels were different between groups, the PHAGE mean was lower for albumin (PHAGE=2.65 g/dL; CON=2.76g/dL) ($P=0.004$) and higher for urea (PHAGE=21.4 mg/dL; CON=17.82mg/dL) ($P<0.0001$). Other parameters, GGT (PHAGE=297.51 U/L; CON=295.08 U/L), total proteins (PHAGE=6.33 g/dL; CON=6.29 g/dL) and cholesterol (PHAGE=87.85 mg/dL; CON=86.63 mg/dL), did not differ ($P>0.1$). All biochemical variables differed considering calves' age ($P<0.0001$), and for albumin and total proteins interaction between experimental groups and age was observed ($P<0.0001$) (Fig. 4).

DISCUSSION

Enteric diseases are common on the first weeks of calves' lives, negatively affecting ADG, and consequently increasing female heifers' age at first conception (Svensson et al, 2003; Botteon et al., 2008). Additionally, diseases occurrence in neonatal period may cause future additional health problems, decreasing quality and quantity of milk production (Kozasa et al., 2005; Laureyns et al., 2009). Therefore, reduction in morbidity and duration of diarrhea can determine higher ADG and, consequently, improvements in animal health and production. In this study, we observed a shorter duration of diarrhea in the PHAGE group in the neonatal period and, consequently, a greater weight gain during the pre-weaned period.

Reduction in the diarrhea period and fewer calves with diarrhea up to 13 d of life in PHAGE may be associated with the control of bacterial proliferation provided by bacteriophage use. Co-infection is commonly seen in diarrheic calves (Cho and Yoon, 2014), and an intestinal dysbiosis and immune suppression may facilitate the proliferation of opportunistic bacteria increasing the duration and severity of the disease (Cho et al., 2013). Such fact was reported by Smith et al (1987) and Sulakvelidze et al. (2001), that confirmed the phage effect on reducing the magnitude of infectious processes. Higher albumin and total protein noticed in PHAGE calves may be explained due to less aggravated diarrheic episodes, as albumin could be lost through feces (Choi et al., 2021). Also, calves supplemented with phage presented faster clinical recovery, less loss of protein, and possibly greater food intake, with better absorption of nutrients that could be the result of smaller extent of intestinal damage (Allison, 2017).

Loss of fluids and less nutrient absorption, associated with inflammatory process caused during diarrhea has a negative correlation with animals' weight gain (Barrington et al., 2002; Gifford et al., 2012). Thus, a shorter diarrhea duration may explain the higher ADG of calves that received milk replacer added with phages. It is noted that this difference is even greater considering only the neonatal period, which is the moment of the highest incidence of diarrhea (García et al., 2000).

In a study conducted with 122 dairy herds in Sweden, calves with neonatal diarrhea were more likely to develop respiratory infections up to six months of age (Hultgren et al., 2008). However, this association was not found in some herds in the USA and in Canada (Windeyer et al., 2014). Besides climate differences, in these countries, individual or collective pens are the most commonly used housing conditions in the first months of life. But, in this study, calves were maintained outdoors and did not had contact with each other, possibly reducing pathogens load and infection between calves. In the present study, the use of phages was not able to reduce respiratory disease morbidity rate but it was not possible to establish if the dose and administration route were able to reach respiratory organs or to determine the infectious agent..

Phage therapy probably acted on bacterial co-infection and secondary bacterial enteritis (Cho et al., 2013), however neonatal diarrhea may be caused by other factors or agents not controlled by phage use, such as nutrition, environment and virus (Cho and Yoon,

2014). Regarding causes of neonatal diarrhea, Santin et al. (2014) observed that 50.3% of fecal samples of calves between 5 days to 2 months of age were positive for *Cryptosporidium* spp., increasing to 66.7% when considering only animals up to 2 weeks of age. Thus, diarrhea morbidity rate did not differ between groups, probably due to primary causing factors or agents other than bacteria. However, phage therapy was able to decrease diarrhea duration by controlling opportunistic bacterial infections.

In vitro studies have shown satisfactory effects on the reduction of *E. coli* O: 157:H7 occurrence (Niu et al., 2009; Coffey et al., 2011; Carter et al., 2012), however, in vivo experiments show contradictory results. The phage therapy was effective in controlling *E. coli* intestinal population in cattle (Sheng et al., 2006; Niu et al., 2008) and sheep (Callaway, 2008). Nonetheless, other studies did not find the same effect (Bach et al., 2003; Raya et al., 2006; Rivas et al. 2010). Bacteria quantification was not possible in this study.

Still, the success of the phage therapy may be associated with administration route. Ronzema et al. (2009) demonstrated that oral route is superior to intra-rectal. Sheng et al. (2006) observed that the rectal application (4 doses of 25 ml with 1×10^{10} CFU) was sufficient to control bacterial population, although in this study continuous administration of phages in lower concentrations in the water was maintained (1.8×10^6 CFU). Also, Anand et al. (2015) found that broad-spectrum phage was able to kill 47.3% of *E. coli* isolates from diarrheic calves.

In this study, lytic bacteriophage added to milk replacer managed to control diarrhea, as observed by the decrease of days with diarrheic feces during neonatal period. Additionally, greater weight gain in the first 80 d of life was obtained for calves supplemented with bacteriophage.

DECLARATIONS

Funding

Commercial bacteriophage used in this study was provided by Bayer Animal Health, Brazil. This work was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) [Finance code 001]; and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil) [grant number 130783/2017-0].

Competing interests statement

The authors declare there are no competing interests.

Availability of data and material

Data will be made available on request

REFERENCES

- Anand, T., R. K. Vaid, B. C. Bera, S. Barua, T. Riyesh, N. Virmani, N. Yadav, and P. Malik. 2015. Isolation and characterization of a bacteriophage with broad host range, displaying potential in preventing bovine diarrhoea. *Virus genes*. 51(2):315-321. <https://doi.org/10.1007/s11262-015-1222-9>
- Allison, R.W. 2017. Avaliação laboratorial das proteínas do plasma e do soro sanguíneo. Pages 398-411 in *Hematologia e Bioquímica Clínica Veterinária*. 2nd ed. Thrall, M.A. Weiser, G., Allison, R.W. and Campbell, T.W. ed. Guanabara Koogan, Rio de Janeiro, Brazil.
- Bach, S. J., T. A. McAllister, D. M. Veira, V. P. Gannon, and R. A. Holley. 2003. Effect of bacteriophage DC22 on *Escherichia coli* O157: H7 in an artificial rumen system (Rusitec) and inoculated sheep. *Anim. Res.* 52(2):89-101. <https://doi.org/10.1051/animres:2003009>
- Bäckhed, F., R. E. Ley, J. L. Sonnenburg, D. A. Peterson, and J. I. Gordon. 2005. Host-bacterial mutualism in the human intestine. *Science*. 307(5717):1915-1920. <https://doi.org/10.1126/science.1104816>
- Botteon, R. C. C. M., P. D. T. L. Botteon, J. D. C. B. S. Júnior, M. H. Pinna, and Z. G. Lóss. 2008. Frequência de diarréia em bezerros mestiços sob diferentes condições de manejo na região do médio Paraíba Rio de Janeiro e Minas Gerais. *Braz. J. Vet. Res. Anim. Sci.* 45(2):153-160. <https://doi.org/10.11606/issn.1678-4456.bjvras.2008.26712>
- Barrington, G. M., J. M. Gay, and J. F. Evermann. 2002. Biosecurity for neonatal gastrointestinal diseases. *Vet. Clin. North Am. Food Anim. Pract.* 18(1):7. [http://doi.org/10.1016/S0749-0720\(02\)00005-1](http://doi.org/10.1016/S0749-0720(02)00005-1)
- Barrow, P., M. Lovell, and A. Berchieri. 1998. Use of lytic bacteriophage for control of experimental *Escherichia coli* septicemia and meningitis in chickens and calves. *Clin. Diagn. Lab. Immunol.* 5(3):294-298. <http://doi.org/10.1128/CDLI.5.3.294-298.1998>
- Callaway, T. R., T. S. Edrington, A. D. Brabban, R. C. Anderson, M. L. Rossman, M. J. Engler, M. A. Carr, K. J. Genovese, J. E. Keen, M. L. Looper, E. M. Kutter, and D. J. Nisbet. 2008. Bacteriophage isolated from feedlot cattle can reduce *Escherichia coli*

- O157: H7 populations in ruminant gastrointestinal tracts. *Foodborne Pathog. Dis.* 5(2):183-191. <https://doi.org/10.1089=fpd.2007.0057>.
- Canadian Council on Animal Care. 2021. Guidelines on the care and use of farm animals in research, teaching and testing. Canadian Council on Animal Care, Ottawa, ON, Canada. ISBN: 978-0-919087-89-7.
- Carter, C. D., A. Parks, T. Abuladze, M. Li, J. Woolston, J. Magnone, A. Senecal, A. M. Kropinski, and A. Sulakvelidze. 2012. Bacteriophage cocktail significantly reduces *Escherichia coli* O157: H7 contamination of lettuce and beef, but does not protect against recontamination. *Bacteriophage.* 2(3):178-185. <https://doi.org/10.4161/bact.22825>.
- Chen, J., and M. W. Griffiths. 1998. PCR differentiation of *Escherichia coli* from other Gram-negative bacteria using primers derived from the nucleotide sequences flanking the gene encoding the universal stress protein. *Lett. Appl. Microbiol.* 27(6):369-371. <https://doi.org/10.1046/j.1472-765x.1998.00445.x>.
- Cho, Y. I., J. I. Han, C. Wang, V. Cooper, K. Schwartz, T. Engelken, and K. J. Yoon. 2013. Case-control study of microbiological etiology associated with calf diarrhea. *Vet. Microbiol.* 166(3-4):375-385. <https://doi.org/10.1016/j.vetmic.2013.07.001>.
- Cho, Y. I., and K. J. Yoon. 2014. An overview of calf diarrhea-infectious etiology, diagnosis, and intervention. *J. Vet. Sci.* 15(1):1-17. <https://doi.org/10.4142/jvs.2014.15.1.1>.
- Choi, K. S., Kang, J. H., Cho, H. C., Yu, D. H., and Park, J. 2021. Changes in serum protein electrophoresis profiles and acute phase proteins in calves with diarrhea. *Can. J. Vet. Res.* 85(1):45-50. PMID: 33390652.
- Coffey, B., L. Rivas, G. Duffy, A. Coffey, R. P. Ross, and O. McAuliffe. 2011. Assessment of *Escherichia coli* O157: H7-specific bacteriophages e11/2 and e4/1c in model broth and hide environments. *Int. J. Food Microbiol.* 147(3):188-194. <https://doi.org/10.1016/j.ijfoodmicro.2011.04.001>.
- Coura, F. M., M. D. Freitas, J. Ribeiro, R. A. de Leme, C. de Souza, A. A. Alfieri, E. J. Facury Filho, A. U. Carvalho, M. X. Silva, A. P. Lage, and M. B. Heinemann. 2015. Longitudinal study of *Salmonella* spp., diarrheagenic *Escherichia coli*, Rotavirus, and Coronavirus isolated from healthy and diarrheic calves in a Brazilian dairy herd. *Trop. Anim. Health Pro.* 47(1):3-11. <https://doi.org/10.1007/s11250-014-0675-5>.

- Craciunas C., A. L. Keul, M. Flonta, and M. Cristea. 2012. Diagnostic tests for *Salmonella* strains targeting *hlaA*, *agfA*, *spvC* and *sefC* genes. *J. Environ. Manage.* 95:512-218. <https://doi.org/10.1016/j.jenvman.2010.07.027>.
- Dall Agnol, A.M., Lorenzetti, E. and Leme, R.A. et al. 2021. Severe outbreak of bovine neonatal diarrhea in a dairy calf rearing unit with multifactorial etiology. *Braz. J. Microbiol.* <https://doi.org/10.1007/s42770-021-00565-5>.
- Foster D. M. and G. W. Smith. 2009. Pathophysiology of diarrhea in calves. *Vet. Clin. North Am. Food Anim. Pract.* 25:13–36. <https://doi.org/10.1016/j.cvfa.2008.10.013>.
- Garcia A., J. A. Ruiz-Santa- Quiteria, J. A. Orden D. Cid., R. Sanz, M. Gómez-Bautista, and R. de la Fuente. 2000. Rotavirus and concurrent infections with other enteropathogens in neonatal diarrheic dairy calves in Spain. *Comp. Immunol. Microb.* 23(3):175-183. [https://doi.org/10.1016/S0147-9571\(99\)00071-5](https://doi.org/10.1016/S0147-9571(99)00071-5).
- Gifford, C. A., B. P. Holland, R. L. Mills, C. L. Maxwell, J. K. Farney, S. J. Terrill, D. L. Step, C. J. Richards, L. O. Burciaga Robles, and C. R. Krehbiel. 2012. Growth and development symposium: Impacts of inflammation on cattle growth and carcass merit. *J. Anim. Sci.* 90(5):1438-1451. <https://doi.org/10.2527/jas.2011-4846>.
- Green, M. R. and J. Sambrook. 2012. *Molecular Cloning: A Laboratory Manual*. 4th ed. Cold Spring Harbor Laboratory Press, New York, NY, USA.
- Harper, D. R. and M. C. Enrigh T. 2011. Bacteriophages for the treatment of *Pseudomonas aeruginosa* infections. *J. Appl. Microbiol.* 111(1):1-7. <https://doi.org/10.1111/j.1365-2672.2011.05003.x>.
- Hultgren, J., C. Svensson, D. O. Maizon, and P. A. Oltenacu. 2008. Rearing conditions, morbidity and breeding performance in dairy heifers in southwest. *Prev. Vet. Med.* 87(3-4):244-260. <https://doi.org/10.1016/j.prevetmed.2008.04.003>.
- Kertz, A. F. and H. Chester-Jones. 2004. Invited review: Guidelines for measuring and reporting calf and heifer experimental data. *J. Dairy Sci.* 87(11):3577-3580. [https://doi.org/10.3168/jds.S0022-0302\(04\)73495-5](https://doi.org/10.3168/jds.S0022-0302(04)73495-5).
- Kozasa, T., M. Tajima, I. Yasutomi, K. Sano, K. Ohashi, and M. Onuma. 2005. Relationship of bovine viral diarrhea virus persistent infection to incidence of diseases on dairy farms based on bulk tank milk test by RT-PCR. *Vet. Microbiol.* 106(1-2):41-47. <https://doi.org/10.1016/j.vetmic.2004.12.009>.

- Laureyns, J., S. Ribbens, and A. De Kruif. 2010. Control of bovine virus diarrhoea at the herd level: Reducing the risk of false negatives in the detection of persistently infected cattle. *Vet J.* 184(1):21-26. <https://doi.org/10.1016/j.tvjl.2008.11.014>.
- Lei, Y. M. K., L. Nair, and M. L. Alegre. 2015. The interplay between the intestinal microbiota and the immune system. *Clin. Res. Hepatol. Gastroenterol.* 39(1):9-19. <https://doi.org/10.1016/j.clinre.2014.10.008>.
- Müller, D., L. Greune, G. Heusipp, H. Karch, A. Fruth, H. Tschäpe, and M. A. Schmidt. 2007. Identification of unconventional intestinal pathogenic *Escherichia coli* isolates expressing intermediate virulence factor profiles by using a novel single-step multiplex PCR. *Appl. Environ. Microbiol.* 73(10):3380-3390. <http://doi.org/10.1128/AEM.02855-06>.
- Niu, Y. D., T. A. McAllister, Y. Xu, R. P. Johnson, T. P. Stephens, and K. Stanford. 2009. Prevalence and impact of bacteriophages on the presence of *Escherichia coli* O157:H7 in feedlot cattle and their environment. *Appl. Environ. Microbiol.* 73(5):1271-1278. <http://doi.org/10.1128/AEM.02100-08>.
- Niu, Y. D., Y. Xu, T. A. McAllister, E. A. Rozema, T. P. Stephens, S. J. Bach, R. P. Johnson, and K. Stanford. 2008. Comparison of fecal versus rectoanal mucosal swab sampling for detecting *Escherichia coli* O157:H7 in experimentally inoculated cattle used in assessing bacteriophage as a mitigation strategy. *J. Food Prot.* 71(4):691-698. <https://doi.org/10.4315/0362-028X-71.4.691>.
- Paton, A. W., and J. C. Paton. 1998. Detection and characterization of Shiga toxigenic *Escherichia coli* by using multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli* *hlyA*, *rfbO111*, and *rfbO157*. *J. Clin. Microbiol.* 36:598–602. <http://doi.org/10.1128/JCM.36.2.598-602.1998>.
- Quigley, J. D., Lago, A., Chapman, C., Erickson, P., and Polo, J. 2013. Evaluation of the Brix refractometer to estimate immunoglobulin G concentration in bovine colostrum. *J. Dairy Sci.* 96(2):1148-1155. <https://doi.org/10.3168/jds.2012-5823>.
- Rakoff-Nahoum, S., J. Paglino, F. Eslami-Varzaneg, and R. Medzhitov. 2004. Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell* 118(2):229-241. <https://doi.org/10.1016/j.cell.2004.07.002>.

- Raya R. R., P. Varey, R. A. Oot, M. R. Dyen, T. R. Callaway, T. S. Edrington, E. M. Kutter, and A. D. Brabban. 2006. Isolation and characterization of a new T-even bacteriophage, CEV1, and determination of its potential to reduce *Escherichia coli* O157:H7 levels in sheep. *Appl. Environ. Microbiol.* 72(9):6405-6410. <http://doi.org/10.1128/AEM.03011-05>.
- Rivas, L., B. Coffey, O. McAuliffe, M. J. McDonnell, C. M. Burgess, A. Coffey, R. P. Ross, and G. Duffy. 2010. In vivo and ex vivo evaluations of bacteriophages e11/2 and e4/1c for use in the control of *Escherichia coli* O157:H7. *Appl. Environ. Microbiol.* 76(21):7210-7216. <http://doi.org/10.1128/AEM.01530-10>.
- Rozema, E. A., T. P. Stephens, S. J. Bach, E. K. Okine, R. P. Johnson, K. Stanford and T. A. Mcallister. 2009. Oral and rectal administration of bacteriophages for control of *Escherichia coli* O157: H7 in feedlot cattle. *J. Food Prot.* 72(2): 241-250. <https://doi.org/10.4315/0362-028X-72.2.241>.
- Smith, B. P. 2009. *Large Animal Internal Medicine*. 4th ed. Mosby Elsevier, St. Louis, MO.
- Tyler, J. W., D. D. Hancock, H. M. Parish, D. E. Rea, T. E. Besser, S. G. Sanders and L. K. Wilson. 1996. Evaluation of 3 assays for failure of passive transfer in calves. *J. Vet. Intern. Med.* 10(5): 304-307. <https://doi.org/10.1111/j.1939-1676.1996.tb02067.x>.
- Santin, M., J. M. Trout, L. Xiao, L. Zhou, E. Greiner and R. Fayer. 2004. Prevalence and age-related variation of *Cryptosporidium* species and genotypes in dairy calves. *Vet. Parasitol.* 122(2):103-117. <https://doi.org/10.1016/j.vetpar.2004.03.020>.
- Santos, S. B., C. Carvalho, J. Azeredo and E. C. Ferreira. 2014. Population dynamics of a *Salmonella* lytic phage and its host: implications of the host bacterial growth rate in modelling. *PLoS one.* 9(7):e102507. <https://doi.org/10.1371/journal.pone.0102507>.
- Sheng, H., H. J. Knecht, I. T. Kudva and C. J. Hovde. 2006. Application of bacteriophages to control intestinal *Escherichia coli* O157:H7 levels in ruminants. *Appl. Environ. Microbiol.* 72(8):5359-5366. <https://doi.org/10.1128/AEM.00099-06>.
- Smith, G. W. 2009. Treatment of calf diarrhea: oral fluid therapy. *Vet. Clin. North Am. Food Anim. Pract.* 25(1):55-72. <https://doi.org/10.1016/j.cvfa.2008.10.006>.

- Smith, H. W., M. B. Huggins and K. M. Shaw. 1987. The control of experimental *Escherichia coli* diarrhoea in calves by means of bacteriophages. *Microbiology*. 133(5):1111-1126. <https://doi.org/10.1099/00221287-133-5-1111>.
- Sulakvelidze, A., Z. Alavidze and J. G. Morris. 2001. Bacteriophage therapy. *Antimicrob. Agents Chemother.*, 45(3): 649-659. <https://doi.org/10.1128/AAC.45.3.649-659.2001>.
- Svensson, C, K. Lundborg, U. Emanuelson and S. Olsson. 2003. Morbidity in Swedish dairy calves from birth to 90 days of age and individual calf-level risk factors for infectious diseases. *Prev. Vet. Med.* 58(3-4):179-197. [https://doi.org/10.1016/s0167-5877\(03\)00046-1](https://doi.org/10.1016/s0167-5877(03)00046-1).
- Volpato, A., A. A. Tonin, G. Machado, L. M. Stefani, G. Campigotto, P. Glombowsky, G. M. Galli, J. F. Favero and A. S. Silva. 2017. Gastrointestinal protozoa in dairy calves: identification of risk factors for infection. *Rev. MVZ Córdoba* 22(2):5910-5924. <https://doi.org/10.21897/rmvz.1027>.
- Windeyer, M. C., K. E. Leslie, S. M. Godden, D. C. Hodgins, K. D. Lissemore and S. J. Leblanc. 2014. Factors associated with morbidity, mortality, and growth of dairy heifer calves up to 3 months of age. *Prev. Vet. Med.* 113(2):231-240. <https://doi.org/10.1016/j.prevetmed.2013.10.019>.

Table 1 Sequence of primers and conditions used in PCR for genes of *Escherichia coli* and *Salmonella* spp. in feces samples of Holstein × Gyr crossbred female calves with diarrhea.

Gene	Protein	Primer sequences	amplicon size (pb)	No. of cycles	annealying temperature (°C)	Reference
<i>hlyA</i>	Hyper Invasibility	F- GCGAGATTGTGAGTAAAAACACC R- CTGCCCCGAGATATAATAATCG	413	35	63	Craciunas et al. (2012)
<i>uspA</i>	Universal stress protein	F-CCGATACGCTGCCAATCAGT R-ACGCAGACCGTAGGCCAGAT	884	30	70	Chen; Griffiths (1998)
<i>eae</i>	Intimin	F-GACCCGGCACAAGCATAAGC R-CCACCTGCAGCAACAAGAGG	384	35	60	Paton; Paton (1998)
<i>stx1</i>	Shiga toxin 1	F-ATAAATCGCCATTTCGTTGACTAC R-AGAACGCCCACTGAGATCATC	180	35	60	Paton; Paton (1998)
<i>est1a</i>	Thermoestable toxin A	F- CCTCTTTTAGCAGACACTGAATCATTG R-CAGGCAGGATTACAACAAAGTTCACAG	157	30	63	Mueller et al. (2007)
<i>hlyA</i>	Hemolysin	F- GCATCATCAAGCGTACGTTCC R- AATGAGCCAAGCTGGTTAAGCT	534	35	60	Paton; Paton (1998)

Table 2 Mortality, morbidity, case fatality rate and relapse rates of Holstein × Gyr crossbred female calves with (PHAGE; n=100) or without (CON; n=100) the addition of bacteriophage in milk replacer during pre-weaning (70 days).

Parameter	Group		P value
	CON	PHAGE	
Morbidity¹			
Diarrhea morbidity (%)	70% (70/100)	63% (63/100)	0.29
Respiratory disease morbidity (%)	29% (29/100)	26% (26/100)	0.66
Mortality²			
Diarrhea mortality (%)	2% (2/100)	1% (1/100)	0.57
Respiratory disease mortality (%)	3% (3/100)	5% (5/100)	0.47
Case fatality rate³			
Diarrhea fatality (%)	2.9% (2/70)	1.6% (1/63)	0.62
Respiratory disease fatality (%)	10.3% (3/29)	13% (5/26)	0.57
Relapse⁴			
Diarrhea relapse (%)	24.3% (17/70)	25.4% (16/63)	0.88
Respiratory disease relapse (%)	13.8% (4/29)	11.53% (3/26)	0.80

1 - Morbidity: number of ill calves/total number of calves; **2 - Mortality:** number of dead calves/total number of calves; **3 - Case fatality:** number of calves that died from diarrhea or respiratory disease/ total number of calves that presented diarrhea or respiratory disease; **4 - Relapse:** number of animals that got sick twice or more.

Table 3 Number of calves Holstein × Gyr crossbred female calves with diarrhea receiving (PHAGE; n=100) or not (CON; n=100) the addition of bacteriophage in milk replacer during pre-weaning (70 days).

AGE	Number of calves with diarrhea		P value
	CON	PHAGE	
1	2/48	2/47	0.9828
2	4/48	8/47	0.2025
3	9/48	2/47	0.0273*
4	9/48	8/47	0.826
5	16/48	5/47	0.0077*
6	15/48	6/47	0.03*
7	16/48	9/47	0.1165
8	25/48	12/47	0.008*
9	24/48	11/46	0.0089*
10	22/48	11/45	0.0312*
11	27/48	9/45	0.0003*
12	18/48	4/45	0.0012*
13	11/48	3/45	0.0285*
14	2/48	3/45	0.5932
15	1/48	1/44	0.9504
16	1/48	1/44	0.9504
17	1/48	2/44	0.5066
18	2/48	0/44	0.171
19	5/48	1/44	0.114
20	4/48	2/44	0.4623
21	2/48	2/44	0.9291
22	5/48	2/44	0.2887
23	7/48	3/44	0.232
24	5/48	0/44	0.0277*
25	3/48	5/44	0.3846
26	8/48	5/44	0.4657
27	3/48	4/44	0.6077
28	2/48	2/44	0.9291
29	6/48	2/44	0.1762
30	2/48	2/44	0.9291

*Statistical difference detected by Fisher's exact test (P<0.05)

Table 4 Performance parameters of Holstein × Gyr crossbred female calves up to 80 days of life with (PHAGE; n = 83) or without (CON; n = 92) the addition of bacteriophage in milk replacer during pre-weaning (70 days).

Parameter	Group				P value		
	CON		PHAGE		Group	Age	Group*Age ²
	Mean	SEM ¹	Mean	SEM ¹			
Thoracic perimeter (cm)	77.67	0.21	78.30	0.26	0.06 [†]	<0.001*	0.26
Withers height (cm)	77.29	0.18	77.44	0.13	0.51	<0.001*	0.81
Croup width (cm)	18.26	0.29	18.53	0.15	0.42	<0.001*	0.32
Weight (kg)	47.34	0.25	49.51	0.26	<0.001*	<0.001*	0.09 [†]
ADG (kg)	0.432	0.016	0.491	0.017	0.011	<0.001*	0.47

¹Standard error of the mean

²Interaction between group and age (days) and group

*Statistical difference identified by Tukey-Kramer test ($P < 0.05$)

[†]Tendency identified by Tukey-Kramer test ($0.05 < P > 0.1$)

Fig. 1 Period (days) of duration of the first diarrheic episode (A) and duration of diarrheic feces (B) in the neonatal period of Holstein × Gyr crossbred female calves with (PHAGE; n = 42) or without (CON; n = 50) the addition of bacteriophage in milk replacer during pre-weaning (70 days) Difference between groups by One-way ANOVA ($P < 0.05$) Bars indicate standard error of the means

Fig. 2 Number of Holstein $\times$ Gyr crossbred female calves with diarrheic and normal feces from days 3 to 13 of life with (PHAGE) or without (CON) the addition of bacteriophage in milk replacer during pre-weaning (70 days)

*Comparison between groups by Fisher's exact test ($P=0.05$)

Fig. 3 Average daily gain (ADG) at the end of the neonatal period (30 days), at the beginning of step-down weaning (42 days) and after weaning (80 days) of calves with (PHAGE; n = 83) or without (CON; = 92) the addition of bacteriophage in milk replacer during pre-weaning (70 days) Comparison between groups by Tukey-Kramer test (* $P<0.01$ ** $P=0.01$)

Fig. 4 Mean value of albumin (A) and total protein (B) of calves rotating with (PHAGE; n = 43) or without (CON; n = 41) the addition of bacteriophage in milk replacer during pre-weaning (70 days)

*Indicates statistical difference by the Tukey-Kramer test ($p < 0.05$)

Interaction between age (days) group and Tukey-Kramer test ($p < 0.05$)