Canadian Journal of Physics

Dynamo seeds from Gravitational Torsional Anomalies and De Sitter Magnetized Metrics

Journal:	<i>Canadian Journal of Physics</i>
Manuscript ID	cjp-2021-0243.R2
Manuscript Type:	Article
Date Submitted by the Author:	15-Oct-2021
Complete List of Authors:	de Andrade, L.C. Garcia; UERJ,
Keyword:	Gravitational anomaly, Teleparallelism, Emergent Torsion, De Sitter space, einstein-cartan
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Dynamo Seeds from Gravitational Torsional Anomalies and De Sitter Magnetized Metrics

By L.C. Garcia de Andrade¹

Abstract

Recently gravitational and Nieh-Yan chiral anomalies have been obtained in Riemann-Cartan spacetime [Class and Quantum Gravity 38 (2021)]. In this example, electrodynamics is encoded in the metric geometry. In this paper we follow this pathway by obtaining a class of deformed de Sitter metrics in teleparallelism. The existence of the unmagnetized De Sitter metric (DSMM) without axial anomalies is proved. Unified theories à la Einstein, Eddington and Schroedinger, called modified de Sitter metric (MDSM) presents some novel features: First, we show that a pure DS unmagnetized metric in T_4 does not induce gravitational anomalies. This is a motivation to study modifications of DSM. Nieh-Yan (NY) torsional anomaly in (DSMM) teleparallel T_4 geometry which is shown to vanish in all cases. Gravitational non-trivial anomalies are obtained from these metrics. Torsional anomaly, much used in condensed matter physics, does not vanish. From these deformed DSMM, we show that dynamo equation with torsional gradients sources is valid from class III of the metrics but is torsionless sourced in second class. We show that in the gravitational anomaly of new deformed de Sitter metric one may cancel the gravitational anomaly, by a proper choice of the metric function. The axial anomaly is obtained for some metric deformation as well. A simple deformation leads to the existence of the NY density in the case of DSMM. This would be class IV of DSMM.

Key-words: Gravitational anomaly, Nieh-Yan anomaly, Teleparallelism, Emergent Torsion

¹Departamento de Física Teórica - IF - UERJ - Rua São Francisco Xavier 524, Rio de Janeiro, RJ, Maracanã, CEP:20550-003 and Institute for Cosmology and Philosophy of Nature, Trg, Florjana 16, Krizvic, Croatia. e-mail:luizandra795@gmail.com

1 Introduction

Recently Laurila and Nissinen [1] made use of emergent torsion in superfluids and Nieh-Yan topology [2] with axial anomalies. Several other models of topological materials [3] have benefit from the application of quantum field theory [4] models such as gravitational anomalies and Pontryagin and Nieh-Yan densities so well-used in cosmological Higgs inflation [5]. Very recently, G Volovik [6] have used the idea of successful application of Cartan torsion [7] to chiral Weyl equations in semimetals to obtain the gravitational anomaly produced in neutral Weyl superfluids by the analog of gravitational instanton. Chiral torsional effects actually [8] have been applied to several examples, such as Casimir effects [9] axial electrodynamics with Hall anomalous effect [10], and in cosmological cyclic models [11]. Earlier an intense activity in analog models of gravity [12] in condensed matter physics, including superfluids [12] and non-Riemannian vortex acoustics have been presented in literature [13]. More recently, K Krasnov [14] has investigated Einstein-Cartan and chirality in nonlinear electrodynamics and gravity unified theory in terms of chiral effect [15]. Motivated by all these facts, we propose in this paper, a new generalization of De Sitter cosmological metric [16] in T_4 teleparallel geometry. This metric is constrained by T_4 vanishing of Riemann-Cartan (RC) curvature. Moreover, the investigations performed in this paper are useful to build unified theories a la Einstein [17]. In addition it possesses magnetic and electric fields encoded in the metric. The magnetic field appears connected to torsion, only when the chiral chemical potential does not vanish and only the magnetic monopoles do not exist. From the $\text{div} B = 0$ the $f(y)$ can be determined in terms of the y-component component of magnetic induction field. Since in this geometry, the RC full curvature vanishes, this allows us to obtain the Riemann curvature in terms of the torsion tensor components. We pointed out that, to obtain the gravitational anomaly, one needs not only the skew torsion component but also the trace torsion vector. The paper is organised as follows: In section 2 we present de Sitter modified metric and the chiral magnetic effect (CME) associated with it. Section 3 addresses topological densities associated to gravitational anomaly of the DSMM. In this same section we compute the NY and chiral gravitational asymmetry due to the non-conservation of current in this torsionful background T_4 spacetime. This is called a torsional anomaly. In section 4 we show that a simple second deformation of de Sitter metric, leads to the

existence of dynamo action with torsion source. Nevertheless, NY anomaly still vanishes. A new feature of this new gravitational anomaly, is that now, anomaly is cancelled by an oscillatory choice of new metric functions. In section 5 we show that the non-conformal original DSM may be deformed in order to obtain the NY and gravitational anomalies. Torsional anomalies however, seems not to exist. Conclusions and discussions are left to section 6.

2 CME and deformations of De Sitter T_4 metric

In this section we present deformations of de Sitter cosmological metric which in general relativity (GR) are conformally flat. This modification is given by

$$ds_I^2 = e^{2\Lambda t}[(dt + \mathbf{B} \cdot d\mathbf{x})^2 - dx^2 - f^2(y)(dy^2 + dz^2)] \quad (1)$$

Note that in this expression, if the $f(y)$ function, equals one, and the magnetic field $\mathbf{B}$ vanishes, the metric reduces to the GR usual cosmological de Sitter metric

$$ds^2 = e^{2\Lambda t}[dt^2 - dx^2 - dy^2 - dz^2] \quad (2)$$

written in terms of one forms basis components

$$e^{(0)} = e^{\Lambda t}(dt + \mathbf{B} \cdot d\mathbf{x}) \quad (3)$$

$$e^{(1)} = e^{\Lambda t}dx \quad (4)$$

$$e^{(2)} = e^{\Lambda t}f(y)dy \quad (5)$$

$$e^{(3)} = e^{\Lambda t}f(y)dz \quad (6)$$

which from the structure Cartan equations of differential forms calculus

$$R^i_j = R^i_{jkl}(\Gamma)e^k \wedge e^l = d\omega^i_j + \omega^i_k \wedge \omega^k_j \quad (7)$$

and

$$T^i = de^i + \omega^i_j \wedge e^j \quad (8)$$

where Γ is the RC spacetime connection. Here we adopt the teleparallel geometry where the Cartan connection vanishes

$$\omega^i_j = 0 \quad (9)$$

where $(i, j = 0, 1, 2, 3)$. From the second structure Cartan equation, the curvature forms R^i_j vanish and consequently all RC curvature tensor components vanish as well. Hence

$$R^i_{jkl}(T_4) = 0 \quad (10)$$

which is the teleparallel geometry condition. From our construction of the metric in terms of Cartan's calculus of differential forms we note that

$$T^1 = de^1 = \Lambda e^{\Lambda t} dt \wedge dx \quad (11)$$

which implies that there is another torsion term T^1_{01} given by

$$T^1_{01} = \Lambda e^{\Lambda t} \quad (12)$$

The remaining components of torsion 2-forms T^i are

$$T^0 = de^0 = e^{\Lambda t} [\partial_{[a} B_{c]} dx^a \wedge dx^c + (\partial_{[0} B_{a]} + \Lambda B_a)] dt \wedge dx^a \quad (13)$$

where indices $(a, c = 1, 2, 3)$. This last expression implies that the corresponding torsion tensor components are

$$T^0_{bc} = e^{\Lambda t} \partial_{[b} B_{c]} \quad (14)$$

Note from this expression, that there are three nonvanishing torsion independent components T^0_{12} , T^0_{13} and T^0_{23} which are linked to $\text{curl} B = J e^{-\Lambda t}$ and therefore represents Ampere-Maxwell law of electrodynamics with a decay factor, or damping of current, lending a geometrical character to torsion T_4 electrodynamics. A very important torsion term, comes from the second torsion component, that although does not contribute to the NY form, because is a trace torsion component, already gives a dynamo amplification [18] and a straightforward relationship to the magnetic field. This expression is

$$T^0_{0a} = [\partial_{[0} B_{a]} + \Lambda B_a] e^{\Lambda t} \quad (15)$$

Note that in this case, the torsion components represent dynamo amplification of the three components, as long as the factor Λ is positive. Note that if one considers that the magnetic field current is chiral, or expressed in terms of the chiral chemical potential μ_5

$$e^{\Lambda t} \partial_{[1} B_{2]} = \mu_5 \epsilon_{012} J^0 \quad (16)$$

and other corresponding currents components. We also

$$e^{\Lambda t} \partial_{[2} B_{3]} = \mu_5 \epsilon_{231} B^1 \quad (17)$$

and so on. Applying the divergent operator on both sides of this expression yields

$$e^{\Lambda t} (\text{div. curl} \mathbf{B}) = \mu_5 \text{div} \mathbf{B} \quad (18)$$

Since div. curl vanishes identically, one obtains $\text{div} B = 0$ or no monopole condition. One still has a CME in this framework of de Sitter modified teleparallelism. This is due to the fact that chiral chemical potential does not necessarily vanish. So our first new result is proved. Performing the analogous computation for T^2 one simply check that T^2 also vanishes by construction of MDSM. The T^3 case yields

$$T^3 = T^3_{ac} dx^a \wedge dx^c = e^{\Lambda t} f_y(y) dy \wedge dz \quad (19)$$

This expression, as we shall see in the next section, is fundamental for the existence of gravitational anomaly, given by the chiral current J_5 non-conservation chiral asymmetry. Therefore, the presence of $f(y)$ is a fundamental trick for the existence of this gravitational anomaly as we shall see in the next section. Now from the last expression one obtains the last component of torsion

$$T^3_{23} = e^{\Lambda t} f_y(y) \quad (20)$$

The sub-index y in $f(y)$ represents the derivative with respect to coordinate- y . One also notices, from this last expression that the torsion component, which shall be proved fundamental to gravitational anomaly, cannot be skew, since there are repeated indices. Hence, besides the presence of totally skew-symmetric torsion axial vector components, one needs also the presence of torsion trace vector.

3 Torsional chiral effects in modified teleparallel de Sitter metric and galactic dynamo seeds

In this section we shall compute the NY and gravitation anomalies for the MDS T_4 metric of last section, in terms of the magnetic field. We also show

that the in principle, arbitrary function $f(y)$ is not arbitrary and can be determined here. Let us now investigate the NY anomaly given by

$$N = \partial_i(\epsilon^{ijkl}T_{jkl}) \quad (21)$$

From the above torsion components one obtains

$$N = -\partial_3(T_{012}) + \partial_1(T_{023}) + \partial_2(T_{013}) \quad (22)$$

Therefore, the NY density vanishes. NY density, vanishes in general when we have no boundaries. Hence the three components of torsion which contribute to the NY topological torsion invariant is given by

$$T^0_{12} = e^{\Lambda t} \partial_1 B_2 \quad (23)$$

and

$$T^0_{13} = e^{\Lambda t} \partial_1 B_3 \quad (24)$$

and

$$T^0_{23} = e^{\Lambda t} \partial_2 B_3 \quad (25)$$

substituting these three last expressions into the expression for NY invariant N , and a simple calculus show that each term vanishes separately, because even being axial totally skew-symmetric they involve skew symmetrization of the partial second derivatives of the components of the magnetic field. Hence, also in this case, the NY form N vanishes. In the case of black holes in RC spacetime it was shown previously that NY form does not vanish. Now let us show that the presence of T^0_{23} torsion component, is fundamental to the gravitational anomaly. Nevertheless, it is not enough to avoid the vanishing of the gravitational anomaly. To achieve this task in a simple manner, we assume linear approximation of the Riemann curvature in terms of torsion. This takes into consideration that only in early universe, and inside black holes, torsion has a strong appreciable value which allows us to neglect products of torsion components. Riemann-Christoffel connections multiplied by torsion and by themselves may be neglected. This, results in a Riemann curvature tensor components coming from the teleparallel condition above as

$$R_{ijkl} = 2\partial_{[i}T_{j]kl} \quad (26)$$

Therefore, by substituting the T^0_{23} and T^1_{23} on the last expressions one obtains the following components of the Riemann tensor

$$R_{0123} = \partial_1 T_{023} - \partial_0 T_{123} \quad (27)$$

Since T_{123} vanishes, the only hope remains in T_{023} term. Therefore, the partial derivative of the T_{023} yields

$$\partial_1 T_{023} = e^{\Lambda t} \partial_1 \partial_{[2} B_{3]} \quad (28)$$

which by the expression of the chiral asymmetry current non-conservation gravitational anomaly yields

$$\partial_i J_5^i \sim \epsilon^{ijkl} R^j{}_{mn} R^{mn}{}_{kl} \sim \epsilon^{0123} R_{0123} R^{23}{}_{23} \quad (29)$$

which shows that the gravitational anomaly for the above modified de Sitter metric vanishes. Now let us write down the torsional anomaly provided by Laurilia and Nissinem as

$$\partial_i J_5^i \sim \epsilon^{ijkl} T_{aij} T^a{}_{kl} \quad (30)$$

Since torsion does not need to be a partial derivative, then one notices immediately that it does not vanish. The expression for T^0 form reads

$$\partial_i J_5^i \sim \epsilon^{0abc} [T^0{}_{0a} T^0{}_{bc}] \quad (31)$$

Let us now consider the above expression for T^0 emergent torsion form components. From Cartan structure equations one obtains

$$T^0{}_{0a} = e^{\Lambda t} [\partial_0 B_a + \Lambda B_a] \quad (32)$$

Note as we mentioned before, this shows that some trace torsion vectorial components are a sort of dynamo amplification of previous magnetic fields B_a . This could be consider as seeds. By rewriting this expression in the magnetogenesis format

$$B_x^{tors} = e^{\Lambda t} [\partial_0 + \Lambda] B_x^{seeds} \quad (33)$$

one obtains

$$B_x^{tors} \sim \Lambda B_x^{seeds} \quad (34)$$

which by considering $\Lambda = H \sim 100 \text{ km s}^{-1} \text{ Mpc}^{-1} = 10^{-9} \text{ s}^{-1}$, H being the Hubble constant. Therefore, from this last expression one obtains a galactic magnetic field of the order of $B_{gal} = B^{tors} = 10^{-6} G$ and a galactic dynamo seed of $10^3 G$. Going back to the chiral torsional anomaly, the other needed component of torsion to obtain this torsional anomaly is

$$T^0_{23} = e^{\Lambda t} \partial_x B_y \quad (35)$$

From expressions (14) and (15) allows one to obtain chiral torsional effect or anomaly, like in the condensed matter Volovik case [6] where

$$\partial_i J_5^i \sim \Lambda e^{\Lambda t} \epsilon^{0abc} [\partial_0 B_a + \Lambda B_a] \partial_b B_c \sim \mu_5 \frac{B^2}{L} e^{\Lambda t} + E \cdot B e^{\Lambda t} \quad (36)$$

Here, the $E \cdot B$ axial chiral term appears, as long as the Faraday equation $\text{curl} B \sim i\omega E \sim \mu_5 B$ is valid. Note that the conclusion is that this chiral asymmetry is proportional to the magnetic energy B^2 plus axial anomaly chiral term. One also notes that Hubble constant can be negative when the expansion is replaced by a big bounce, where a universe phase contracts instead of expanding. Then, one may consider the chiral torsional asymmetry could be cancelled even when the universe undergoes a big bounce. An interesting implication from expression (32) is that it yields a dynamo equation sourced by torsion, and besides the time derivative of this component leads to

$$\partial_0 T^1_{01} \sim e^{Ht} [2H \partial_t B_x + \partial_t^2 B_x] \quad (37)$$

Since H is small we have neglected terms of second order in H . Therefore the expression inside the brackets must decay so the first and second time derivatives of magnetic field are negative and B decays with expansion of the universe as well as torsion. Therefore as predicted by SenGupta et al [?] in the case of expanded braneworld universe torsion is highly suppressed by the end of inflation. In the particular case where the time derivative of torsion component vanishes one obtains

$$H e^{Ht} [H B_x + \partial_t B_x] = 0 \quad (38)$$

which is the dynamo equation in Friedmann like GR spacetime. Note that if torsion is non-constant But, let us now check what happens when the expression inside the square brackets is negative when the universe undergoes expansion. This means first that the H is positive and therefore expression on the LHS of equation (37) is negative as well and this correspond to a damping in the torsion component. Substitution of this expression into the chiral anomaly expression (36) we note that the expression for torsional anomaly vanishes in general relativity Riemannian manifold. If the universe has a bounce and H is negative we might find an expression where the torsion grows and

$$\partial_0 T^1_{01} \sim e^{Ht} [2H\partial_t B_x + \partial_t^2 B_x] \geq 0 \quad (39)$$

One then notes that H being negative we have an inequality that

$$2H\partial_t B_x \leq \partial_t^2 B_x \quad (40)$$

or $2HB_x \leq \partial_t B_x$, which certainly would imply that the magnetic field would grow in this bouncing contracting phase of the universe, ending our proof. Clearly torsion would also grow in this contracting phase of bouncing universe.

4 Gravitational anomaly from new deformations of De Sitter T_4 metric

Let us now deform or modify the de Sitter metric by introducing into the metric, a function $g = g(x, y, z)$ in place of $f(y)$. This deformation yields the new metric

$$ds_{II}^2 = e^{2Ht} [(dt + \mathbf{B} \cdot d\mathbf{x})^2 - f^2(x)dx^2 - g^2(x, y, z)(dy^2 + dz^2)] \quad (41)$$

of class II, where we have also introduced function $f(x)$ as the metric coefficient g_{11} . Now the new one-form basis are

$$e^{(0)} = e^{Ht}(dt + \mathbf{B} \cdot d\mathbf{x}) \quad (42)$$

$$e^{(1)} = e^{Ht}f(x)dx \quad (43)$$

$$e^{(2)} = e^{Ht} g(x, y, z) dy \quad (44)$$

$$e^{(3)} = e^{Ht} g(x, y, z) dz \quad (45)$$

From the structure Cartan equations of differential forms

$$R^i_j = R^i_{jkl}(\Gamma) e^k \wedge e^l = d\omega^i_j + \omega^i_k \wedge \omega^k_j \quad (46)$$

and

$$T^i = de^i + \omega^i_j \wedge e^j \quad (47)$$

where Γ is the RC spacetime connection obtained from torsion tetrad components in T_4 . From our construction of the metric in terms of Cartan's calculus of differential forms one obtains

$$T^1 = de^1 = e^{Ht} [f H dt \wedge dx] \quad (48)$$

which implies that another torsion term T^1_{01} is

$$T^1_{01} = H f(x) e^{Ht} \quad (49)$$

which yields

$$\partial_1 T^1_{01} = H f_x(x) e^{Ht} \quad (50)$$

This time, the torsion gradient source of dynamo equation above, does not vanish. The remaining components of torsion 2-forms T^i are

$$T^0 = de^0 = e^{Ht} [\partial_{[a} B_{c]} dx^a \wedge dx^c + (\partial_0 B_a + H B_a) dt \wedge dx^a] \quad (51)$$

where indices $(a, c = 1, 2, 3)$. This last expression implies that the corresponding torsion tensor component is

$$T^0_{ab} = e^{Ht} \partial_{[a} B_{c]} \quad (52)$$

Note from this expression that there are three nonvanishing torsion independent components T^0_{12} , T^0_{13} and T^0_{23} . They already generates a dynamo amplification [18] and a straight relationship with the magnetic field. This expression is

$$T^0_{0a} = e^{Ht} (\partial_0 B_a + H B_a) \quad (53)$$

Now T^2 and T^3 forms yields

$$T^3_{23} = e^{Ht} \partial_{[2} B_{3]} \quad (54)$$

and

$$T^3_{03} = e^{Ht} H g(x, y, z) \quad (55)$$

These two expressions are enough to obtain the two Riemann curvature components which are needed to find the now nonvanishing gravitational anomaly. Then

$$R_{0123} = -\partial_1 T_{023} = e^{Ht} \partial_1 \partial_{[2} B_{3]} \quad (56)$$

and the other component of the Riemann tensor reads

$$R_{2323} = -\partial_2 T_{323} = e^{Ht} g_{yy} \quad (57)$$

This time gravitational anomaly of T_4 de Sitter deformed metric does not vanish as can be seen from

$$\partial_i J_5^i \sim \epsilon^{0123} R^{01}_{23} R^{23}_{23} \quad (58)$$

which yields

$$\partial_i J_5^i \sim (g_{yy} - g_{zz}) e^{2Ht} \partial_1 \partial_{[2} B_{3]} \quad (59)$$

Note that in this case, anomaly could be cancelled by choosing the relation $g_{yy} = g_{zz}$. A simple solution of this PDE yields oscillatory behavior for function $g(y, z)$.

5 Nieh-Yan anomaly from deformations of non-conformal DSM T_4 metric

Let us now deform the non-conformal de Sitter metric, by introducing into the metric a function $f = f(x, y, z)$ to replace $f(y)$. This deformation yields the new metric

$$ds_{III}^2 = [dt + e^{Hx} B_a dx^a]^2 - e^{2H(t+x)} d\mathbf{x}^2 \quad (60)$$

of class III. In this section we shall prove that this metric, contrary to other deformations of classes I and II in previous sections, does

induce the NY anomaly. To achieve this task, let us build from this class III metric the one-form basis of Cartan's calculus

$$e^{(0)} = (dt + e^{Hx} \mathbf{B} \cdot d\mathbf{x}) \quad (61)$$

$$e^{(1)} = e^{H(t+x)} dx \quad (62)$$

$$e^{(2)} = e^{H(t+x)} dy \quad (63)$$

$$e^{(3)} = e^{H(t+x)} dz \quad (64)$$

which yields the following set of torsion 2-forms

$$T^0 = de^0 = T^0_{db} dx^d \wedge dx^b = (e^{Hx} [\partial_{[d} B_{b]} + x_{,d} B_b]) dx^d \wedge dx^b \quad (65)$$

where in this expression $x_{,a} = \partial_a x$. The other torsion forms are given by

$$T^a = de^a = T^a_{0b} dt \wedge dx^b = e^{H(t+x)} [\partial_0 B_b + H B_b] dt \wedge dx^b \quad (66)$$

From these last two expressions we obtain T^4 torsion components in tetrad form as

$$T^0_{db} = (e^{Hx} [\partial_{[d} B_{b]} + H x_{,d} B_b]) \quad (67)$$

and

$$T^a_{0a} = e^{H(x^b)} [\partial_{[0} B_{a]} + H B_a] \quad (68)$$

From expression (67) one obtains the NY form

$$N \sim \epsilon^{c0db} \partial_c T_{0db} \quad (69)$$

Substitution of the expression (67) into this N expression one obtains

$$N \sim e^{Hx} \epsilon^{c0db} H (\partial_d x) \partial_c B_d \quad (70)$$

where some terms have vanish due to symmetry between the skew-symmetric Levi-Civita symbol ϵ^{c0db} . Since the last term on the RHS of this expression is proportional to magnetic field B in the CME case, or $\text{curl} B = J_5 = \mu_5 B$, one may reexpress the NY anomaly in terms of chemical chiral potential as

$$N \sim e^{Hx} H (\partial_d x) \mu_5 B^d \quad (71)$$

So we may reach an important conclusion, namely that the NY anomaly vanishes when $x = \text{constant}$ surface or when dynamos are non-chiral. Other class called IV metric is such that

$$ds_{IV}^2 = [dt + e^{Hx} B_a dx^a]^2 - e^{2Ht} f^2(\mathbf{x}) d\mathbf{x}^2 \quad (72)$$

from which we easily show to possess gravitational anomaly but no NY anomaly. Several other DS metrics in T_4 with distinct new physical properties may be computed in a future work.

6 Conclusions

Previously Laurila and Nissinen have investigated the torsional Landau levels and geometric anomalies in condensed matter Weyl systems with coordinate dependent tetrads and Fermi velocities, momentum space geometry and low-energy weyl quasi-particles. With the universal coupling of tetrads and momentum, they obtained a chiral and axial anomalies by gravitational analogy. At low energy, they found background fields for fermions as RC spacetime, so common in real gravitation in Einstein-Cartan gravity with spin-spin contact interaction. On those backgrounds they were able to find out Nieh-Yan anomaly which appears in condensed matter with UV parameter. Moreover, they applied these ideas to superfluids, as we have done here following the Volovik idea of using gravitational anomaly analogs. In this paper NY anomaly was found to be very rare in the de Sitter class of metrics in teleparallelism discussed here. Several magnetized and unmagnetized metrics with implications to CME and chiral dynamos with torsion have been presented, in analogy to chiral dynamos in bouncing model recently presented by the author [19]. Their torsion and chiral chemical potential link has also been obtained in the context of cosmology. Several torsional chiral and axial anomalies in the T_4 DS framework have been presented in this paper. More detailed investigations were left for future publications. Recently, I was told about a very recent paper by Ze-Min Huang et al [20] where torsion, energy magnetization and thermal Hall effect have been addressed. In this paper, the teleparallel geometry has been used with fully implications to condensed matter physics. This paper lends more support for the interest in the research of the common foreground between cosmology and condensed matter applications. Important to say that in their reference, they also use the spin

connection ω^i_j free condition to obtain T_4 exactly as we did here in the cosmological and magnetogenesis frameworks. Actually, magnetism seems to be an important point of contact between microscopic and macroscopic worlds of condensed matter and cosmology. A method of encoding physical information into the metrics in teleparallelism presented recently by the author [21] has been used here throughout the paper.

7 Acknowledgements

I would like to thank G Volovik for many enlightning discussions on the subject of this paper. Financial support from University of State of Rio de Janeiro (UERJ) are grateful acknowledged.

References

- [1] S Laurila and J Nissenin, Torsional Landau Levels and geometric anomalies in condensed matter Weyl systems, ArXives: 2007.10682v2 [cond-mat. str-el] 2020. Phys Rev B.
- [2] The Nieh-Yan Anomaly: Torsional Landau Levels, Central Charge and Anomalous thermal Hall Effect, Z-M Huang, B Han, M Stone, arXive: 1911.00174. H T Nieh, ML Yan, Annals of Phys 138, 2 237 (1982). Garcia de Andrade, Teleparallel Pseudo-electromagnetic fields and pseudo-magnetic monopoles in topological materials, preprint (2021).
- [3] J L Manes, M Valle, M A Vazques-Mozo, Chiral torsional effects in anomalous fluids, arXiv: 2012.08449v1 (2020) JHEP in press.
- [4] F W Hehl, Phys Lett 36 A(1971) 225.
- [5] M Shaposhnikov , A Shkerin, I Timiryasov, S Zell, Higgs Inflation in Einstein-Cartan Gravity, Submitted to JCAP, arXiv: 2007.14978v1, (2020). Phys Rev Lett 126, 16130, (2021).
- [6] G E Volovik, JETP letters, Analog of gravitational anomaly in topological chiral superconductors, arXives: 2014.01020v1 (cond-mat.supr-con) March (2021).

- [7] E Cartan, Sur les equations de la gravitation dEinstein, (1922) Paris, Gauthier-Villars. E Cartan, Comptes Rendus (Paris), 174 (1922). E Cartan, Ann Inst Henri Poincare (1922). For a historical perspective see A Einstein and E Cartan, Letters on Absolute Parallelism (1987) Ed R Debever, Ed Princeton University Press.
- [8] S Imaki and Z Qiu, Phys Rev D 102, 016001 (2020).
- [9] Yago Ferreira and K Landsteiner, On chiral responses to geometric torsion, arXiv:2011.10535v2 (2020). Lan-Lan Gao, S Kaushik, D E Kharzeev and E J Philip, Chiral Kinetic theory of anomalous transport induced by torsion, arXiv: 2010. 07123v2 [cond-mat.mes-hall] 23 Nov 2020.
- [10] L Garcia de Andrade, Class and Quantum Gravity 18, No 18, 3907 (2001). Chiral anomalies and helical dynamos in non-Riemannian spaces, Can J Physics, in press (2020). L C Garcia de Andrade. Cosmic magnetism in modified theories of gravity, (2017) Editions europeennes universitaires.
- [11] L Garcia de Andrade, Eur Phys J C 78: 254 (2018) and Eur Phys J C 77: 401 (2017). L C Garcia de Andrade, Eur Phys J C 78: 530 (2018). N Mavromatos, private communication (2020). T Bossingham, N Mavromatos and S Sarkar, The role of temperature dependent string-inspired CPT violation backgrounds in leptogenesis and the chiral magnetic effect, arXiv:1810.13384. published in Eur Phys J C (2019).
- [12] G Volovik, JETP Letters 113, issue 8 (2021). G Volovik The Universe in a Helium Droplet, Oxford university Press (2003).
- [13] Garcia de Andrade, Phys Rev D 70 (2004) 064004.
- [14] K Krasnov, Formulations of General Relativity: Gravity, Spinors and Differential Forms (Cambridge monographs on Mathematical Physics) Cambridge UK (202).
- [15] Y Herfray, K Krasnov, C Scarini, Yuri Shtanov, Class Quantum Gravity 33 (2016) 235001.
- [16] W de Sitter, MNRAS 76, (9) : 699-728 (1916).

- [17] E Schroedinger, Structure of Spacetime, Cambridge University Press.
- [18] V Arnold and , Topological Methods in Hydrodynamics, (1989(**19** S Gera and S SenGupta, Invisible magnetic monopoles and spacetime geometry, arXiv: 2004.13083 (2020).
- [19] L Garcia de Andrade, Eur Phys J C 78: 254 (2018) and Eur Phys J C 77: 401 (2017).
- [20] Ze Min Huang, Bo Han and Xiao Qi Sun, Torsion, Energy Magnetization and Thermal Hall Effect (2021) arXiv: 2105.01600 v2.
- [21] L C Garcia de Andrade, Class and Quantum Gravity 38, (2021).