Bioline International

[HOME](#) [JOURNALS](#) [REPORTS](#) [NEWSLETTERS](#) [BOOKS](#) [SAMPLE PAPERS](#) [RESOURCES](#) [FAQ](#)

African Journal of Reproductive Health, ISSN: 1118-4841
Women's Health and Action Research Centre

African Journal of Reproductive Health, Vol. 7, No. 2, Aug, 2003 pp. 89-99

Gender Issues in the Prevention and Control of STIs and HIV/AIDS: Lessons from Awka and Agulu, Anambra State, Nigeria

Nkoli N Ezumah¹

¹Department of Sociology/Anthropology, University of Nigeria, Nsukka, Nigeria
Correspondence: 63 Nike Lake Expressway, Trans-Ekulu, P.O. Box 4235, Enugu. E-mail:
helitze@infoweb.abs.net Tel: 234 42 551980, 234 42 255301, 234 42 255671

Code Number: rh03026

ABSTRACT

The study examined perceptions, practices and norms underlying sexuality and gender relations that constrain the prevention and control of STIs and HIV/AIDS. It was carried out among the Igbo of Awka and Agulu in Anambra State of Nigeria. Both quantitative and qualitative methods were used for data collection. Findings indicate that cultural practices that encourage the establishment of sexual networks by men persist in the study communities. Some married women who are not able to achieve pregnancy with their husbands get involved in such networks to have children. Some parents who do not have male children encourage their unmarried daughters to have children out of wedlock so as to perpetuate the lineage. Inequality in gender relations and fear of repercussions constrain women from negotiating safe sex. Sexuality education is, therefore, necessary to improve the knowledge base, perceptions and sexual behaviours of the study communities. Women should be empowered to make informed decisions about sexuality and childbearing. Male responsibility in reproductive health should also be encouraged. (*Afr J Reprod Health* 2003; 7[2]: 89-99)

RÉSUMÉ

Questions de genre dans la prévention et la lutte contre les IST et le VIH/SIDA: leçons tirées d'Awka et d'Agulu, Etat d'Anambra, Nigeria. L'étude a examiné les perceptions, les pratiques et les normes qui sous-tendent la sexualité et les relations de genre qui limitent la prévention et la lutte contre les IST et le VIH/SIDA. L'étude a été menée auprès des Ibos d'Awka et d'Agulu dans l'Etat d'Anambra au Nigéria. Pour collecter les données, nous avons employé les méthodes quantitative et qualitative. Les résultats montrent que les pratiques culturelles qui encouragent l'établissement des réseaux sexuels par les hommes, demeurent toujours dans les communautés qui ont fait l'objet de l'étude. Certaines femmes mariées qui n'arrivent pas à devenir enceintes de leurs maris, deviennent impliquées dans de tels réseaux pour avoir des enfants. Certains parents qui n'ont pas d'enfants de sexe masculin encouragent leurs filles célibataires d'avoir des enfants hors mariage afin de prolonger la lignée. L'inégalité dans les relations de genre et la peur des repercussions obligent les femmes à ne pas négocier la sexualité sans danger. L'éducation sexuelle est donc nécessaire pour améliorer la base de la connaissance, les perspectives et les comportements sexuels des communautés qui ont fait l'objet de l'étude. Il faut donner le pouvoir aux femmes qui leur permettra de prendre des décisions en connaissance de cause sur la sexualité et l'accouchement. Il faut également encourager la responsabilisation de la part des hommes en matière de la santé reproductive. (*Rev Afr Santé Reprod* 2003; 7[2]: 89-99)

KEY WORDS: *Gender, sexuality, STIs, HIV/AIDS, prevention, safe sex*

INTRODUCTION

STIs constitute a reproductive health problem for men and women with many complications including infertility. It has been reported that 85% of primary infertility in women and 20% in men in sub-Saharan Africa is due to previous genital tract infections.¹ In the case of HIV/AIDS, about 60% of the 33 million infected people worldwide live in Africa.² In Nigeria, there has been a dramatic increase in the rate of HIV/AIDS infection from 1.8% in 1990 to 5.4% in 1999.³ In some states that constitute the hot spots of Nigeria the rate is 6-10%.⁴

Research has established a link between risky sexual behaviour, especially sexual networking, and the transmission of HIV/AIDS. About 80% of infections in Africa are as a result of heterosexual transmission.⁵ There is a higher rate of infection in women than men, and several factors contribute to this. Gender issues are of prime importance among such factors. Due to gender inequality and women's low socio-economic status, they are less likely than men to exercise control over the timing, frequency and conditions of sex.⁶⁻⁸ The vulnerability of women to STIs and HIV/AIDS is also aggravated by sexual exploitation, coercion, rape and commercial sex work. Cubis⁹ argues that without considering the impact of gender relations and strategies for dealing with women's low socio-economic status, the distribution of condoms and education will not be effective in

dealing with the spread of HIV/AIDS.

In addition, ideologies about sexuality tend to predispose men and women to indulge in risky sexual behaviours and unprotected sex. Women are often expected to be passive in sexual relations. Their sexual expression is assumed to be reactive or in response to male advances, whereas men's sexuality is assumed to be active. People's adherence to these notions about sexuality explains why a double standard of sexual morality persists in some societies.¹⁰ This has been identified as one of the predisposing factors to STIs in Nigeria.¹¹ The belief that women should be passive in sexual relationships also makes them to be ignorant and misinformed about the sexual behaviour of their husband and the consequences of such actions.¹²

Gender roles and cultural values related to marriage and childbearing also predispose men and women to STIs and HIV/AIDS. Marriage and childbearing provide social status and respectability for women in Nigeria as in other African countries.^{13,14} In fact, procreation is perceived as the major role of Igbo women in Nigeria.¹⁵ Consequently, the society tends to treat childless women with disdain. Although the increasing phenomenon of "outside wives" who serve as mistresses to married men who are still living with their "inside wives" was attributed to women's quest for materialism,¹³ such a development could also be related to women's desire to bear children. This paper examines how perceptions, practices and norms underlying sexuality and gender relations among the Igbo constrain men and women from taking appropriate measures to protect themselves as well as their partners from STIs including the HIV/AIDS.

THEORETICAL FRAMEWORK

The sexuality-gender framework propounded by Dixon-Mueller⁷ provides a useful theoretical orientation for the analysis of the effect of gender issues and sexuality in the prevention and control of STIs and HIV/AIDS. She identified four dimensions of sexuality-gender framework, namely, sexual behaviour, nature of sexual acts, social construction of sexuality, and nature of sexual drives and enjoyment.

The first dimension, which focuses on sexual behaviour, deals with the issue of timing, duration, number, choice and identity of sexual partners. The conditions that influence the initiation of sexual activities have important influence on peoples' subsequent attitudes and behaviour. In some societies, a double standard of morality exists with respect to gender differences in sexual behaviour. It has been pointed out that sexual relationships between older men and younger women are acceptable in many societies.¹⁷ Extramarital relationships by married men are also tolerated in Nigerian communities.^{16,17} Similarly, the practice of patronising commercial sex workers by married men is often pervasive.^{10,11}

The second aspect focuses on sexual acts. These include the nature, frequency and conditions of choice of sexual partners as well as the types of sexual practices individuals engage in. Dixon-Mueller⁷ argues that because heterosexual practices tend to be more dominant in societies, the existence of homosexual practices might appear invisible in apparently heterosexual populations. Moreover she raises the issue of certain sexual practices such as oral and anal sex, masturbation, use of devices to enhance sexual pleasure that may have implications for disease transmission and prevention. That dimension also addresses the implications of gender-based violence and coercive sex for disease transmission and unwanted pregnancies.

The third dimension, which focuses on the social construction of sexuality, deals with the cultural interpretations and meanings that are ascribed to sexual thoughts and behaviours. These reflect the beliefs and ideas people have about "the nature of the body". Cultural taboos and sanctions are established to control undesirable sexual behaviours of males and females. However, it has been observed that due to cultural inhibitions, the "culture of silence" in sexuality has persisted in some societies among couples and between parents and children.¹⁸ In particular, women's compliance with their expected norms in sexual relationships (i.e., passivity in sexual relationships) is often responsible for their inability to participate in meaningful communication about sexuality. Consequently, some women dwell on misinformation and ignorance about the sexual behaviour of their partners and the likely consequences of such actions. Concomitantly, they are not able to protect themselves from infection especially if their partners indulge in multiple sexual networking.^{12,16} The lack of communication about sexuality is aggravated in some societies like Nigeria by the use of poignant language that stigmatises women as disease vectors.¹

The fourth dimension refers to sexual drives and pleasures. It deals with how individuals perceive their sex drives and the meanings they attribute to sexual pleasure. The perception that sexual pleasure is desirable for males predisposes them to early involvement in sexual experience. This often starts during adolescence. In the case of women, the expression of sexual pleasure is hardly thought of. However, for many Nigerian women the consideration of sexuality as a source of pleasure still remains beyond imagination.¹⁴

Due to the interrelationship between gender, sexual behaviours and practices that enhance the transmission of STIs and HIV/AIDS as demonstrated in this theoretical orientation, one joins in the call for a better understanding of sexual behaviour as well as its determinants in different cultural settings.^{9,19} Such information would be indispensable in ensuring that culturally specific and appropriate programmes are instituted to bring about desired positive changes in sexual behaviour as a measure towards prevention and control of STIs and HIV/AIDS.

MATERIALS AND METHODS

The study was conducted in two Igbo communities in Anambra State of Nigeria, namely, Awka and Agulu. Awka is the capital city of the state. The 1991 National Census puts the population of Awka at 58,225 persons, comprising 28,335 males and 29,890 females. On the other hand, Agulu is a rural community situated about 30km from Awka. The 1991 National Census puts the population at 49,310 persons, comprising 23,573 males and 25,737 females.

Data were collected from five randomly selected communities - three in Awka (Ifite, Amenyi and Ezioka) and two in Agulu (Obeagu and Nkita-aku). Eight and six enumeration areas (EAs) (based on the 1991 census figures) were randomly selected from Awka and Agulu respectively. Both quantitative and qualitative methods were used for data collection. Survey instruments were administered to a total of 1000 randomly selected respondents, 600 from Awka and 400 from Agulu. The distribution pattern was based on the proportion of the two communities in relation to the total population. The sample distribution ratio (3:2 in favour of women) was a reflection of the male:female ratio in the general population of the two communities.

Eighteen focus group discussions (FGDs), made up of 12 in Awka (comprising six female and six male groups) and six in Agulu (comprising three male and three female groups) were administered. The participants were purposively selected to fall within three age groups -15-24 years, 25-44 years and 45 years and above. Other criteria used for the selection of FGD participants included education and occupation. Due to greater migration of young men from Agulu to Awka and other urban areas, it was difficult to get up to six male focus groups of the required age, education and occupational categories at the time of investigation. We therefore had to scale down the number of FGDs in Agulu to three male and three female groups to enhance effective gender comparison of responses. Each focus group comprised 8-10 participants. Sixteen respondents were selected for the in-depth interviews, eight in Akwa (four men and four women) and eight in Agulu (four men and four women). The participants included leaders of organisations, community leaders, youth leaders, teachers, public servants, farmers and housewives.

RESULTS AND DISCUSSION

Cultural Practices Associated with Childbearing that Predispose Women to STIs and HIV/AIDS

Although the Igbo frown at premarital sex,¹⁶ the study identified circumstances that the community approves of women having children out of wedlock. About 63.4% of the survey respondents indicated that where a couple does not have a male child, there could be a tendency to coerce one of their daughters to remain unmarried and have male

children for the family in order to perpetuate the lineage. Also, 20.6% indicated that if a man dies without having children from his wife, or if he is impotent or unable to father a child, his wife would be encouraged to have a lover with whom she can have children. Both male and female respondents expressed similar views on these issues ([Table 1](#)).

The establishment of sexual networks by widows and childless women to have children was corroborated by findings from the FGDs and IDIs. A male community leader at Awka referred to the practice as "taking shelter". He explained that through this practice the woman is trying to save herself from the embarrassment and scorn that childless women usually experience in the Igbo society. One of the problems associated with the practice of "taking shelter" was, however, indicated by a respondent as follows:

After going to take shelter, she would continue to have sexual relationship with her husband so that if she became pregnant it would appear as if he was responsible for it, even though she knew where the pregnancy came from.

The practice of having multiple sexual partners in order to have children and of couples coercing their unmarried daughters to bear children for their natal families expose women and girls to the risk of contracting STIs including HIV/AIDS. This therefore supports the finding that HIV/AIDS seroprevalence is higher among women of childbearing age in Nigeria.⁷

Perceptions of People's Vulnerability to STIs

Results show that 81.5% of the survey respondents held the view that only promiscuous women and commercial sex workers are vulnerable to STIs. Only 14.2% indicated that any woman who has unprotected sex is at risk of contracting STIs. Male and female respondents held similar views irrespective of their marital status ([Table 2](#)). It is important to note that most of the responses from the IDIs and FGDs reflected the ideas expressed by survey respondents. For example, male respondents in the FGDs (25-44 years) at Awka pointed out that only commercial sex workers and unmarried promiscuous women are at risk of contracting STIs.

These findings demonstrate that ignorance and misinformation about sexual relationships are still serious issues to be addressed in any effort to prevent STIs and HIV/AIDS in these communities. This is because married women who might not be promiscuous but whose husbands might be having unprotected sex with multiple partners are also at risk of contracting STIs. In fact, it has been pointed out that the single most important risk factor for women in the transmission of HIV/AIDS in sub-Saharan Africa is being married.²⁰

However, 81.6% of male respondents rightly indicated that any man who patronises

commercial sex workers or has many sexual partners is at risk of contracting STIs. To buttress that point, a male respondent in an FGD (45 years and above) at Agulu said:

Men who travel frequently are also at risk because they tend to make love with women before coming home.

PERCEPTIONS OF THE USE OF CONDOM BY MARRIED COUPLES

Some 62.6% of the respondents agreed that married couples could use condoms ([Table 3](#)) to avoid unwanted pregnancies and for child spacing. Among those who disagreed that couples should use condoms, 52.6% said married couples should accept the number of children God has given them because condoms are meant for unmarried people who desire to avoid pregnancy. About 31% expressed their fears that the use of condoms could create such problems as infection, bleeding, barrenness and cancer of the womb for women. The remaining 16.4% indicated that if women know their safe periods they and their partners would be in a position to practice periodical abstinence as a natural method of family planning.

When women were asked if they would discuss the use of condoms with their husband if they suspect that he indulges in extramarital affairs, only 54% of the married women said they would do so. Several reasons were given by those who said they would not discuss the use of condoms with their husband even when they suspect that he has multiple sexual partners. About half (49.6%) said such a discussion would encourage them to have outside lovers, 24.8% said they did not know about the use of condoms, and 12.4% said their husbands would not listen to them and that it might even worsen the situation. Seven per cent said it is not their business to make such suggestion while 6.2% said it is immoral to use condoms. These responses also highlight the fact that fear of repercussions constrains women from negotiating safe sex practices.

The views expressed by participants at the FGDs and IDIs on the use of condoms reflected both inter-generational gap and gender differences. A female vice principal (above 50 years) in Awka said it is all right for a woman to advise her husband to use condoms if she finds out that he is unfaithful. At the same time, she quickly pointed out thus:

In the real sense, it is not wise for a woman to advise her husband to use condoms, because that amounts to providing support for his waywardness.

Similarly, a retired female headmistress (above 60 years) in Agulu said although a woman could suggest the use of condoms to her husband, in practice men are those who would be expected to bring up such issues for discussion because of their constant urges for sex.

Some of the males in the IDIs, both at Agulu and Awka, were opposed to the idea of a woman suggesting the use of condoms to her husband regardless of the circumstances. For example, a male youth leader at Agulu said:

A woman should not say so with her mouth. It is the responsibility of the man. It is the man who should put up such a suggestion for discussion because it is the man's prerogative, not that of the woman.

It was only in the male FGDs (15-24 years) in both communities that there was a consensus that women could suggest the use of condoms to their husbands. In fact, one of them indicated:

If the man does not know about condoms but the wife knows, she should let him know that it is important to use it.

On the other hand, a male in one of the FGDs (25-44years) at Awka insisted:

It is not good for any married woman to do that because it is the man who is marrying her. She is not marrying the man.

Similarly, a male respondent in an FGD (25-44 years) at Agulu said:

A woman has no authority to do anything like that. It is the duty of a man. A woman has no right to ask the husband to use condoms. It only concerns the man.

These responses show that male control is a fundamental problem confronting women in exercising their rights to protect themselves from infection. The fact that the opposing views were expressed by some men and women clearly reflect the prevailing notion that males should have control of heterosexual relationships. For many of them the issue was who has the prerogative to control decision-making in sexuality. This finding corroborated reports from other studies, which identified lack of power as a serious impediment to women's ability to negotiate safe sex.^{21, 22}

People's Perception of Whose Responsibility it is to ensure that one does not contract STIs

A majority of the respondents (86.4%) indicated that both men and women are responsible for ensuring that one does not contract STIs. Only 10.7% indicated that it is the responsibility of men, 2.7% said it is the responsibility of women, while 2% had no clear views on the issue. These views were supported by responses from the IDIs and FGDs. A male community leader at Awka argued:

A man may contract STIs and extend to his wife, and a woman may contract them and extend to the husband. So it is their joint responsibility to ensure that they do not contract that.

A male teacher in the IDI at Awka pointed out:

If the man takes adequate care of his wife, gives her due respect, abstains from meeting with other women outside, the wife is likely to remain faithful in the home. It is the man that should care for the woman.

Similarly, women at the FGD for those 45 years and above at Agulu pointed out that if a husband and wife are able to avoid outside lovers, they would have no problems of contracting STIs.

Some respondents said it is the responsibility of the man to ensure that he and his wife do not contract STIs. Sixty eight per cent of the women and 55.6% of the men who gave this response attributed their view to the prevalence of male promiscuity. A female respondent in one of the FGDs (15-24 years) at Awka added that males tend to be promiscuous because "there is no punishment for them for flirting". These responses underscore the need to emphasise male responsibility in ensuring the sexual health of their partners.

Responses by those who said it is the responsibility of women to ensure that couples do not contract STIs demonstrate how the prevailing language of sexuality stigmatises women. About 28.6% of the men and 16.7% of women perceived women as disease vectors. Also, more men (35.7%), compared to women (16.7%), felt that it is the responsibility of women to monitor their husbands' activities in order to know whether they are indulging in extramarital affairs. Similarly, a male youth leader in the IDI at Agulu argued that it is the responsibility of women (to ensure that couples do not contract STIs) because they tend to be promiscuous. He emphasised:

You may be saying that you have a wife, not knowing that she has outsmarted you. If the woman says the home will be good, it will be.

Perceptions on whether it is Appropriate for Women to Refuse to have Sex with their Husbands when they do not desire it, or when they feel that it is not Safe

Findings reveal that 81.6% of respondents agreed that it is appropriate for women to refuse to have sex with their husband if they do not desire it. They said the woman might be indisposed due to ill health, she may be weak, pregnant, menstruating or still weaning an infant (78.3%). Others said she could refuse if she knows that her husband is having an extramarital affair (10.4%), she might be unhappy with the husband especially if he is not taking good care of her (7.0%), if the woman does not want to be pregnant (3.3), or if she

is fasting or meditating. The fact that only 10.4% raised the issue of fear for disease transmission shows that many of the respondents are still ignorant of the adverse consequences of the sexual behaviours of their partners. This highlights the gravity of the danger of disease transmission that risky sexual behaviours of husbands could pose on their wives. Similar responses were obtained from IDIs and FGDs as with the questionnaire survey.

Some of the respondents said a woman cannot refuse to have sex with her husband because they believe the males should have control over female sexuality and that women should remain passive in sexual relationships. A male youth leader in the IDI at Agulu said:

It is better for any woman that refuses to have sex with her husband to go back to her father's house.

Similarly a female Christian woman leader in Agulu said if a woman refuses to have sex with her husband and the later starts flirting she is to be blamed for his action. A male participant in one FGD (45 years and above) at Awka also said:

It is not the right thing to do. Women should be ready all the time.

Even a female in the FGD for the 45 years and above group at Agulu said:

Wives should be ready to answer their husbands whenever they call on them.

Possible Consequences of Refusing to have Sex with the Husband

Majority of the respondents said the man would be angry with his wife if she refuses to have sex with him, he may also beat her, reject her food, refuse to provide financial support for her, and he may start having extramarital affair. These responses were consistent among both men and women and among the married and unmarried ([Table 4](#)). The responses were also corroborated by participants in the FGDs and IDIs. Participants at the female and male FGDs (45 years and above) in Awka and Agulu specifically emphasised that a husband who has been denied sex by his wife would become suspicious. This could lead to fighting and quarrelling, making the man to deny her of housekeeping allowance. One of the females pointed out that Arefusal is giving the husband a license to seek an outside lover.

A female vice principal in the IDIs at Awka responded:

Do you need to ask? Some beat their wives, force them into having sex...

A female leader of a Christian women organisation in the IDIs at Agulu also emphasised that refusal of sex by a wife could lead to trouble in the home. A male teacher at Awka pointed out:

If she refuses her husband sex how can she stop him from going out?

These responses demonstrate that fear of repercussions still poses a serious constraint to women's ability to exercise their rights in negotiating sexual relationships with their spouses. There was a consensus among both males and females, indicating group norms that emanated from their socialisation, a situation in which female passivity as well as male control in sexuality is still predominant. The persistence of such views is a serious impediment to social change. This consequently continues to pose a threat to the prevention of STIs and HIV/AIDS among married women in the study communities.

Respondents' Perceptions about Categories of Women who would be able to Refuse Sex with their Husbands when they Suspect that it is not Safe

Majority (81.3%) of the survey respondents felt that mostly breadwinners, especially educated and highly placed women who know the implication of STIs, would be able to refuse sex with their husbands when they suspect that it is not safe ([Table 5](#)). The importance of economic independence in enhancing women's ability to be assertive and to refuse sex when it is not safe was also highlighted in the FGDs and IDIs. A female teacher in the IDI at Awka indicated that women who cannot refuse sex are immature wives who do not know their rights. She identified other categories such as illiterates, the jobless who are dependent on their husbands, and those who could be easily intimidated by their husbands among those who cannot refuse their husbands' sexual advances. Female participants in the FGDs (15-24 and 25-44 years) at Agulu and Awka emphasised the issue of fear and intimidation. One of them said:

The inability of women to refuse sex is due to fear of violence on their husbands' part because some men beat fear into their wives. So, because of fear the women may not voice out even when they suspect that their men may be infected.

CONCLUSION AND RECOMMENDATIONS

This paper makes some important contributions to knowledge and research in reproductive health issues. It identified the cultural practices, perceptions and norms about gender relations and sexuality that influence the sexual behaviours of men and women, thus increasing their chances of contracting STIs including HIV/AIDS in the communities studied. Findings from the study can be summarised as follows:

- In their effort to have male children and perpetuate their lineage, some parents in

the study communities encourage their unmarried daughters to bear children for them outside wedlock.

- Some married women who cannot achieve pregnancies with their husbands establish sexual networks (taking shelter) in order to fulfil their perceived gender roles of childbearing.
- Due to ignorance and misconception, a majority of the respondents perceived that only promiscuous women and commercial sex workers are vulnerable to STIs.
- A majority of the respondents felt that condom is only meant for preventing pregnancy, consequently they oppose its use for religious and moral reasons.
- Married women who are of the low socioeconomic status cannot suggest the use of condom to their husband or refuse his sexual advances because of the fear of repercussions.
- Male dominance over female sexuality is a serious impediment to achieving social change in peoples' attitudes to sexuality and sexual behaviour.
- Male sexual promiscuity is still pervasive because it is condoned in the communities.
- Lack of inter-spousal communication arising from the 'culture of silence' in sexuality is perpetuated by women's adherence to the notion that females ought to be passive in sexual relationships.
- Economic empowerment of women is very important if their ability to exercise their sexual rights is to be achieved.

These findings raise important policy issues that must be addressed in any effort to prevent the spread of STIs and HIV/AIDS. The intervention strategies suggested are two-fold. The first entails the provision of culturally specific and appropriate sexuality educational programmes in these communities to improve their knowledge and perceptions of sexual behaviours. Such programmes should highlight the reproductive health hazards involved in the harmful cultural practices that encourage the involvement of men and women in risky sexual behaviours. Specifically, women should be empowered to make informed decisions about their sexuality and childbearing. Male responsibility in reproductive health should also be encouraged in order to promote the sexual health of men and the wellbeing of their partners.

The second is the need to embark on empowerment strategies that would deal with issues of male dominance in sexual relationship, women's passivity and the fear of reprisals, which arise as a result of women's low socio-economic status. Women should be educated on the need to improve their self-image, they should be given opportunities for skills training and assisted to engage in income generating activities that would reduce their dependence on men. If women are empowered economically they will be better able to overcome the fear of reprisals when they need to take decisions on issues that affect their sexuality.

ACKNOWLEDGEMENT

The author acknowledges the support of the John D. and Catherine T. MacArthur Foundation, which provided fund for the project under the Fund for Leadership Development (FLD) in Nigeria. The support of field staff in data gathering is also acknowledged.

REFERENCES

1. Olukoya Adepeju. Reproductive tract infections in Nigeria. In: Jennifer Grant and Diana M Measham (Eds.). *Reproductive Tract Infection: Lessons Learned from the Field*. New York: The Population Council, 1996, 21-22.
2. UNAIDS/WHO. AIDS Epidemic Update. December 1999.
3. National AIDS and STDs Control Programme (NASCP). HIV/syphilis sentinel seroprevalence survey in Nigeria, Abuja, Nigeria, 1999.
4. Bawakira C. Role of CSOs in massive awareness raising campaign (MARC): adolescents and young people. Paper presented at a dialogue on comprehensive sexuality curriculum organized by Action Health Incorporated, at Sheraton Hotel and Towers, Abuja, Nigeria, October 26, 2001.
5. United Nations. *World Population Monitoring*. New York: United Nation's Department of International Economic and Social Affairs, 1990.
6. Rivers Kim and Aggleton Peter. *Men and the HIV Epidemic*. New York: UNDP, 1999.
7. Dixon-Mueller Ruth. The sexuality connection in reproductive health. In: Sondra Zeidenstein and Kirsten Moore (Eds.). *Learning About sexuality: A Practical Beginning*. New York: The Population Council, 1996, 137-157.
8. Hankins Catherine. Sexuality issues, reproductive choices and gynaecological concerns affecting women living with HIV. In: Lorraine Sherr Catherine (Eds.). *AIDS as a Gender Issue: Psychological Perspectives*. London: Taylor and Francis, 1996, 1-15.
9. Cubis Jacinta. Australia's international response to HIV/AIDS. *Development Bulletin* 2000; 52: 26B29.
10. Havanon Napapon. Talking to men and woman about their sexual relationships: insights from a Thai study. In: Sondra Zeidenstein and Kirsten Moore (Eds.). *Learning About Sexuality: A Practical Beginning*. New York: The Population Council, 1996, 110B118.
11. Orubuloye IO. Patterns of sexual behaviour of high risk populations and the implications for STDs as HIV/AIDS transmission in Nigeria. In: Richard G Parker and John H Gagnon (Eds.). *Conceiving Sexuality*. New York: Rutledge, 1995, 235-246.
12. Gammage Sarah. Women's role in household decision-making: a case study of Nigeria. International Center for Research on Women, Washington DC, USA, 1997.
13. Karanja WW. Outside wives and inside wives in Nigeria: a study of changing

- perceptions in marriage. In: David Parkin and David Nyamwaya (Eds.). *Transformations of African Marriage*. Manchester: Manchester University Press, 1987, 247.
14. Grace Osakue, Bene Madunagu and Hajira Usman (Eds.). *Voices: Findings of a Research into Reproductive Rights of Women in Nigeria*. Benin City: International Reproductive Rights Research Action Group (IRRAG), 1995.
  15. Amadiume Ifi. *Male Daughters, Female Husbands: Gender and Sex in an African Society*. London: Zed Books, 1987.
  16. Ezumah Nkoli N. Perceptions of sexuality and gender relations among, and implications for the reproductive health of men and women: selected findings from Awka and Agulu, Anambra State, Nigeria. In: Elaine Salo and Helen Moffet (Eds.). *Associate Publication*. Cape Town: University of Cape Town, 2000, 30-62.
  17. Aral SO. Sexual behaviour as a risk factor for sexually transmitted diseases. In: Adrienne Germain, King K Holmes, Peter Piot and Judith W Wasseerheit (Eds.). *Reproductive Tract Infections: Global Impact and Priorities for Women's Reproductive Health*. New York: Plenum Press, 1992, 185-198.
  18. Smyke Patricia. *Women and Health*. London: Zed Books Ltd., 1991.
  19. Carael Michel. Sexual behaviour. In: John Cleland and Benoit Ferry (Eds.). *Sexual Behaviour and AIDS in the Developing World*. London: Taylor and Francis Ltd., 1995, 75-123
  20. Moodie Rob. Should HIV be on the development agenda? *Development Bulletin* 2000; 52: 6-8.
  21. Richardson Diane. AIDS education and women: sexual and reproductive issues. In: Peter Aggleton, Peter Davies and Graham Hart (Eds.). *AIDS: Individual, Cultural and Policy Dimensions*. London: The Falmer Press, 1990, 169-179.
  22. Sherr Lorraine. Tomorrow's era: gender, psychology and HIV infection. In: Lorraine Sherr, Catherine Hankins and Lydia Bennett (Eds.). *AIDS as a Gender Issue: Psychological Perspectives*. London: Taylor and Francis, 1996, 16-45.

Copyright 2003 - Women's Health and Action Research Centre

Contact: [African Journal of Reproductive Health](#)

The following images related to this document are available:

Photo images

[\[rh03026t1.jpg\]](#) [\[rh03026t2.jpg\]](#) [\[rh03026t3.jpg\]](#) [\[rh03026t4.jpg\]](#) [\[rh03026t5.jpg\]](#)

[ABOUT BIOLINE](#)

[NEWS](#)

[FORUM](#)

[MAILING LIST](#)

[EMAIL BIOLINE](#)

[HOME](#)

powered by **<XML>@CRIA**

Table 1   **Circumstances in which the Society Approves of a Woman Having Children out of Wedlock by Gender (%)**

Response	Male (N = 423)	Female (N = 576)	Mean
Only when there is no male child in the family and her parents encourage her to stay back and have children	65.2	62.2	63.4
If the husband is dead or impotent/unable to father a child	17.7	22.7	20.6
Under no circumstances	9.5	8.3	8.9
Don't know	4.5	4.9	4.7
If the husband suffers from a mental problem	4.5	4.9	4.7
Where the woman is advanced in age and has no marriage prospects	1.4	0.5	0.9
It depends on what the woman wants	1.2	0.9	1.0
Total	100	100	100

Number of missing observation = 1

Table 2 Perceptions of Women's Vulnerability to STIs (%)

Response	A (Gender)		Mean	Married N = 501	B (Marital status)		Mean
	Male N = 423	Female N = 576			Separated N = 48	Never Married N = 451	
Promiscuous women and those involved in commercial sex work	82.3	80.9	81.5	79.0	89.6	83.4	81.5
Any woman who has unprotected sex, who has multiple sexual partners, or whose husband is promiscuous	13.2	14.9	14.2	16.6	8.3	12.1	14.0
Don't know	1.7	0.7	1.1	2.0	0.0	0.7	1.3
Women with low immunity	1.4	1.1	1.3	1.8	0.0	1.8	1.7
Dirty women	1.2	2.1	1.7	0.6	2.1	1.6	1.1
Lesbians	0.2	0.7	1.1	0.0	0.0	0.4	0.2
Total	100	100	100.0	100	100	100	100

In (A), number of missing observation = 1

"Separated" refers to all previously married who are separated, divorced or widowed

Table 3   **Should Couples Use Condoms? (%)**

Response	Married (N = 501)	Separated (N = 48)	Never married (N = 451)	Mean
Yes	63.5	70.8	60.8	62.6
No	34.5	27.1	29.7	32.0
Don't know	2.0	2.1	9.5	5.4
Total	100	100	100	100

"Separated" refers to all previously married who are separated, divorced or widowed.

Table 4 Possible Consequences for Refusing to have Sex with the Husband (%)

Response	A (Gender)		Mean	B (Marital status)			Mean
	Male N = 423	Female N = 576		Married N = 501	Separated N = married	Never N = 451	
He will be angry with her, may beat her, reject her food and refuse to provide financial support for her	61.7	60.4	61.0	61.6	62.5	60.1	61.0
He may start having extramarital affairs	20.3	24.7	22.8	22.8	25.0	22.6	22.8
He will do nothing	0.7	1.2	1.0	1.4	2.1	0.5	1.0
He may report her to his family or hers, or send her away	5.9	5.0	5.4	5.0	2.1	6.2	5.4
He may force her to have sex with him	11.3	8.7	9.8	9.2	8.3	10.6	9.8
Total	100	100	100	100	100	100	100

In (A) number of missing observation = 1

The category separated refers to all previously married who are separated, divorced or widowed

Table 5 Perceptions about the Categories of Women who can Refuse Sexual Advances with their Husbands when they think it is not Safe (%)

Response	Male (N = 423)	Female (N = 576)	Mean
Breadwinners, especially educated and highly placed women who know the implications of STIs	81.8	80.9	81.3
Any woman	13.0	11.6	12.2
Women who can intimidate their husbands, also unfaithful and wicked ones	4.7	5.0	4.9
Those who do not use condoms	0.2	0.5	0.4
Don't know	0.2	1.9	1.2
Total	100	100	100

Number of missing observations = 1