1 **Title:** Reaching to recover balance in unpredictable circumstances: is online
2 visual control of the reach-to-grasp reaction necessary or sufficient?
3 (revised February 23, 2012)

4
5 **Authors:** Kenneth C. Cheng (MAsc)¹⁻³, Sandra M. McKay (PhD)^{1,2}, Emily C.
6 King (MAsc)^{1,2,4} and Brian E. Maki (PhD, PEng)¹⁻⁵

7 **Affiliations:** ¹Centre for Studies in Aging, Sunnybrook Health Sciences Centre;
8 ²Toronto Rehabilitation Institute;
9 ³Institute of Medical Science,
10 ⁴Institute of Biomaterials and Biomedical Engineering, and
11 ⁵ Department of Surgery, University of Toronto.

12 **Correspondence:** Brian E. Maki, PhD, PEng
13 Centre for Studies in Aging, Sunnybrook Health Sciences Centre,
14 2075 Bayview Avenue, Toronto, Ontario, CANADA M4N 3M5
15 416-480-6100 ext. 3513
16 brian.maki@sri.utoronto.ca

17 **Running head:** Visual control of perturbation-evoked reach-to-grasp

18 **Keywords:** Arm movements; Postural balance; Spatial working memory;
19 Triggered reactions; Vision

20
21
22
23

1 **ABSTRACT**

2 Reaching to grasp an object for support is a common and functionally important
3 response to sudden balance-perturbation. The need to react very rapidly (to prevent falling)
4 imposes temporal constraints on acquisition and processing of the visuospatial information
5 (VSI) needed to guide the reaching movement. Previous results suggested that the CNS may
6 deal with these constraints by using VSI stored in memory proactively, prior to perturbation-
7 onset; however, the extent to which online visual control is necessary or sufficient to guide
8 these reactions has not been established. This study examined the speed, accuracy and
9 effectiveness of perturbation-evoked reach-to-grasp reactions when forced to rely entirely on
10 either online or stored VSI, by using liquid-crystal goggles to occlude vision either before or
11 after perturbation-onset. The reactions were evoked, in twelve healthy young-adults, via
12 sudden unpredictable antero-posterior platform-translation (barriers deterred stepping
13 reactions). Prior to perturbation-onset, a small cylindrical handhold was positioned
14 unpredictably (by a motor-driven device) at one of four locations in front of the subject.
15 Results indicated that equilibrium could be recovered successfully by grasping the handhold
16 using either online-VSI or stored-VSI to guide the arm reaction; however, both sources of
17 VSI were required for optimal performance. Reach initiation and arm movement were
18 slowed when dependent on online-VSI, whereas reach-accuracy and grip-formation were
19 impaired when dependent on stored-VSI. Comparison with normal-VSI trials suggests that
20 both sources of VSI are utilized when grasping a small handhold for support under normal
21 visual conditions, with stored-VSI predominating during initiation/transport and online-VSI
22 contributing primarily to final target acquisition/prehension.

23

1 INTRODUCTION

2 Rapid reach-to-grasp reactions are a prevalent response to sudden balance
3 perturbation, and can play an important role in preventing falls (McIlroy and Maki 1995;
4 Maki and McIlroy 1997; Maki et al. 1998; Bateni et al. 2004; Maki and McIlroy 2005;
5 Marigold and Misiasek 2009). To grasp an object, it is necessary to identify and locate a
6 suitable handhold and to use this visuospatial information (VSI) to guide the reaching
7 movement. Studies of volitional goal-directed arm movements have clearly revealed a
8 natural propensity for online visual control, wherein visual fixation of the target is used to
9 guide the hand toward it (Prablanc et al. 1979; Carnahan and Marteniuk 1991; Abrams 1992;
10 Haycoe and Ballard 2005; Land 2006). It is not clear, however, the extent to which online
11 visual control can contribute to guiding the rapid reach-to-grasp balance-recovery reactions
12 evoked by a sudden unpredictable balance perturbation, which are initiated and executed
13 much more rapidly than volitional arm movements (McIlroy and Maki 1995; Maki and
14 McIlroy 1997; Gage et al. 2007). The urgent need to react very rapidly in order to prevent a
15 fall imposes temporal constraints that may limit the capacity to acquire and process the
16 needed VSI online, i.e. after the onset of the balance perturbation (Maki and McIlroy 2005;
17 King et al. 2011).

18 Recent studies of arm reactions evoked by a truly-unexpected balance perturbation
19 while ambulating in an unfamiliar environment did, in fact, show that the reach-to-grasp
20 movements were invariably executed in the absence of concurrent visual fixation of the
21 handrail (King et al. 2009; King et al. 2011). Most subjects did, however, fixate briefly on
22 the handrail one or more times upon first entering the environment. This has led to
23 suggestions that the CNS may guide these arm reactions using VSI stored in working spatial
24 memory. The needed VSI, regarding the location of potential handholds (as well as other

1 salient objects), would be acquired and stored automatically as a contingency, through
2 natural exploratory gaze behaviour, as the person enters or moves through the environment
3 (King et al. 2009; King et al. 2011). This stored-VSI would then allow the hand to be moved
4 very rapidly toward the nearest available handhold, if and when an unexpected balance
5 perturbation occurs.

6 The capacity to use stored-VSI to guide effective reach-to-grasp reactions is
7 supported by a study in which liquid-crystal goggles were used to occlude vision for a 200ms
8 interval at time of balance-perturbation onset (Ghafouri et al. 2004). It was shown that the
9 initiation of the grasping reaction was unaffected by the blocking of online visual feedback.
10 However, because vision was occluded only during the initiation of the arm movement, this
11 study did not address potential contributions of online visual control during the target
12 acquisition phase, nor did it address the capacity of the CNS to utilize online visual control to
13 plan and execute the initiation of the reach. Such online control could occur, even in the
14 absence of visual fixation of the target handhold, by utilizing VSI from the peripheral visual
15 field (King et al. 2009; King et al. 2010; King et al. 2011).

16 The objective of the current study was to determine the extent to which online visual
17 control is necessary, or sufficient in itself, to execute rapid, accurate and effective reach-to-
18 grasp reactions in response to sudden unpredictable balance perturbation. To force reliance
19 on online visual control, liquid-crystal goggles were used to occlude vision prior to
20 perturbation onset, and the location of a small graspable handhold was varied unpredictably
21 while vision was occluded. These “online-VSI” trials were compared to “stored-VSI” trials
22 in which vision was occluded at time of perturbation onset (thereby forcing a reliance on VSI
23 acquired prior to perturbation onset) and to “normal-VSI” trials in which vision was not
24 occluded.

1 In view of the temporal constraints noted above, we anticipated that response
2 initiation would be governed primarily using stored VSI. Therefore, we hypothesized that: 1)
3 there would be little difference between normal-VSI and stored-VSI trials in the timing and
4 spatial features of the early portion of the reaching movement; and 2) response initiation
5 would be delayed and arm movement would be slowed in online-VSI trials. The latter
6 effects would presumably serve to “buy” more time to acquire and process the VSI needed to
7 select the appropriate limb and to plan the initial limb trajectory. Conversely, we anticipated
8 that target acquisition would be governed primarily using online VSI, as a consequence of
9 the rapid decay in the accuracy of VSI stored in working spatial memory (Elliott and Calvert
10 1990; Westwood et al. 2003) and the need for online visual control to improve reach
11 accuracy (Prablanc et al. 1979). Therefore, we further hypothesized that mean endpoint
12 accuracy would be reduced, endpoint variability would be increased and prehension errors
13 would occur more frequently in stored-VSI trials, in comparison to online-VSI and normal-
14 VSI trials.

15

1 **METHODS**

2 *Participants*

3 Twelve naïve, right-handed healthy young adults were tested [six male, six female;
4 age 19-29 years (mean=23); mass 48-86kg (mean=66); height 159-186cm (mean=172)].
5 Exclusion criteria included diabetes, neurological or sensory disorders, recurrent dizziness or
6 unsteadiness, use of medications that may affect balance, joint replacement, medical
7 conditions interfering significantly with daily activities, or functional limitations of limb use.
8 All subjects had a Snellen visual acuity of 20/30 or better, without wearing spectacles. Each
9 subject provided written informed consent to comply with ethics approval granted by the
10 institutional review board. All subjects participated in a short and separate balance study
11 involving three gait-perturbation trials (King et al. 2009; King et al. 2011) immediately prior
12 to the start of the present study.

13 *Protocol*

14 Balance-recovery reach-to-grasp reactions were evoked by sudden forward (0.12m,
15 0.41m/s, 1.4m/s²) or backward (0.18m, 0.6m/s, 2.0m/s²) translation of a 2m×2m computer-
16 controlled motion-platform (Maki et al. 1996). Each platform translation comprised an
17 approximately-square 300ms acceleration pulse followed immediately by an equal and
18 opposite deceleration pulse. The focus of the study was on the grasping reactions evoked
19 during the backward falling motion induced by forward platform translations; however, to
20 increase unpredictability and deter anticipatory arm movements, we also included backward
21 translations (~30% of trials) and “catch” trials (~15% of trials) in which the platform
22 remained stationary for the duration of the trial (~10s). For safety, subjects wore a harness
23 designed to prevent impact between body and floor without restricting movement or
24 providing somatosensory feedback that could aid in control of balance.

At the start of each trial, a motor-driven device mounted on the platform (Cheng et al. 2009) controlled a cylindrical handhold (length=20cm, diameter=3.8cm) to move along a transverse axis in front of the subject (distance from handhold to back of heels=33% of body-height; handhold height=60% of body-height) and to stop unpredictably at one of four locations (left or right of midline; distance from mid-sagittal plane=50% or 75% of shoulder-width, SW); see Figure 1. Perturbation-onset was then triggered after a random delay (2-5s), or not at all (“catch” trials). To provide moderate visual contrast, the handhold was covered with medium-gray grip tape, while the visual background (floor and wall) was black, with an ambient room-illumination level of ~230lux.

Subjects began each trial facing forward (looking at the handhold, in normal- and stored-VSI trials), with arms resting at sides. They were instructed to recover balance by grasping a marked (with red tape) “target” section of the handhold (length=125% of hand-width) as quickly as possible after onset of platform motion (overhand grasp with hand closest to target), but to refrain from moving their arms if the platform did not move. They were also told that a \$50 prize would be awarded to the subject with the quickest responses, but that penalties for premature initiation of arm movements (prior to onset of platform motion, or in the absence of any platform motion) would reduce their chances of winning the prize. Instructions not to move the feet, plus foam-rubber barriers (height=30cm), were used to deter foot motion and reinforce reliance on reach-to-grasp reactions.

Three visual conditions were tested, using computer-controlled translucent liquid-crystal goggles (Translucent Technologies, Toronto, ON) to occlude vision: 1) prior to perturbation-onset (forcing use of online-VSI), 2) after perturbation-onset (forcing use of stored-VSI), or 3) not at all (normal-VSI). The goggles were custom-modified to hold the liquid-crystal element flush against the orbit of each eye, so as to completely block both

central and peripheral vision when activated (opaque) and to permit near-complete field-of-view when de-activated (Scovil et al. 2008). Goggle activation/de-activation was synchronized with perturbation-onset (platform acceleration $>0.1\text{m/s}^2$). Subjects were instructed to look at the handhold

For each visual condition, each subject performed two blocks of fourteen trials comprising (in random order) two forward-translation trials at each of the four handhold locations, two catch (no-perturbation) trials, and four backward-translation trials. Subjects were informed of the visual condition prior to starting each block. Order of the blocks was balanced across and within subjects. To dampen learning effects, seven practice trials were performed each time a new visual condition was introduced. In addition, five unrestricted-vision trials were performed prior to starting the main protocol.

Data collection and analysis

Video recordings from four cameras were used to determine which arm was used to grasp the handhold, whether a full grasp was achieved (all digits wrapped around the handhold), whether a collision error occurred (contact with back of wrist, hand or digits), and whether the subject attempted to step (by kicking the foam-rubber barriers) or fell into the safety harness (confirmed by load cell on harness cable). Surface electrodes were used to record electromyographic (EMG) activity bilaterally in the anterior deltoid, lateral deltoid and biceps muscles (band-pass filtered, 10-500Hz; sampling-rate=1000Hz). Reaction time was defined as the earliest EMG onset latency in any of these muscles, as determined by a computer algorithm (McIlroy and Maki 1993) and confirmed by visual inspection. Contact time was detected by force-sensing resistors mounted on the front, back and top of the handhold (sampling-rate=200Hz), and confirmed using a hand-velocity criterion ($<5\%$ peak velocity), as determined by the motion-analysis system described below. Timing measures

1 were defined relative to onset of platform acceleration ($>0.1\text{m/s}^2$), determined by an
2 accelerometer mounted on the motion platform.

3 A three-dimensional motion-analysis system (Vicon-Peak Performance; Englewood,
4 CO) collected kinematic data (sampling rate=200Hz; low-pass filtered at 6Hz using a dual-
5 pass fourth-order filter). Analysis focussed on the location and trajectory of each hand
6 (marker on third-metacarpal knuckle), relative to markers mounted at each end of the
7 handhold. The hand-marker data were used to determine the maximum resultant velocity,
8 time-to-peak-velocity (relative to the reaction-onset time) and time-after-peak-velocity (time
9 from peak velocity to handhold contact). The location of the hand marker relative to the
10 center of the handhold target area, at time of handhold contact, was used to define the reach
11 error in each coordinate direction.

12 Data from the hand marker were also used to describe the trajectory of the reach in
13 the transverse plane. Trajectory data were used to determine: a) the angle of the initial reach
14 trajectory; and b) the deviation from the 'direct-path' to the handhold, using a method
15 adapted from studies of volitional reaching (Khan et al. 2003; Khan et al. 2006). The 'direct-
16 path' was defined as the straight-line path connecting the hand position at movement onset
17 (hand velocity $>5\%$ of peak velocity) to the center of the target region of the handhold.
18 Orthogonal deviation from this direct-path was calculated at increments of 5% of the direct-
19 path distance. The initial trajectory angle was based on the line segment connecting the
20 movement-onset hand position to the position of the hand when it had moved 5% of the
21 direct-path distance to the handhold. The angle of this line segment was determined in
22 relation to a line segment connecting the movement-onset hand position to the 50%SW
23 handhold position (see inset in Figure 4A).

Repeated-measures analysis of variance (ANOVA) and post-hoc Tukey multiple comparisons were performed to test the hypotheses. The primary dependent variables were: 1) reach timing (reaction-onset, movement-time and contact-time); 2) initial trajectory angle; 3) reach velocity (peak velocity, time-to-peak-velocity and time-after-peak-velocity); 4) reach accuracy; 5) variability in reach accuracy; and 6) frequency of full-grasp. The reach accuracy and variability variables were analyzed separately in each coordinate direction, and were expressed as a proportion of subject height prior to analysis. The primary focus was on effects due to visual condition. Handhold side (left or right) and eccentricity (50% or 75% of SW from midline) were also included as factors to account for variation due to these factors, and any significant interactions with visual condition were noted. All data were rank-transformed prior to analysis to avoid errors arising from potential violations of the assumptions underlying the ANOVA (Conover and Iman 1981). The criterion level of significance was 0.05.

For frequency variables, the proportion of trials in which the event was observed was calculated within each subject, for each of the twelve experimental conditions (three visual conditions $\times$ four handhold locations), and the ANOVA was performed on the rank-transformed proportions. To analyze accuracy variability, the standard deviation of the grasp error was determined within each subject, for each of the twelve experimental conditions, and the ANOVA was performed on the rank-transformed standard deviations. For all of the other variables, rank-transformed data from individual trials were used. Each subject performed 48 trials; therefore, there were a total of 576 trials potentially available for analysis. Trials in which subjects initiated the arm movement prematurely (EMG latency $<70\text{ms}$), failed to react rapidly to the perturbation (EMG latency $>250\text{ms}$, movement-time $>550\text{ms}$, or contact-time $>750\text{ms}$) or did not reach for the handhold were excluded from the

1 analyses (n=11). In addition, due to technical problems with the motion-analysis system, one
2 subject was excluded from the accuracy analyses and two subjects were excluded from the
3 trajectory analyses.

4

5

1 **RESULTS**

2 All twelve subjects completed the protocol and all were well able to recover balance
3 by reaching to grasp the handhold. None of the subjects fell into the safety harness
4 (maximum harness loading <5% body-weight) or attempted to step to recover balance (by
5 kicking the foam barriers out of the way). The reach failed to contact the handhold in only
6 0.71% (4/565) of trials, and collision between the back of the hand and the handhold
7 occurred in only 0.88% (5/565) of trials (these missed- or inadvertent-contact events were
8 invariably followed by a trajectory correction and successful prehension). Subjects never
9 reached with the “wrong” arm (contralateral to the handhold location); however, six of the
10 twelve subjects occasionally raised both arms prior to completing the reach with the
11 “correct” arm (1 of 48 trials in four subjects, 3 trials in one subject and 9 trials in one
12 subject). This tendency was dependent on visual condition [$F(2,22)=4.04$, $p=0.032$], and was
13 more likely to occur in online-VSI trials (6.4% of trials, versus 2.1% of normal-VSI and
14 0.0% of stored-VSI trials). Examples of typical reach trajectories are shown in Figure 2.

15 **Timing and speed of motion:** Mean reaction-onset time, movement time and contact
16 time were all affected significantly by the visual condition [$F(2,22)>15.33$, p 's<0.0001; see
17 Figure 3A]. As hypothesized, movement times were prolonged when dependent on online-
18 VSI (369ms vs 347ms for normal-VSI and 345ms for stored-VSI). Reaction-onset times
19 were also slower when dependent on online-VSI, in comparison to stored-VSI trials (158ms
20 vs 143ms), but did not differ when compared to normal-VSI trials (162ms). As a
21 consequence of the slowing in both reaction-onset time and movement time, handhold
22 contact occurred significantly later in online-VSI trials, in comparison to both of the other
23 visual conditions, whereas the earliest contact times occurred in stored-VSI trials (527ms for
24 online-VSI vs 509ms for normal-VSI vs 487ms for stored-VSI). Visual condition also had a

1 significant effect on peak velocity [$F(2,18)=4.10$, $p=0.034$], time-to-peak-velocity
2 [$F(2,18)=3.85$, $p=0.041$], and time-after-peak-velocity [$F(2,18)=5.57$, $p=0.013$]; see Figure
3 3B. Again, the timing was prolonged during online-VSI trials (time-to-peak-velocity: 210ms
4 vs 203ms for stored-VSI and 201ms for normal-VSI; time-after-peak-velocity: 153ms vs
5 146ms for normal-VSI and 143 for stored-VSI). The velocity was slower in online-VSI trials
6 (3.14m/s) in comparison to normal-VSI trials (3.30m/s), but not in comparison to stored-VSI
7 trials (3.14m/s).

8 **Reach trajectory:** Although there was no main effect of visual condition on the angle
9 of the initial reach trajectory in the transverse plane [$F(2,18)=0.18$, $p=0.84$], there was a
10 significant interaction between handhold eccentricity and visual condition [$F(2,18)=9.5$,
11 $p=0.0016$]; Figure 4A. Specifically, online-VSI trials appeared to exhibit less handhold-
12 eccentricity-dependent modulation of the initial trajectory direction. In these trials, the mean
13 difference in initial trajectory angles between the 50%SW and 75%SW handhold positions
14 was 8° , versus 14° for normal-VSI and 21° for stored-VSI trials. Note that the 8° difference
15 in online-VSI trials was considerably smaller than the mean angular difference in the actual
16 handhold locations (16°). Consistent with these findings, analysis of deviations from the
17 direct-path to the handhold showed a tendency for medial deviation during online-VSI trials
18 involving the more lateral handhold positions (75%SW from midline); see Figure 4B. Such
19 medial deviation would tend to cause the trajectory to be more similar to the trajectory used
20 when the handhold was in the more medial position (50%SW from midline). The apparent
21 tendency to adopt a “generic” initial trajectory in some online-VSI trials is also evident in the
22 example data shown in Figure 2; these example data also illustrate the subsequent trajectory
23 corrections that occurred.

Reach accuracy and variability: Analysis of mean error at hand contact revealed a significant effect due to visual condition in the antero-posterior [$F(2,20)=22.76$, $p<0.0001$] and vertical [$F(2,20)=9.09$, $p=0.0016$] directions; see Figure 5A. Specifically, in stored-VSI trials, the hand tended to contact the handhold with antero-posterior undershoot (-30mm vs -9mm for normal-VSI and -10mm for online-VSI) and reduced elevation (51mm vs 58mm for online-VSI and normal-VSI). Although there was also a main effect due to visual condition on medio-lateral accuracy [$F(2,20)=4.91$, $p=0.018$], the effect was not consistent across handhold locations [visual-condition $\times$ handhold-eccentricity interaction; $F(2,20)=3.87$, $p=0.038$]: lateral error was greatest in stored-VSI trials when the handhold was in the 50%SW position (11mm, vs 5-7mm for normal- and online-VSI), but the hand tended to land medially with respect to the handhold center (in all visual conditions) in the 75%SW trials (-9mm for normal- and stored-VSI, -19mm for online-VSI). Analysis of variability in hand-landing position revealed a significant effect of visual condition in the vertical direction [$F(2,20)=14.92$, $p=0.0001$], with the greatest variability occurring in stored-VSI trials (9.5mm vs 5.5mm for normal-VSI and 6.0mm for online-VSI); see Figure 5B. There were no significant vision effects for landing variability in the medio-lateral or antero-posterior directions [$F(2,20)$'s <0.58 , p 's >0.57].

Prehension: Analysis of the type of grasp revealed a small but near-significant effect due to visual condition [$F(2,22)=3.30$, $p=0.056$], with subjects least likely to achieve a full grasp (all digits wrapped around the handhold) when forced to rely on stored-VSI (80% of trials vs 91% for normal-VSI and 86% for online-VSI; see Figure 6). The failure to achieve a full grasp in some of the stored-VSI trials appeared to be associated with the aforementioned tendency to undershoot the handhold in the antero-posterior direction. The most common consequence of the undershoot was a hook grip, in which the fingers hooked onto the top of

1 the handhold (without palmar contact) and the thumb failed to wrap around the undersurface
2 of the handhold (11% of trials for stored-VSI, vs 5% for online-VSI and 3% for normal-VSI;
3 see Figure 6).

4

1 **DISCUSSION**

2 Subjects were generally well able to achieve a functionally-adequate grasp and
3 prevent themselves from falling despite the experimental manipulations in access to
4 visuospatial information (VSI). These findings thus indicate that either online-VSI or stored-
5 VSI alone can be sufficient to guide functional reach-to-grasp reactions in response to sudden
6 unpredictable balance perturbation. It appears, however, that both sources of VSI are
7 required for optimal performance. As hypothesized, reach initiation and arm movement were
8 slowed when dependent on online VSI (in comparison to stored-VSI trials), whereas reach
9 accuracy and ability to achieve a full grasp were reduced when dependent on stored VSI (in
10 comparison to online-VSI trials). Comparison with the normal-VSI trials suggests that both
11 sources of VSI were utilized under normal visual conditions, whereby stored-VSI facilitated
12 rapid initiation and transport and online-VSI facilitated accurate target acquisition and
13 prehension.

14 The slowing in mean reaction-onset time observed in online-VSI trials presumably
15 served to buy more time to acquire and process the VSI needed to select the appropriate limb
16 and to plan the initial portion of the reach trajectory, whereas the slowing of the arm
17 movement (and prolongation of the time-after-peak-velocity) presumably served to further
18 increase the time available to formulate and execute online trajectory corrections during the
19 later stages of the reach (Khan et al. 2006). There was also evidence of a tendency to preplan
20 a “generic” initial trajectory in online-VSI trials, which may have helped to avoid delays in
21 response initiation in some trials. Use of a generic initial trajectory, in some proportion of
22 online-VSI trials, is supported by the finding that the mean initial trajectory angle in the
23 transverse plane was less affected by changes in handhold location when dependent on
24 online-VSI. Another possible strategy involved initially raising both arms, prior to selecting

1 which arm to use to grasp the handhold. This could potentially serve to buy time for the
2 VSI-processing needed to select the most appropriate limb, without causing delay in
3 initiating arm movement. Based on online-VSI acquired during the initial raising of the arms,
4 the CNS could then terminate the motion of the “incorrect” (contralateral-to-handhold) arm
5 and complete the reach with the “correct” (ipsilateral) arm. Further research is needed to
6 investigate whether bilateral-arm-raise does, in fact, lead to faster reach-to-grasp initiation,
7 whether it has other functional benefits [e.g. helping to “counter-balance” the backward
8 falling motion of the body (Marigold and Misiashzek 2009)], or whether these responses were
9 simply a failure to follow the instructions to reach with one arm.

10 With regard to effects of forced reliance on stored-VSI, the findings supported our
11 hypothesis that this would lead to reduction in mean endpoint accuracy, increase in endpoint
12 variability and less effective grip formation, in comparison to trials where online VSI was
13 available. In addition, we found support for our hypotheses that the timing and spatial
14 features of the reaching movement would be similar in stored-VSI and normal-VSI trials;
15 however, there was one exception: reaction-onset time was significantly faster in stored-VSI
16 trials (by ~20ms, on average). Ghafouri et al (2004) also found that the forced dependence
17 on stored-VSI had no effect on the initial direction or speed of the reach (in comparison to
18 normal-VSI trials), but in contrast to the present study, found very little effect (<5ms, on
19 average) on reaction-onset time. The reason for this discrepancy in findings is not clear, but
20 could possibly be related to the level of postural challenge. Potentially, the larger
21 perturbation magnitudes used by Ghafouri et al, and associated arousal levels (Carpenter et
22 al. 2004), served to evoke the most rapid reactions possible, independent of visual condition,
23 whereas the less challenging perturbations used here may have allowed less rapid responses
24 to suffice in the less challenging visual condition, i.e. normal-VSI trials.

Other than the work by Ghafouri et al (2004), we are not aware of any other studies addressing the contributions of stored- or online-VSI to the control of perturbation-evoked reach-to-grasp reactions. However, this issue has been addressed in the context of perturbation-evoked lower-limb movements, i.e. stepping reactions evoked by sudden platform translation (Zettl et al. 2005; Zettl et al. 2007; Scovil et al. 2008; Zettl et al. 2008). Similar to the present findings, these studies have suggested that the initiation of the limb movement is typically governed by stored-VSI and that forced reliance on online-VSI can adversely affect response initiation. For example, in a study which required the step to be directed through a narrow, unpredictably-located aperture, use of liquid-crystal goggles to force reliance on online visual control led to delay in foot-off time (by ~50ms), as well as increased tendency to step with the “wrong” foot (i.e. contralateral to the aperture location) and to contact with the barrier while moving the foot through the aperture (Scovil et al. 2008). Step-reaction studies have also indicated that stored-VSI can generally suffice in guiding the final phases of the limb movement; however, an increased tendency to look downward when required to land the foot on a target suggests that online-VSI may play a role when accurate end-point control of the foot is required (Zettl et al. 2005; Zettl et al. 2008).

The presently-observed tendency to undershoot targets when dependent on stored-VSI has also been reported to occur in young-adults in previous studies of volitional arm movements (Elliott et al. 1999; Westwood et al. 2003). It has been suggested that this might be a strategy to ensure that the hand does not collide with the target (in the absence of any opportunity for visually-guided online corrections to the reach trajectory) and/or a strategy to avoid time-consuming reversals in reach direction (which would be required in the event of overshoot error) (Elliott et al. 1999). Alternatively, the undershoot error could reflect a

1 failure to fully compensate for the backward body motion induced by the forward platform
2 translation, a misrepresentation of the target position stored in memory (Flanders et al. 1992),
3 or a time-dependent decay in the accuracy of the stored target information (Lemay and
4 Proteau 2002). The latter, however, would be expected to lead to increase in antero-posterior
5 end-point variability (based on studies of volitional pointing and grasping tasks (Khan et al.
6 2006)), but this was not observed in the present study.

7 The mean changes in reaction timing and accuracy due to the visual manipulations
8 were relatively small (~40ms or less and ~30mm or less), and did not prevent subjects from
9 achieving a functionally-adequate grasp and recovering equilibrium; however, the presently-
10 observed errors and delays in contacting the handhold may have a greater functional impact
11 when responding to larger perturbations. The present perturbation magnitudes were selected
12 (based on our previous studies) to be large enough to force a need to grasp for support (but
13 not so large as to evoke stepping reactions); however, because subjects were instructed to
14 grasp in all trials, we cannot ascertain the degree to which grasping was actually needed to
15 augment the stabilization provided by the lower-limb “feet-in-place” postural reactions that
16 occur in conjunction with reach-to-grasp reactions (McIlroy and Maki 1995). Presumably,
17 larger perturbations will require larger stabilizing hand-handhold reaction forces to be
18 generated, and hence may require a stronger grip. Some of the “partial” grips that were
19 sufficient to stabilize the body in the present study may prove to be inadequate in responding
20 to larger perturbations; hence, grasp-accuracy demands may increase. In addition, larger
21 perturbations will demand more rapid responses. Further work is needed to determine
22 whether the increased accuracy and speed-of-response demands imposed by larger
23 perturbations will affect the extent to which functionally-effective reach-to-grasp reactions
24 can be executed using stored- or online-VSI.

Further work is also needed to assess the degree to which the present findings generalize to daily-life situations, which are likely to involve complex environments and balance perturbations that are completely unexpected and highly unpredictable. In contrast, the task conditions in the present study were limited to a finite number of options (two perturbation directions/magnitudes and four handhold locations), and subjects were given prior instruction to respond to the perturbation by grasping the handhold. Such task conditions may have allowed the CNS to take advantage of its capacity to encode motor commands for multiple trajectories/directions simultaneously prior to action selection (Cisek and Kalaska 2010). In addition, more complex environments are likely to require a more extensive visual-search to locate an appropriate handhold, which could further delay and jeopardize successful completion of reach-to-grasp reactions in situations where the CNS is forced on rely on online visual control, especially when the perturbation is truly unexpected.

Further work is also needed to examine situations where the VSI must be stored in spatial memory for some interval of time prior to perturbation onset. Such a situation could occur in daily life, for example, when potential handhold locations are mapped via natural exploratory gaze behaviour upon first entering an environment but the balance perturbation occurs several seconds later. Studies of volitional arm movements indicate that the accuracy of the VSI stored in working spatial memory decays rapidly (Elliott and Madalena 1987; Hu et al. 1999). For example, a 2s delay between vision-occlusion and onset of an auditory cue to initiate arm movement led to increased endpoint variability, undershoot error, reaction time and movement time (Hu et al. 1999; Lemay and Proteau 2002; Westwood et al. 2003). Finally, it will also be of interest to determine whether visual fixation of the handhold occurred during online-VSI trials (subsequent to the offset of visual occlusion) or whether these reactions were guided by peripheral vision. The wearing of the liquid-crystal goggles

1 precluded use of an eye-tracker in the present study; however, it may be possible to address
2 this question using other methods to record eye movements or by using special contact lenses
3 to force reliance on peripheral or central vision (Sivak and MacKenzie 1990). Work is in
4 progress to address these issues.

5

1 CONCLUSION

2 The results of this study indicate that either stored-VSI or online-VSI is sufficient to
3 guide rapid perturbation-evoked reach-to-grasp reactions with a level of accuracy that is
4 sufficient to achieve a functional grasp, even when the handhold is small and the location of
5 the handhold is varied in an unpredictable manner prior to perturbation onset. However, it
6 appears that both sources of VSI are required for optimal performance, as reach initiation and
7 arm movement were slowed when dependent on online-VSI and reach-accuracy and grip-
8 formation were impaired when dependent on stored-VSI. Nonetheless, despite these
9 performance decrements, our findings suggest that either stored- or online-VSI alone can be
10 an adequate source of information for guiding functionally-effective reach-to-grasp balance-
11 recovery reactions. Further work is needed to establish whether this remains the case when
12 responding to larger and completely unexpected perturbations in more complex and
13 unpredictable environments and situations, and to determine the effects of age-related
14 impairments in the capacity to rapidly process online VSI or to accurately store VSI in
15 working spatial memory.
16

1 **FUNDING**

2 Supported by grant #MOP-13355 from the Canadian Institutes of Health Research
3 (CIHR) and by scholarships from CIHR, the Ontario Neurotrauma Foundation, the Vision
4 Science Research Program and the Ontario Graduate Scholarship Program.

6 **ACKNOWLEDGMENTS**

7 Assistance from Sandy Wong is gratefully acknowledged. Preliminary results were
8 presented at the 19th Conference of the International Society for Postural and Gait Research
9 (Bologna, Italy; July/2009) and the Symposium on Basic Mechanisms Underlying Balance
10 Control under Static and Dynamic Conditions (Pavia, Italy; June/2009).

13 **CONFLICT OF INTEREST**

14 The authors declare that they have no conflict of interest.

1 **REFERENCES**

- 2 Abrams RA (1992) Coordination of eye and hand for aimed limb movements. In: Proteau L,
3 Elliot D (eds) Vision and Motor Control. Elsevier, Amsterdam, pp 129-152
- 4 Bateni H, Zecevic A, McIlroy WE, Maki BE (2004) Resolving conflicts in task demands
5 during balance recovery: does holding an object inhibit compensatory grasping? Exp
6 Brain Res 157:49-58
- 7 Carnahan H, Marteniuk RG (1991) The temporal organization of hand, eye, and head
8 movements during reaching and pointing. J Motor Behav 23:109-119
- 9 Carpenter MG, Frank JS, Adkin AL, Paton A, Allum JH (2004) Influence of postural anxiety
10 on postural reactions to multi-directional surface rotations. J Neurophysiol 92:3255-
11 3265
- 12 Cheng KC, McKay SM, King EC, Tung JY, Lee TA, Scovil CY, Maki BE (2009) The
13 moveable handhold: a new paradigm to study visual contributions to the control of
14 balance-recovery reactions. Gait Posture 29:339-342
- 15 Cisek P, Kalaska JF (2010) Neural mechanisms for interacting with a world full of action
16 choices. Annu Rev Neurosci 33:269-298
- 17 Conover WJ, Iman RL (1981) Rank transformations as a bridge between parametric and
18 nonparametric statistics. Am Stat 35:124-133
- 19 Elliott D, Calvert R (1990) The influence of uncertainty and premovement visual information
20 on manual aiming. Can J Psychol 44:501-511

- 1 Elliott D, Heath M, Binsted G, Ricker K, Roy EA, Chua R (1999) Goal-directed aiming:
2 correcting a force-specification error with the right and left hands. *J Motor Behav*
3 31:309-324
- 4 Elliott D, Madalena J (1987) The influence of premovement visual information on manual
5 aiming. *Q J Exp Psychol A* 39:541-559
- 6 Flanders M, Helms-Tillery SI, Soechting JF (1992) Early stages in a sensorimotor
7 transformation. *Behav Brain Sci* 15:309-362
- 8 Gage WH, Zabjek KF, Hill SW, McIlroy WE (2007) Parallels in control of voluntary and
9 perturbation-evoked reach-to-grasp movements: EMG and kinematics. *Exp Brain Res*
10 181:627-637
- 11 Ghafouri M, McIlroy WE, Maki BE (2004) Initiation of rapid reach-and-grasp balance
12 reactions: is a pre-formed visuospatial map used in controlling the initial arm
13 trajectory? *Exp Brain Res* 155:532-536
- 14 Haycoe M, Ballard D (2005) Eye movements in natural behavior. *Trends Cogn Sci* 9:188-
15 194
- 16 Hu Y, Eagleson R, Goodale MA (1999) The effects of delay on the kinematics of grasping.
17 *Exp Brain Res* 126:109-116
- 18 Khan MA, Franks IM, Elliott D, Lawrence GP, Chua R, Bernier PM, Hansen S, Weeks DJ
19 (2006) Inferring online and offline processing of visual feedback in target-directed
20 movements from kinematic data. *Neurosci Biobehav Rev* 30:1106-1121

- 1 Khan MA, Lawrence GP, Franks IM, Elliott D (2003) The utilization of visual feedback in
2 the control of movement direction: evidence from a video aiming task. *Motor Control*
3 7:290-303
- 4 King EC, Lee TA, McKay SM, Scovil CY, Peters AL, Pratt J, Maki BE (2011) Does the
5 "eyes lead the hand" principle apply to reach-to-grasp movements evoked by
6 unexpected balance perturbations? *Hum Mov Sci* 30:368-383
- 7 King EC, McKay SM, Cheng KC, Maki BE (2010) The use of peripheral vision to guide
8 perturbation-evoked reach-to-grasp balance-recovery reactions. *Exp Brain Res*
9 207:105-118
- 10 King EC, McKay SM, Lee TA, Scovil CY, Peters AL, Maki BE (2009) Gaze behavior of
11 older adults in responding to unexpected loss of balance while walking in an
12 unfamiliar environment: a pilot study. *J Optom* 2:119-126
- 13 Land MF (2006) Eye movements and the control of actions in everyday life. *Prog Retin Eye*
14 *Res* 25:296-324
- 15 Lemay M, Proteau L (2002) Effects of target presentation time, recall delay, and aging on the
16 accuracy of manual pointing to remembered targets. *J Motor Behav* 34:11-23
- 17 Maki BE, McIlroy WE (1997) The role of limb movements in maintaining upright stance: the
18 "change-in-support" strategy. *Phys Ther* 77:488-507
- 19 Maki BE, McIlroy WE (2005) Change-in-support balance reactions in older persons: an
20 emerging research area of clinical importance. *Neurol Clin* 23:751-783

- 1 Maki BE, McIlroy WE, Perry SD (1996) Influence of lateral destabilization on compensatory
2 stepping responses. *J Biomech* 29:343-353
- 3 Maki BE, Perry SD, McIlroy WE (1998) Efficacy of handrails in preventing stairway falls: a
4 new experimental approach. *Safety Sci* 28:189-206
- 5 Marigold DS, Misiaszek JE (2009) Whole-body responses: neural control and implications
6 for rehabilitation and fall prevention. *Neuroscientist* 15:36-46
- 7 McIlroy WE, Maki BE (1993) Changes in early 'automatic' postural responses associated
8 with the prior-planning and execution of a compensatory step. *Brain Res* 631:203-211
- 9 McIlroy WE, Maki BE (1995) Early activation of arm muscles follows external perturbations
10 of upright stance. *Neurosci Lett* 184:177-180
- 11 Prablanc C, Echallier JE, Jeannerod M, Komilis E (1979) Optimal response of eye and hand
12 motor systems in pointing at a visual target. I. Spatio-temporal characteristics of eye
13 and hand movements and their relationships when varying the amount of visual
14 information. *Biol Cybern* 35:113-124
- 15 Scovil CY, Zettel JL, Maki BE (2008) Stepping to recover balance in complex environments:
16 is online visual control of the foot motion necessary or sufficient? *Neurosci Lett*
17 445:108-112
- 18 Sivak B, MacKenzie CL (1990) Integration of visual information and motor output in
19 reaching and grasping: the contributions of peripheral and central vision.
20 *Neuropsychologia* 28:1095-1116

1 Westwood DA, Heath M, Roy EA (2003) No evidence for accurate visuomotor memory:
2 systematic and variable error in memory-guided reaching. *J Motor Behav* 35:127-133

3 Zettel JL, Holbeche A, McIlroy WE, Maki BE (2005) Redirection of gaze and switching of
4 attention during rapid stepping reactions evoked by unpredictable postural
5 perturbation. *Exp Brain Res* 165:392-401

6 Zettel JL, McIlroy WE, Maki BE (2008) Effect of competing attentional demands on
7 perturbation-evoked stepping and associated gaze behavior in young and older adults.
8 *J Gerontol A Biol Sci Med Sci* 63:1370-1379

9 Zettel JL, Scovil CY, McIlroy WE, Maki BE (2007) Gaze behavior governing balance
10 recovery in an unfamiliar and complex environment. *Neurosci Lett* 422:207-212

11

FIGURE CAPTIONS

Figure 1: Motion platform and moveable handhold systems. Dashed lines in the insert indicate the four handhold positions that were tested.

Figure 2: Example transverse-plane hand trajectories from a representative subject. In the trials shown, the handhold was located to the right of midline. Each panel corresponds to one of the three visual conditions, and shows the trajectories for the four trials where the handhold location was 50% of shoulder-width (SW) from midline (thin grey lines) and the four trials where the handhold location was 75% of SW from midline (thicker black lines). The trajectories have been adjusted so that the starting position of the hand marker (at time of movement onset) is at the origin of the coordinate system; the center of the handhold relative to this coordinate system is indicated, for each trial, by an encircled (and shaded) number (1-8) which matches the unshaded number displayed at the end-point of the corresponding trajectory. The square or triangular symbol superimposed on each trajectory indicates the hand-marker position 100ms after onset of hand movement. Note that the direction of this initial 100ms of hand movement was modulated according to handhold location in normal- and stored-VSI trials, whereas online-VSI trials showed a tendency for the hand to move initially toward the 50%SW handhold location regardless of the actual handhold location (necessitating a subsequent trajectory correction in trials where the handhold was actually located at the 75%SW position).

Figure 3: Effect of visual condition on mean reach-to-grasp speed: **A.** timing (reaction-onset, movement and handhold-contact times); and **B.** hand velocity (peak velocity, time-to-peak-velocity and time-after-peak-velocity). Note the slowing of the reaction in

online-VSI trials, in comparison to stored-VSI and/or normal-VSI trials. * indicates a significant difference between means ($\alpha=0.05$); whiskers indicate standard deviations.

Figure 4: Effect of visual condition on mean hand trajectory (transverse plane): **A.** initial-trajectory angle; and **B.** orthogonal deviation from a direct-path trajectory (see insets for definitions). Data are shown separately for the two handhold eccentricities (distance to midline = 50% or 75% of shoulder-width, SW); **B** also shows data separately for the left and right handhold locations. In **A**, * indicates a significant difference between means ($\alpha=0.05$); whiskers indicate standard deviations. Indicators of significance ($\alpha=0.05$) in **B**: online-VSI different from stored-VSI and normal-VSI (*); online-VSI different from stored-VSI (¶); online-VSI different from normal-VSI (†); no significant differences between means (ns).

Figure 5: Effect of visual condition on reach-to-grasp accuracy, for each coordinate axis: **A.** systematic error; **B.** variable error. Note the increased tendency for reduced elevation and antero-posterior undershoot in stored-VSI trials, as well as the greater tendency for lateral error (note, however, that the latter effect was dependent on handhold eccentricity; see Results for details). Note also the increased vertical variable error in stored-VSI trials. * indicates a significant difference between means ($\alpha=0.05$); whiskers indicate standard deviations.

Figure 6: Effect of visual condition on grasping. The different types of grasping behavior that were observed are illustrated by the photographs. The bar graphs indicate the mean and standard deviation for the percentage of trials in which each grasping behavior occurred. Note that the reduced tendency to achieve a full grip in stored-VSI trials was nearly statistically significant ($p=0.056$).