Title: Redirection of gaze and switching of attention during rapid stepping reactions evoked by unpredictable postural perturbation (revised February 7, 2005)

Authors: John L. Zettel^{1,2,3}, Andrea Holbeche¹, William E. McIlroy^{1,4} and Brian E. Maki^{1,2,3,5}

Affiliations: ¹Centre for Studies in Aging, Sunnybrook and Women's College Health Sciences Centre;

²Institute of Medical Science,

³Institute of Biomaterials and Biomedical Engineering,

⁴Graduate Department of Rehabilitation Science and

⁵Department of Surgery, University of Toronto

Correspondence: John L. Zettel

Centre for Studies in Aging

Sunnybrook and Women's College Health Sciences Centre

2075 Bayview Avenue, Toronto, Ontario, CANADA M4N 3M5

Tel: (416) 480-5858; Fax: (416) 480-5856

e-mail: john.zettel@sw.ca

Running Head: Gaze behavior during triggered stepping

Meetings: Presented (in part) at the 16th International Symposium on Posture and Gait (Sydney, Australia; March, 2003), the 13th Biennial Conference of the Canadian Society for Biomechanics (Halifax, Canada; August 2004), and the 34th Annual Meeting of the Society for Neuroscience (San Diego, California; October, 2004).

Keywords: attention, balance, environmental constraints, eye movements, postural control, stepping, triggered reactions, vision

Abstract

In many situations, successful execution of a balance-recovery reaction requires visual information about the environment. In particular, reactions that involve rapid limb movements, such as stepping, must be controlled to avoid obstacles and accommodate other constraints on limb trajectory. However, it is unknown whether the CNS can acquire the necessary visuospatial information prior to perturbation onset, or must instead redirect gaze at the floor during the execution of the stepping reaction. To study this, we examined gaze behavior during rapid forward-directed stepping reactions triggered by unpredictable platform perturbation, in 12 healthy young adults. We also monitored switching of attention, as inferred from onset of significant error in performing a concurrent visuomotor tracking task. Obstacles and/or step targets were used as constraints, to increase demands for accurate foot movement. Downward gaze shifts toward the floor almost never occurred during stepping reactions when foot motion was unconstrained, but did occur more frequently as the demands for accurate foot movement increased. Nonetheless, even in the most challenging condition (target plus obstacle), downward redirection of gaze occurred in less than 40% of the trials, and subjects were commonly well able to clear the obstacle and land the foot on the target without redirecting gaze toward the floor. An apparent switching of attention, subsequent to perturbation onset, occurred frequently (>80% of trials) in all task conditions, independent of the gaze shifts. The findings indicate that visual fixation of the foot or floor was not essential for accurate control of the foot movement, nor was the apparent switching of attention that followed perturbation onset linked, in any consistent way, to overt changes in visual fixation. Spatial features of the support surface were apparently “remembered” prior to perturbation onset, thereby allowing both vision and attention to be directed to other demands during the execution of the balance reaction.

1 **Introduction**

2 Numerous studies have demonstrated that vision contributes to the control of postural
3 stability, predominantly affecting slow or low-frequency balance corrections (Diener et al, 1986).
4 However, it would appear that no studies have addressed the potential role of eye movements in
5 acquiring visual information that may be needed to respond effectively to a sudden unexpected
6 loss of balance. Visual information about the spatial features of the nearby environment is likely
7 to be of critical importance when the balancing reaction involves limb movements, such as
8 stepping. Rapid, triggered compensatory stepping reactions must be directed and scaled
9 appropriately in order to arrest the center-of-mass motion suddenly evoked by an unexpected
10 perturbation (Maki and McIlroy, 1997; Hsiao and Robinovitch, 1999; Maki and McIlroy, 1999;
11 Pai, 2003), but the control of these reactions must also take into account the location of obstacles
12 and other environmental constraints on foot trajectory and step placement (Zettel et al, 2002a;
13 Zettel et al, 2002b; Maki et al, 2003).

14 These demands suggest a critical role for visual attention, which involves the allocation of
15 cognitive and/or perceptual resources to objects or events in the visual field (Harris and Jenkin,
16 2001). Clearly, it is important for the central nervous system (CNS) to direct visual attention to
17 the immediate environment in order to acquire the spatial information needed to execute
18 compensatory stepping reactions successfully. It is not known, however, whether the CNS is able
19 to acquire the requisite visual information in “real time”, after onset of postural perturbation. An
20 alternate possibility is that this information is acquired prior to perturbation onset and
21 subsequently integrated with on-line sensory feedback about the body motion induced by the
22 perturbation. This control strategy would have the advantage of avoiding potential delays in
23 initiating the response (associated with time required to plan and execute eye and/or head
24 movements to acquire the visual information and then process this information), but would

1 instead necessitate reliance on a pre-formed egocentric spatial map of the immediate environment,
2 which would have to be updated automatically as we move about in our daily lives (Colby, 1998;
3 Ghafouri et al, 2004).

4 It is also possible that the CNS uses eye movements toward the anticipated landing site to
5 help guide the foot trajectory and step placement, during the execution of the stepping reaction.
6 This would be consistent with findings from studies of targeted volitional stepping movements,
7 which suggest that a saccade toward the step target is used to guide feedforward control of the
8 foot movement (Hollands and Marple-Horvat, 2001; Di Fabio et al, 2003). Interestingly, the
9 timing of this saccade, which typically occurred prior to foot-off during these self-initiated
10 volitional movements (Hollands et al, 1995; Hollands and Marple-Horvat, 1996), appears to be
11 similar to the timing at which we have observed an apparent switching of attention during rapid
12 stepping reactions that are triggered by postural perturbation (Maki et al, 2001a). However, we
13 did not record eye movements in that study. As a result, it is not clear whether the switching of
14 attention was associated with redirection of gaze and acquisition of new visual information, or
15 whether instead this reflected reallocation of cognitive resources for other purposes (e.g. to
16 process sensory information related to the perturbation-induced body motion and to plan the
17 stabilizing motor response).

18 In the present study, we examined whether gaze is redirected downwards, toward the feet
19 or floor, during the execution of forward-directed compensatory stepping reactions evoked by
20 unpredictable postural perturbation. We also examined the timing of any such gaze shifts in
21 relation to the apparent switching of attention that typically occurs prior to foot-off during such
22 reactions (Maki et al, 2001a). We sought to determine whether: 1) the CNS can acquire the
23 necessary visual information about environmental constraints prior to perturbation onset, or must
24 instead redirect gaze at the floor surface during the execution of the reaction; 2) any such need to

1 redirect gaze during the reaction is influenced by increasing demands for accurate control of the
2 foot trajectory; and 3) the switching of attention that typically precedes foot-lift is associated with
3 changes in gaze direction. The findings will be shown to support the view that on-line visual
4 control is not required to meet demands imposed by nearby constraints, and that the switching of
5 attention that follows onset of postural perturbation is not related to on-line acquisition of
6 visuospatial constraint information. Rather, it appears that spatial information about
7 environmental constraints can be acquired beforehand, stored in memory and then used after
8 perturbation onset to aid in guiding the foot trajectory.

10 **Methods**

11 *Subjects*

12 Twelve healthy, naive, young adults were tested (5 male, 7 female; ages 22-29, height
13 160-181cm, mass 54-93kg). All were right-hand and right-leg dominant (as determined by
14 preferred limb for writing and kicking) and reported no medical conditions affecting balance.
15 Subjects were required to have a minimum uncorrected visual acuity of 20/40 (Snellen chart);
16 those who required corrective lenses were excluded, to avoid potential problems in using the eye
17 tracking apparatus. Each subject provided written informed consent to comply with ethics approval
18 granted by the institutional review board.

20 *Protocol*

21 Compensatory stepping reactions were evoked by sudden, unpredictable horizontal
22 movement of a large (2mx2m), computer-controlled moveable platform ((Maki et al, 2003); Fig.
23 1a). Subjects began all trials in a standardized comfortable foot position (McIlroy and Maki,
24 1997) at the center of the platform. For safety, they wore a harness, designed to prevent falls

1 without providing somatosensory feedback that might influence balance control, and safety
2 guardrails and walls were mounted around the platform perimeter. The focus of the study was on
3 forward stepping reactions evoked by large backward platform translations (average acceleration
4 3.0m/s^2 ; peak velocity 0.9m/s , duration 0.6s); however, other perturbation directions and
5 magnitudes were included for purposes of unpredictability. Each block of trials comprised, in
6 random order: six large backward platform translations, two “catch” trials (no platform motion)
7 and eight additional platform translations of various direction (forward, backward, left or right)
8 and magnitude ($0.13\text{--}3.0\text{m/s}^2$; $0.2\text{--}0.9\text{m/s}$).

9 A visuomotor tracking task was performed concurrently in order to monitor switching of
10 attention during the perturbation reactions (McIlroy et al, 1999; Maki et al, 2001b; Norrie et al,
11 2002). This task involved pursuit tracking of a continuously-moving, pseudorandom target (sum
12 of sines: 0.24 , 0.28 , 0.42 Hz; see Fig. 2a.1, 2b.1) displayed for 20s on a computer screen (1.2m in
13 front of the subject). The pursuit cursor was controlled by using the right thumb to rotate a
14 potentiometer (splinted to the wrist and supported by a sling and restraining strap that held the
15 hand comfortably in front of the abdomen). Onset of significant error in tracking performance,
16 subsequent to perturbation onset, was inferred to reflect switching of attention from the tracking
17 task to the balance-recovery task. The timing of perturbation onset was varied unpredictably;
18 however, to facilitate accurate detection of perturbation-related tracking deviation, the
19 perturbations were always delivered during relatively straight sections of the target waveform,
20 where unperturbed tracking accuracy is highest. Practice “tracking-only” trials ($n=6$) preceded the
21 perturbation trials. An additional 16 tracking-only trials (performed at start, midpoint and end of
22 session) were used to determine the thresholds for onset of significant tracking deviation (see
23 Maki et al, 2001b for details).

Blocks of trials were performed for each of four environmental-constraint conditions (Fig. 1): 1) *no-constraint*, 2) *obstacle-only*, 3) *target-only* and 4) *obstacle-plus-target*. The obstacle was a styrofoam-covered metal bar (3.5cm diameter, 150cm length) that was mounted transversely on the moving platform, in front of the subject (2.5% of body height from the toes) at a height equal to 12.5% of body height; a block of foam rubber (2.4cm thick) filled the gap between the bar and the platform surface. In the target tasks, a line of red tape (2cm x 75cm), extending forward in front of each foot, was used to indicate the required medio-lateral (m-l) step placement (forward step length was left unrestricted so as not to compromise ability to arrest the forward perturbation-induced body motion). At the start of each trial block, the subject's attention was directed to the constraint conditions when the task instructions were given. The subject was also free to look at the floor during the interval between trials (approximately 30 seconds). However, once a trial had started and the subject was performing the tracking task, it was not possible to see the obstacle or step targets (in the central or peripheral fields) without redirecting gaze downward, away from the tracking-task display.

Each subject performed two blocks of randomized trials (16 per block) for each of the four constraint conditions, the order counterbalanced both within and across subjects. Two large backward-translation trials preceded the first block of randomized trials. Trials alternated between tracking and no tracking (subjects looked straight ahead at a static visual target on the computer screen). The no-tracking trials were included to allow comparison with previous studies and will not be discussed further here. The protocol yielded 25 large-backward-translation tracking trials per subject, all of which involved forward stepping reactions. These trials were the focus of the analysis.

Vigilance to the tracking task was encouraged by a monetary reward. Subjects were told to do whatever came naturally to prevent falling, but they invariably stepped in the large-

1 perturbation trials. For constrained trials, they were told to avoid contacting the obstacle and/or to
2 direct the step (if they needed to step) so as to land the great toe on the target tape, and that failure
3 to do so would be penalized (reducing their chances of winning the monetary reward for best
4 tracking performance).

5 6 *Measurements and analysis*

7 A lightweight, video-based eyetracker (Applied Science Laboratories, model 501;
8 Bedford, MA) was worn on the head and used to record movements of the left eye, at a sampling
9 rate of 60Hz. This system uses infrared corneal reflections to determine the gaze direction,
10 relative to the head. It displays this information as crosshairs (indicating the horizontal and
11 vertical position of the point of gaze) which are superimposed on the video images recorded by a
12 “scene camera” mounted rigidly on the head (see Figs. 2c,d). The scene camera records changes
13 in the field of view resulting from changes in head position and orientation, and the crosshairs
14 displayed on the video indicate the object in the environment at which gaze is directed. By
15 playing back these video images frame by frame, we were able to determine whether the subject
16 looked downward toward the floor following perturbation onset (using head and/or eye
17 movements), as well as the onset time of each such downward gaze shift (to within 16ms). The
18 same method was used by Patla and Vickers (1997) in a study of obstacle avoidance during gait.

19 Onset of significant error in the tracking task was determined by comparing the tracking
20 performance to the thresholds determined from the tracking-only trials (see (Maki et al, 2001b)
21 for details). Two forceplates, embedded flush with the surface of the moving platform, were
22 sampled at 200 Hz to determine time of foot-off and foot-contact (swing-foot unloading or
23 reloading < or > 5% of body weight). Video recordings from four overhead cameras were used to
24 identify obstacle contact, to measure step-to-target accuracy (by resolving, to within 1cm, the

1 position of a reflective marker on the great toe relative to a grid marked on the platform surface),
2 and to verify foot-off and foot-contact times determined from the forceplates. All timing values
3 were defined relative to onset of platform acceleration (0.1m/s^2), as recorded by an accelerometer.
4 Only gaze and tracking deviations that occurred prior to foot-contact were considered. Example
5 tracking, gaze and force-plate data are shown in Fig. 2.

6 For five subjects, electromyographic (EMG) activity was recorded by surface electrodes
7 placed over the right extensor digitorum muscle, in order to check for the presence of
8 perturbation-evoked activation that could potentially interfere with the performance of the
9 visuomotor tracking task (McIlroy and Maki, 1995). The EMG signal was bandpass filtered (10-
10 500Hz) and sampled at a rate of 1000 Hz. Onset of significant activation was detected using a
11 computerized algorithm (rectified magnitude > 3 standard deviations above pre-perturbation level
12 for $>25\text{ms}$) and confirmed by visual inspection.

13 Repeated-measures analysis of variance was performed to assess the effect of constraint
14 condition on frequency and onset timing of downward gaze shift and tracking deviation during
15 stepping. For the frequency analyses, the dependent variable was the percentage of trials (as
16 determined within each subject, for each of the four task conditions) in which gaze shift or
17 tracking deviation occurred. Of the 300 trials potentially available for analysis, two trials were
18 excluded because a subject misunderstood instructions, and another trial was dismissed because a
19 subject stepped high as if to clear an obstacle even though no obstacle was present. In addition,
20 due to technical problems, gaze data were not available for five trials and tracking data were not
21 available for twelve trials.

Results

Gaze redirection

The redirection of gaze toward the floor, during the stepping reactions, was highly dependent on the constraint condition, and occurred more frequently as the demands for accurate control of the foot movement increased ($F(3,47)=8.04$, $p=0.0004$; Fig. 3). In the most demanding task, involving the obstacle and step target, there was a gaze shift toward the floor, prior to foot-contact, in 37% (27/73) of trials. This rate dropped to 24% (16/68) when only the step target was present. Downward gaze shift almost never occurred in the absence of the step target (3 of 70 *obstacle-only* trials; 1 of 69 *no-constraint* trials). The tendency to look down was primarily limited to six subjects, who each looked down in four or more of the twelve step-target trials. Of the remaining subjects, four looked down in only one of these trials and two did not look down at all. There was no evidence of substantive changes in gaze behavior due to increasing familiarity or “practice” effects: subjects were equally likely to look downward in their first three trials involving the step target as in their last three step-target trials (in each case: 12 of 36 trials).

The timing of the downward gaze shift varied widely, with onset times ranging from 140 to 710ms after perturbation onset. On average, however, the change in gaze direction occurred around the time of foot-off, nearly 300ms prior to foot-contact (Fig. 4a,b; see also Fig. 2a.2). In relation to perturbation onset, the gaze shift actually began more than 100ms earlier, on average, in the *target-only* trials, in comparison to the *obstacle-plus-target* trials: 337 ± 119 ms (mean $\pm$ s.d.) versus 449 ± 159 ms ($F(1,14)=5.83$, $p=0.02$). In relation to foot contact, the difference in gaze-shift timing was somewhat less pronounced: onset of gaze shift preceded foot contact by 224 ± 131 ms in *target-only* trials, versus 287 ± 157 ms in *obstacle-plus-target* trials ($F(1,14)=1.67$, $p=0.20$). This latter finding reflects the fact that foot-contact occurred much later in the *obstacle-plus-*

1 *target* trials (560 ± 76 ms versus 737 ± 130 ms; $F(1,14)=24.4$, $p<0.0001$), due to the large increase in
2 swing duration required to step over the obstacle (Fig. 4b).

3 Gaze redirection had no effect on the ability to avoid the obstacle. The obstacle was
4 cleared successfully (without contact) in 97% of trials regardless of whether subjects did or did
5 not look downwards (successful clearance in 29 of 30 gaze-shift trials and 110 of 113 no-shift
6 trials). On the other hand, downward gaze deviation did appear to improve ability to land the foot
7 on the step target. For *target-only* trials, the success rate was 81% (13/16) when subjects looked
8 down prior to foot-contact, versus 52% (27/52) when they did not look down. There was a
9 similar improvement in success rate during *obstacle-plus-target* trials: 74% (20/27) when looking
10 down versus 48% (22/46) when no downward gaze shift occurred. Mean absolute step-to-target
11 error ($|m-l \text{ distance}|$ from great toe to target tape) improved by 2cm when downward gaze shift
12 did occur (21 ± 20 mm versus 41 ± 39 mm). “Practice” appeared to lead to some small improvements
13 in ability to land on the target. In comparing the first three step-target trials versus the last three,
14 success rates were 75% (9/12) versus 83% (10/12) in trials where gaze shift occurred, and 29%
15 (7/24) versus 46% (11/24) in trials where subjects did not look downwards.

16 17 *Attention switching*

18 An apparent switching of attention from tracking task to balance-recovery task (as
19 reflected by sudden significant deviation in tracking performance, e.g. Fig. 2a.3, 2b.3) followed
20 perturbation onset in 84% (235/280) of trials, while 9% (26/280) of trials had no tracking
21 deviation (in 19 trials, tracking performance was too inconsistent to allow onset of perturbation-
22 related tracking deviation to be determined reliably; these trials were omitted from the tracking-
23 related analyses). Although tracking deviation occurred with high frequency in all constraint
24 conditions (Fig. 3), there was a small but significant constraint-related effect ($F(3,47)=3.05$,

p=0.04), with tracking deviation occurring most consistently in the most demanding task (69 of 70 of *obstacle-plus-target* trials; 99%) and least consistently in the least demanding task (51 of 63 *no-constraint* trials; 81%). The frequency of tracking deviation in the other two conditions was at an intermediate level (91%, 59/65, for *obstacle-only* trials; 89%, 56/63, for *target-only* trials).

The average onset time of tracking deviation was 231 ± 107 ms, and did not differ significantly between constraint conditions ($F(3,46)=0.94$, $p=0.43$; Fig. 4). As in our previous study of attention switching during compensatory stepping (Maki et al, 2001a), the tracking deviation almost always began prior to foot-off (217 of 235 cases; 92%). The average interval between onset of tracking deviation and foot-off was 165 ± 110 ms.

An absence of attention switching, as inferred from an absence of tracking deviation, did not appear to compromise ability to contact the step target or avoid the obstacle. Average absolute step-to-target error was 34 ± 34 mm in trials with tracking deviation ($n=125$) versus 35 ± 29 mm in the eight step-target trials where tracking deviation did not occur. For obstacle trials, successful obstacle clearance occurred in 98% (125/128) of trials with tracking deviation versus 100% (7/7) of trials where tracking deviation did not occur.

The immobilization of the tracking arm appeared to be effective in inhibiting perturbation-evoked reactions in the distal arm muscles; hence it is unlikely that the attention-switching results were confounded by “motor interference”. Performance of the tracking task required continuous activation of the extensor digitorum muscle, which we monitored in five of the subjects. Previous studies have shown that postural perturbation can evoke a burst of activity in this muscle (Maki and McIlroy, 1997); however, this was typically absent in the present trials. Within the 111 trials where extensor digitorum EMG was monitored and tracking deviation occurred, there was no perturbation-related activation in 86 cases (76%). For the 25 trials where the perturbation did evoke a burst of extensor digitorum activity, the onset of the burst occurred

after the onset of tracking deviation in five cases. This leaves only 20 of 111 trials (18%) where the perturbation-evoked reaction could have potentially caused the tracking deviation.

Relation between gaze redirection and attention switching

There were a total of 261 trials where the presence or absence of tracking deviation and gaze shift could both be determined reliably. Evidence of attention switching (i.e. significant tracking deviation) occurred in 90% (235/261) of these trials, but gaze shift was detected in only 17% (45/235) of the tracking-deviation trials. While aversion of gaze from the computer screen would obviously be a cause of subsequent tracking error, the start of the gaze shift preceded (or coincided with) the onset of tracking deviation in only ten trials. In the remaining 35 trials, the tracking deviation preceded gaze deviation (mean latency: 220 ± 116 ms versus 420 ± 150 ms); however, there was little evidence of the consistent temporal coupling that would be expected if the attention switching was related to the motor control of the gaze shift or the subsequent acquisition of new visual information. The time interval between the onset of tracking deviation and subsequent gaze shift was not at all consistent and varied widely among trials (s.d. 127ms, range 6 to 503ms). See Fig. 5.

Discussion

The present study is, to our knowledge, the first to examine the potential role of eye movements in guiding the selection and execution of balance-recovery reactions. The findings indicate that visual fixation of the foot or floor is not required for accurate control of the foot movement during the execution of rapid, perturbation-triggered stepping reactions. Even when demands for accurate limb movement were increased by means of an obstacle and/or step target, subjects were commonly able to avoid the obstacle and/or land on the target without looking downward. Although the CNS is able to acquire substantial information about objects that are

located in the peripheral visual field (Lyon, 1990), it was not possible for subjects in the present study to see the obstacle or step targets, in any part of the visual field, without redirecting gaze downward (away from the eye-level tracking display). The finding that gaze redirection commonly did not occur during the stepping reactions is therefore consistent with the view that the required spatial information about environmental constraints can be acquired beforehand, stored in memory and then used after perturbation onset to aid in guiding the foot trajectory. It should be noted, however, that these conclusions are specific to the current experimental conditions, i.e. relatively simple environmental constraints that remained static over the course of repeated trials. Ongoing work is addressing whether similar gaze behavior occurs when responding to postural perturbation in more complex, dynamic environments.

The findings also indicate that the apparent switching of attention that follows onset of postural perturbation is not related to redirection of gaze. Evidence of attention switching (i.e. significant tracking deviation) occurred in 85% of trials, but gaze shift was detected in only 17% of these trials. When gaze shift did occur, it usually began after the onset of tracking deviation, which would indicate that eye or head movement *per se* was not the cause of the tracking error. Furthermore, the absence of any consistent temporal coupling between the tracking and gaze deviations would appear to rule out the possibility that the switching of attentional resources was associated with the motor control of the gaze shift or the subsequent acquisition of new visual information. Rather, it would seem that the attentional demands were primarily associated with other aspects of balance control, likely related to the processing of the perturbation-induced sensory discharge (Shumway-Cook and Woollacott, 2000; Redfern et al, 2001; Quant et al, 2004) and/or the planning of the foot trajectory (Maki et al, 2001a) or other aspects of the motor response (Rankin et al, 2000; Brauer et al, 2002). A possible relation to foot-trajectory planning is supported by the present findings that the attention switching almost always preceded foot-lift,

1 and that attention switching was most likely to occur when the constraint conditions demanded
2 the most accurate trajectory control. In any case, the fact that there was seldom a complete
3 cessation of efforts to track the target and that the eye movements often persisted in following the
4 target even after the onset of significant error in tracking (e.g. see Fig. 2b.2) would suggest that
5 the attention switching involved a partial, rather than complete, reallocation of cognitive
6 resources to the balance-recovery task.

7 It has been suggested that the tracking-task paradigm may yield delayed estimates for the
8 onset of attention switching if the tracking involves any feedforward (predictive) control elements
9 (Redfern et al, 2002). Any such feedforward control could cause the appearance of tracking error
10 to lag behind the true onset of attention switching. The pseudorandom target waveforms that were
11 used did, in fact, have certain predictable features that may have allowed subjects to use
12 predictive control over short intervals (e.g. the fact that the target motion must necessarily reverse
13 direction whenever the target approaches the limits of the display monitor). The fact that some
14 subjects were able, in a small number of trials, to track successfully for 50-250ms after looking
15 away from the target display (see Fig. 5) may reflect a capacity for short-term predictive control.
16 We are currently performing experiments to document the extent to which different subjects are
17 able to use predictive control in our tracking task and to quantify the degree to which the tracking
18 deviation may lag behind the true onset of attention switching. However, it is important to
19 recognize that these issues are unlikely to invalidate the main conclusions of the present study, i.e.
20 that gaze redirection was not required for accurate stepping movement and that the initial
21 switching of attention that followed perturbation onset was typically not related to redirection of
22 gaze. In fact, if the true onset of attention switching did occur earlier than the estimates derived
23 from the tracking-deviation data, this would serve to further weaken the evidence for any causal

1 association between the gaze shifts and the attention switching, by further reducing the number of
2 trials where the gaze shift preceded attention switching.

3 It is perhaps surprising that saccades were not consistently used, after perturbation onset,
4 to assist in on-line control of the foot movement, particularly when required to step to a target.
5 Studies of volitional limb movement would suggest that such eye movements could potentially
6 help to guide the limb toward the target during the perturbation reactions. For example, studies of
7 volitional upper-limb reaching or pointing movements have demonstrated that visual fixation of
8 the target improves movement accuracy (Abrams, 1992), and studies of volitional step-to-target
9 movements have suggested that a saccade toward the target, completed prior to foot-off or early
10 in the swing phase, is used to control the foot movement (Hollands et al, 1995; Hollands and
11 Marple-Horvat, 1996; Hollands and Marple-Horvat, 2001; Di Fabio et al, 2003). For obstacle
12 clearance, there is apparently less need for precise foot-movement control. Studies of volitional
13 stepping indicate that step height can be exaggerated to guarantee clearance (Chou et al, 1997),
14 and that the required visual information can be acquired prior to initiating the obstacle-clearing
15 step (Patla and Vickers, 1997). The present results would suggest that a similar obstacle-
16 avoidance strategy can be used during stepping reactions that are triggered by sudden
17 unpredictable postural perturbation.

18 There are a number of potential reasons why visual fixation of the target did not occur
19 consistently during the perturbation-triggered stepping reactions. The CNS may actually act to
20 inhibit eye and head movements during stabilizing reactions, since such movements could
21 exacerbate instability (Hunter and Hoffman, 2001) or impede ability to acquire spatial orientation
22 or self-motion information from the visual flow (Brandt et al, 1973). It is also possible that
23 difficulty in disengaging attention from the tracking task impeded the redirection of attentional
24 focus to the step target, which some researchers have postulated must precede the saccade to a

1 target (McFadden and Wallman, 2001). Although subjects were told that inaccurate step-to-target
2 performance would be penalized (reducing the changes of winning the monetary reward for most
3 accurate tracking), they may have nonetheless given top priority to the tracking task, and the fact
4 that the tracking target was not extinguished after perturbation onset may have further impeded
5 saccade execution (McFadden and Wallman, 2001). As noted earlier, there was, in fact, a
6 tendency for the eye movements often to persist in following the tracking target throughout the
7 trial, even after the onset of significant error in thumb-tracking performance (see Fig. 2b.2).

8 Because of the rapid speed at which the triggered stepping reactions were completed, one
9 might also anticipate that temporal constraints could impede effective use of visual fixation to
10 guide the foot movement; however, it appears that the CNS is, in fact, capable of initiating the eye
11 movements with sufficient speed. In the *target-plus-obstacle* trials, gaze shifts were initiated
12 nearly 300ms prior to foot-contact, on average. Even in the *target-only* trials, in which foot-
13 contact occurred much earlier (560ms after perturbation onset, as compared to 740ms in *target-*
14 *plus-obstacle* trials), the gaze shifts were still initiated nearly 250ms prior to foot-contact, on
15 average. In individual trials, gaze shifts were initiated as early as 595ms prior to foot-contact.
16 Such “lead times” would appear to be sufficient to enable useful visual information to be
17 acquired. It is well known that the saccadic eye movement itself could be completed in about
18 50ms, whereas an interval on the order of 300ms might be needed for the CNS to extract the
19 required visual information after completion of the saccade (Harris and Jenkin, 2001). The
20 functional value of the gaze shifts observed in the present study is supported by the fact that the
21 mean step-to-target accuracy did appear to improve, to some extent, in trials where downward
22 gaze deviation occurred. However, it should be noted that at least some of the gaze shifts that
23 were recorded very likely occurred too late to provide useful information for online control of the

1 limb trajectory. In these trials, it is possible that the gaze shifts served another purpose, e.g. efforts
2 of subjects to ascertain whether they had been successful in landing on the target.

3 Despite the apparent use of gaze shifts to help guide the foot movement in some of the
4 step-target trials, it should be reiterated that subjects were, in the majority of trials, able to avoid
5 the obstacle and/or land on the target in the absence of any downward gaze redirection. This
6 would indicate that “remembered” visuospatial information about the support surface, acquired
7 prior to perturbation onset, was incorporated into the control. The static, unchanging obstacles
8 and targets used in this study may have facilitated the capacity to “remember” this spatial
9 information; however, it is important to recognize that the unpredictable variation in perturbation
10 magnitude and direction precluded the possibility of “memorizing” the required limb trajectories.
11 In order to direct the motion of the swing foot appropriately, it was necessary for the CNS to
12 combine on-line sensory feedback about the perturbation-induced body motion with the
13 “remembered” visual support-surface information. This information is presumably
14 “remembered” within the framework of an egocentric visual map, formed by combining
15 retinotopic information with extraretinal cues pertaining to the orientation of the eyes with
16 respect to the head (Colby, 1998). We have proposed that the CNS automatically formulates
17 such a map in daily life, as a contingency in the event that rapid reaction to loss of balance is
18 suddenly required (Ghafouri et al, 2004). This egocentric mapping may occur in parallel with
19 the allocentric mapping that guides navigational behavior (Colby, 1998). Ongoing work,
20 involving intermittent and unpredictable motion of multiple obstacles, is examining factors that
21 affect how frequently the postulated egocentric map is updated, and how long the map can be
22 “remembered” without updating.

1 **Conclusions**

2 Visual fixation of the foot or floor was not essential for accurate control of the foot
3 movement during the execution of rapid, perturbation-evoked stepping reactions, nor was the
4 apparent switching of attention that followed perturbation onset linked, in any consistent way, to
5 overt changes in visual fixation. Spatial features of the support surface were apparently
6 “remembered” prior to perturbation onset and used to help plan or guide the foot trajectory and
7 step placement, thereby allowing both vision and attention to be directed to other demands during
8 the execution of the balance reaction.

1 **Acknowledgments**

2 Supported by grants from the Canadian Institutes of Health Research (CIHR). BEM was a CIHR
3 Senior Investigator and WEM held a Canada Research Chair in Neurorehabilitation. JLZ held
4 scholarships from the Natural Sciences and Engineering Research Council and from the Health
5 Care, Technology and Place Program (University of Toronto).

References

- Abrams RA (1992) Coordination of eye and hand for aimed limb movements. In: L Proteau, D Elliot (eds) *Vision and Motor Control*, pp 129-152. Elsevier, Amsterdam
- Brandt TH, Dichgans J, Koenig E (1973) Differential effects of central versus peripheral vision on egocentric and exocentric motion perception. *Exp Brain Res* 16: 476-491
- Brauer SG, Woollacott M, Shumway-Cook A (2002) The influence of a concurrent cognitive task on the compensatory stepping response to a perturbation in balance-impaired and healthy elders. *Gait Posture* 15: 83-93
- Chou LS, Draganich LF, Song SM (1997) Minimum energy trajectories of the swing ankle when stepping over obstacles of different heights. *J Biomech* 30: 115-120
- Colby CL (1998) Action-oriented spatial reference frames in cortex. *Neuron* 20: 15-24
- Di Fabio RP, Zampieri C, Greany JF (2003) Aging and saccade-stepping interactions in humans. *Neurosci Lett* 339: 179-182
- Diener HC, Dichgans J, Guschlbauer B, Bacher M (1986) Role of visual and static vestibular influences on dynamic postural control. *Human Neurobiology* 5: 105-113
- Ghafouri M, McIlroy WE, Maki BE (2004) Initiation of rapid reach-and-grasp balance reactions: is a pre-formed visuospatial map used in controlling the initial arm trajectory? *Exp Brain Res* 155: 532-536
- Harris LR, Jenkin M (2001) Vision and attention. In: M Jenkin, LR Harris (eds) *Vision and Attention*, pp 2-17. Springer, New York
- Hollands MA, Marple-Horvat DE (1996) Visually guided stepping under conditions of step cycle-related denial of visual information. *Exp Brain Res* 109: 343-356
- Hollands MA, Marple-Horvat DE (2001) Coordination of eye and leg movements during visually guided stepping. *J Motor Behav* 33: 205-216

- 1 Hollands MA, Marple-Horvat DE, Henkes S, Rowan AK (1995) Human eye movements during
2 visually guided stepping. *J Motor Behav* 27: 155-163
- 3 Hsiao ET, Robinovitch SN (1999) Biomechanical influences on balance recovery by stepping. *J*
4 *Biomech* 32: 1099-1106
- 5 Hunter MC, Hoffman MA (2001) Postural control: visual and cognitive manipulations. *Gait*
6 *Posture* 13: 41-48
- 7 Lyon D (1990) Large and rapid improvement in form discrimination accuracy following a
8 location precue. *Acta Psychologica* 73: 69-82
- 9 Maki BE, McIlroy WE (1997) The role of limb movements in maintaining upright stance: the
10 "change-in-support" strategy. *Phys Ther* 77: 488-507
- 11 Maki BE, McIlroy WE (1999) The control of foot placement during compensatory stepping
12 reactions: does speed of response take precedence over stability? *IEEE Trans Rehabil*
13 *Eng* 7: 80-90
- 14 Maki BE, McIlroy WE, Fernie GR (2003) Change-in-support reactions for balance recovery:
15 control mechanisms, age-related changes and implications for fall prevention. *IEEE*
16 *Engineering in Medicine and Biology Magazine* 22: 20-26
- 17 Maki BE, Norrie RG, Zecevic A, Quant S, Kirshenbaum N, Bateni H, McIlroy WE (2001a)
18 Initiation and execution of rapid postural reactions and stepping movements: which
19 phases require visuospatial attention? In: J Duysens, BCM Smits-Engelsman, H Kingma
20 (eds) *Control of Posture and Gait*, pp 573-576. International Society for Postural and
21 *Gait Research*, Maastricht, Netherlands
- 22 Maki BE, Zecevic A, Bateni H, Kirshenbaum N, McIlroy WE (2001b) Cognitive demands of
23 executing rapid postural reactions: does aging impede attentional switching?
24 *NeuroReport* 12: 3583-3587

- 1 McFadden S, Wallman J (2001) Shifts of attention and saccades are very similar: are they
2 causally linked? In: M Jenkin, LR Harris (eds) Vision and Attention, pp 19-39.
3 Springer, New York
- 4 McIlroy WE, Maki BE (1995) Early activation of arm muscles follows external perturbations of
5 upright stance. *Neurosci Lett* 184: 177-180
- 6 McIlroy WE, Maki BE (1997) Preferred placement of the feet during quiet stance: development
7 of a standardized foot placement for balance testing. *Clin Biomech* 12: 66-70
- 8 McIlroy WE, Norrie RG, Brooke JD, Bishop DC, Nelson AJ, Maki BE (1999) Temporal
9 properties of attention sharing consequent to disturbed balance. *NeuroReport* 10: 2895-
10 2899
- 11 Norrie RG, Maki BE, Staines WR, McIlroy WE (2002) The time course of attention shifts
12 following perturbation of upright stance. *Exp Brain Res* 146: 315-321
- 13 Pai YC (2003) Movement termination and stability in standing. *Exercise and Sports Sciences*
14 *Reviews* 31: 19-25
- 15 Patla AE, Vickers JN (1997) Where and when do we look as we approach and step over an
16 obstacle in the travel path? *Neuroreport* 8: 3661-3665
- 17 Quant S, Adkin A, Staines WR, Maki BE, McIlroy WE (2004) The effect of a concurrent
18 cognitive task on cortical potentials evoked by unpredictable balance perturbations.
19 *BioMed Central Neuroscience* 5: 18
- 20 Rankin J, Woollacott MH, Shumway-Cook A, Brown L (2000) Cognitive influence on postural
21 stability: a neuromuscular analysis in young and elders. *J Gerontol* 55A: M112-M119
- 22 Redfern MS, Jennings JR, Martin C, Furman JM (2001) Attention influences sensory integration
23 for postural control in older adults. *Gait Posture* 14: 211-6.

- 1 Redfern MS, Muller ML, Jennings JR, Furman JM (2002) Attentional dynamics in postural
- 2 control during perturbations in young and older adults. *J Gerontol* 57: B298-B303
- 3 Shumway-Cook A, Woollacott M (2000) Attentional demands and postural control: the effect of
- 4 sensory context. *J Gerontol* 55A: M10-16
- 5 Zettel JL, McIlroy WE, Maki BE (2002a) Can stabilizing features of rapid triggered stepping
- 6 reactions be modulated to meet environmental constraints? *Exp Brain Res* 145: 297-308
- 7 Zettel JL, McIlroy WE, Maki BE (2002b) Environmental constraints on foot trajectory reveal the
- 8 capacity for modulation of anticipatory postural adjustments during rapid triggered
- 9 stepping reactions. *Exp Brain Res* 146: 38-47

Captions for figures

Figure 1: Schematic drawings of the obstacle and step targets, as seen from a side view (A) and overhead view (B) of the moving platform. The obstacle height and the location of the obstacle and target lines, in relation to the subject, are drawn approximately to scale. The large arrow denotes the backward direction of platform motion used to evoke the forward-step reactions.

Figure 2: Example data from one subject. Data are shown for an obstacle-plus-target trial (A) and an obstacle-only trial (B), where significant tracking deviation was (A) or was not (B) accompanied by downward gaze shift toward the floor. Plots A.1 and B.1 each display the recorded point-of-gaze and thumb-tracking signals, superimposed on the tracking target waveform, for the full 20-second duration of the trial. The target moved vertically up and down on the computer screen located in front of the subject; the plots display the vertical position of the point-of-gaze and thumb-tracking cursor in relation to the screen. The rectangular box in plots A.1 and B.1 indicates a 1-second time window during which the platform perturbation was delivered. Plots A.2-A.4 and B.2-B.4 are enlargements of this 1-second time window. Plots A.2/B.2, A.3/B.3 and A.4/B.4 display the vertical point-of-gaze signal, vertical tracking (cursor position) signal and vertical ground-reaction forces, respectively. The calibration (scaling) markings displayed along the right edge of panel B also apply to panel A. Vertical broken lines marked P, TD, GD, FO and FC indicate time of perturbation onset, onset of significant tracking deviation, onset of downward gaze deviation, foot-off and foot-contact, respectively. Note that the onset of tracking deviation, prior to foot-off, occurred in the complete absence of any downward gaze shift throughout the duration of the stepping reaction (trial depicted in panel B), or preceded the gaze shift by a substantive (~300ms) time interval (trial depicted in panel A). For purposes of displaying the point-of-gaze data in panels A and B, kinematic measurements of

head position and orientation (Flock of Birds; Ascension Technology Corp.; Burlington, VT) were integrated with the corneal reflection signals indicating gaze direction relative to the head. However, as detailed in the Methods, a simpler video-based method was used in the actual analysis of the gaze-shift data. Panels C and D illustrate the video-based approach: crosshairs indicating the point of gaze are superimposed on the video recorded by a head-mounted “scene” camera. The example video frames shown in C and D illustrate the gaze behavior occurring shortly after foot-off for the trials depicted in panels A and B, respectively. Note that gaze remains directed at the tracking display in D, whereas in C, gaze has begun to move downward due to a combination of head movement (indicated by the change in scene) and eye movement (indicated by the change in crosshair position relative to the scene-camera field of view).

Figure 3: Relative frequency of downward gaze shift and tracking deviation, according to constraint condition. Each bar represents the percentage of trials in which a downward gaze shift or tracking deviation occurred. Note the high frequency of tracking deviation, across all conditions, and the much lower frequency of gaze shifts. Note also that the gaze shifts were almost entirely limited to the conditions that involved the step target.

Figure 4: Timing of key events. In Panel A, mean time of foot off, foot contact, onset of downward gaze shift and onset of tracking deviation is shown for each of the four constraint conditions, with error bars indicating the standard deviation. Note that tracking deviation consistently began prior to foot-off, in all constraint conditions. No gaze-timing data are shown for the no-target tasks because gaze shift was almost exclusively limited to the two tasks involving the step target (even in these tasks, gaze shift occurred in a minority of trials, as indicated in Fig. 3). In the few trials where downward gaze shift did occur, it began 112ms earlier, on average, in the *target-only* task, in comparison to the *obstacle-plus-target* task (*

p<0.05). Panel B shows the time interval by which the onset of gaze shift preceded foot-contact, as well as the swing duration (time interval between foot-off and foot-contact). Note that foot-contact was delayed in the obstacle trials (see panel A; § p<0.05) because a much prolonged swing duration was needed to clear the obstacle (panel B; ¶ p<0.05).

Figure 5: Scatterplot showing the absence of a consistent temporal relationship between gaze shift and switching of attention (as inferred from onset of significant tracking deviation). Onset of tracking deviation is plotted against onset of downward gaze shift, for the 45 trials where both gaze shift and tracking deviation were observed to occur. For data values that fall below the superimposed line (79 % of trials), the tracking deviation began before the start of the downward gaze shift; hence, the tracking error cannot be explained by the aversion of gaze away from the tracking display in these trials. More generally, the wide variation in the relative timing of the two events would appear to rule out any consistent causal link.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5