Local Environmental Impacts of Alternative Forms of Residential Development

Tenley Conway
Department of Geography
University of Toronto, Mississauga
3359 Mississauga Rd N
Mississauga, ON L5L 1C6
Canada
T:1-905-828-3928
tenley.conway@utoronto.ca

Local Environmental Impacts of Alternative Forms of Residential Development

Abstract

New Urbanism is often presented as an improvement over conventional suburban development along economic, social, and environmental lines. While the economic and social claims of new urbanism have been investigated, relatively little work has examined the potential environmental impacts of New Urbanism as compared to conventional forms. One of the challenges of studying the environmental impacts of New Urbanism is that it is still too early to adequately evaluate many sites as most New Urban development is less than a decade old, but it often takes several decades for vegetation and related ecological processes to establish. This paper attempts to address the gap in our understanding of New Urban environmental conditions by examining the relationship between residential parcel vegetation abundance and specific components of residential form across the Greater Toronto Area (Ontario, Canada). While not explicitly comparing New Urban sites to other locations, this analysis illuminates how many New Urban design principles are related to vegetation conditions by focusing on those aspects of residential form that are addressed by New Urbanism— road pattern, neighborhood land use, and site-level characteristics— across a region with heterogeneous form. The results indicate that locations aligned with New Urbanism design principles do not necessarily support more vegetation than those areas based on more conventional suburban patterns, with many design principles having no relationship to vegetation abundance. The paper concludes by examining the planning implications of the results, and outlines future research needed to more fully understand the long-term impact of New Urbanism on local environmental conditions.

1 Introduction

Since the start of the Twentieth Century, urban growth in North America has increasingly taken the form of single use, auto-dependent, low density development (Jackson, 1987; Squires, 2002). As a result, cities are often described as polycentric entities sprawling in fractal and spider-like configurations (Batty and Xie, 1996; Berry and Kim, 1993; Gottdiener and Klephart, 1991; Longley and Mesev, 2000; White and Engelen, 1993). In response to the ‘sprawling’ pattern that now dominates many metropolitan areas, alternative forms of development are being discussed by both academics and planners, with much of the attention given to New Urbanism (Young, 1995). Proponents argue that unlike recent urban forms, New Urbanism integrates traditional urban morphological characteristics to create vibrant, pedestrian friendly communities that have a relatively low environmental impact (Congress for the New Urbanism, 2000). Although carried out in a variety of ways, New Urbanism is often implemented through traditional neighborhood developments, which include high building densities, mixed uses, grid street patterns, narrow streets, and short setbacks (Talan, 2001; Thompson-Fawcett and Bond, 2003).

Social and economic conditions have received significant attention in the literature on suburban sprawl, New Urbanism and other urban forms (c.g. Bjelland et al., 2006), but considerably less research has investigated conditions of the physical environment (Berke, 2002). While adoption of New Urbanism and related development types may present a better environmental option than conventional suburban development (Ellis, 2002)— due to lower per person land consumption and reduction of auto dependency— most New Urbanism developments are still located on greenfields (Berke et al., 2003). Conversion of greenfield sites to even to a “smart” urban form radically impacts local conditions through increased water pollution, heat

island effects, and destruction and modification of vegetation (Sukopp and Hejný, 1990). In addition, the intensity of localized environmental impacts has the potential to be much greater in the higher density developments associated with New Urbanism (Nilon et al., 1995).

One of the challenges of studying the local environmental impacts of New Urbanism is that it is still too early to adequately evaluate many sites (Ellis, 2002). Most New Urban developments are less than a decade old, but it often takes several decades for vegetation to establish, making it premature to assess vegetation conditions and related processes (micro-climate, surface hydrology, pollution filtration). As a result, an approach like the paired-case method, where each New Urbanism community is evaluated along side a comparable conventional suburban development (Berke et al., 2003; Youngentob and Hostetler, 2005), is not necessarily appropriate when considering local environmental impacts. However, in the absence of empirical studies, we have no idea if the touted environmental benefits of New Urbanism exist, or if these communities perform no differently than conventional suburbs. While difficult, the need to better understand these issues in the short-term is imperative as more and more communities adopt New Urbanism principles in an attempt to meet a variety of goals, including protection of the physical environment.

This paper begins to address the gap in our understanding of local environmental conditions in New Urban and conventional suburban developments by examining the relationship between vegetation abundance and specific aspects of residential form across the Greater Toronto Area (Ontario, Canada). While not explicitly comparing New Urban sites to other locations, this analysis will illuminate how many New Urban design principles are related to vegetation conditions by investigating the specific aspects of residential form that are addressed by New

Urbanism— road pattern, neighborhood land use, and site-level characteristics— across a region with heterogeneous form. A similar approach has been employed by Greenwald (2003) to determine the potential influence of new urban design principles on travel mode choice. Given that many New Urban communities directly model themselves on historical development patterns, exploring existing areas with similar design features is reasonable.

The focus of the study is on vegetation because of the important role urban vegetation plays in local environmental conditions; vegetation contributes to air filtration, micro-climate regulations, noise reduction, rainwater drainage, sewage treatment, habitat availability, and recreation opportunities (Bolund and Hunhammar, 1999). A simple vegetation index (NDVI- the Normalized Difference Vegetation Index) is used to represent vegetation abundance given the large spatial extent of the study area. The following sections describe the environmental impact of New Urbanism and the analysis of the Toronto case study. The paper concludes with a discussion of the planning implications of the analysis.

2 New Urbanism and Environmental Impacts

The principles of New Urbanism have been outlined in a series of documents (Calthorpe, 1993; Congress for the New Urbanism, 2000; Duany and Plater-Zyberk, 1991) and implemented in a growing number of communities throughout North America (Bressi, 2002; Grant, 2006). New Urbanism is presented as an alternative to conventional suburbs, which are characterized as having relatively low densities, separation of land uses, a lack of housing variety, and reduced road connectivity.

To date, most evaluations of New Urbanism have focused on social and economic goals (Ellis, 2002; Grant, 2006). In addition to whole hearted supporters

(c.g. Bressi, 2002), several studies have identified potential shortcomings of New Urbanism, including evidence that counters its central goals: New Urbanism may not create to a stronger sense of community (Nasar, 2003), tends to support very homogenous populations (Grant, 2002; Talen, 1999), does not include housing for poor people and difficult to house populations (Al-Hindi, 2001), and mixed development in practice includes a relatively narrow array of uses (Grant, 2002).

In addition to social and economic benefits, environmental benefits are often tied to the New Urbanism design approach (Ellis, 2002; c.g. Till, 2001). In many cases, environmental benefits are fore fronted in the selling of such development (Till, 2001), with ecological analogies used throughout the promotional literature describing New Urban communities (Grant, 2006). Calthorpe (1993) states that “nature should provide the order and underlying structure of the metropolis”, while the Congress for the New Urbanism (2000) argues development should respect the local climate and ecology. New Urbanism is often presented as a reaction to wasteful consumption of natural resources (Calthorpe, 1993), and the lack of connection between local ecology and the built environment typical of conventional suburbs (Hough, 1995; Kunstler, 1996). Some have suggested that New Urbanism is more sustainable than conventional suburban developments (c.g. Zimmerman, 2001), and many argue that it ensures resources are preserved for future generations (Calthorpe, 1993; Duany and Plater-Zyberk, 1991; Katz, 1994; Kuntsler, 1996; Landgon, 1994), echoing an oft stated goal of environmental sustainability.

When the actual environmental benefits of New Urbanism are more closely examined, a mixed picture begins to emerge. Studies on environmental impacts have addressed regional and local-level effects. Touted regional benefits associated with New Urbanism are primarily (1) the potential to lower land consumption and create/

maintain more public green space and (2) reduce transportation associated pollution and energy use, through pedestrian friendly design and (in some cases) transit access (Duany and Plater-Zyberk, 1991). These claims have received some criticism, with Skaburskis (2006) arguing that when New Urban developments are examined in the context of regional dynamics, presence of such communities does not necessarily lead to more green space (or a smaller urban footprint) in the long run. Because most New Urban development occurs on greenfields, Leung (1995) concludes it is nothing more than pretty sprawl.

Studies examining impacts from travel behaviour are also mixed. While Shay and Khattak (2005) found lower automobile ownership and usage in New Urban communities, several others did not (Crane, 1996; Crane and Crepeau, 1998; Hygeia and REIC, 1995; Pogharian, 1996). The lack of difference in travel behaviour may be due to factors that cannot be address through local community design, the focus of New Urbanism (Crane and Crepeau, 1998); a result of most New Urbanism communities lacking sufficient or suitable stores and employment to meet residents needs (Crane, 1996; Grant, 2002; Steiner, 1998); and/or because many New Urban developments occur on the urban edge and are not well linked to transit.

Local environmental conditions that may be effected by urban form are hydrology (including water quality), micro-climate, vegetation patterns, and site-level biodiversity (Alberti, 1999). Few studies have compared conditions between conventional suburban development, New Urbanism, and other forms along these lines. Those that have, tend to focus on the analysis of plans given that the actual communities were often still under construction.

Gordon and Tamminga (2002) evaluated 10 New Urbanism communities' secondary plans in Markham (Ontario, Canada) in light of a recently completed

Natural Features study conducted by the municipality. Markham, located in the northern section of the Greater Toronto Area, is the largest adopter of New Urbanism principles in North America. When development is complete, over 20,000 dwelling units will be constructed across several communities based on New Urbanism designs.

In the study, Gordon and Tamminga (2002) identified several strategies developers were encouraged, but were not required to, adopt as a result of the Natural Features study. Their analysis indicated that the majority of communities only minimally or somewhat incorporated the eight strategies under examination. The following strategies were least often adopted: ecological restoration of significant areas, having the urban edge influenced by natural feature patterns, using open space to buffer natural features, and linking internal parks to natural vegetation corridors. While these results may be better than those for conventional development plans—which were not examined in this study—there was clearly not wholesale support for the ecological principles articulated in the Natural Features study.

Berke et al. (2003), who compared site plans for conventional and New Urban developments, found that the plans adopting New Urbanism principles did a better job of protecting local hydrology, with the exception that the higher road densities would likely contribute to increased storm water run-off. However, Brander et al. (2006) argue that adoption of specific mitigation practices, compatible with New Urbanism designs, can minimize run-off and increase infiltration. While micro-climate and New Urbanism has not been directly studied, extensive research has investigated urban micro-climatic conditions, primarily temperature fluxes and water balance. Not surprisingly, micro-climate varies with urban form (c.g. Arnfield, 2006), and higher

building densities, like those found in New Urbanism communities, are usually associated with higher temperatures (Bonan, 2000; Golany, 1996).

Finally, studies examining vegetation conditions in New Urban communities are rare, although this issue has been indirectly considered. Grant (2006) shows that over time houses in New Urbanism developments have become larger, and are sometimes even larger than in surrounding conventional suburbs (Brown and Cropper, 2001). Large houses on small lots may mean less area is available for vegetation, as well as a higher consumption of fossil fuels (for heating and cooling) and a greater contribution to heat island effects.

Studies more generally evaluating the influence of alternative urban forms on vegetation patterns are also scarce. Those that exist tend to focus on the single variable of distance from city center (Medley et al., 1995). Such studies are based on the assumption that metropolitan areas are linear gradients of urban (population, building, employment) densities, which decrease as one moves away from the city centre (McDonnell and Pickett, 1990). Interestingly, this approach is parallel to the transect concept that has been proposed within the New Urbanism literature, which attempts to create linear patterns of decreasing density as a way to separate rural and urban activities (Duany and Talen, 2002). While the urban-rural gradient research has illuminated ecological variations that exist along such gradients, the role of other urban form components (street pattern; housing type; etc.) is not addressed through this approach.

Recently a few studies have considered other aspects, like street pattern (Conway and Hackworth, 2007; Stone, 2004), lot density (Stone, 2004), zoning class (Wilson et al., 2003), current and past land use (Dow, 2000), and housing age (Hope et al., 2003), indicating that other urban form components are related to urban

vegetation conditions. This study more fully explores the relationship between urban form and vegetation, to highlight any differences between conventional suburbs and New Urban design features.

3 Methods

3.1 Study area

The study area includes the five regional municipalities that make up the 5,902 km² Greater Toronto Area (GTA) in Canada (Figure 1). The original city settlement and several other small towns were located on a sandy plain that stretches from the shore of Lake Ontario to 10 km inland. The majority of the region was cleared for agriculture in the 19th Century, with only about six percent of the land in woodlots by the 1940s (Chapman and Putnam, 1984). The 20th century saw rapid urban development expanding outwards from Toronto, converting agricultural lands to the North and engulfing previously separate small towns.

By 2001, the study area had a population just over five million people. With an annual growth rate of 1.9 percent (Statistics Canada, 2001), the GTA is one of the fastest growing large metropolitan regions in North America. In the study area, approximately 13 percent of the land is used for urban residential purposes, eight percent for commercial and industrial activities, and three percent is protected parkland. The remaining 77 percent is agricultural, rural residential development, forests, and wetlands. Today, the region includes several commercial centres, often remnants of previously separate urban settlements; significant industrial and commercial strip development along highways and major roads; traditional older residential areas with neighborhood commercial establishments; and extensive areas of conventional suburban development.

3.2 Urban form variables

Fourteen variables were included in the analysis to reflect neighborhood or site-level design components associated with New Urbanism (Table 1). Several variables representing urban density were included because of New Urbanism's focus on increasing density. The first of the density variables is parcel size, with New Urbanism favouring relatively small residential parcels. Hough (1995) argues that parcel size is likely a key factor limiting vegetation abundance in built-up environments due to physical space limitations. However, Robbins' et al. (2001) study of parcel size effects in suburban Ohio found that as parcel size increases, so does building size, suggesting that there may not be proportionally more room for vegetation on larger parcels.

New Urbanism writings are not as explicit about population density, although one may assume that as the density of building lots increases so would the number of people. However, this is not always the case, as some studies indicate that single people or couples without children are more likely to live in New Urbanism communities than in conventional suburbs (Bookout, 1992). As a result, population density is examined separately from building parcel size, represented by the number of people per square kilometer in each census dissemination area. Dissemination areas contain a population of 400 to 700 people, and are the smallest geographical unit for which all census data is released by Statistics Canada (2001).

In North America, road patterns within residential neighborhoods have changed from the basic grid pattern primarily used in the early 20th Century to the curvilinear streets and cul-de-sacs that dominate post-war development. This change is associated with a decrease in road length and intersection density, leading to less connectivity (Southworth and Owens, 1993). A central tenet of New Urbanism is an

increase in connectivity through the use of a grid road network. Because increased connectivity is usually associated with an increase in road and intersection density this measure was included. Road density was defined as the length of road, in kilometers. Neighborhoods were defined using a 1.0 km radius, following Geoghegan et al. (1997), to represent a walkable distance around each residential parcel. As the variable is defined, most grid networks will have a higher neighborhood road density than curvilinear and cul-de-sac patterns.

Several other variables are also included in the analysis to represent additional aspects of neighborhood land use pattern. The number of different land use classes within a neighborhood is a broad measure of the diversity of functions, with a higher number of land use classes suggesting a mixed use development pattern over a more segregated configuration. This variable reflects the emphasis on mixed use communities in New Urbanism.

Even when mixed uses are emphasized, in practice this can still mean a variety of land use patterns. As a result, the percent of different land uses within a 1.0 km neighborhood were also considered: (1) percent of neighborhood in parkland; (2) percent of neighborhood green space (open area and parkland); and (3) percent of neighborhood used for industrial/commercial purposes.

The percent of renter-occupied dwelling units and two measure of housing structure type (percent of dwellings that are detached or semi-detached single family houses and percent of dwellings in apartment buildings five stories or higher) were included as measures of diversity in residential options. New Urbanism writing often promotes variations in housing styles as a way to support a diverse community, although housing types are often limited to single family or low rise structures.

At the site level, several variables reflecting view from residential properties were identified. View variables were included because previous research in urban landscapes has suggested that residents prefer a view of certain features over others (Kaplan, 2001). Communities with mixed uses or integrated green space provide different views than areas that consist only of residential development. Given the lack of topographic variation across much of the region, viewsheds were defined using a 0.1 km neighborhood around each sample point, following Geoghegan et al. (1997). Similar to neighborhood uses, views of both parkland and green space (parkland and open land) were examined.

3.3 Vegetation data

The Normalized Difference Vegetation Index (NDVI) was chosen to represent vegetation abundance because it is a good measure of the amount of green biomass present (Tucker, 1979), and correlated with urban land cover type (Fung and Siu, 2000) and leaf area index (Xavier and Vettorazzi, 2004). The index is a ratio of near infrared (NIR) and visible red (R) reflective values defined by the following formula:

$$NDVI = \frac{NIR - R}{NIR + R}$$

Heavily vegetated areas have an NDVI value closer to 1, while locations dominated by water, impervious surfaces, or bare soil have values closer to -1.

In this study, NDVI was calculated from Landsat 7 ETM+ satellite imagery. The imagery has a cell size of 30 m, an appropriate resolution for metropolitan-wide studies of NDVI variation (Wilson et al., 2003). Two images were needed to cover the study area. Images collected two weeks apart were used to minimize cloud coverage: Path 18 Row 29 on September 3, 1999 and Path 18 Row 30 on September

19, 1999. The images were georeferenced using the roads data layer from the CanMap Route Logistics 8.2 dataset¹.

3.4 Analysis

For this study, only areas of residential land uses were included in the analysis. Hough (1995) notes that age of construction is one of the most important factors in determining urban vegetation because the construction process usually represents the last major disturbance, with most vegetation removed from the site and new vegetation planted. When trees and other woody species are considered, vegetation levels may continue to increase for the first few decades after development due to natural growth processes. To try to account for these growth processes two analyses were conducted: (1) in residential neighborhoods where the dominate age of construction was prior to 1946 and (2) where the dominate age of construction was between 1946 and 1970 (Figure 1). The first time period likely represents only those neighborhoods with mature vegetation, while the second analysis is able to include the post war period where significant suburban development occurred in the GTA. The first time period included 6,723 data points the second time period includes 26,733 data points. These numbers reflect the relative portion of residential areas in the two time periods, with specific sites selected based on a random sample spatial separated above the level of spatial autocorrelation for NDVI.

The dominate age of construction for each neighborhood was determined based on Statistics Canada 2001 census data at the dissemination level. The percent of dwellings constructed prior to 1946, from 1946 to 1960, and in 10 year spans through 2001 was first calculated. The most active period of construction was then

¹ All data layers were projected in UTM's, and, where necessary, georectified to align with the CanMap Route Logistics 8.2 dataset.

assigned as the dominate age of construction for each dissemination area, and subsets of pre-1946 and 1946-1970 neighborhoods were created. For each sample of residential parcels, rights-of-ways and water bodies were removed to ensure that NDVI values reflect conditions on residential property.

In both samples, residential areas with very low population densities were excluded. Thus, the analyses were conducted within only those areas that have a population density above 400 people per square kilometre, Statistics Canada's definition of an urban area.

A correlation analysis between NDVI and the urban form variables was conducted. All variables listed in Table 1 were used in this analysis. The first step was to examine the relationship between the urban form variables, to determine if certain features (i.e. high road density and low green space) are found in the same location to aid with the interpretation of the vegetation results. The second step looked at the relationship between NDVI and each urban form variable. Based on hypothesized relationships between the urban form variables, partial correlations were calculated to control for certain variables. Specially four questions were addressed: (1) what is the relationship between neighbourhood percent of single family homes and NDVI when controlling for parcel size, (2) what is the relationship between neighbourhood percent of renter-occupied dwelling units and NDVI when controlling for road density, (3) what is the relationship between the industrial- commercial land uses variables and NDVI when controlling for neighbourhood housing type, (4) what is the relationship between neighbourhood green space and NDVI when controlling for road density?

4 Results

Table 2 shows the summary statistics for the two time periods. Based on the Kolmogorov-Smirnov test, all variables have significantly different means and distribution shapes between the two time periods. The most notable differences are for average parcel size and population density, with both variables having higher means in pre-1946 neighborhoods. These seemingly contradictory conditions exist because there are fewer detached single family houses and more (low-rise) apartment buildings in pre-1946 areas. Thus, the average pre-1946 parcel is more intensely used.

Tables 3 and 4 show the correlations between the urban form variables. As might be expected, the strongest correlations for both time periods are between the three housing type variables (percent renter-occupied, percent of dwelling units that are single family houses, percent of dwellings in apartments with five stories or more). Interestingly, population density and parcel size are not correlated with the housing variables or each other. The housing type variables are also correlated with neighborhood industrial/ commercial and views of those land uses. A strong negative relationship exists between road density and neighborhood green space, suggesting that the road density variable is not only picking up changes in road patterns but also differences in areas used more or less intensity for urban purposes. Finally, there were only weak correlations between neighborhood land use variables and the corresponding view variables, suggesting that land use patterns are not related at these two scales (0.1 km and 1.0 km).

The correlations between NDVI and the urban form variables are given in Table 5. In residential areas primarily built before 1946, parcel size and percent single family homes are positively correlated with NDVI. When parcel size is

controlled for, the relationship between NDVI and single family homes remains (0.414, $p < 0.001$). Road density and percent neighborhood industrial/commercial land use are negatively correlated. Weaker correlations also exist for three other variables: percent of neighborhood green space is positively correlated and percent neighborhood industrial/ commercial, view of industrial/ commercial, and the percent of renters are negatively correlated.

When housing type is controlled for the relationship between view of industrial/ commercial and NDVI stays moderate (-0.313, $p < 0.001$) but only a weak relationship exist for the percent of neighborhood industrial/ commercial land uses (-0.206, $p < 0.001$). Furthermore, when the percent of single family homes is controlled for, there is no significant relationship between NDVI and percent renter-occupied. The same is true for NDVI and percent of neighborhood green space when controlling for road density.

The primary difference between the two time periods is that weaker relationships exist for the 1946-1970 residential areas, perhaps due to the younger age of the some of the residential areas leading to less mature vegetation.

5 Discussion

The analysis indicates that a few of the design principles associated with New Urbanism are related to variations in NDVI. One of the factors most strongly correlated with NDVI is neighborhood road patterns, where higher densities are associated with lower levels of vegetation on residential property. Higher road density is a key component of New Urbanism, as it is seen as a way to meet social goals through increased connectivity. Based on this analysis, adoption of a highly connected road pattern, like a grid, may actually increase negative environmental

impacts. Gordon and Tamminga (2002) acknowledge the clear conflict between higher built connectivity (i.e. a grid road pattern) and natural ecosystem connectivity, and suggest that a cul-de-sac design may actually better minimize that conflict. Alternatively, a modified grid pattern, using pedestrian pathways not roads to maintain high levels of built connectivity (CMHC, 2002; Southworth and Ben-Joseph, 2004), could be a compromise solution.

The positive correlation between NDVI and parcel size indicates another dilemma that arises through New Urbanism's design principles. On the one hand larger building lots can support more vegetation, and the positive ecological services (micro-climate regulation, pollution filtration, habitat, etc.) associated with that vegetation. On the other hand, larger parcels mean that less land is available for other uses, including protection of ecological features.

The very weak correlation between population density and NDVI may point to an alternative to this problem. While New Urbanism is usually associated with higher density building lots, most of the residents are still single family structures (detached, semi-detached, or townhouses). Replacement of single family homes with multi-family medium or high rise buildings could potentially allow for a larger percent of land to be kept in a vegetated state. However, this type of approach, which includes the modernist design of the 'high rise in the park', has been criticized by advocates of New Urbanism because it is not seen as aesthetically pleasing or supportive of social interaction (Calthorpe, 1993; Grant, 2006). However, medium or high rise buildings may be a more efficient use of space that still allows for abundant vegetation.

The results do suggest the housing structure and occupant type are related to NDVI. Neighborhoods with a larger percentage of single family dwellings are associated with higher vegetation levels, even when controlling for parcel size. This

is one example where a common feature of New Urbanism is associated with a positive environmental situation. Of course this is also a dominant feature of conventional suburbs. Interestingly, larger apartment buildings (greater than five stories) are not associated with less (or more) vegetation, again suggesting that the 'high rise in the park' style development many supporters of New Urbanism scorn may have some environmental benefits over New Urbanism. However, the negative relationship between percent renter-occupied dwellings and vegetation abundance appears to primarily be due to the influence of fewer single family homes. Thus, while high levels of vegetation abundance may be possible with apartment buildings, these conditions are more likely to occur in areas with high concentrations of single family homes.

The relationship between vegetation and views of industrial/commercial land suggests the influence of these types of uses extends beyond the site they occupy, with industrial/commercial land and surrounding residential parcels tending to have low levels of vegetation. One possible explanation may be that fewer single family homes and more rental units tend to occur in neighborhoods with higher levels of industrial and commercial uses, thus housing type may be influencing the relationship. However, the partial correlations suggest that this is not the case for residential properties with a view of industrial and commercial uses. Alternatively, the positive effects of neighboring green space seem to merely be an artefact of lower road densities in those neighborhoods.

Interestingly, while specific neighborhood land use types are related to residential vegetation, the total number of land uses in a neighborhood is not. Thus, a mixed use design, which is promoted by New Urbanism, in and of itself, may not lead

to lower levels of vegetation, but that certain uses (industrial/commercial) and/or the type of dwellings that usually occur near these uses do.

5.1 Urban environmental impacts

There is growing evidence that New Urban communities do not necessarily support healthier natural systems or reduced local environmental impacts as compared to conventional suburban design. In addition, the relatively small scale of New Urbanism within a metropolitan context, and tendency for such development to be located on the urban fringe, has the potential for most New Urbanism communities to not only be associated with more intensive local environmental impacts, but also to not necessarily encompass the benefits of compact forms- cheaper infrastructure and reduced travel distances. Thus, New Urbanism in practice may represent no more than the process of converting greenfields along the urban edge to high density sprawl (Grant 2006).

To date New Urbanists have focused on creating the illusion that there are no negative environmental effects, but actually draw on a very narrow understanding of ecology in their design principles (Zimmerman, 2001). In particular, New Urbanism glosses over the inherent conflict between built and natural systems. Calthorpe (1993:44) notes that “environmentally sound communities need parks, regional greenbelts, and high-quality open spaces, but they also need density and street-life,” yet the principles associated with New Urbanism do not provide a clear pathway to reconcile these needs.

New Urbanism also ignores what Neuman (2005) calls the compact city paradox: a compact city (i.e. high density along the lines of New Urbanism) is not necessarily a more sustainable one as such cities are associated with their own set of

environmental problems (i.e. pollution, heat islands, little room for vegetation). In fact, minimizing these problems has historically been one of the push factors associated with the development of conventional suburbs (Neuman, 2005).

One way to overcome the environmental shortcomings of New Urbanism may be better integration of landscape ecology, which involves the study of ecological processes across settled landscapes (Turner et al., 2001), into urban design principles. Gordon and Tamminga (2002) point out that there is a wide gap between landscape ecology and New Urbanism, but landscape ecologists have begun to develop a set of strategies to protect natural systems in urbanizing landscapes (MacCracken and Bignal, 1998; Odell et al., 2003; Opdam et al., 2002). Joining landscape ecology principles and a neighborhood-level urban design approach, like New Urbanism, may help elucidate a way to balance built and natural systems in an urban context, ensuring that the environmental promises of New Urbanism can be met.

5.2 Limitations and next steps

This study was an initial attempt to better understand the relationship between urban vegetation in residential neighborhoods based on conventional suburban and New Urban design principals. A major limitation of the study is that the vegetation measure was limited to abundance, and does not take into account species composition. Native species are likely to require less water and other inputs, while more likely to provide habitat to other species. The benefits of planting native species may be even greater when it replaces conventionally managed lawn grass, due to reduction in chemical inputs (Robbins et al., 2001). Site-level surveys of species composition in New Urban and conventional suburbs would begin to determine if developers of New Urbanism are more likely to incorporate native vegetation and

non-lawn alternatives, and if property owners retain or replace that vegetation over time.

A second limitation is that New Urbanism communities were not actually examined in this study due to the relatively recent creation of many of these developments. While analysis of locations that do and do not have some of the same design components as New Urbanism may begin to illuminate conditions associated with this form, this approach does not enable a study of the ways in which different design features work together to influence the local environment and residents' property management. A long-term study tracing conditions in New Urbanism communities would provide a more robust understanding of the environmental impacts of this type of urban form. However, the type of approach used in this study provides some basic insight into the impacts of New Urban design elements.

A study focusing on specific New Urbanism communities would also enable neighborhood boundaries tailored to each site. In this analysis, variables were summarized reflecting using either a standard distance to reflect a 'walkable' neighbourhood or one defined by the census data (i.e. dissemination areas), rather than functional neighborhoods. This was a result of data availability and the large size of the study area. If actually New Urbanism communities were studied, then functional neighbourhoods specific to each site could be identified (c.g. Song and Knaap, 2004), better reflecting actual conditions.

Overall, the generally weak relationship between the urban form variables and NDVI suggests that factors other than design elements included in this analysis are primarily driving urban vegetation conditions. For example the socioeconomic status of households, cultural influences, and municipal or regional urban forestry policy, which have already been shown to play some role in urban vegetation conditions

(Conway and Urbani, 2007; Heynen et al., 2006; Heynen and Lindsay, 2003; Martin et al. 2004; Stone, 2004; Wilson et al., 2003), may play a greater role than urban form. Further research needs to more thoroughly address the relative influence of, and interactions between, each of these broader factors.

6. Conclusions

New Urbanism is often associated with a number of social, economic, and environmental goals. While the social and economic impacts of New Urbanism have been well studied, relatively little attention has been given to the environmental claims. As an initial attempt to better understand the environmental implications of New Urbanism, this study compared vegetation abundance on residential property in relation to urban form variables that differ between conventional suburbs and New Urbanism. The results indicate that these design principles are not necessarily associated with higher levels of vegetation, and may actually be associated with less. These results are inline with a growing discussion that New Urbanism is not necessarily an improvement over conventional suburbs in terms of local environmental conditions, and is far from an environmentally sustainable form of residential development.

References

- Alberti M, 1999, "Urban patterns and environmental performance: what do we know?" *Journal of Planning Education and Research* **19** 151-163
- Al-Hundi KF, 2001, "The New Urbanism: where and for whom? investigation of a new paradigm" *Urban Geography* **22(3)** 202-219
- Arnfield AJ, 2006, "Micro- and mesoclimatology" *Progress in Physical Geography* **30(5)** 677-689
- Batty M, Xie Y, 1996, "Preliminary evidence for a theory of the fractal city" *Environment and Planning A* **28(10)** 1745-1762
- Berke P, 2002, "Does sustainable development offer a new direction for planning? Challenges for the twenty first century" *Journal of Planning Literature* **17(1)** 21-36
- Berke PR, MacDonald J, White N, Holmes M, Line D, Oury K, and Ryznar R, 2003, "Greening development to protect watersheds: does New Urbanism make a difference?" *Journal of the American Planning Association* **69(4)** 397-314
- Berry B, Kim H, 1993, "Challenges to the monocentric model" *Geographical Analysis* **25** 1-4
- Bjelland MD, Maley M, Cowger L, Barajas LB, 2006, "The quest for authentic place: The production of suburban alternatives in Minnesota's St. Croix Valley" *Urban Geography* **27(3)** 253-270
- Bolund P, Hunhammar S, 1999, "Ecosystem services in urban areas" *Ecological Economics* **29(2)** 293-301
- Bonan GB, 2000, "The microclimates of a suburban Colorado (USA) landscape and implications for planning and design" *Landscape and Urban Planning* **49** 97-114

- Bookout LW, 1992, "Neotraditional town planning: the test of the marketplace"
Urban Land **51** 12-17
- Brander KE, Owen KE, Potter KW, 2006, "Modelled impacts of development type on runoff volume and infiltration performance" *Journal of the American Water Resources Association* **30(4)** 961-969
- Bressi TW, 2002 *The Seaside Debates: A Critique of the New Urbanism* (Rizzoli International Publications Inc., New York)
- Brown B, Cropper V, 2001, "New urban and standard suburban subdivisions: evaluating psychological and social goals" *Journal of the American Planning Association* **67(4)** 402-419
- Calthorpe P, 1993 *The Next American Metropolis: Ecology, Community, and the American Dream* (Princeton Architectural Press, New York)
- Chapman LJ, Putman DF, 1984 *The Physiography of Southern Ontario. Ontario Geological Survey, Special Volume 2* (Government of Ontario, Toronto)
- CMHC, 2002, "Residential street pattern design", in *Research Highlights, Socio-economic Series, No. 75*, (Canadian Mortgage and Housing Corporation, Ottawa)
- Congress for the New Urbanism, 2000 *Charter of the New Urbanism: Region/Neighborhood, District, and Corridor/Block, Street, and Building* (McGraw Hill, New York)
- Conway TM, Hackworth J, 2007, "Urban pattern and land cover variation the Greater Toronto Area" *The Canadian Geographer* **51(1)** 43-57
- Conway TM, Urbani L, 2007 "Variations in municipal urban forestry policies: A case study of Toronto, Canada" *Urban Forestry and Urban Greening* **6** 181-192

- Crane R, 1996, "Cars and drivers in the new suburbs: linking access to travel in neotraditional planning" *Journal of the American Planning Association* **62(1)** 51-65
- Crane R, Crepeau R, 1998, "Does neighborhood design influence travel? A behavioral analysis of travel diary and GIS data" *Transportation Research Part D: Transport and Environment* **3(4)** 225-238
- Dow K, 2000, "Social dimensions of gradients in urban ecosystems" *Urban Ecosystems* **4** 255-275
- Duany A, and Plater-Zyberk E, 1991 *Towns and Town-Making Principles* (Rizzoli, New York)
- Duany A, Talen E, 2002, "Transect planning" *Journal of the American Planning Association* **68(3)** 245-265
- Ellis C, 2002, "The New Urbanism: critiques and rebuttals" *Journal of Urban Design* **7(3)** 261-291
- Fung T, Sui W, 2000, "Environmental quality and its changes, an analysis using NDVI" *International Journal of Remote Sensing* **21** 1011-1024
- Geoghegan J, Wainger LA, Bockstael NE, 1997, "Spatial landscape indices in a hedonic framework: An ecological economics analysis using GIS" *Ecological Economics* **23** 251-264
- Golany GS, 1996, "Urban design morphology and thermal performance" *Atmospheric Environment* **30(3)** 455-465
- Gordon DLA, Tamminga K, 2002, "Large-scale traditional neighborhood development and pre-emptive ecosystems planning: the Markham experience, 1989-2001" *Journal of Urban Design* **7(2)** 321-340

- Gottdiener M, Klephart G, 1991, "The multinucleated metropolitan region: a comparative analysis", in *Postsuburban California: The Transformation of Orange County Since World War II* Eds R Kling, S Olin, and M Poster (University of California Press, Berkeley, CA)
- Grant J, 2006 *Planning the Good Community: New Urbanism in Theory and Practice* (Routledge, London)
- Grant J, 2002, "Mixed use in theory and practice: Canadian experience with implementing a planning principle" *Journal of the American Planning Association* **68(1)** 71-84
- Greenwald MJ, 2003, "The road less traveled: New Urban inducements to travel mode substitution for nonwork trips" *Journal of Planning Education and Research* **23** 39-57
- Heynen NC, Lindsay G, 2003, "Correlates of urban forest canopy cover: implications for local public works" *Public Works Management and Policy* **8(1)** 33-47
- Heynen, N, Perkins HA, Roy P, 2006, "The political ecology of uneven green space: the impact of political economy on race and ethnicity in producing environmental inequality in Milwaukee" *Urban Affairs Review* **42(1)** 3-25
- Hope D, Gies C, Zhu W, Fagab WF, Redman CL, Grimm NB, Nelson AL, Martin C, Kinzig A, 2003, "Socioeconomics drive urban plant diversity" *Proceedings in the National Academy of Science* **100(15)** 8788-8792
- Houg M, 1995 *Cities and Natural Process* (Routledge, New York)
- Hygeia, REIC, 1995 *Changing Values, Changing Communities: A Guide to the Development of Healthy, Sustainable Communities* (Canada Mortgage and Housing Corporation, Ottawa)

- Jackson KT, 1987 *Crabgrass Frontier: the Suburbanization of the United States* (Oxford University Press, New York)
- Kaplan R, 2001, "The nature of the view from home: psychological benefits" *Environment and Behaviour* **33(4)** 507-542
- Katz P, 1994 *The New Urbanism: Towards an Architecture of Community* (McGraw-Hill, New York)
- Kunstler JH, 1996 *The Geography of Nowhere: The Rise and Decline of America's Man-made Landscape* (Simon and Schuster, New York)
- Landgon P, 1994 *A Better Place to Live: Reshaping the American Suburb* (Amherst, MA: University of Massachusetts Press)
- Leung H-L, 1995, "A new kind of sprawl" *Plan Canada* **35(5)** 4-5
- Longley PA, Mesev V, 2000, "On the measurement and generalization of urban development" *Environment and Planning A* **32(3)** 473-488
- McCracken DI, Bignal EM, 1998, "Applying the results of ecological studies to land-use policies and practices" *Journal of Applied Ecology* **35** 961-967
- McDonnell MJ, Pickett STA, 1990, "Ecosystem structure and function along urban-rural gradients: an unexploited opportunity for ecology" *Ecology* **71(4)** 1232-1237
- Martin CA, Warren PS, Kinzig AP, 2004, "Neighborhood socioeconomic status is a useful predictor of perennial landscape vegetation in residential neighborhoods and embedded small parks of Phoenix, AZ" *Landscape and Urban Planning* **69** 355-368
- Medley KE, McDonnell MJ, Pickett STA, 1995, "Forest-landscape structure along an urban-to-rural gradient" *Professional Geographer* **47(2)** 159-168
- Nasar J, 2003, "Building in affordability" *Urban Land* **62(5)** 64-65

- Neuman M, 2005, "The compact city fallacy" *Journal of Planning Education and Research* **25** 11-26
- Nilon CH, Long CN, Zipperer WC, 1995, "Effects of wildland development on forest bird communities" *Landscape and Urban Planning* **32(2)** 81-92
- Odell EA, Theobald DM, Knight RL, 2003, "Incorporating ecology into land use planning: the songbirds' case for clustered development" *Journal of the American Planning Association* **69** 72-82
- Opdam P, Foppen R, Vos C, 2002, "Bridging the gap between ecology and spatial planning in landscape ecology" *Landscape Ecology* **16** 767-779
- Pogharian S, 1996, "Street design: learning from suburbia" *Plan Canada* **36(5)** 41-42
- Robbins P, Polderman A, Birkenholtz T, 2001, "Lawns and toxins: an ecology of the city" *Cities* **18(6)** 369-380
- Shay E, Khattak AJ, 2005, "Automobile ownership and use in neotraditional and conventional neighborhoods" *Transportation and Land Development* **1902** 18-25
- Skaburskis A, 2006, "New urbanism and sprawl - A Toronto case study" *Journal of Planning Education and Research* **25(3)** 233-248
- Song Y, Knapp G-J, 2004, "Measuring urban form: is Portland winning the war on sprawl?" *Journal of the American Planning Association* **70(2)** 210-225
- Southworth M, Ben-Joseph E, 2004, "Reconsidering the Cul-de-sac" *Access* **24** 28-33
- Southworth M, Owens PM, 1993, "The evolving metropolis: studies of community, neighborhood, and street form at the urban edge" *Journal of the American Planning Association* **59(3)** 271-284
- Squires GD, 2002 *Urban Sprawl: Causes, Consequences, and Policy Responses* (Urban Institute Press, Washington, D.C)

- Statistics Canada, 2001. Population and Dwelling Counts for Census Divisions, Census Subdivisions (municipalities), and Designated Places (DPL). Last updated October, 17 2005
- Steiner R, 1998, "Traditional shopping centers" *Access* **12** 8-13
- Stone BJ, 2004, "Paving over paradise: How land use regulations promote residential imperviousness" *Landscape and Urban Planning* **69** 101-113
- Sukopp H, Hejný S, 1990 *Urban Ecology* (SPB Academic Publishing, The Hague)
- Talen E, 1999, "Sense of community and neighborhood form: an assessment of the social doctrine of New Urbanism" *Urban Studies* **36(8)** 1361-1379
- Till KE, 2001, "New Urbanism and nature: green marketing and the neotraditional community" *Urban Geography* **23(3)** 220-248
- Thompson-Fawcett M, Bond S, 2003, "Urban intentions for the built landscape: examples of concept and practice in England, Canada, and New Zealand" *Progress in Planning* **60** 147-234
- Tucker C, 1979, "Red and photographic infrared linear combination for monitoring green vegetation" *Remote Sensing of Environment* **8** 127-150
- Turner MG, Gardner RH, O'Neill RV, 2001 *Landscape Ecology in theory and Practice* (Springer, New York)
- White R, Engelen G, 1993, "Cellular-automata and fractal urban development: a cellular modeling approach to the evolution of urban land-use patterns" *Environment and Planning A* **25(8)** 1175-1199
- Wilson JS, Clay M, Martin E, Stuckey D, Vedder-Risch K, 2003, "Evaluating environmental influences of zoning in urban ecosystems with remote sensing" *Remote Sensing of the Environment* **86** 303-321

- Xavier AC, Vettorazzi CA, 2004, "Mapping leaf area index through spectral vegetation indices in a subtropical watershed" *International Journal of Remote Sensing* **25(9)** 1661-1672
- Young D, 1995 *Alternatives to Sprawl* (Lincoln Institute of Land Policy, Cambridge, MA)
- Youngentob K, Hostetler M, 2005, "Is a new urban development model building greener communities?" *Environment and Behavior* **37(6)** 731-759
- Zimmerman J, 2001, "The "nature" of urbanism on the New Urban frontier: sustainable development, or defence of the suburban dream?" *Urban Geography* **22(3)** 249-267

Table 1. Urban development pattern and socioeconomic variables.

Variable Name	Description
Parcel_Size	The size of each parcel in meters squared, using Teranet's 2002 Property Tax Parcel Map Data for Halton, Peel, York, Durham, and Toronto Regions and The City of Toronto's 2002 Parcel data for City of Toronto.
Pop_Denisty	Population density, by dissemination area from 2001 Census
Roads_0.1	Road density within a 0.1 km neighborhood; proxy for view, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Road_density	The density of roads within a 1.0 km neighborhood, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Land_Diversity	Number of land uses present within a 1.0 km neighborhood, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset. Maximum value = 7
Park_0.1	Percent of parkland within a 0.1 km neighborhood; proxy for view, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Park_1.0	Percent of parkland within a 1.0 km neighborhood, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Green_0.1	Percent of land that is open or parkland within a 0.1 km neighborhood; proxy for view, derived from DMTI's 2004

	CanMap Route Logistics 8.2 Ontario dataset
Green_1.0	Percent of land that is open or parkland within a 1.0 km neighborhood, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Ind-Comm_0.1	Percent of land used for industrial and commercial purposes within a 0.1 km neighborhood; proxy for view, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Ind-Comm_1.0	Percent of land used for industrial and commercial purposes within a 1.0 km neighborhood, derived from DMTI's 2004 CanMap Route Logistics 8.2 Ontario dataset
Per_Renter_Occ	Percent of dwelling units occupied by renters, by dissemination area from 2001 Census
Per_Single_Fam	Percent of dwelling units that are detached, single family homes, by dissemination area from 2001 Census
Apt $\geq$ 5	Percent of dwelling units in apartment building five stories or higher, by dissemination area from 2001 Census

Table 2. Descriptive statistics for NDVI and urban form variables^a.

	Residential Neighborhoods Constructed Prior to 1946			Residential Neighborhoods Constructed 1946 to 1971		
	Min.	Mean	Max.	Min.	Mean	Max.
NDVI	-0.27	0.05	0.49	-0.29	0.07	0.45
Parcel_Size	6	7,442	435,057	4	5,222	383,545
Pop_Density	401	5,798	150,200	417	3,841	150,300
Roads_0.1	1	18	42	1	17	42
Roads_density	1	545	815	6	510	900
Land_Diversity	2	6	7	2	5	7
Park_0.1	0	3	94	0	4	100
Park_1.0	0	7	53	0	8	69
Green_0.1	0	4	100	0	6	100
Green_1.0	0	14	97	0	15	95
Ind-Comm_0.1	0	2	73	0	1	90
Ind-Comm_1.0	0	3	30	0	3	60
Per_Renter_Occ	0	32	100	0	22	100
Per_Single_Fam	0	54	100	0	73	100
Apt $\geq$ 5	0	22	100	0	14	100

a. All variables significantly different between the two time periods based on the Kolmogorov-Smirnow test.

Table 3. Spearman's Rho for NDVI and urban form variables in neighborhoods dominated by residential development built before 1946^a.

[illegible]

Per_ Apt≥5														

a. All values are significant at the 0.001 p-level.

Table 4. Spearman's Rho for NDVI and urban form variables in neighborhoods dominated by residential development built between 1946 and 1970^a.

[illegible]

Per_ Apt $\geq$ 5														

a. All values are significant at the 0.001 p-level.

Table 5. Spearman's correlations between NDVI and urban form variables^a.

Variable Name	Prior to 1946	1946 to 1970
Parcel_Size	0.395	0.251
Pop_Denisty	-0.142	-0.179
Roads_0.1	-0.062	-0.032
Roads_density	-0.374	-0.119
Land_Diversity	-0.117	-0.136
Park_0.1	0.092	-0.075
Park_1.0	0.074	0.001
Green_0.1	0.179	0.100
Green_1.0	0.308	0.079
Ind-Comm_0.1	-0.372	-0.203
Ind-Comm_1.0	-0.394	-0.185
Per_Renter_Occ	-0.303	-0.214
Per_Single_Fam	0.434	0.262
Apt $\geq$ 5	-0.210	-0.171

a. All values are significant at the 0.001 p-level.

Figure

Figure 1. Urban residential development in the Greater Toronto Area, Ontario, Canada.

Residential Neighborhoods

N

10 0 10 20 km