Canadian Journal of Forest Research
Revue canadienne de recherche forestière

Total and epiphytic litter under the canopy of *Acer macrophyllum* in an old-growth temperate rainforest, Washington State, USA

Journal:	<i>Canadian Journal of Forest Research</i>
Manuscript ID	cjfr-2014-0492.R2
Manuscript Type:	Article
Date Submitted by the Author:	01-Jun-2015
Complete List of Authors:	Tejo, Camila; Universidad Austral de Chile, Instituto de Conservacion, Biodiversidad y Territorio Zabowski, Darlene; University of Washington Nadkarni, Nalini; University of Utah, Department of Biology
Keyword:	<i>Acer macrophyllum</i>, carbon cycle, epiphytic litterfall, forest productivity, old-growth forest

SCHOLARONE™
Manuscripts

Total and epiphytic litter under the canopy of *Acer macrophyllum* in an old-growth temperate rainforest, Washington State, USA

Authors: Camila F. Tejo ^{a*}, Darlene Zabowski ^b, Nalini M. Nadkarni ^c

^a Instituto de Conservacion, Biodiversidad y Territorio, Universidad Austral de Chile, Casilla 567, Valdivia, Chile.

^b School of Environmental Forest Sciences, University of Washington, Seattle, WA, 98195, USA.

^c Department of Biology, University of Utah, Salt Lake City, UT 84112 USA

*Corresponding author. Tel.: +56 9 6215 2781. Email address: cftejo@gmail.com

Abstract

The amounts and ecological importance of epiphytic litterfall has often been overlooked in forest ecosystem studies. However, epiphytes participate in whole-ecosystem dynamics by capturing and retaining nutrients from atmospheric sources, and transferring these nutrients to other ecosystem components. We quantified epiphytic litterfall under the canopy of mature bigleaf maples (*Acer macrophyllum* Pursh) and compared it to other litter components in an old-growth temperate rainforest in Washington State. Total litterfall during one year was 4760 kg ha⁻¹, with the greatest contribution from bigleaf maple leaves. Of the total litter input, 546 kg ha⁻¹ consisted of epiphytic litter, equivalent to 12% of total fine litter input, the highest contribution of epiphyte litterfall documented for this type of forest. Compared with other studies in the Pacific Northwest, our estimations of conifer needle inputs relate to the location of the littertraps. Bigleaf maple leaves dominated C and N returns in litter; epiphytic litterfall contributed 240 kg ha⁻¹ yr⁻¹ of C (~11% of total C inputs) and 5.7 kg ha⁻¹ yr⁻¹ of N (~11% of total N inputs) to the forest floor. Inputs of N from epiphytes and bigleaf maple litter under the canopy of this tree could be important in augmenting N in this old-growth ecosystem.

Acer macrophyllum, carbon cycle, epiphytic litterfall, forest productivity, nitrogen cycle, old-growth forest.

Introduction

Epiphytes, plants that derive support but not nutrients from their host trees, are a component of many forest ecosystems (Lowman and Nadkarni 1995). Whereas terrestrially rooted vegetation (trees, shrubs and herbs) obtain their nutrients predominantly from the soil, epiphytic plants (e.g., mosses, epiphytic ferns) and lichens garner their nutrients from wind- and rain-borne sources or arboreal soil (Enloe et al. 2006; Hietz et al. 2002; Knops et al. 1996; Nadkarni and Matelson 1991). Because epiphytic material (plants, lichens plus canopy soil, hereafter EM) captures allochthonous nutrients that settle within the canopy, the deposition of epiphytic litterfall to the forest floor can increase the total amount of nutrients (particularly N) intercepted and retained within the forest. The contribution of EM to litterfall might be important in forest ecosystems where N limits productivity, a common situation in Pacific Northwest forests (Edmonds et al. 1989; Fried et al. 1990; O'Keefe and Naiman 2006).

Litterfall provides a major pathway for nutrient and energy transfer from plants to soil (Coxson and Nadkarni 1995; Lindo and Winchester 2007). This flux has been related to forest productivity, nutrient cycling, and the dynamics of soil organic matter (Edmonds and Murray 2002; O'Keefe and Naiman 2006). Litterfall quantity and quality is influenced by a variety of factors including climate, seasons, forest type, successional stage and site productivity (Berg and McClaugherty 2008; Edmonds 1980; Edmonds and Murray 2002; Perez et al. 1991; Perez et al. 2003).

There is extensive information on the quantity of litterfall from terrestrially rooted plants, but very few studies have addressed the contribution of epiphytic litterfall in forests where EM is present (e.g., Kohler et al. 2008; Nadkarni and Matelson 1992; O'Keefe and Naiman 2006; Veneklaas 1991). In those forests, EM litterfall can be a substantial contributor to total litterfall

biomass. For instance, in a Colombian montane forest, Veneklaas (1991) estimated the total litterfall as $4.3 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{yr}^{-1}$, with $\sim 5\%$ of this litter attributed to epiphytes (vascular plants, bryophytes, and lichens). In an upper montane forest of Costa Rica, Kohler et al. (2008) estimated the contribution of epiphytic litterfall in an old-growth tropical forest as 6% of total litterfall. Collection of EM litterfall is challenging because it is extremely variable spatially and temporally. In the Monteverde cloud forest, over 50% of EM litterfall was collected in less than 2% of the collection containers (Nadkarni and Matelson 1992). A study of an old-growth forest in Costa Rica also indicated that epiphytic litterfall had high spatial and temporal variability (Kohler et al. 2008).

In the Pacific Northwest, total fine litterfall has been estimated (Abee and Lavender 1972; Edmonds and Murray 2002; Harmon et al. 2004; Klopatek 2007; O'Keefe and Naiman 2006), but only a few studies have quantified the contribution of EM to total litter biomass (Abee and Lavender 1972; McShane et al. 1983; O'Keefe and Naiman 2006). McShane et al. (1983) in a Douglas-fir/ western hemlock forest, estimated epiphytic litterfall between 46 to $115 \text{ kg} \cdot \text{ha}^{-1} \cdot \text{yr}^{-1}$. O'Keefe and Naiman (2006) noted a significant relationship between epiphytic litterfall and the basal area of Sitka spruce (*Picea sitchensis* (Bong.) Carrière) and bigleaf maple (*Acer macrophyllum* Pursh) that correlated with the large biomass of epiphytes held by individual trees of these species. Most of these studies, however, did not specifically focus on assessing epiphyte litterfall input. Therefore, the contribution of epiphytes may have been underestimated, as EM often fall as large clumps and by “riding down” large and small branches and on whole fallen trees, which would not be captured in traditional tree litterfall collectors.

Bigleaf maple is a native tree of the Pacific Northwest characterized by large accumulations of epiphytic biomass (Nadkarni 1984). Bigleaf maple litterfall can enhance soil

fertility through the supply of nutrients to the forest floor by providing litter that is rich in N, Ca and K (Turk et al. 2008). Moreover, Chandler et al. (2008) reported that over 80% of bigleaf maple litter is deposited directly under bigleaf maple canopy. The combined effects of epiphytic biomass, nutrient-rich litter, and high bigleaf maple litter inputs suggest that the presence of bigleaf maples and their epiphytes could enhance the capture of atmospheric N and its movement from the canopy to the forest floor of temperate rainforests.

The objective of this study was to determine the contributions of biomass, C, and N derived from epiphytic litterfall underneath bigleaf maples in an old-growth temperate rainforest in Washington State. In the study area, epiphytic biomass is extremely high: large trees can hold over 500 kg of dry epiphytic biomass (Naiman et al. 2010). Specifically we: i) determined monthly and seasonal litter inputs under the canopy of bigleaf maple during one year; ii) quantified annual C and N returns from litter components, iii) related total and EM litterfall to temperature and precipitation; and iv) compared EM and tree litter inputs under bigleaf maple to what has been documented for this and other temperate forests.

91 **Methods**

92 *Study Site*

93 The study was conducted from June 2010 to November 2012 in an old-growth forest
94 located at the Queets River watershed on the western side of Olympic National Park,
95 Washington State (47.34 N, 124.09 W). The area has a temperate climate, with cool-wet winters
96 and warm-dry summers. The mean annual precipitation is ~3000 mm, with most precipitation
97 falling from November to June. Mean annual air temperature is 14.7 °C, mean winter and
98 summers temperatures are 7.3°C and 22°C, respectively (Bechtold and Naiman 2009). Wind
99 speed ranges from 6.3 km h⁻¹ to 10.5 km h⁻¹, with highest speeds between December to March
100 (Quillayute station, Western Regional Climate Center 2013).

101 The forest is described by Van Pelt et al. (2006) and is dominated by Sitka spruce (*Picea*
102 *sitchensis*). Occasional western hemlock (*Tsuga heterophylla* (Raf.) Sarg.) and Douglas-fir
103 (*Pseudotsuga menziesii* (Mirb.) Franco var. *menziesii*) also occur. Bigleaf maple (*Acer*
104 *macrophyllum*) is the dominant hardwood species and its crown coverage corresponds to 34% of
105 total forest cover, with coverage determined using the line intercept technique (Canfield 1941).
106 Other hardwood species present are red alder (*Alnus rubra* Bong.) and vine maple (*Acer*
107 *circinatum* Pursh). Understory vegetation is dominated by sword fern (*Polystichum munitum*
108 (Kaulf.) C. Presl) and redwood sorrel (*Oxalis oregana* Nutt.) (Van Pelt et al. 2006). In the
109 canopy, the predominant epiphytes are two bryophyte species: icicle moss (*Isothecium*
110 *myosuroides* Brid.) and hanging moss (*Antitrichia curtispindula* (Hedw.) Brid.). The epiphytic
111 vascular plants licorice fern (*Polypodium glycyrrhiza* D.C. Eaton) and Oregon selaginella
112 (*Selaginella oregana* D.C. Eaton) are common in the canopy; epiphytic lichens such as lettuce

lung (*Lobaria oregana* (Tuck.) Müll. Arg.) and lungwort (*L. pulmonaria* (L.) Hoffm.) are also common.

Sampling and Analysis

In June 2010, four 30 m x 30 m plots were installed in the study site, based on the presence of at least two mature bigleaf maple trees. Within each plot, crown cover of bigleaf maple was estimated, and ranged between 39 and 52%. We used two methods to estimate tree and epiphytic litter inputs to the forest floor. For smaller materials (epiphyte fragments <100 cm² and litter from trees), we installed three 50 cm x 50 cm litter traps per plot (hereafter, “small traps”) lined with 1.5 mm nylon mesh across wooden frames. Each trap was 20 cm high and placed on the forest floor to reduce disturbance from elk. To collect inputs of larger EM, we installed two litter traps in each plot that consisted of 1.5 mm nylon mesh covering 2.7 m² and 5.4 m² of the ground (hereafter, “large traps”). Litter collected from large traps only included clumps of epiphytes ($\geq 100\text{cm}^2$), epiphytes attached to fallen branches, and branches and twigs (all woody material from large traps was >1 cm in diameter). Within each plot, all littertraps were randomly distributed under the canopy of bigleaf maple, and were emptied monthly for 16 months (August 2010 to November 2011). The sample design did not include epiphytic litterfall from fallen trees, so total epiphytic litterfall of the forest is still an underestimate.

Collected litterfall was taken to the laboratory, air-dried for 48 hr, sorted, and weighed. Air-dried weights were compared with oven-dry weights (65°C for 48 hr) and were within 1% of oven-dry weight. Litter collected in the smaller traps was separated into the following categories: bigleaf maple leaves, other deciduous leaves (vine maple, red alder, and black cottonwood), conifer needles (spruce, western hemlock, and Douglas fir), woody materials

(twigs, branches, and bark), EM (bryophytes, lichens, licorice ferns, and canopy soil), and miscellaneous material (reproductive material, understory litter, dead insects, and unidentifiable fragments). The EM collected from the larger littertraps was separated to bryophytes, lichens, licorice fern, canopy soil; woody material (twigs, branches and bark) was separated from the epiphytic material. Monthly litter inputs of each trap for each litter category were standardized to kg ha^{-1} then averaged per plot. Monthly litter collected in small and large traps (EM and woody material) was standardized to kg ha^{-1} , sum, and then averaged per plot.

The relationship between litter inputs and climatic variables (precipitation and temperature) was examined using Pearson's correlation coefficient. Precipitation and temperature data for the Queets River watershed during the sampling period was obtained from the PRISM Climate Group database from Oregon State University (<http://prism.oregonstate.edu>, created May 14, 2013).

Total C and N were determined by dry combustion using a Perkin-Elmer Model 2400 CHN analyzer (Perkin-Elmer, Wellesley, MA) in the Analytical Lab, School of Environmental and Forest Sciences, University of Washington. The sorted and weighed litterfall of each plot was grouped by season. A mixed litter category was created that included conifer needles, other deciduous species, and miscellaneous material. Litterfall C and N inputs for each litter category (epiphytes, bigleaf maple, woody material and mixed litter) were determined by multiplying the nutrient concentration of each category by the mass of that litter fraction for each season. The C/N ratio of epiphytic litterfall was calculated in proportion to the biomass and C and N concentration of each epiphytic component. To determine annual C and N inputs in litter for each category, total C and N from September 2010 through August 2011 were summed. All the data were analyzed using R software version 2.14.1 (R Development Core Team 2012).

Results

Litterfall biomass and seasonality

The mean total overstory litterfall under the canopy of bigleaf maple was 4760 kg ha⁻¹ during one year (Table 1). Bigleaf maple leaves were the dominant litter fraction and accounted for over one-third of total litterfall. In decreasing order, woody material, conifer needles and miscellaneous material were the next largest fractions of litter biomass. Other deciduous species contributed only 5% of total litterfall. Inputs from epiphytes were 12% of total litter inputs during one year (546 kg ha⁻¹ yr⁻¹). Surprisingly, no canopy soil litterfall was collected during the sampling period even though it is a large component of the arboreal canopy ecosystem. No larger branches were collected in either the small or large litter plots, and the only canopy soil that was noted on the forest floor was attached to larger branches. Determining the amount of canopy soil litter may require a longer sampling period, more litter collectors, or larger plots, as it appears to be a rare component of litterfall.

Litterfall inputs showed a distinct seasonality, with highest monthly inputs in October and November (Fig. 1). Leaves from bigleaf maple and other deciduous species were present mainly during the autumn (September - November), and were almost completely absent between March and June. Conifer needle litter had a maximum in November 2010 (183 kg ha⁻¹ mo⁻¹), while for the rest of the year, needle litter biomass averaged 48 kg ha⁻¹ mo⁻¹. Woody litterfall showed high spatial and temporal variability, particularly during the fall and winter months, when it reached 142 kg ha⁻¹ mo⁻¹ (Fig. 1). When relating inputs of total litterfall with climatic variables, we found a positive correlation between total litterfall with precipitation during the autumn months (September to November; $r = 0.98$, $p = 0.0001$) (Fig 2), whereas total litter

inputs and temperature were positive correlated only during the winter months (December to February; $r = 0.99$, $p \leq 0.05$).

Epiphytic litterfall

The inputs from epiphytic litterfall were highly variable over time, with biomass ranging from 4 to 90 kg ha⁻¹ mo⁻¹ during the sampling period (Fig. 1 and 3). The lowest and highest inputs of epiphyte biomass were in August and November 2010, respectively. In July 2011, the contribution of EM litterfall corresponded to 26% of the total litter inputs for that month. This contribution of EM to total litterfall was mainly litter from licorice fern.

Overall, bryophytes contributed 90% of total EM litterfall, whereas lichens and licorice fern were 5% and 6% of total EM litterfall, respectively (Table 3, Fig. 3). Bryophyte litterfall showed a strong seasonal pattern, and particularly between October and December 2010, bryophyte litterfall was derived mainly from larger clumps of mosses in the large littertraps. Nevertheless, small clumps and small fragments of bryophytes were continuously present in littertraps during the entire collection period (Fig. 3). Lichen litter was also collected every month during the sampling period but was greatest between January and June 2011 (ca. 2 - 3.5 kg ha⁻¹ mo⁻¹). Licorice fern litterfall contributed the least of all three epiphyte categories and there was no licorice fern litterfall during some months (Fig. 3). Highest inputs of licorice fern were between July and October 2011 (12 and 8 kg ha⁻¹ mo⁻¹, respectively) with small amounts scattered throughout the other months. Epiphytic litterfall was positive correlated with precipitation during the sampling period ($r = 0.53$, $p = 0.02$).

Carbon and Nitrogen in litterfall

Carbon concentration across litter categories ranged between 37 to 49%, whereas N concentration ranged between 1% and 2% (Table 2). Nitrogen concentration of litter components followed the sequence: lichens > bigleaf maple leaves > bryophytes and mixed litter > licorice fern, and woody material.

The largest quantity of C and N inputs to the forest floor was bigleaf maple leaf litter, followed by mixed litter, woody material, and bryophytes (Table 2). Lichens and licorice fern had the lowest C and N deposition during one year, respectively. Bigleaf maple leaves and mixed litter (conifer needles, other deciduous and miscellaneous category) contributed nearly 80% of total litter N during one year. Overall, epiphytic litterfall (bryophytes, lichens and licorice fern) and woody material each contributed ~10% of total N in litter (Table 3).

Total C and N inputs varied among seasons (Fig. 4). Bigleaf maple leaves were present only during the autumn, with C and N inputs from bigleaf maple leaf litter higher in November 2010 than in November 2011. Woody material had higher C and N transfer during the winter and spring, which coincides with high precipitation in December 2010, and January and March of 2011 (Fig. 2). The mixed litter category had higher C and N inputs during the autumn, which corresponds to a higher deposition of litter from non-bigleaf maple deciduous species. On average, C/N ratios ranged from 29 for lichens to 71 for woody material (Table 2).

Total C and N inputs from epiphyte litter were mainly derived from bryophytes. However, total N inputs from epiphytic litterfall during the spring correspond to lichen litterfall, which had the highest N concentration (Table 2). The C/N ratio for epiphytic litterfall was 42 due to the large biomass of bryophytes, the dominant epiphytic litterfall component. Throughout the seasons, changes in the C/N ratio of EM occurred. The lowest C/N ratio for epiphytic litterfall (36) occurred during autumn 2010, when bryophytes were the major epiphytic litter

227 component. However, during the summer 2010 C/N ratio of the epiphytic litterfall had a high
228 C/N ratio (71) and was dominated by licorice fern.

Draft

229 Discussion

230 Although the presence of EM has been increasingly acknowledged as an important
231 component of forest ecosystems (Cardelus 2010; Lindo and Whiteley 2011; Nadkarni et al.
232 2011), collections of epiphytic litterfall have often been included with miscellaneous litter or
233 mixed with other litter categories (Abee and Lavender 1972; Edmonds and Murray 2002).
234 Studies that have specifically measured inputs of EM relative to total litterfall, have estimated
235 EM litterfall contribution between 4-7% of total litterfall (Table 3). At the Queets River
236 watershed, O'Keefe and Naiman (2006) documented epiphytic litter between 4-6% of total litter
237 inputs. Although we did not collect epiphytic litter from whole trees or large fallen branches, our
238 estimation of epiphytic litterfall biomass under the canopy of bigleaf maple was 12% of the total
239 litter biomass, the highest contribution of epiphytic litterfall reported for this forest type. Using
240 larger littertraps may have allowed us to collect a more accurate measure of EM litterfall,
241 particularly inputs from canopy soils, as larger clumps were more likely to be collected in the
242 larger traps. Collection of canopy soils is challenging since canopy soils developed underneath
243 epiphytes are attached to branches and crotches of the host tree (Haristoy et al. 2014). When
244 found on the ground, canopy soils are part of the EM conglomerate, or remain attached to the
245 fallen branch or tree. On the other hand, some lichen litter could have been lost from littertraps
246 due to animal predation of alectoroid lichens (Esseen and Renhorn 1998; McCune and Daly
247 1994).

248 Litterfall under the canopy of bigleaf maple during one year was similar to litter biomass
249 sampled from the entire forest around the bigleaf maple plots include coniferous and other
250 hardwood trees (O'Keefe and Naiman 2006) as well as to other old-growth forest of the region
251 (Abee and Lavender 1972; Edmonds and Murray 2002). Maple leaves were the dominant litter

component under bigleaf maple trees which indicates that bigleaf maple leaves are predominantly deposited directly under bigleaf maple canopy similar to the finding of Chandler et al. (2008). Hirabuki (1991) also determined that litter distribution corresponded with a tree's canopy distribution.

In previous studies, the dominant component of litterfall in this region was conifer needles (Edmonds and Murray 2002; O'Keefe and Naiman 2006). At the Queets river watershed, the dominant conifers reach heights above 80 m and their needles are carried by the wind to the area covered by bigleaf maple. Whereas O'Keefe and Naiman (2006) reported 2980 kg ha⁻¹ yr⁻¹ of annual conifer needle inputs, in our study, we only collected 25% of that value under the canopy of bigleaf maple. This lower biomass of conifer needles is undoubtedly related to the location of the littertraps under maple but shows that needles are still a major contributor to litter biomass under bigleaf maple.

Litterfall seasonality in the Pacific Northwest has been linked with temperature and precipitation patterns (Abee and Lavender 1972; Edmonds and Murray 2002). The difference in the total litter inputs between autumn 2010 and autumn 2011 may be attributed to a later start of autumn rains in 2011 compared with 2010. In 2010, the beginning of the rainy season occurred in August, but in 2011, the rain began to increase in September (Fig. 2). Furthermore, while the dry period in 2010 lasted from July to August, in 2011 the dry period extended from June through August, which could also have affected the peak in litterfall.

Nitrogen inputs from total litterfall under the canopy of bigleaf maple were much higher than reported by other studies in the Pacific Northwest (Abee and Lavender 1972; Edmonds and Murray 2002) (Table 3). High N inputs are related to the high proportion of bigleaf maple biomass in the litter, which has a high N concentration (Chandler et al. 2008; Turk et al. 2008).

Total N inputs from EM litterfall under bigleaf maple during one year are the highest recorded for temperate ecosystems, corresponding to 11% of total N inputs. A previous study in tropical ecosystems estimated the N contribution of epiphytic litterfall as 7% of total N in litterfall (Nadkarni and Matelson 1992).

Total N input from epiphytic litterfall ($5.7 \text{ kg ha}^{-1} \text{ yr}^{-1}$) to the forest floor was almost double the annual atmospheric N deposition with precipitation in the region ($3.1 \text{ kg ha}^{-1} \text{ yr}^{-1}$; (Edmonds et al. 1998). Because of their location in the canopy, epiphytes capture and retain allochthonous nutrients that become potentially available to the whole forest via litter deposition, throughfall or stemflow. Furthermore, N-fixing lichens such as *L. oregana* can contribute up to 16 kg ha^{-1} of N in Pacific Northwest forests, increasing the N pool of EM (Antoine 2004). The contribution of epiphytic N inputs to the forest floor should be considered in future studies to understand nutrient cycling in these ecosystems.

The contribution of bigleaf maple trees at the ecosystem level might have a disproportionate effect on N cycling compared with litter inputs of conifer trees in the stand. In this area, bigleaf maple form clusters surrounded by the conifer-dominant canopy (Van Pelt et al. 2006). The inputs of litterfall under the canopy of bigleaf maple are nearly equivalent to average litter inputs of the whole forest, as estimated by O'Keefe & Naiman (2006) (Table 3), making bigleaf maple a rich focal point of nutrient deposition. Furthermore, since bigleaf maple drops its leaves directly under its canopy, and considering the biomass and litterfall of EM associated with bigleaf maple, the influence that bigleaf maple has on the surrounding vegetation may be directly limited to the area where bigleaf maples are present until whole trees and its associated EM biomass falls into the ground, decomposers and other organisms transfer it to other areas of the forest, or the surrounding vegetation takes up this pool of nutrients through their root system.

Conclusions

Litterfall estimations are critical for understanding nutrient returns to the forest floor, especially in areas where nutrients such as N are a limiting factor to forest productivity. Under the canopy of bigleaf maple, maple leaves dominated total litter inputs, and epiphytic litterfall under these trees was higher than the average epiphytic litterfall for the whole area at the same location. Annual variation in litter deposition correlates with precipitation from late summer and early autumn. Nitrogen inputs to the forest floor from EM litterfall could have a positive influence on N availability to the surrounding vegetation as this N is released during decomposition. Furthermore, as EM captures and retains nutrients in the canopy mats, litter deposition from EM is enhancing the sources of these nutrients to the forest floor. The combined effect of nutrient-rich litter and large accumulations of EM within bigleaf maple canopies suggests that bigleaf maple is a focal point for nitrogen movement to the forest floor, and its conservation could be critical for long-term contributions to forest productivity.

Acknowledgements

We thank Dr. Jerry Freilich and the Olympic National Park for access to and protection of study sites (study project OLYM-00309). Climbing equipment came from the International Canopy Network and Gear for Good (Ingrid Gordon), and is gratefully recognized. Thanks also to Tim Miller, Colton Miller, Deanne Greaves, Korena Mafune and Georgia Seyfried for Laboratory and field assistance. Partial funding was provided by a research grant from the National Science Foundation (DEB-09-56301).

Draft

References

- 320 Abee, A., and Lavender, D. 1972. Nutrient cycling in throughfall and litterfall in 450-year-old
321 Douglas-fir stands. Pages 133-143 in Proceedings of Symposium on Research on Coniferous
322 Forest Ecosystems: First year Progress in the Coniferous Forest biome. US International
323 Biological Program, Bellingham, Washington.
- 324 Antoine, M.E. 2004. An ecophysiological approach to quantifying nitrogen fixation by *Lobaria*
325 *oregana*. *The Bryologist* **107**(1): 82-87.
- 326 Bechtold, J.S., and Naiman, R.J. 2009. A quantitative model of soil organic matter accumulation
327 during floodplain primary succession. *Ecosystems* **12**(8): 1352-1368.
- 328 Berg, B., and McClaugherty, C. 2008. Plant litter decomposition, humus formation and Carbon
329 sequestration. Springer, Berlin. 338p.
- 330 Callaway, R.M., and Nadkarni, N.M. 1991. Seasonal patterns of nutrient deposition in a *Quercus*
331 *douglasii* woodland in central California. *Plant and Soil* **137**(2): 209-222.
- 332 Canfield, R. 1941. Application of the line interception method in sampling range vegetation.
333 *Journal of Forestry* **39**(4): 388-394.
- 334 Cardelus, C.L. 2010. Litter decomposition within the canopy and forest floor of three tree species
335 in a tropical lowland rain forest, Costa Rica. *Biotropica* **42**(3): 300-308.
- 336 Chandler, J.R., Schmidt, M.G., and Dragicevic, S. 2008. Spatial patterns of forest floor
337 properties and litterfall amounts associated with bigleaf maple in conifer forests of southwestern
338 British Columbia. *Canadian Journal of Soil Science* **88**(3): 295-313.
- 339 Coxson, D.S., and Nadkarni, N.M. 1995. Ecological Roles of Epiphytes in Nutrient Cycles of
340 Forest Ecosystems. *In Forest Canopies. Edited by M. D. Lowman, and N. M. Nadkarni.*
341 Academic Press. pp. 495-543.

- 342 Edmonds, R.L. 1980. Litter decomposition and nutrient release in Douglas-fir, red alder, western
343 hemlock, and Pacific silver fir ecosystems in western Washington. *Canadian Journal of Forest*
344 *Research* **10**(3): 327-337.
- 345 Edmonds, R.L., Binkley, D., Feller, M.C., Sollins, P., Abee, A., and Myrold, D.D. 1989. Nutrient
346 cycling: effects on productivity of northwest forests. Maintaining the long-term productivity of
347 Pacific northwest forest ecosystems. Edited by D. Perry, R. Meurisse, B. Thomas, R. Miller, J.
348 Boyle, C. Perry, and R. Powers. Timber Press, Portland, Oregon: 17-35.
- 349 Edmonds, R.L., Blew, R.D., Marra, J.L., Blew, J., Barg, A.K., Murray, G., and Thomas, T.B.
350 1998. Vegetation patterns, hydrology, and water chemistry in small watersheds in the Hoh River
351 Valley, Olympic National Park: National Park Service Scientific Monograph
352 NPSD/NRUSGS/NRSM-98/02. 131p.
- 353 Edmonds, R.L., and Murray, G.L.D. 2002. Overstory litter inputs and nutrient returns in an old-
354 growth temperate forest ecosystem, Olympic National Park, Washington. *Canadian Journal of*
355 *Forest Research* **32**(4): 742-750.
- 356 Enloe, H.A., Graham, R.C., and Sillett, S.C. 2006. Arboreal Histosols in old-growth redwood
357 forest canopies, northern California. *Soil Science Society of America Journal* **70**(2): 408-418.
- 358 Esseen, P.-A., and Renhorn, K.-E. 1998. Mass loss of epiphytic lichen litter in a boreal forest. *In*
359 *Annales Botanici Fennici*. Helsinki: Societas Biologica Fennica Vanamo, 1964-. pp. 211-217.
- 360 Fried, J.S., Boyle, J.R., Tappeiner, J.C., and Cromack, K. 1990. Effect of bigleaf maple on soils
361 in Douglas-fir forests. *Canadian Journal of Forest Research* **20**(3): 259-266.
- 362 Haristoy, C.T., Zabowski, D., and Nadkarni, N. 2014. Canopy Soils of Sitka Spruce and Bigleaf
363 Maple in the Queets River Watershed, Washington. *Soil Sci. Soc. Am. J.* **78**: S118–S124. doi:
364 10.2136/sssaj2013.07.0300nafsc.

- 365 Harmon, M.E., Bible, K., Ryan, M.G., Shaw, D.C., Chen, H., Klopatek, J., and Li, X. 2004.
366 Production, respiration, and overall carbon balance in an old-growth *Pseudotsuga-tsuga* forest
367 ecosystem. *Ecosystems* **7**(5): 498-512.
- 368 Hietz, P., Wanek, W., Wania, R., and Nadkarni, N.M. 2002. Nitrogen-15 natural abundance in a
369 montane cloud forest canopy as an indicator of nitrogen cycling and epiphyte nutrition.
370 *Oecologia* **131**(3): 350-355.
- 371 Hirabuki, Y. 1991. Heterogeneous dispersal of tree litterfall corresponding with patchy canopy
372 structure in a temperate mixed forest. *Vegetatio* **94**(1): 69-79.
- 373 Klopatek, J.M. 2007. Litterfall and fine root biomass contribution to nutrient dynamics in
374 second- and old-growth Douglas-fir ecosystems. *Plant and Soil* **294**(1-2): 157-167.
- 375 Knops, J.M.H., Nash, T.H., and Schlesinger, W.H. 1996. The influence of epiphytic lichens on
376 the nutrient cycling of an oak woodland. *Ecological Monographs* **66**(2): 159-179.
- 377 Kohler, L., Holscher, D., and Leuschner, C. 2008. High litterfall in old-growth and secondary
378 upper montane forest of Costa Rica. *Plant Ecology* **199**(2): 163-173.
- 379 Lebrete, M., Nys, C., and Forgeard, F. 2001. Litter production in an Atlantic beech (*Fagus*
380 *sylvatica* L.) time sequence. *Annals of Forest Science* **58**(7): 755-768.
- 381 Lindo, Z., and Whiteley, J.A. 2011. Old trees contribute bio-available nitrogen through canopy
382 bryophytes. *Plant and Soil* **342**(1-2): 141-148.
- 383 Lindo, Z., and Winchester, N.N. 2007. Oribatid mite communities and foliar litter decomposition
384 in canopy suspended soils and forest floor habitats of western redcedar forests, Vancouver
385 Island, Canada. *Soil Biology and Biochemistry* **39**(11): 2957-2966.
- 386 Lowman, M., and Nadkarni, N. 1995. *Forest Canopies*. Academic Press, San Diego.

- 387 McCune, B., and Daly, W.J. 1994. Consumption and decomposition of lichen litter in a
388 temperate coniferous rainforest. *The Lichenologist* **26**(01): 67-71.
- 389 McShane, M.C., Carlile, D.W., and Hinds, W.T. 1983. The effect of collector size on forest
390 litter-fall collection and analysis. *Canadian Journal of Forest Research* **13**(6): 1037-1042.
- 391 Nadkarni, N.M. 1984. Biomass and mineral capital of epiphytes in an *Acer macrophyllum*
392 community of a temperate moist coniferous forest, Olympic Peninsula, Washington State.
393 *Canadian Journal of Botany* **62**(11): 2223-2228.
- 394 Nadkarni, N.M., and Matelson, T.J. 1991. Fine litter dynamics within the tree canopy of a
395 tropical cloud forest. *Ecology* **72**(6): 2071-2082.
- 396 Nadkarni, N.M., and Matelson, T.J. 1992. Biomass and nutrient dynamics of epiphytic litterfall
397 in a neotropical montane forest, Costa Rica. *Biotropica* **24**(1): 24-30.
- 398 Nadkarni, N.M., Parker, G.G., and Lowman, M.D. 2011. Forest canopy studies as an emerging
399 field of science. *Annals of Forest Science* **68**(2): 217-224.
- 400 Naiman, R.J., Bechtold, J.S., Beechie, T.J., Latterell, J.J., and Van Pelt, R. 2010. A Process-
401 Based View of Floodplain Forest Patterns in Coastal River Valleys of the Pacific Northwest.
402 *Ecosystems* **13**(1): 1-31.
- 403 O'Keefe, T.C., and Naiman, R.J. 2006. The influence of forest structure on riparian litterfall in a
404 Pacific Coastal rain forest. *Canadian Journal of Forest Research* **36**(11): 2852-2863.
- 405 Perez, C., Armesto, J.J., and Ruthsatz, B. 1991. Litter decomposition, root biomass and soils of
406 mixed broadleaf-conifer forests in Chiloe National Park, southern Chile. *Revista Chilena De*
407 *Historia Natural* **64**(3): 479-490.

- 408 Perez, C.A., Armesto, J.J., Torrealba, C., and Carmona, M.R. 2003. Litterfall dynamics and
409 nitrogen use efficiency in two evergreen temperate rainforests of southern Chile. *Austral*
410 *Ecology* **28**(6): 591-600.
- 411 R Development Core Team. 2012. R: A language and environment for statistical computing. R
412 Foundation for Statistical Computing, Vienna, Austria.
- 413 Turk, T.D., Schmidt, M.G., and Roberts, N.J. 2008. The influence of bigleaf maple on forest
414 floor and mineral soil properties in a coniferous forest in coastal British Columbia. *Forest*
415 *Ecology and Management* **255**(5–6): 1874-1882.
- 416 Van Pelt, R., O'Keefe, T.C., Latterell, J.J., and Naiman, R.J. 2006. Riparian forest stand
417 development along the Queets River in Olympic National Park, Washington. *Ecological*
418 *Monographs* **76**(2): 277-298.
- 419 Veneklaas, E.J. 1991. Litterfall and nutrient fluxes in two montane tropical rainforest, Colombia.
420 *Journal of Tropical Ecology* **7**: 319-336.
- 421 Western Regional Climate Center. 2013. Cooperative Climatological Data Summaries [accessed
422 November 23, 2013].
- 423

Tables

Table 1: Mean total litterfall (SD) by litter type under the canopy of bigleaf maple between September 2010 and August 2011 at the Queets River watershed, Washington State.

Material	Litterfall (kg·ha ⁻¹)	% Total litterfall
Bigleaf Maple	1820 (690)	38
Epiphytic material	546 (333)	11.5
Bryophytes	495 (317)	10.4
Lichens	23 (25)	0.5
Licorice fern	28 (27)	0.6
Conifer Needles	748 (412)	16
Other Deciduous	238 (174)	5
Miscellaneous	580 (429)	12.1
Wood	826 (481)	17.4
Total	4762 (1331)	100

Table 2: Carbon and nitrogen concentrations (%), annual returns (kg ha^{-1}) between October 2010 and September 2011, and C/N ratios of litter collected under the canopy of bigleaf maple at the Queets River watershed, Washington State. Values were calculated by multiplying the nutrient concentration of each litter fraction by the total accumulated biomass of that fraction in each plot. Values are means with standard deviation in parenthesis ($n = 18$).

Litter component	Carbon		Nitrogen		C/N
	(%)	(kg ha^{-1})	(%)	(kg ha^{-1})	
Bigleaf maple	47 (0)	875 (324)	1.6 (0)	23 (9)	36 (5)
Epiphytic material		240 (144)		5.7 (3)	
Bryophytes	43 (1)	215 (137)	1.1 (0)	5 (3)	42 (3)
Lichens	41 (6)	10 (11)	1.7 (1)	0.5 (1)	29 (7)
Licorice fern	37 (11)	13 (12)	0.7 (0)	0.2 (0)	48 (10)
Mixed Litter*	48 (1)	775 (437)	1.1 (0)	17 (9)	47 (7)
Woody	49 (1)	405 (234)	0.7 (0)	5.4 (3)	71 (9)
Total		2290 (634)		51 (18)	

*This category includes conifer needles, other deciduous species and miscellaneous material.

Table 3: Mean annual biomass and N inputs (kg ha⁻¹) in total and epiphytic litterfall under the canopy of bigleaf maple between October 2010 and September 2011, at the Queets River watershed (this study) in comparison with other studies from Washington State and tropical forests that reported epiphytic litterfall.

Ecosystem/Source	—Total Litterfall—		—Epiphytic material—			
	Biomass	N inputs	—Biomass—		—N inputs—	
	(kg ha ⁻¹)		(kg ha ⁻¹)	% total	(kg ha ⁻¹)	% total
Temperate forest						
This study	4760	51	546	11.5	5.7	11
O’Keefe and Naiman 2006	5613		238	4.2		
Lebret et al 2001	4710		171	3.6		
Callaway and Nadkarni*	3495	42	166	4.7	2.2	5.2
McShane et al 1983 ⁺	2941		62	2.1		
Abee and Lavender 1972	6325	27			1.4	5.2
Tropical Forest						
Kohler et al 2008	17200		790	6		
Nadkarni and Matelson 1992	7500	100	500	7	7.5	7
Veneklaas 1991	4300		230	5		

*Includes only data from June 1986 - May 1987

⁺Values are the mean of their data

Figures

Fig. 1: Mean and SD of litterfall inputs (kg·ha⁻¹) per month of bigleaf maple leaves (A), conifer needles (B), epiphytes (C), other deciduous species leaves (D), woody material (E) and miscellaneous material (F) collected under the canopy of bigleaf maples at the Queets River watershed, Washington State between August 2010 to November 2011. Note the scale of the y-axis for bigleaf maple and conifer needles is 8x and 2x, respectively, of other categories of litter.

Fig. 2: Mean and SD of total and epiphytic litterfall inputs (kg·ha⁻¹) per month plotted against modeled maximum and minimum temperatures (A) and precipitation (B) data for the area from August 2010 to November 2011.

Fig. 3: Mean and SD of epiphytic litter biomass (kg ha^{-1}) per month by component, bryophytes (A), licorice fern (B), and lichens (C), collected under the canopy of bigleaf maple between August 2010 to November 2011 at the Queets River watershed, Washington state. Note the scale of the y-axis for bryophytes and licorice fern is 30x and 3x, respectively, of the lichens.

Fig. 4: Mean and SD of total litter carbon (A) and nitrogen (B) per season from September 2010 to November 2011 by litter type under the canopy of bigleaf maple at the Queets River watershed, WA. Mixed litter category includes conifer needles, other deciduous species, and miscellaneous material.