Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

**The contribution of fecundity and embryo quality to
reproductive potential of eastern Bering Sea snow crab
(*Chionoecetes opilio*)**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2015-0485.R1
Manuscript Type:	Article
Date Submitted by the Author:	28-Apr-2016
Complete List of Authors:	Webb, Joel; Alaska Department of Fish and Game, Division of Commercial Fisheries Slater, Laura; Alaska Department of Fish and Game, Division of Commercial Fisheries Eckert, Ginny; University of Alaska Fairbanks, Juneau Center, School of Fisheries and Ocean Sciences Kruse, Gordon; University of Alaska Fairbanks, School of Fisheries and Ocean Sciences
Keyword:	CRABS < Organisms, REPRODUCTION < General, FECUNDITY < General, FISHERY MANAGEMENT < General, LIFE HISTORY < General

SCHOLARONE™
Manuscripts

1 The contribution of fecundity and embryo quality to reproductive potential of eastern Bering Sea
2 snow crab (*Chionoecetes opilio*)

3 Joel B. Webb^{1,2*}, Laura M. Slater³, Ginny L. Eckert¹, and Gordon H. Kruse¹

4 ¹ University of Alaska Fairbanks, School of Fisheries and Ocean Sciences, 17101 Point Lena
5 Loop Road, Juneau, AK 99801

6 ² Alaska Department of Fish and Game, Division of Commercial Fisheries, PO Box 115526,
7 Juneau, AK 99811-5526

8 ³ Alaska Department of Fish and Game, Division of Commercial Fisheries, 351 Research Court,
9 Kodiak, AK 99615

10 Corresponding Author: Joel Webb (e-mail: joel.webb@alaska.gov)

11 *Present Address: Alaska Department of Fish and Game, Division of Commercial Fisheries, PO
12 Box 115526, Juneau, AK 99811. Phone: 907-465-6129, Fax: 907-465-6402,

Abstract

Development of refined indices of female reproductive potential is needed for estimation of alternative biological reference points for the eastern Bering Sea (EBS) snow crab, *Chionoecetes opilio*, fishery, which is managed with large male-only harvest regulations. Females were collected from 2007 to 2009 to investigate seasonal and interannual variation in fecundity with maternal size, shell condition (a proxy for age post maturity), and recent mating and again in 2010 to examine biochemical measures (CHN) of embryo quality. Mean model-adjusted fecundity was highest for primiparous and young multiparous females and declined with advancing shell condition, presumably from senescence. This pattern was also found for clutch fullness indices evaluated from 20 years of stock assessment survey data. Indicators of low female sperm reserves were associated with decreased (~10%) fecundity for multiparous females. Seasonal comparison of size-fecundity relationships suggested that embryo loss during brooding was minimal and embryo quality analyses suggested that strong variation with maternal characteristics was unlikely. Finally, fecundity-at-size of EBS females may be lower than that of conspecifics in Japan and eastern Canada.

28 Introduction

29 Large-male only harvest regulations are used to manage many crab fisheries worldwide. A
30 challenge to crab fisheries management is that information is often lacking to inform
31 relationships between male harvest and female reproductive potential. Peak harvests in the
32 eastern Bering Sea (EBS) snow crab, *Chionoecetes opilio*, fishery were among the largest crab
33 harvests in the world but declined rapidly in the late 1990s and have varied substantially under a
34 more conservative management regime to the present (Fig.1). Biomass of mature males is
35 currently used as an index of stock reproductive potential for stock assessment and management
36 (Turnock and Rugolo 2014). Information is needed to develop a refined index of female
37 reproductive potential, intuitively preferable to a male index for estimation of biological
38 reference points, and to evaluate whether trends are similar between indices.

39 Female reproductive output provides the initial input (egg production) from which density-
40 dependent and density-independent processes ultimately regulate cohort strength (Caputi 1993;
41 Émond et al. 2015; Houde 2008; Lipcius and Stockhausen 2002). Density-independent variables,
42 including climate regime impacts on larval advection and predation of juveniles, are recognized
43 to influence recruitment for EBS snow crab (Burgos et al. 2013; Parada et al. 2010; Szuwalski
44 and Punt 2013; Zheng and Kruse 2006). However, it has also been argued that strong recruitment
45 pulses for the EBS stock are serially linked (Ernst et al. 2012), and reproductive potential was
46 positively associated with recruitment under at least one historic climate regime (Szuwalski and
47 Punt 2013). Major geographic shifts in reproductive potential also occurred coincident with
48 climate regime shifts (Orensanz et al. 2004; Parada et al. 2010), such that disentangling the
49 influence of these factors, particularly from aggregate indices, may be challenging. Biomass-
50 based indices of reproductive potential assume that stock egg production is proportional to

biomass (Rothschild and Fogarty 1989). However, this relationship may diverge for species in which fecundity or embryo quality varies strongly with maternal factors (Lambert 2008), potentially biasing estimation of biological reference points and perceptions of stock resiliency to fishing (Brooks 2013; Murawski 2001; Marshall et al. 2006). Incorporation of detailed information on reproduction (e.g. maturity, sex ratio, and egg production) can improve estimation of relationships between reproductive potential and recruitment, particularly for species with highly variable reproductive traits (Morgan et al. 2011), but can also increase uncertainty in the estimation of reference points (Kell et al. 2016). Management strategy evaluation and simulation indicated that the current reference point framework, based on a proxy maximum fishing mortality rate ($F_{35\%}$), is relatively robust in the face of environmental regime shift impacts on recruitment (Punt et al. 2014; Szuwalski and Punt 2012), but also emphasized that defining an index of effective spawning biomass including female reproductive potential is a priority research objective (Punt et al. 2014).

Among decapod crustaceans, fecundity and embryo quality can vary with maternal characteristics and environmental conditions. Patterns of increasing fecundity with increasing maternal size are generally conservative (Hines 1982; Reid and Corey 1991), and egg clutch size is constrained allometrically by both the volume of the body cavity and variation in egg size (Hines 1982). Fecundity-at-size can vary substantially among stocks and years for commercially exploited crabs (Haynes et al. 1976; Prager et al. 1990; Swiney et al. 2012). Factors potentially influencing size-specific fecundity include maternal age relative to maturity (Sainte-Marie 1993), variation in maternal condition through bottom-up processes (Wells 2009), density-dependence (Prager et al. 1990), variation in the physical environment (Hannah et al. 1995), brood sequence in species with multiple annual broods (Darnell et al. 2009; Verisimo et al. 2011), mating

74 dynamics/fertilization success (Rondeau and Sainte-Marie 2001; Sainte-Marie et al. 2002), and
75 embryo loss during brooding (Kuris 1990). Similarly, embryo quality can vary with both
76 maternal and environmental factors (Amsler and George 1984; Gimenez and Anger 2001;
77 Ouellet and Plante 2004; Verisimo et al. 2011) influencing larval fitness and survival (Racotta et
78 al. 2003; Gimenez 2006).

79 Factors known to influence the size-fecundity relationship for snow crab include many of the
80 same factors described above for decapod crustaceans. Female snow crab have a terminal (final
81 lifetime) molt to maturity (Watson 1970). Duration of elapsed time relative to terminal molt is
82 indexed by assignment of a subjective shell condition index (SC) based on the accumulation of
83 epibionts and wear on the exoskeleton (Jadamec et al. 1999). Shell condition index information
84 collected annually during stock assessment surveys is useful for distinguishing primiparous
85 from multiparous females (Ernst et al. 2005; Fonseca et al. 2008). On an annual reproductive
86 cycle, primiparous females (carrying the first egg clutch of ontogeny) are SC2 (new shell),
87 whereas young, multiparous females likely bearing the second or third clutch of ontogeny are
88 SC3 (old shell), and old, multiparous females carrying the fourth or greater clutch are generally
89 classified as SC4 (very old shell) or SC5 (very, very old shell) (Jadamec et al. 1999; Ernst et al.
90 2005). Exposure, early in development, of embryos brooded by female snow crab to
91 temperatures less than approximately 1° C induces a protracted ~2 y duration (biennial) of
92 embryo development (Moriyasu and Lanteigne 1998; Sainte-Marie et al. 2008; Webb et al.
93 2007). Snow crab fecundity increases with female size (Haynes et al. 1976), and primiparous
94 females have reduced fecundity-at-size compared to young, multiparous females (Sainte-Marie
95 1993). Among EBS females, variability in mean clutch fullness (CFI) increases with increasing
96 shell condition index, and mean CFI is lower for SC4 and SC5 than for younger females

(Orensanz et al. 2005). Reductions in fecundity with advanced shell condition may be due to reduced energy allocation to reproduction (Kon and Adachi 2006; Kon et al. 2010). The occurrence of primiparous females with relatively low fecundity may be due to egg loss during post-oviposition mating at male-biased sex ratios or sperm limitation at female-biased sex ratios (Sainte-Marie et al. 2002). Primiparous females with very low sperm reserves extruded unfertilized egg clutches in the laboratory (Rondeau and Sainte-Marie 2001). Estimates of embryo loss during brooding for snow crab are highly variable, ranging from negligible to a 50% decrease, with greater decreases for larger females (Kon 1974; Sainte-Marie 1993; Comeau et al. 1999).

Improved understanding of the relationship between male and female reproductive potential and the primary factors influencing female reproductive potential is needed to clarify the relationship between reproductive potential and recruitment for this stock. The objective of this study was to provide information for improved assessment of reproductive potential of EBS snow crab by: 1) estimating functional relationships between maternal characteristics (size, shell condition, and mating success) and fecundity; 2) evaluating the efficacy of a subjective index of clutch size routinely collected during assessment surveys as a predictor of differences in fecundity; 3) characterizing seasonal changes in the size-fecundity relationship to investigate embryo loss during brooding; and 4) estimating functional relationships between maternal characteristics (size and shell condition) and embryo quality.

Methods and Materials

Specimen Collection

Mature female snow crab, identified by a disproportionately large abdomen relative to immature females, were sampled in summer (June through early August) of 2007-2009 during

annual stock assessment bottom trawl surveys conducted by the US NOAA National Marine Fisheries Service (NMFS; Chilton et al. 2011). Females were sampled from across their surveyed distribution. Sampled locations and sample size at each location varied annually (Fig. A1). Crab were collected, tagged, maintained live or frozen depending on condition, and shipped to laboratory facilities in Juneau or Kodiak, Alaska. Crab that died during collection or shipping were frozen whole until processing. Live females were maintained in large aquaria (>700 L) in the laboratory at temperatures similar to those *in situ* (0.0 to 3.0 °C) for periods of one to four weeks until processing. Because bottom trawl surveys occur only during the summer, ovigerous female snow crab were opportunistically collected during the winter and spring of 2008 (n=102) and 2010 (n=66) as bycatch in groundfish fisheries by the North Pacific Groundfish Observer Program. Collection locations were constrained to the area fished by observed vessels in the southern and central portion of the snow crab distribution, and females were frozen aboard the fishing vessel and shipped to Juneau.

For evaluation of embryo quality, females (22 SC2, 25 SC3, and 17 SC4) with embryos in the early stages of development (gastrula or earlier; Moriyasu and Lanteigne 1998) were sampled from the EBS in June 2010 during the annual NMFS bottom trawl survey. Females were held live in flow-through seawater tanks aboard the survey vessel, and shipped via airfreight to the University of Alaska Fairbanks laboratory in Juneau, Alaska, where they were held in 700 L aquaria with flow-through seawater chilled to 1.5 - 2.0° C.

Clutch fullness index (CFI)

Weighted mean CFI by shell condition group and year was estimated from data (n=187,848 CFI observations) collected at-sea during annual NMFS trawl surveys from 1994 to 2014 (data available from R. Foy, NMFS Kodiak Laboratory, Kodiak, Alaska). CFI is a categorical

assignment of egg clutch size based on a visual index of the volume of the egg clutch relative to the female abdomen with levels of 6 – full, 5 – 0.75, 4 – 0.5, 3 – 0.25, 2 – 0.125, 1 – empty or trace (Jadamec et al. 1999; Orensanz et al. 2005).

Laboratory data collection

Laboratory data collected from live females included carapace width (measured to the nearest 0.1 mm), SC, egg clutch color (Pantone™), grasping marks on legs, ovary color (Pantone™), ovary % fullness (25% increments relative to examples in Jadamec et al. 1999), ovary dry weight (1 in 3 sampling rate), egg viability based on the presence of symmetric cleavage planes or embryo structure, embryo development stage (Moriyasu and Lanteigne 1998), spermathecal fullness (25% intervals, where a 100% full spermathecae was defined as the absence of a visible empty portion of the spermathecae, and empty was defined as a lack of visible material in the spermathecae), presence of fresh ejaculate (Duluc et al. 2005), and CFI. The same data were collected from frozen specimens from survey and observer collections except when degraded condition of the ovary or embryos precluded reliable assessment of embryo viability or ovary fullness and weight. Fecundity was estimated by division of the dry weight of the egg clutch stripped from the pleopods by mean embryo dry weight (determined from two replicate subsamples of >150 embryos). All embryos were dried at 60° C for >48 h in a drying oven and weighed to the nearest 0.1 mg with an Ohaus® DV215CD or AV264 analytical microbalance.

Females collected in 2010 were processed as above with additional data collected on embryo volume and proximate biochemical composition. Mean embryo dry weight was estimated from six subsamples of ~50 embryos per sample, and fecundity was calculated as the dry weight of the egg clutch divided by the mean embryo dry weight. Embryo volume was estimated by measurement of 10 live embryos per female photographed at 2.5x magnification on a Leica M80

166 stereomicroscope. The diameter of each embryo was estimated as the mean of its major and
167 minor axis (Sainte-Marie 1993) as measured using ImageJ image analysis software, and embryo
168 volume was calculated as a spheroid $V = \frac{4}{3}\pi 1/2\bar{d}^3$, where $\bar{d}$ was the mean of major and minor
169 axes (Brante et al. 2003). Subsamples of the dried embryo clutches (~80 mg) of 16 SC2, 17 SC3
170 and 15 SC4 were randomly selected and analyzed for proximate biochemical composition by
171 measurement of elemental composition of carbon, hydrogen, and nitrogen (CHN) by combustion
172 methods (Gnaiger and Bitterlich 1984).

173 *Statistical analysis*

174 Data included in analyses of fecundity and embryo quality were screened to limit analysis to
175 females with egg clutches in the early stages of development and without evidence of
176 hybridization. Snow crab in the EBS can hybridize with its congener, Tanner crab (*C. bairdi*).
177 Tanner crab have smaller embryos than snow crab (Haynes et al. 1976) and morphologically
178 identified hybrids can have mixed proportions of eggs the size of either snow or Tanner crab (L.
179 Slater, unpublished data). Females identified morphologically as snow crab, but with mean
180 embryo weights similar to Tanner crab, indicating they were possibly *C. opilio* X *C. bairdi*
181 hybrids, were excluded from comparisons using a threshold mean embryo weight of 4.80×10^{-5}
182 g (J. Webb and L. Slater, unpublished data). Females in the intermediate year of biennial
183 reproduction were excluded based on embryo development (post-gastrula) and ovary fullness
184 (>50%). Final datasets included 453 SC2, 235 SC3, 136 SC4, and 29 SC5 (n=853) females for
185 fecundity analyses and 22 SC2, 15 SC3, and 17 SC4 females for embryo quality.

186 Potential differences in fecundity and clutch dry weight as a function of female carapace
187 width, mean embryo dry weight, shell condition, year of collection, and location (random

covariate) were evaluated using linear mixed models for females collected in summer from 2007 to 2009. Variability in mean embryo dry weight was evaluated by the same approach, as a function of female size, shell condition, and year of collection. Model-adjusted means of the dependent variables are reported throughout. Quantile regression (Koenker 2006) by shell condition group at the 5th and 95th percentiles (proxies for minimum and maximum fecundity at size) were used to evaluate possible changes in fecundity-at-size between shell condition groups with years pooled. ANOVA was used to evaluate differences in standardized fecundity residuals from the linear mixed model by shell condition group for females with or without fresh ejaculate in the spermathecae and among spermathecal fullness classes for SC2 and SC3 females. SC4 and SC5 females were excluded from comparisons of fecundity residuals among spermathecal fullness classes due to low sample sizes. Differences in size-fecundity relationships between summer (early in the reproductive cycle) and spring (late in the reproductive cycle) were evaluated by ANCOVA using only females collected south of 58° N in the summers of 2007 and 2009, since spring collections were confined to the southern portion of the snow crab distribution. The fixed effect year was defined as the year of embryo extrusion for this analysis to facilitate comparisons (i.e. females collected in the spring were grouped with females collected the prior summer). A linear mixed model with carapace width and CFI as predictors was also used to estimate the slopes and intercepts and their standard errors of the size-fecundity relationship by CFI group.

Relationships among mean embryo volume, mean embryo dry weight, elemental composition (% C, % H, % N, and C:N ratio), and elemental weight per embryo (C, H, and N in $\mu\text{g} \cdot \text{embryo}^{-1}$), maternal carapace width, maternal shell condition, and fecundity of females collected in 2010 were evaluated by analysis of covariance (ANCOVA). Female carapace width, shell condition,

211 and their interaction were treated as fixed effects. Mean embryo dry weight, elemental
212 composition (% C, % H, % N, and C:N ratio), and elemental weight per embryo were modeled
213 as dependent variables. Associations among embryo characteristics (volume and weight),
214 elemental composition (% C, % H, and % N), and model-adjusted fecundity residuals were
215 evaluated with estimates of pairwise Pearson correlation (r).

216 Mean female CFI by shell condition and year was estimated from the NMFS survey database
217 as the mean CFI among all surveyed stations weighted by the mean density of female snow crab
218 for each combination of shell condition and clutch fullness index within each year. Differences
219 in the weighted mean CFI and the coefficient of variation of CFI among shell condition groups
220 were tested with a Kruskal-Wallis test and Dunn's post-hoc test (Zar 1999).

221 A top-down procedure was used for the mixed-effects model selection (Zuur et al. 2009).
222 Normality of model residuals was evaluated by normal-quantile plots and the Shapiro-Wilk test.
223 Homogeneity of variance was evaluated by plots of model residuals versus predicted values.
224 When necessary, square root or natural logarithm transformations were applied to the dependent
225 variable to achieve homogeneity of variance of the residuals. Influential outliers were identified
226 by Cook's distance, and model robustness to influential outliers was evaluated by fitting models
227 both including and excluding outliers. The threshold significance level for all statistical analyses
228 was $\alpha=0.05$. Post-hoc Tukey HSD tests of the model-adjusted mean were conducted for
229 significant results with more than two comparisons. Quantile regression was conducted using the
230 R package quantreg in R 2.15.0 (R Development Core Team 2012) with standard errors of the
231 slopes and intercepts estimated by bootstrap. All linear model and CFI analyses were conducted
232 in R 2.15.0 or JMP 9.02 (SAS Institute, Cary, NC).

Results

Size-fecundity relationships among shell condition classes and years

A high proportion of the overall variability in snow crab fecundity was explained by a linear mixed model ($n=853$, $R^2=0.80$, $RMSE=3895$; Table A1) including female size, shell condition, and year of collection as fixed effects and location of collection as a random effect. Fecundity was positively associated with female size and varied among shell condition groups and years (Table A1; Fig. 2A). SC3 females had the highest fecundity when pooled among years, but this group had lower fecundity in 2008 compared to 2007 and 2009. Significantly lower fecundity for SC3 compared to primiparous (SC2) females was observed in 2008 (Fig. 2A). SC4 and SC5 females generally had lower fecundity than SC2 and SC3 females. Fecundity of SC5 females was ~50% of the fecundity of SC3 females in 2007 and 2008, but this comparison was limited in 2009 owing to a small sample size ($n=3$) for SC5 females (Fig. 2A). Collection location contributed ~21% of the residual variance to the size-fecundity relationship (Table A1).

Variability in fecundity-at-size differed among shell condition groups. Variance in fecundity-at-size was greater for SC3 and SC4 than SC2 females (Table A2; Fig. 3). Quantile regression at the 95th percentile showed increased maximum fecundity-at-size from SC2 to SC3 and decreased maximum fecundity at size for shell conditions SC4 and SC5 (Table A2; Fig. 3). Minimum fecundity at size (5th percentile) decreased markedly with advancing shell condition, indicating a weakening of the positive relationship between size and fecundity (Table A2; Fig. 3).

Mean embryo and clutch dry weight

Among all females in this study, mean ($\pm$ SE) embryo dry weight was $66.92 \pm 4.43 \mu\text{g}$. Estimated by a linear mixed model ($n=853$, $R^2=0.52$, $RMSE= 5.53 \times 10^{-6}$; Table A3), mean

embryo dry weight also differed significantly among years and shell condition groups but not with maternal size (Table A3; Fig. 2B). Clutch dry weight, estimated from a linear mixed model ($n=853$, $R^2=0.73$, $RMSE=0.656$; Table A4), differed significantly among years and shell condition groups (Table A4; Fig. 2C). Carapace width and shell condition were the strongest predictors of clutch dry weight, followed by year and mean embryo dry weight (Table A4). Mean ($\pm$ SE) clutch dry weight of all females was 2.10 ± 0.04 g.

Sperm reserves and fecundity

Indicators of greater or more recent sperm reserves, measured as % spermathecal fullness and presence of fresh ejaculate, respectively, were positively associated with fecundity residuals (Fig. 4). Spermathecal fullness and the presence of fresh ejaculate were positively related among all females, as the proportion of females classified with fresh ejaculate increased with increasing spermathecal fullness from 13% in the 0-25% fullness category to 94% in the 75-100% fullness category. Model standardized fecundity residuals pooled among years were higher for SC3 than SC2 females (ANOVA, $F=5.02$, $p=0.002$) and for SC3 females with relatively full spermathecae (25%-50%, 50-75%, and 75 -100% categories) compared to SC3 females with relatively empty spermathecae (0-25% category; ANOVA, $F_{7,658} = 5.93$, $p<0.0001$; Tukey HSD $p<0.05$; Fig. 4A). Model-standardized fecundity residuals did not vary significantly with spermathecal fullness for SC2 females. Model standardized fecundity residuals also varied among shell condition groups with the presence or absence of fresh ejaculate (ANOVA; $F_{7,830} = 4.94$, $p<0.0001$). The presence of fresh ejaculate was associated with a significant (Tukey HSD, $p<0.05$) increase in residual fecundity, a difference in residuals of 3,000-4,000 embryos, for SC3 and SC4 females but not for SC2 or SC5 females (Fig. 4B).

Clutch fullness index

Similar to the amount of variation explained by a suite of biological variables, 80% of the variation in fecundity was explained by female size and CFI ($n=853$, $R^2=0.80$, $RMSE=469.6$; Table A5; Fig. 5A). Pooled among years (1994-2014), weighted mean CFI varied significantly among shell condition groups, with greater mean CFI for SC2 than SC5 females, SC3 females than SC4 and SC5 females, and SC4 than SC5 females. Weighted mean CFIs were similar between SC2 and SC3 and SC2 and SC4 females (Kruskal Wallis; $\chi^2 = 57.60$, $df = 3/83$; $p < 0.001$; Dunn; $p < 0.05$; Table 3). The coefficient of variation (CV) of CFI also differed among shell condition groups with a significantly higher CV for SC4 than SC2 and SC3 females and SC5 females versus all other groups (Kruskal-Wallis; $\chi^2 = 72.35$, $df = 3/83$; $p < 0.001$; Dunn; $p < 0.05$; Table 3). Temporal variation in the time series of weighted mean clutch fullness index by shell condition was also qualitatively quasi-cyclic with regular peaks and troughs for multiparous (SC3-SC5) but not primiparous (SC2) females (Fig. 5B).

Embryo quality

Embryo quality, determined by embryo volume, dry weight, and elemental composition did not vary as a function of female size. Significant differences ($p < 0.05$) were observed among shell condition groups (Table 1 & A7), but were limited in magnitude. Mean embryo dry weight, C weight per embryo, and H weight per embryo were significantly lower for SC4 than SC2 females, but a similar trend was not detected in embryo volume. Significant, but generally weak ($r = 0.3-0.4$), positive correlations were observed among embryo volume, weight, and elemental composition (Table 2; Fig. A2). Mean embryo volume was positively correlated with mean embryo dry weight (Fig. A2). Embryo % C was positively correlated with embryo % N and fecundity residuals (Fig. A2).

300 *Seasonal comparisons of size-fecundity relationships and reproductive status*

301 Fecundity was similar within shell condition group for SC3 and SC4 females in both summer
302 and spring, but very oldshell (SC4) females had significantly lower fecundity at size, by
303 approximately 4,500 embryo, than SC3 females in both summer and spring (Table A6; Fig. 6).
304 Most (90%) females collected in the spring were brooding clutches with eyed embryos, and 19%
305 of females had egg clutches with evidence of hatching (egg cases mixed with eyed eggs) at the
306 time of collection. Evidence of hatching was observed from March to June but not in February.
307 Most (97%) females collected in the spring were SC3 or SC4.

308 **Discussion**

309 Fecundity of sampled EBS snow crab varied among years and as a function of female size,
310 shell condition, and sperm reserves. Comparison of size-fecundity relationships by shell
311 condition group suggested that average fecundity was generally highest for both primiparous
312 (SC2) and young, multiparous (SC3) females, and decreased thereafter with increasing duration
313 of female reproductive history. Estimation of variability in maximum and minimum fecundity-at-
314 size by quantile regression provided finer scale resolution and demonstrated 26% greater
315 maximum fecundity at size of SC3 versus SC2 females. In the Gulf of Saint Lawrence and Japan,
316 young multiparous snow crab had 17-30% greater mean fecundity than did primiparous females
317 (Sainte-Marie 1993; Kon and Adachi 2006), although not consistently across years (Sainte-Marie
318 et al. 2008). The lower maximum fecundity of SC2 compared to SC3 females is likely
319 attributable to limited body volume available for ovary maturation and energy allocation to both
320 growth and reproduction prior to the molt to maturity (Sainte-Marie 1993), as in the congener,
321 Tanner crab (Somerton and Meyers 1983). The decreasing trend in both minimum and mean
322 fecundity-at-size from SC3 to SC5 is likely a result of female senescence. The size-fecundity

relationship of primiparous females was relatively stable through the duration of this study, while multiparous female fecundity varied among years in response to variability in embryo size/weight, possible sperm limitation, and senescence. Pooled mean embryo dry weight of multiparous females (SC3, SC4, and SC5) in 2008 was ~22 % greater than in 2007, while clutch dry weight was similar or decreased, indicating a potential change in per embryo energy allocation. Per embryo energy allocation may be affected by maternal condition/nutrition or environmental factors during oocyte formation and maturation, but these patterns have not been comprehensively studied for female snow crab (Sainte-Marie et al. 2008). Variability in fecundity related to differences in mean embryo weight is unlikely to be detected by the current CFI monitoring protocol. Annual monitoring of the fecundity of SC2 and SC3 female snow crab has continued since 2009 in the EBS and will likely provide further insights into interannual variability in mean embryo weight between these two groups.

Indicators of low sperm reserves were associated with reduced fecundity of multiparous females in this study. Lower mean residual fecundity, equivalent to ~ 10% reduction at mean female size, was observed for SC3 females with low spermathecal fullness and with an absence of fresh ejaculate for SC3 and SC4 females (Fig. 4). Fresh ejaculate is an indicator of recent mating (Duluc et al. 2005) and increased sperm reserves (Webb and Bednarski 2010). Sperm limitation may occur for multiparous snow crab if sperm reserves stored from previous mating are absent or low and females are not mated prior to oviposition of the subsequent clutch (Rondeau and Sainte-Marie 2001; Sainte-Marie et al. 2002; Slater et al. 2010). Sperm viability also decreases with extended duration of storage (Sainte-Marie et al. 2008). Sperm reserves of primiparous female EBS snow crab are lower than those typically observed for conspecifics from eastern Canada, with a high proportion classified with empty spermathecae by visual indicators

in this study (Fig. 4). There was no evidence of retention of mature oocytes in the ovary or high proportions of abnormally developing (e.g. unfertilized) embryos in the egg clutches of females with low residual fecundity and low spermathecal fullness (J. Webb and L. Slater, unpublished data). Association between lower fecundity and reduced sperm reserves could be a result of sloughing of unfertilized embryos due to sperm limitation or embryo loss due to post-oviposition mating (Rondeau and Sainte-Marie 2001; Sainte-Marie et al. 2002). Operational sex ratio and the maturity status (i.e. adolescent versus adult) of males present at the time of mating likely differ between primiparous and multiparous females in the EBS (Somerton 1981; Nichol and Somerton 2015). Further evaluation of relationships between the quantity and quality of available males, measures of female sperm reserves in the EBS (c.f. Slater et al. 2010), and the potential influence of fishery removals of large males will be valuable to provide further insight into the mating dynamics of this stock.

CFI could serve as a useful index of reproductive potential for fishery management. In the spatiotemporally comprehensive CFI time series collected during stock assessment surveys, patterns of variability in fecundity were well approximated. In the fecundity analysis, a statistical model with only CFI and female size as predictors explained the same high proportion (~80%) of the overall variation in fecundity as the more complex model integrating a suite of biological factors and structured by sampling period (Table A1). Trends towards increasing (SC2-SC3) and then decreasing mean fecundity and increased variance in fecundity (SC3-SC5) were also similar between the CFI time series and fecundity analyses. The quasi-cyclic peaks and troughs in mean-weighted CFI (Fig. 6B) observed in this study were also noted in other studies and attributed to interactions among fluctuating recruitment, mature female movement, and declining

female fecundity during periods of increasing average mature female shell condition (Orensanz et al. 2005; Ernst et al. 2012).

For primiparous EBS snow crab variances around both the size-fecundity relationship and temporal variation in mean CFI were lower than that of multiparous females (Fig. 3; Fig. 6B). In addition to the effects of mating and senescence already described, both spatial dynamics and other life history characteristics may also influence this pattern. Primiparous females are distributed in the middle domain of the EBS at 50 -100 m depth (Zheng et al. 2001; Orensanz et al. 2004), a region interannually and seasonally dominated by cold bottom water (-1.5 to 4°C ; Orensanz et al. 2007; Mueter and Litzow 2008). Following the molt to maturity, females undertake a cross shelf ontogenetic migration into the deeper, warmer (~ 2 to 5°C) waters of the outer domain (Orensanz et al. 2004; Ernst et al. 2005). A prolonged duration of ovarian maturation (~ 3 y; Alunno-Bruscia and Sainte-Marie 1998) precedes the production of the primiparous egg clutch, while ovarian maturation takes place in one (annual) or two (biennial) years for multiparous females. Most multiparous females in the outer domain of the EBS are likely on an annual reproductive cycle (Armstrong et al. 2008). Associated with a shorter duration of ovarian maturation (~ 1 y), the fecundity and oocyte/embryo characteristics (e.g. mean dry weight) of multiparous females may exhibit greater variability in relation to the influence of varying environmental or primary production dynamics (Iverson et al. 1979) on maternal condition and energy allocation.

Substantial embryo loss during brooding for female snow crab in the EBS was not detected in this study. Size-fecundity relationships were similar between seasons regardless of shell condition. In support of this conclusion, fecundity was $\sim 13\%$ lower for SC4 versus SC3 females

390 in both summer and spring. Differences in the size-fecundity relationship are likely established
391 prior to oviposition or early in embryo development and persist through embryo incubation
392 without substantial modification due to embryo loss or embryo mortality for females of both
393 shell condition groups. This finding is consistent with that of Sainte-Marie (1993) and contrasts
394 with estimates of reductions in fecundity of 20% to 50% during incubation in other snow crab
395 stocks (Kon 1974; Comeau et al. 1999).

396 Our observation of empty egg cases mixed with pre-hatch stage embryos, indicative of larval
397 release during the spring (March to May) was consistent with previous studies, in which peak
398 snow crab larval abundance was observed in April (Incze et al. 1987), and springtime hatching
399 was inferred from staging of ovarian maturation and embryo development (Reppond et al. 2008).
400 None of the females collected in the spring in our opportunistic samples were SC2, which could
401 be due to sampling outside the geographic distribution of SC2 females (Zheng et al. 2001, Ernst
402 et al. 2005) or perhaps the transition from SC2 to SC3 occurred before the spring sampling
403 (Ernst et al. 2005).

404 Quality of early stage embryos did not vary with maternal size, but some differences were
405 observed among shell condition groups for EBS snow crab. Stability in % elemental composition
406 with significant variation in mean embryo dry weight (~8% reduction for SC4 relative to SC2
407 females; Table 1) implied that differences in mean embryo dry weight are likely to indicate real
408 differences in embryo energy content a difference of in the embryo quality sample. This
409 difference could be interpreted to indicate a trend toward reduced embryo quality with increasing
410 age post-maturity (e.g. elapsed reproductive cycles). However, the pattern of mean embryo dry
411 weight by shell condition observed in the fecundity analysis (Fig. 2B) contrasted with that
412 observed in the embryo quality analysis (Table 1). Values of % C (47-55), % H (7-10), and % N

(8-10) estimated for early stage snow crab embryos were similar to those estimated for early embryo stages of caridean shrimps (Clarke 1993; Anger et al. 2002; Urzúa et al. 2012), and other brachyuran crabs (Fischer et al. 2009). Interestingly, fecundity residuals for females of all shell conditions collected in 2010 were positively correlated with % C (Table 2; Fig. A2), suggesting a potential relationship between higher embryo energy content and higher than average fecundity for females regardless of shell condition. Linkages among environmental variability, maternal condition, and embryo size and energy content have been demonstrated for other brachyuran crabs (Brante et al. 2003; Gimenez and Anger 2001; Bas et al. 2007; Wells 2009) and may also influence snow crab fecundity and embryo quality.

Qualitative comparison of best-fit size-fecundity relationships among snow crab stocks imply that both primiparous and multiparous EBS females may have lower mean fecundity-at-size than conspecifics from Wakasa Bay, Japan and the northwestern Gulf of Sainte-Lawrence, Canada (Fig. 9). Estimated fecundity-at-size for primiparous females sampled in the EBS in 1968 (Haynes et al. 1976) and Chukchi Sea in 1976 (Jewett 1981) were comparable to that of primiparous females sampled in this study (Fig. 7). Potential mechanisms underlying differences in the size-fecundity relationship among stocks have not been assessed for snow crab. However, the possibility of stock-specific differences in reproductive potential per unit of mature female biomass highlight the importance of studies to quantitatively evaluate reproductive parameters and develop stock-specific understanding of the life history processes influencing stock productivity and persistence (Orensanz et al. 2004; Parada et al. 2010; Ernst et al. 2012; Burgos et al. 2013).

In support of the development of biological reference points based on female reproductive potential for EBS snow crab, this study developed quantitative descriptions of relationships

among female size, shell condition, CFI, and fecundity necessary to estimate an index of fertilized egg production for EBS snow crab. This index, estimated from female size and shell condition or size and CFI data collected during annual stock assessment surveys, has the advantage of explicitly accounting for the pronounced demographic trends in size and shell condition observed for this stock (Ernst et al. 2005; Orensanz et al. 2007). Such an index is also unlikely to be biased by either embryo loss during brooding or strong influences of maternal factors on embryo quality, which were not detected in this study.

Relative to spatiotemporally comprehensive collection of CFI data, studies of size-fecundity relationships have limited ability to describe stock-level variability in reproductive potential for high-abundance, broadly-distributed crab stocks. Nevertheless, such studies are necessary to provide contextual insight into factors influencing the size-fecundity relationship, including variability in embryo characteristics, embryo viability, and sperm limitation. Our results suggest that variability in reproductive potential per unit biomass or abundance is lower for primiparous than multiparous females. If spatiotemporally persistent, the patterns of variability in embryo dry weight and carbon content observed between SC2 and SC3/SC4 females in both the fecundity and embryo quality components of this study may warrant further attention due to the potential for “carry-over” effects of embryo quality to the larval and juvenile stages. Improved understanding of life-stage specific (primiparous and multiparous) mating dynamics *in situ* and pre-oviposition factors influencing ovarian maturation and oocyte production are needed to clarify the relative importance of these processes in shaping female reproductive potential. Monitoring of female sperm reserves (e.g. Slater et al. 2010), egg fertilization, and quantitative evaluation of sperm reserves relationships between primiparous and multiparous females similar to efforts for snow crab in the Gulf of Saint Lawrence, Canada (Sainte-Marie et al. 2008), may

provide further insight into factors mediating the probability and magnitude of sperm limitation with female size and shell condition. Biennial reproduction was not addressed in this study, but consideration of this factor will be a key aspect of future efforts to define biological reference points for fishery management based on interannual variability in female reproductive potential and relationships to the male-only fishery (e.g. sperm demand). Monitoring of female fecundity and sperm reserves has continued in the EBS as a long-term study and will provide further insight into the temporal dynamics for this stock.

Draft

Acknowledgements

This research was supported by funding from the North Pacific Research Board (NPRB Project #714). This is NPRB publication number 594 and Alaska Department of Fish and Game professional publication PP-281. Funding for L. Slater’s contributions to this work came from federal grants NA06NMF4370073, NA07NMF4370268, and NA09NMF4370166. Special thanks are extended to Robert Foy of the US NMFS, Kodiak Laboratory for providing data from the EBS annual stock assessment survey, to all personnel who assisted with sample collection and processing during the survey, and to three anonymous reviewers whose advice improved the manuscript.

Draft

References

- Alunno-Bruscia, M., and Sainte-Marie, B. 1998. Abdomen allometry, ovary development, and growth of female snow crab, *Chionoecetes opilio* (Brachyura, Majidae), in the northwestern Gulf of Sainte-Lawrence. Can. J. Fish. Aquat. Sci. **55**: 459-477. doi:10.1139/cjfas-55-2-459.
- Amsler, M.O., and George, R.Y. 1984. Seasonal variation in the biochemical composition of the embryos of *Callinectes sapidus* Rathbun. J. Crust. Biol. **4**(4): 546-553. doi:10.2307/1548068.
- Anger, K., Moriera, G.S., and Ismael, D. 2002. Comparative size, biomass, elemental composition (C, N, H), and energy concentration of caridean shrimp eggs. Invertebr. Reprod. Dev. **42**(2-3): 83-93. doi:10.1080/07924259.2002.9652765.
- Armstrong, D.A., Burgos, J., Ernst, B. and Orensanz, J.M. 2008. Female effective reproductive output of the snow crab stock in the eastern Bering Sea. North Pacific Research Board, Project 508. Final Report. Available from: http://doc.nprb.org/web/05_prjs/508_Final%20Report.pdf. Accessed 21 December 2013. 165 p.
- Bas, C.C., Spivak, E.D., and Anger, K. 2007. Seasonal and interpopulational variability in fecundity, egg size, and elemental composition (CHN) of eggs and larvae in a grapsoid crab, *Chasmagnathus granulatus*. Helgol. Mar. Res. **61**: 225-237. doi:10.1007/s10152-007-0070-y.
- Brante, A., Fernández, M., Eckerle, L., Mark, F., Pörtner, H-O, and Arntz, W. 2003. Reproductive investment in the crab *Cancer setosus* along a latitudinal cline: egg production, embryo losses, and embryo ventilation. Mar. Ecol. Prog. Ser. **251**:221-232. doi:10.3354/meps251221.

- 496 Brooks, E.N., 2013. Effects of variable reproductive potential on reference points for fishery
497 management. *Fish. Res.* **138**:152-58. doi:10.1016/j.fishres.2012.06.003
- 498 Burgos, J., Ernst, B., Armstrong, D., and Orensanz, J.M., 2013. Fluctuations in range and
499 abundance of snow crab (*Chionoecetes opilio*) from the eastern Bering Sea: What role for Pacific
500 cod (*Gadus macrocephalus*) predation? *Bull. Mar. Sci.* **89**:57-81. doi:10.5343/bms.2011.1137
- 501 Caputi, N. 1993. Aspects of spawner-recruit relationships, with particular reference to crustacean
502 stocks: a review. *Aust. J. Mar. Fresh. Res.* **44**: 589-607. doi:10.1071/MF9930589.
- 503 Daly, B.J., Armistead, C.E., and Foy, R.J. 2014. The eastern Bering Sea continental shelf trawl
504 survey: results for commercial crab species. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-
505 AFSC-282.167 p.
- 506 Clarke, A. 1993. Egg size and composition in polar shrimps (Caridea; Decapoda). *J. Exp. Mar.*
507 *Biol. Ecol.* **168**: 189-203. doi:10.1016/0022-0981(93)90259-Q.
- 508 Comeau, M., Starr, M., Conan, G.Y., Rochibaud, G., and Therriault, J.-C. 1999. Fecundity and
509 duration of egg incubation for multiparous female snow crabs (*Chionoecetes opilio*) in the fjord
510 of Bonne Bay, Newfoundland. *Can. J. Fish. Aquat. Sci.* **56**: 1088-1095. doi:10.1139/f99-044.
- 511 Darnell, M.Z., Rittschof, D., Darnell, K.M., and McDowell, R.E. 2009. Lifetime reproductive
512 potential of female blue crabs *Callinectes sapidus* in North Carolina, USA. *Mar. Ecol. Prog. Ser.*
513 **394**: 153-163. doi:10.3354/meps08295.

- 514 Duluc, C., Sainte-Marie, B., and Brethes, J.-C. 2005. Value of indicators for assessing recent
515 mating in brachyuran crabs. J. Mar. Biol. Ass. UK. **85**: 129-135.
516 doi:10.1017/s0025315405010945h.
- 517 Émond, K., Sainte-Marie, B., Galbraith, P.S. , and J. Bêty. 2015. Top-down vs. bottom-up
518 drivers of recruitment in a key marine invertebrate: investigating the early life stages of the snow
519 crab. ICES J. Mar. Sci. doi: 10.1093/icesjms/fsu240.
- 520 Ernst, B., Armstrong, D.A., Burgos, J., and Orensanz, J.M. 2012. Life history schedule and
521 periodic recruitment of female snow crab (*Chionoecetes opilio*) in the eastern Bering Sea. Can. J.
522 Fish. Aquat. Sci. **69**(3): 532-550. doi:10.1139/f2011-173.
- 523 Ernst, B., Orensanz, J.M., and Armstrong, D.A. 2005. Spatial dynamics of female snow crab
524 (*Chionoecetes opilio*) in the eastern Bering Sea. Can. J. Fish. Aquat. Sci. **62**(2): 250-268. doi:
525 10.1139/f04-201.
- 526 Fischer, S., Thatje, S., and Brey, T. 2009. Early egg traits in *Cancer setosus* (Decapoda,
527 Brachyura): effects of temperature and female size. Mar. Ecol. Prog. Ser. **377**: 193-202.
528 doi:10.3354/meps07845.
- 529 Gimenez, L. 2006. Phenotypic links in complex life cycles: conclusions from studies with
530 decapod crustaceans. Integr. Comp. Biol. **46**(5): 615-622. doi:10.1093/icb/icl010.
- 531 Gimenez, L., and Anger, K. 2001. Relationships among salinity, egg size, embryonic
532 development, and larval biomass in the estuarine crab *Chasmagnathus granulata* Dana, 1851. J.
533 Exp. Mar. Biol. Ecol. **260**: 241-257.

- 534 Gnaiger, E., and Bitterlich, G. 1984. Proximate biochemical composition and caloric content
535 calculated from elemental CHN analysis: a stoichiometric concept. *Oecologia* **62**: 289-298.
536 doi:10.1007/BF00384259.
- 537 Hannah, R.W., Jones, S.A., and Long, M.R. 1995. Fecundity of ocean shrimp (*Pandalus*
538 *jordani*). *Can. J. Fish. Aquat. Sci.* **52**: 2098-2107. doi:10.1139/f95-803.
- 539 Haynes, E., Karinen, J.F., Watson, J., and Hopson, D.J. 1976. Relation of eggs and egg length to
540 carapace width in the brachyuran crabs *Chionoecetes bairdi* and *C. opilio* from the southeastern
541 Bering Sea and *C. opilio* from the Gulf of St. Lawrence. *J. Fish. Res. Board Can.* **33**: 2592-2595.
542 doi:10.1139/f76-304.
- 543 Hines, A.H. 1982. Allometric constraints and variables of reproductive effort in brachyuran
544 crabs. *Mar. Biol.* **69**: 309-320. doi:10.1007/BF00397496.
- 545 Houde, E.D. 2008. Emerging from Hjort's Shadow. *J. Northwest Atl. Fish. Sci.* **41**: 53-70.
546 doi:10.2960/J.v41.m634.
- 547 Incze, L.S., Armstrong, D.A., and Smith, S.L. 1987. Abundance of larval Tanner crabs
548 (*Chionoecetes* spp.) in relation to adult females and regional oceanography of the southeastern
549 Bering Sea. *Can. J. Fish. Aquat. Sci.* **44**: 1143-1156. doi:10.1139/f87-137.
- 550 Iverson, R.L., Coachman, L.K., Cooney, R.T., English, T.S., Goerling, J.J., Hunt, G.L., Jr.,
551 Macauley, M.C., McRoy, C.P., Reeburg, W.S., and Whiteledge, T.E. 1979. Ecological
552 significance of fronts in the southeastern Bering Sea. *In* *Ecological processes and marine*
553 *systems*. Edited by R.J. Livingston. Plenum Press, New York. pp. 437-466.

- 554 Jadamec, L.S., Donaldson, W.E., and Cullenberg, P. 1999. Biological field techniques for
555 *Chionoecetes* crabs. University of Alaska Sea Grant College Program. AK-SG-99-02. Fairbanks,
556 AK. 80 p. doi:10.4027/bftcc.1999. 80 p.
- 557 Jewett, S.C. 1981. Variations in some reproductive aspects of female snow crabs *Chionoecetes*
558 *opilio*. J. Shellfish Res. **1**: 95-99.
- 559 Kell, L.T., Nash, R.D.M., Dickey-Collas, M., Mosqueira, I. and Szuwalski, C. 2015. Is spawning
560 stock biomass a robust proxy for reproductive potential?. Fish Fish. doi:10.1111/faf.12131
- 561 Koenker, R. 2006. Quantile regression in R: A Vignette. Available from [http://cran.r-](http://cran.r-project.org/web/packages/quantreg/vignettes/rq.pdf)
562 [project.org/web/packages/quantreg/vignettes/rq.pdf](http://cran.r-project.org/web/packages/quantreg/vignettes/rq.pdf).
- 563 Kon, T. 1974. Fisheries biology of the Japanese Tanner crab - VI. On the number of ovarian eggs
564 and eggs held in the pleopods. Bull. Jap. Soc. Sci. Fish. **40**(5): 465-469.
- 565 Kon, T., and Adachi, T. 2006. Fecundity of snow crab *Chionoecetes opilio* off Wakasa Bay in
566 the Sea of Japan. Nippon Suisan Gakkaishi. **72**(4): 673-680. doi:10.2331/suisan.72.673.
- 567 Kon, T., Ono, M., and Honma, Y. 2010. Histological studies on the spent ovaries of aged snow
568 crabs *Chionoecetes opilio* caught in the Sea of Japan. Fish. Sci. **76**: 227-233.
569 doi:10.1007/s12562-010-0215-9.
- 570 Kuris, A.M. 1990. A review of patterns and causes of crustacean brood mortality. In Crustacean
571 Egg Production. Edited by A. Wenner and A.M. Kuris. A. A. Balkema, Brookfield, Vermont. pp.
572 117-142.

- 573 Lambert, Y. 2008. Why should we closely monitor fecundity in marine fish populations? J.
574 Northwest Atl. Fish. Sci. **41**: 93-106. doi:10.2960/J.v41.m628.
- 575 Lipcius, R.N., and Stockhausen, W.T. 2002. Concurrent decline of the spawning stock,
576 recruitment, larval abundance, and size of the blue crab *Callinectes sapidus* in Cheseapeake Bay.
577 Mar. Ecol. Prog. Ser. **226**: 45-61. doi:10.3354/meps226045.
- 578 Marshall, C.T., Needle, C.L., Thorsen, A., Kjesbu, O.S., and Yaragina, N.A. 2006. Systematic
579 bias in estimates of reproductive potential of an Atlantic cod (*Gadus morhua*) stock: implications
580 for stock-recruit theory and management. Can. J. Fish. Aquat. Sci. **63**(5): 980-994.
581 doi:10.1139/f05-270.
- 582 Moran, A.L., and McAlister, J.S. 2009. Egg size as a life history character of marine
583 invertebrates: Is it all it's cracked up to be? Biol. Bull. (Woods Hole). **216**: 226-242.
- 584 Morgan, M.J., Perez-Rodriguez, A., Saborido-Rey, F., and Marshall, C.T. 2011. Does increased
585 information about reproductive potential result in better prediction of recruitment? Can. J. Fish.
586 Aquat. Sci. **68**(8): 1361-1368. doi:10.1139/f2011-049.
- 587 Moriyasu, M., and Lanteigne, C. 1998. Embryo development and reproductive cycle in the snow
588 crab, *Chionoecetes opilio* (Crustacea: Majidae), in the southern Gulf of Sainte-Lawrence. Can. J.
589 Zool. **76**: 2040-2048. doi:10.1139/z98-147.
- 590 Mueter, F.J., and Litzow, M.A. 2008. Sea ice retreat alters the biogeography of the Bering Sea
591 continental shelf. Ecol. Appl. **18**(2): 309-320. doi:10.1890/07-0564.1.

- 592 Murawski, S. 2001. Impacts of demographic variation in spawning characteristics on reference
593 points for fishery management. ICES J. Mar. Sci. **58**(5): 1002-1014.
594 doi:10.1006/jmsc.2001.1097.
- 595 Nichol, D.G., and Somerton, D.A. 2015. Seasonal migrations of morphometrically mature male
596 snow crab in the eastern Bering Sea in relation to mating dynamics. Fish. Bull. **113**:313-326.
597 doi:10.7755/FB.113.3.7
- 598 Orensanz, J.M., Ernst, B., and Armstrong, D.A. 2007. Variation of female size and stage at
599 maturity in snow crab (*Chionoecetes opilio*)(Brachyura: Majidae) from the eastern Bering Sea. J.
600 Crust. Biol. **27**: 536-591. doi:10.1651/S-2790.1.
- 601 Orensanz, J.M., Ernst, B., Armstrong, D.A., and Parma, A.M. 2005. Detecting early warnings of
602 recruitment overfishing in male-only crab fisheries: an example from the snow crab fishery. *In*
603 Fisheries assessment and management in data-limited situations. Edited by G.H. Kruse, V.F.
604 Galluci, D.E. Hay, R.I. Perry, R.M. Peterman, T.C. Shirley, P.D. Spencer, B. Wilson and D.
605 Woodby. Alaska Sea Grant College Program AK-SG-05-02, University of Alaska Fairbanks. pp.
606 267-288.
- 607 Orensanz, J.M., Ernst, B., Armstrong, D.A., Stabeno, P.J., and Livingston, P. 2004. Contraction
608 of the geographic range of distribution of snow crab (*Chionoecetes opilio*) in the eastern Bering
609 Sea: an environmental ratchet? CalCOFI Reports. **45**: 65-79.
- 610 Ouellet, P., and Plante, F. 2004. An investigation of the sources of variability in American
611 lobster (*Homarus americanus*) eggs and larvae: female size and reproductive status, and
612 interannual and interpopulational comparisons. J. Crust. Biol. **24**: 481-495. doi:10.1651/C-2467.

- 613 Parada, C., Armstrong, D.A., Ernst, B., Hinckley, S., and Orensanz, J.M. 2010. Spatial dynamics
614 of snow crab (*Chionoecetes opilio*) in the eastern Bering Sea - Putting together the pieces of the
615 puzzle. Bull. Mar. Sci. **86**(2): 413-437.
- 616 Prager, M.H., McConaughy, J.R., Jones, C.M., and Geer, P.J. 1990. Fecundity of blue crab,
617 *Callinectes sapidus*, in Cheseapeake Bay: Biological, statistical and management considerations.
618 Bull. Mar. Sci. **46**: 170-179.
- 619 Punt, A.E., Szuwalski, C.S., and Stockhausen, W. 2014. An evaluation of stock-recruitment
620 proxies and environmental change points for implementing the US Sustainable Fisheries Act.
621 Fisheries Research **157**:28-40. doi:10.1016/j.fishres.2194.03.015
- 622 Racotta, I.S., Palacios, E., and Ibarra, A.M. 2003. Shrimp larval quality in relation to broodstock
623 condition. Aquaculture **227**(1-4): 107-130. doi:10.1016/s0044-8486(03)00498-8.
- 624 Reid, D.M., and Corey, S. 1991. Comparative fecundity of decapod crustaceans, II. The
625 fecundity of fifteen species of anomuran and bracyuran crabs. Crustaceana. **61**: 175-189.
626 doi:10.1163/156854091X00669.
- 627 Reppond, K., Rugolo, L, and de Oliveira, A.C.M. 2008. Change in biochemical composition
628 during development of snow crab, *Chionoecetes opilio*, embryos. J. Crust. Biol. **28**(3): 519-527.
629 doi:10.1651/07-2862r1.1
- 630 Rondeau, A., and Sainte-Marie, B. 2001. Variable mate-guarding time and sperm allocation by
631 male snow crabs (*Chionoecetes opilio*) in response to sexual competition, and their impact on the
632 mating success of females. Biol. Bull. (Woods Hole) **201**: 204-217. doi:10.2307/1543335.

- 633 Rothschild, B., and Fogarty, M. 1989. Spawning-stock biomass: A source of error in
634 recruitment/stock relationships and management and advice. ICES J. Mar. Sci. **45**: 131-135.
635 doi:10.1093/icesjms/45.2.131.
- 636 Sainte-Marie, B. 1993. Reproductive cycle and fecundity of primiparous and multiparous female
637 snow crab, *Chionoecetes opilio*, in the northwest Gulf of Saint Lawrence. Can. J. Fish. Aquat.
638 Sci. **50**: 2147-2156. doi:10.1139/f93-240.
- 639 Sainte-Marie, B., Gosselin, T., Sévigny, J.-M., and Urbani, N. 2008. The snow crab mating
640 system: opportunity for natural and unnatural selection in a changing environment. Bull. Mar.
641 Sci. **83**: 131-161.
- 642 Sainte-Marie, B., Sévigny, J.-M., and Carpentier, M. 2002. Interannual variability of sperm
643 reserves and fecundity of primiparous females of the snow crab (*Chionoecetes opilio*) in relation
644 to sex ratio. Can. J. Fish. Aquat. Sci. **59**: 1932-1940. doi:10.1139/f02-162.
- 645 Slater, L.M., MacTavish, K.A., and Pengilly, D. 2010. Preliminary analysis of spermathecal load
646 of primiparous snow crabs (*Chionoecetes opilio*) from the eastern Bering Sea, 2005-2008. In
647 Biology and management of exploited crab populations under climate change. Edited by G.H.
648 Kruse, G.L. Eckert, R.J. Foy, R.N. Lipcius, B. Sainte-Marie, D.L. Stram and D. Woodby. Alaska
649 Sea Grant, University of Alaska Fairbanks. pp. 237-248.
- 650 Somerton, D. A. 1981. Life history and population dynamics of two species of Tanner crab,
651 *Chionoecetes bairdi* and *C. opilio* in the eastern Bering Sea with implications for the
652 management of the commercial harvest. Ph.D. dissertation. University of Washington. Seattle,
653 Wash. 220 pp.

- 654 Somerton, D.A., and W.S. Meyers. 1983. Fecundity differences between primiparous and
655 multiparous female Alaskan Tanner crab (*Chionoecetes bairdi*). J. Crust. Biol. **3**: 183-183.
656 doi:10.2307/1548254.
- 657 Swiney, K.M., Long, W.C., Eckert, G.L., and G.H. Kruse. 2012. Red king crab, *Paralithodes*
658 *camtschaticus*, size-fecundity relationship, and interannual and seasonal variability in fecundity.
659 J. Shellfish Res. **31**(4):925-933. doi:10.2983/035.031.0403.
- 660 Szuwalski, C.S. and Punt, A.E. 2012. Fisheries management for regime-based ecosystems: a
661 management strategy evaluation for the snow crab fishery in the eastern Bering Sea. ICES J.
662 Mar. Sci. **70**:955-967. doi:10.1093/icesjms/fss182
- 663 Szuwalski, C.S. and Punt, A.E. 2013. Regime shifts and recruitment dynamics of snow crab,
664 *Chionoecetes opilio*, in the eastern Bering Sea. Fish. Ocean. **22**(5):345-354.
665 doi:10.1111/fog.12026
- 666 Turnock, B.J., and Rugolo, L.J. 2014. Stock assessment of eastern Bering Sea snow crab. *In*
667 Stock Assessment and Fishery Evaluation for the king and Tanner crab fisheries of the Bering
668 Sea and Aleutian Islands regions. North Pacific Fishery Management Council, Anchorage, AK.
669 pp 41-177.
- 670 Urzúa, Á., Paschke, K., Gebauer, P., and Anger, K. 2012. Seasonal and interannual variations in
671 size, biomass and chemical composition of the eggs of North Sea shrimp, *Crangon crangon*
672 (Decapoda: Caridea). Mar. Biol. **159**: 583-599. 10.1007/s00227-011-1837-x.

- 673 Verisimo, P., Bernardez, C., Gonzalez-Gurriaran, E., Freire, J., Muino, R., and Fernandez, L.
674 2011. Changes between consecutive broods in the fecundity of the spider crab, *Maja*
675 *brachydactyla*. ICES J. Mar. Sci. **68**: 472-478. doi:10.1093/icesjms/fsq164.
- 676 Watson, J. 1970. Maturity, mating, and egg laying in the spider crab, *Chionoecetes opilio*. J.
677 Fish. Res. Board Can. **27**: 1607-1616. doi:10.1139/f70-181.
- 678 Webb, J.B., and Bednarski, J. 2010. Variability in reproductive potential among exploited stocks
679 of Tanner crab (*Chionoecetes bairdi*) in southeastern Alaska. In Management of exploited crab
680 population under climate change. Edited by G.H. Kruse, G.L. Eckert, R.J. Foy, R.N. Lipcius, B.
681 Sainte-Marie, D.L. Stram and D. Woodby. Alaska Sea Grant College Program AK-SG-10-01,
682 University of Alaska Fairbanks. pp. 295-317.
- 683 Webb, J.B., Eckert, G.L., Shirley, T.C., and Tamone, S.L. 2007. Changes in embryonic
684 development and hatching in *Chionoecetes opilio* (snow crab) with variation in incubation
685 temperature. Biol. Bull. (Woods Hole) **213**: 67-75. doi:10.2307/25066619.
- 686 Wells, S. 2009. Changing fecundity and reproductive output in females of a Chesapeake Bay
687 population of blue crab, *Callinectes sapidus*. Ph. D. Dissertation, Old Dominion University,
688 Norfolk, VA. 91 pp.
- 689 Zar, J.H. 1999. Biostatistical Analysis. Prentice-Hall, Upper Saddle River, NJ.
- 690 Zheng, J. and Kruse, G.H. 2006. Recruitment variation of eastern Bering Sea crabs: Climate-
691 forcing or top-down effects. Prog. Ocean. **68**:184-204. doi:10.1016/j.pocean.2006.02.002.

- 692 Zheng, J., Kruse, G.H., and Ackley, D.R. 2001. Spatial distribution and recruitment patterns of
693 snow crabs in the eastern Bering Sea. *In* Spatial processes and management of marine
694 populations. *Edited by* G.H. Kruse, N. Bez, M.W. Booth, M.W. Dorn, S. Hills, R.N. Lipcius, D.
695 Pelletier, C. Roy, S.J. Smith and D. Witherell. Alaska Sea Grant College Program AK-SG-01-02,
696 Fairbanks, AK. pp. 233-255.
- 697 Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., and Smith, G.M. 2009. Mixed effects
698 models and extensions in ecology with R. Springer, New York, NY

Draft

Table 1. Embryo characteristics and embryo elemental composition by shell condition for newshell (SC2), oldshell (SC3), and very oldshell (SC4) female snow crab collected from the eastern Bering Sea in 2010. Values sharing the same letters were not significantly different (Tukey HSD, $p \geq 0.05$).

Embryo Characteristics			
Shell Condition	2	3	4
N	22	18	17
Embryo Volume (mm ³)	0.13 ± 0.00 ^a	0.13 ± 0.00 ^a	0.12 ± 0.00 ^a
Mean Embryo Dry Mass (µg)	68.2 ± 0.97 ^a	65.1 ± 1.32 ^{ab}	63.2 ± 1.24 ^b
Embryo Elemental Composition			
Shell Condition	2	3	4
N	16	10	15
% C	51.1 ± 0.36 ^a	51.4 ± 0.35 ^a	50.8 ± 0.41 ^a
% N	8.96 ± 0.08 ^a	9.18 ± 0.05 ^a	9.23 ± 0.06 ^a
% H	8.60 ± 0.29 ^a	7.64 ± 0.17 ^a	7.98 ± 0.26 ^a
C/N	5.71 ± 0.03 ^a	5.60 ± 0.05 ^{ab}	5.50 ± 0.02 ^b
C (µg/embryo)	34.4 ± 6.01 ^a	33.8 ± 7.90 ^{ab}	32.3 ± 7.19 ^b
N (µg/embryo)	6.04 ± 0.11 ^a	6.04 ± 0.13 ^a	5.86 ± 0.14 ^a
H (µg/embryo)	5.79 ± 0.21 ^a	5.03 ± 0.14 ^b	5.06 ± 0.17 ^b

Table 2. Correlation among indices of embryo quality for eastern Bering Sea snow crab, *Chionoecetes opilio*, sampled in 2010. Pairwise Pearson correlation coefficient (*r*) and *p*-values for one-tailed t-tests among, % C (carbon), % N (nitrogen), C/N (carbon/nitrogen ratio), mean embryo weight (MEW), mean embryo volume (MEV), and RES (residuals of the ANCOVA of fecundity are as a function of size and shell condition) are summarized, respectively, in the upper and lower triangles. Significant relationships (*p*<0.05) are indicated in bold.

	% C	%N	C/N	MEW	MEV	RES
% C	*	0.612	0.353	-0.019	-0.143	0.399
% N	<0.001	*	-0.520	-0.211	-0.182	0.259
C/N	0.024	<0.001	*	0.229	0.054	0.229
MEW	0.909	0.186	0.151	*	0.357	-0.122
MEV	0.374	0.256	0.737	0.008	*	-0.122
RES	0.010	0.102	0.447	0.365	0.367	*

712 Table 3. Weighted mean clutch fullness index (CFI) by shell condition class pooled among years
713 (1994 to 2014) for eastern Bering Sea snow crab. Weighted mean CFI ($\pm$ SD) and coefficient of
714 variation (CV $\pm$ SD) were not significantly different for groups sharing the same letter within
715 each column (Dunn's post-hoc test; $p < 0.05$).

	Weighted Mean CFI	CV of Weighted Mean CFI
SC2	4.7 ± 0.68^{ab}	0.12 ± 0.017^a
SC3	4.7 ± 0.99^a	0.30 ± 0.026^a
SC4	3.8 ± 1.7^b	0.60 ± 0.059^b
SC5	1.3 ± 1.4^c	0.55 ± 0.50^c

716

717

718 **Figure Captions:**

719 Figure 1. Value (nominal USD) and landings of eastern Bering Sea snow crab, 1965-2015.

720 Figure 2. Model-adjusted fecundity (A), mean embryo weight (B), and clutch dry weight (C) of
 721 sampled mature female snow crab from the eastern Bering Sea (n=853) with embryos in the
 722 early stages of development, by year (2007, 2008, 2009) and by shell condition index (SC2 –
 723 Newshell, SC3 – Oldshell, SC4 – Very Oldshell, SC5 – Very, Very Oldshell), presented as mean
 724 $\pm$ SE. Levels sharing the same letter were not significantly different (Tukey HSD; $p < 0.05$).
 725 Sample sizes indicated within each bar.

726 Figure 3. Quantile regression fits (solid black lines) of size-fecundity by shell condition group
 727 (SC2 – Newshell, SC3 – Oldshell, SC4 – Very Oldshell, SC5 – Very, Very Oldshell) at the 95th
 728 (upper line) and 5th (lower line) for mature female snow crab with egg clutches in the early
 729 stages of embryo development collected from the eastern Bering Sea from 2007 to 2009.
 730 Parameter estimates are summarized in Table A2.

731 Figure 4. Mean standardized fecundity residuals and sperm reserves, indexed as the qualitative
 732 assessment of spermathecal fullness (A) or the presence-absence of fresh ejaculate (B), by shell
 733 condition group (SC2 – Newshell, SC3 – Oldshell, SC4 – Very Oldshell, SC5 – Very, Very
 734 Oldshell) for mature female snow crab sampled from eastern Bering Sea from 2007 to 2009.
 735 Residuals (mean $\pm$ SE) were standardized by a linear mixed model and levels sharing the same
 736 letter were not significantly different (Tukey HSD; $p < 0.05$). Sample sizes indicated within each
 737 bar.

738 Figure 5. Size-fecundity (A), with best-fit lines by clutch fullness index (CFI) category estimated
 739 by ANCOVA (Table 10) for females collected 2007 to 2009. Time series of CFI (B) from
 740 eastern Bering Sea annual stock assessment surveys (Daly et al. 2014), with interannual (1994-
 741 2014) variation in weighted mean CFI by shell condition index (SC). CFI ranges from no eggs
 742 (CFI -1) to 100% full egg clutch (CFI – 6), and SC ranges from newshell (SC2) to very, very
 743 oldshell (SC5) for mature female snow crab. The fecundity analysis study period (2007-2009) is
 744 indicated by the vertical lines

745 Figure 6. Size-fecundity (A) early (summer) versus late (spring) in the brooding duration for
 746 female eastern Bering Sea snow crab collected in summer 2007 and 2009 and spring 2008 and
 747 2010 and mean model-adjusted (ANCOVA) fecundity (B) of SC3 and SC4 pooled among years
 748 and seasons. Sample sizes are indicated within the bars, and levels sharing the same letter were
 749 not significantly different (Tukey HSD; $p < 0.05$).

750 Figure 7. Best-fit size-fecundity relationships among snow crab stocks based on literature values
 751 evaluated at 5 mm intervals of female carapace width (mm) for primiparous (P) and multiparous
 752 (M) females from the northwest Gulf of Sainte Lawrence, Canada (Sainte-Marie 1993), Wakasa

753 Bay, Japan (Kon and Adachi 2006), southeastern Bering Sea (Haynes et al. 1976, this study),
754 Chukchi Sea (Jewett 1981), and this study.

755

Draft

756 Figure 1.

757 Figure 2.

Figure 3.

761 Figure 4.

762 Figure 5.

763 Figure 6.

764

Figure 7.

Appendix to:

The contribution of fecundity and embryo quality to reproductive potential of eastern Bering Sea snow crab

Joel B. Webb^{1*}, Laura M. Slater², Ginny L. Eckert¹, and Gordon H. Kruse¹

¹ University of Alaska Fairbanks, School of Fisheries and Ocean Sciences, 17101 Point Lena Loop Road, Juneau, AK 99801

² Alaska Department of Fish and Game, Division of Commercial Fisheries, 351 Research Court, Kodiak, AK 99615

*Corresponding Author: Alaska Department of Fish and Game, Division of Commercial Fisheries, PO Box 115526, Juneau, AK 99811. Phone: 907-465-6129, Fax: 907-465-6402, Email: joel.webb@alaska.gov

Table A1. Results of linear mixed model analysis of variation in female size-fecundity with the effect of shell condition index (SC), year of collection (YEAR), carapace width (CW) nested within shell condition index and year, mean embryo dry weight (MEW) nested within shell condition index and year, and location of collection (LOC) estimated as a random component of the residual variance, on fecundity of eastern Bering Sea snow crab with embryos in the early stages of development from summer 2007 to summer 2009.

	N	R^2	RMSE
WHOLE MODEL FIT	853	0.80	3895
FIXED EFFECT TESTS	df	F	p
CW[SC, YEAR]	12	89.1	<0.0001
MEW[SC, YEAR]	12	19.5	<0.0001
SC	3	15.1	<0.0001
YEAR	2	3.77	0.044
YEAR*SC	6	5.07	<0.0001
COMPONENTS OF RESIDUAL VARIANCE	Variance	SE	% of Total
LOC	1.03×10^7	3.65×10^6	20.8
Error	5.10×10^7	2.57×10^6	79.2

Table A2. Parameter estimates and bootstrap estimated standard errors (SE) of the slope and intercepts of quantile regression fits to the 95% and 5% quantiles of the size-fecundity relationship by shell condition group for eastern Bering Sea snow crab collected from 2007 to 2009. Data and linear fits are presented in Figure 3.

Quantile	95%		5%	
	Estimate	SE	Estimate	SE
Intercept	-77170	4305	-47770	15240
CW	2019	79	1217	283
SC3	-28640	8072	-14740	22840
SC4	-6183	12100	32670	26940
SC5	18720	38010	41080	33560
CW*SC3	709	140	28	473
CW*SC4	148	217	-871	430
CW*SC5	-334	729	-1082	611

Draft

Table A3. Results of linear mixed model analysis of variation in mean embryo weight by shell condition index (SC), year of collection (YEAR), an interaction term, and location of collection (LOC) estimated as a random component of the residual variance on mean embryo dry weight of female eastern Bering Sea snow crab with embryos in the early stages of development from summer 2007 to summer 2009. Carapace width nested within shell condition and year was not a significant covariate ($F = 1.36, p = 0.18$) and was excluded from the final model. Results are summarized in Fig. 2.

WHOLE MODEL FIT		N	R^2	RMSE
		853	0.52	5.53×10^{-6}
FIXED EFFECT TESTS		df	F	p
	SC	3	42.6	<0.0001
	YEAR	2	21.5	<0.0001
	SC*YEAR	6	12.2	<0.0001
COMPONENTS OF RESIDUAL VARIANCE		Variance	SE	% of Total
	LOC	0.005	0.0014	26.0
	Error	0.016	0.0008	74.0

Table A4. Results of linear mixed model analysis of variation in clutch dry weight by shell condition index (SC), year of collection (YEAR), carapace width (CW) nested within shell condition index and year, mean embryo dry weight (MEW) nested within shell condition index and year, and location of collection (LOC) estimated as a random component of the residual variance, for females with embryos in the early stages of development in the eastern Bering Sea from summer 2007 to summer 2009. Results are summarized in Fig. 2.

WHOLE MODEL		N	R^2	RMSE
		853	0.73	0.656
FIXED EFFECT TESTS		df	F	p
	CW[SC, YEAR]	12	76.6	<0.0001
	MEW[SC, YEAR]	12	4.09	<0.0001
	SC	3	16.6	<0.0001
	YEAR	2	4.78	0.044
	YEAR*SC	6	6.41	<0.0001
COMPONENTS OF RESIDUAL VARIANCE		Variance	SE	% of Total
	LOC	0.0055	0.0025	12.5
	Error	0.0375	0.0020	87.5

Table A5. Results of linear mixed model analysis of size-fecundity by clutch fullness index (CFI), female carapace width (CW in mm), an interaction term, and location of collection (LOC) estimated as a random component of the residual variance with fecundity of female eastern Bering Sea snow crab collected from 2007 to 2009. Linear fits are shown in Figure 6.

WHOLE MODEL FIT		N	R^2	RMSE
		853	0.80	469.6
FIXED EFFECT TESTS		df	F	p
	CW	1	133.1	<0.0001
	CFI	4	133.2	<0.0001
	CW*CFI	4	13.7	<0.0001
COMPONENTS OF RESIDUAL VARIANCE		Variance	SE	% of Total
	LOC	178.2	44.7	27.5
	Error	469.8	23.9	72.5

Table A6. Results of analysis of covariance (ANCOVA) of the size-fecundity relationships early versus late in the brooding duration for female snow crab collected from the eastern Bering Sea. Females were collected in the summer with egg clutches in early stages of development or in the spring with egg clutches in the late (i.e. eyed) stages of embryo development. ANCOVA model effects include carapace width (CW) nested by shell condition index (SC), year of embryo extrusion (YEAR), and season (summer or spring), mean embryo dry weight (MEW) nested within the same terms as CW, and the nesting factors as main effects. Interaction terms were not significant ($p>0.05$) and were dropped from the model.

WHOLE MODEL FIT		N	df	<i>F</i>	<i>p</i>	<i>R</i> ²
		375	355	40.2	<0.0001	0.67
FIXED EFFECTS TESTS			df	<i>F</i>	<i>p</i>	
	CW [SC, YEAR, SEASON]		8	70.47	<0.0001	
	MEW [SC, YEAR, SEASON]		8	26.81	<0.0001	
	SC		1	7.60	0.0420	
	YEAR		1	0.05	0.8273	
	SEASON		1	1.74	0.1877	

Table A7. Results of analyses of covariance (ANCOVA) of the embryo characteristics and embryo elemental composition as a function of female carapace width (CW), shell condition (SC), and their interaction for female snow crab, *Chionoecetes opilio*, sampled in the eastern Bering Sea in 2010. The number of error degrees of freedom was 56 for each ANCOVA of embryo characteristics and 37 for embryo elemental composition. Significant factors ($p<0.05$) are indicated in bold.

Effects	<i>df</i>	<i>F</i>	<i>P</i>	<i>df</i>	<i>F</i>	<i>P</i>
Fecundity				Mean Embryo Dry Weight		
CW	1	89.2	<0.001	1	1.24	0.270
SC	2	16.9	<0.001	2	4.74	0.013
CW*SC	2	6.11	0.004	2	1.03	0.365
Embryo Volume				C/N		
CW	1	0.458	0.502	1	1.200	0.281
SC	2	0.151	0.861	2	8.200	<0.001
CW*SC	2	1.07	0.353	2	0.591	0.559
% C				C (µg/embryo)		
CW	1	0.325	0.572	1	0.75	0.393
SC	2	0.462	0.634	2	3.85	0.0308
CW*SC	2	1.145	0.330	2	3.41	0.0443
% H				H (µg/embryo)		
CW	1	0.035	0.852	1	0.05	0.83
SC	2	3.26	0.051	2	6.15	0.01
CW*SC	2	0.221	0.803	2	1.47	0.24
% N				N (µg/embryo)		
CW	1	2.070	0.159	1	1.405	0.244
SC	2	2.750	0.078	2	1.389	0.263
CW*SC	2	0.318	0.731	2	2.597	0.089

Figure A1. Map of sampling locations of female eastern Bering Sea snow crab by shell condition (SC) collected during annual summer stock assessment bottom trawl surveys for investigation of fecundity in 2007 (A), 2008 (B), and 2009 (C) and investigation of embryo quality in 2010 (D). Symbol size (top left of each panel) and symbol hatching (key in panel D) indicate, respectively, the number of females collected and shell condition composition at each location sampled.

Figure A2. Size-fecundity relationship (A), fecundity residuals (standardized for maternal size and shell condition) and embryo % C (B), mean embryo volume and mean embryo weight (C), and embryo % C and embryo % N (D) by shell condition (SC) for female snow crab collected from the eastern Bering Sea in 2010 for investigation of embryo quality.