2011

Temperature-growth divergence in white spruce forests of Old Crow Flats, Yukon Territory, and adjacent regions of northwestern North America

Post-print/ Accepted manuscript

Trevor J. Porter

Michael F.J. Pisaric

Porter, T. J. and Pisaric, M. F. J. (2011). Temperature-growth divergence in white spruce forests of Old Crow Flats, Yukon Territory, and adjacent regions of northwestern North America. *Glob. Change Biol.*, 17: 3418–3430. [doi:10.1111/j.1365-2486.2011.02507.x](https://doi.org/10.1111/j.1365-2486.2011.02507.x)

This is the peer reviewed version of the following article: Porter, T. J. and Pisaric, M. F. J. (2011). Temperature-growth divergence in white spruce forests of Old Crow Flats, Yukon Territory, and adjacent regions of northwestern North America. *Glob. Change Biol.*, 17: 3418–3430, which has been published in final form at [doi:10.1111/j.1365-2486.2011.02507.x](https://doi.org/10.1111/j.1365-2486.2011.02507.x) This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

HOW TO CITE TSPACE ITEMS

Always cite the published version, so the author(s) will receive recognition through services that track citation counts, e.g. Scopus. If you need to cite the page number of the TSpace version (original manuscript or accepted manuscript) because you cannot access the published version, then cite the TSpace version **in addition to** the published version using the permanent URI (handle) found on the record page.

1 Temperature-growth divergence in white spruce forests of Old Crow Flats, Yukon Territory, and
2 adjacent regions of northwestern North America

3
4 Trevor J. Porter* and Michael F.J. Pisaric

5
6 Department of Geography and Environmental Studies, Carleton University, 1125 Colonel By
7 Drive, Ottawa, Ontario, Canada, K1S 5B6

8
9 *Corresponding author: +1-613-520-2600 ext. 1836, +1-613-520-4301 (fax),

10 tjporter@connect.carleton.ca

11
12 Keywords: dendroclimatology, ring-width, divergence, boreal treeline, white spruce, Old Crow
13 Flats, Yukon Territory

Abstract

We present a new 23-site network of white spruce ring-width chronologies near boreal treeline in Old Crow Flats, Yukon Territory, Canada. Most chronologies span the last 300 years, and some reach the mid-16th century. The chronologies exhibit coherent growth patterns before the 1930s. However, since the 1930s they diverge in trend and exhibit one of two contrasting, but well-replicated patterns we call Group 1 and Group 2. Over the instrumental period (1930-2007) Group 1 sites were inversely correlated with previous-year July temperatures while Group 2 sites were positively correlated with growth-year June temperatures. At the broader northwestern North America (NWN) scale, we find that the Group 1 and Group 2 patterns are common to a number of white spruce chronologies, which we call NWN 1 and NWN 2 chronologies. The NWN 1 and NWN 2 chronologies also share a single coherent growth pattern prior to their divergence (~1950s). Comparison of the NWN 1/NWN 2 chronologies against gridded 20th-century temperatures for NWN and reconstructed Northern Hemisphere summer temperatures (A.D. 1300-2000) indicates that all sites responded positively to temperature prior to the mid-20th century (at least back to A.D. 1300), but that some changed to a negative response (NWN 1) while others maintained a positive response (NWN 2). The spatial extent of divergence implies a large-scale forcing. As the divergence appears to be restricted to the 20th century, we suggest the temperature response shift represents a moisture stress caused by an anomalously warm, dry 20th-century climate in NWN, as indicated by paleoclimatic records. However, because some sites do not diverge, and are located within a few kilometres of divergent sites, we speculate that site-level factors have been important in determining the susceptibility of sites to the large-scale drivers of divergence.

Introduction

In high-latitude regions, dendroclimatic studies often report a positive relation between summer temperatures and tree-ring width (e.g., Briffa et al. 1990; Szeicz and MacDonald 1995; Gostev et al. 1996; D'Arrigo et al. 2006; Frank et al. 2007; Wilson et al. 2007; Youngblut and Luckman 2008). This observed relation is intuitive for trees living at the cold northern margins of the boreal forest where tree growth is largely thought to be temperature-dependent; however, this is not always the case. In northwestern North America (NWN), in particular, many white spruce (*Picea glauca* [Moench] Voss) stands are inversely correlated with previous-year summer temperature (e.g., Barber et al. 2000; Lloyd and Fastie 2002; Wilmking et al. 2004; Pisaric et al. 2007; McGuire et al. 2010). At some of these sites, inverse relations may have persisted over the entire 20th century (Lloyd and Bunn 2007), but at others they appear to be a recent phenomenon (Jacoby and D'Arrigo 1995; D'Arrigo et al. 2004). Similar temperature response shifts are also found at high-latitude Eurasian sites (Briffa et al. 1998; Jacoby et al. 2000). Collectively, these instances of transient temperature-growth responses are referred to as the 'Divergence Problem' (DP) (D'Arrigo et al. 2008).

In a paleoclimatology context, DP complicates the use of affected tree-ring chronologies as temperature proxies since reconstructions depend on time-stable proxy-climate relations. However, because high-latitude tree-ring networks are an important data source for centennial- to millennial-length temperature reconstructions (Jansen et al. 2007), it is important to improve understandings of DP and the extent to which past climate-growth relations can be considered time-stable.

The divergence problem is often observed as a low-frequency departure between summer temperature and ring-width/density occurring after the mid- to late-20th century (D'Arrigo et al.

2008), coinciding with the warmest period the Arctic has experienced in the last two millennia (Kaufman et al. 2009). Although the cause(s) of DP are largely unknown, many have suggested the relatively warm late-20th century may be driving this non-linear behaviour by temperature-induced drought stress (Jacoby and D'Arrigo 1995; Barber et al. 2000; Lloyd and Bunn 2007; McGuire et al. 2010) or optimal biological temperature thresholds being surpassed (D'Arrigo et al. 2004; Wilmking et al. 2004). Other possible explanations for DP include late-20th-century changes in snow cover (Vaganov et al. 1999), UV-B radiation (Briffa et al. 2004), and global dimming (D'Arrigo et al. 2008). However, current understandings of DP are based on a small number of study sites, limiting our ability to draw conclusions about its causes and the likelihood that past temperature-growth relations were also impacted.

Here we present a new network of site-averaged ring-width chronologies from 23 white spruce sites in Old Crow Flats, Yukon Territory, Canada, to satisfy 3 objectives: (1) expand the high-latitude tree-ring network into a region where this research has been absent; (2) examine the response of this network to temperature; and (3) determine if trees in this region were impacted by DP. Further, we draw comparisons between these results and a larger-scale network of white spruce sites across NRNA. Old Crow Flats hosts the northernmost extent of boreal treeline in Yukon Territory and is adjacent to interior Alaska, central Yukon, and the Mackenzie Delta, areas where divergence has been identified (D'Arrigo et al. 2004; Wilmking et al. 2004; Pisarcic et al. 2007). Our site chronology network is unique in NRNA due to its very high site density. This characteristic allows us to better identify regionally-significant growth patterns that are more closely linked to regional-scale factors such as climate than any individual site chronology (Hughes 2011).

Materials and methods

Study region

The Old Crow Flats region (Fig. 1) is a low-lying basin complex bounded by mountain ranges in Alaska and Yukon Territory. The region's surficial geology is primarily defined by a thick glaciolacustrine clay unit deposited by Glacial Lake Old Crow when it occupied the area from ~24,000-12,000 years BP (Hughes 1972; Morlan 1980; Thorson and Dixon 1983; Dyke et al. 2002). Much of the region is poorly drained and covered by a vast mosaic of shallow lakes and peatlands (Ovenden and Brassard 1989; Labreque et al. 2009). Meandering channels incise the surficial clay and provide some drainage (Lauriol et al. 2002). Channel floodplains are well drained and covered by thick organic layers underlain by fine-to-coarse fluvial deposits (Hughes, 1971). White spruce forests in the region are generally found on floodplains while black spruce (*Picea mariana* [Mill.] BSP) forests tend to occupy poorly drained areas amongst the lakes.

Old Crow Flats is in the continuous permafrost zone (Heginbottom et al. 1995). Climate is highly seasonal (Fig. 2) with dry, stable Arctic air dominating during winter, and relatively warm, moist air from the North Pacific and Beaufort Sea during summer (Dyke 2000). Mean annual, winter (Dec.-Feb.), and summer (Jun.-Aug.) temperatures at Old Crow are -9.0°C, -28.6°C, and 12.6°C, respectively (www.climate.weatheroffice.gc.ca). Minimum (maximum) temperatures are below freezing for all months except June-August (May-September) (Fig. 2). Despite its abundance of lakes, annual precipitation at Old Crow Flats is low compared to other northern regions with only parts of the Canadian Arctic Archipelago and northern Greenland receiving less precipitation per annum (Serreze and Barry 2005). Old Crow receives ~265 mm of precipitation annually, ~38 mm during winter, and ~119 mm in summer (Fig. 2).

Tree-ring data

Twenty-three white spruce stands were sampled in 2007 and 2008 (Fig. 1a); all sites are within 150 km of latitudinal treeline. General sampling locations were preselected so that sites would be distributed across the region. In the field, mature sites were preferentially sampled to maximize the length of resultant tree-ring chronologies. Mature sites were identified based on tree morphology and abundance of deadwood. All sites shared a similar open canopy structure such that light could easily penetrate to ground level, and a comparable assemblage of shrubs, mosses, grasses, and lichens. All sites except TM2 and SC1 are situated on channel floodplains. TM2 is situated on a hill known locally as Timber Hill and faces south-west; SC1 is a relatively high-elevation site (~647 m a.s.l. compared to ~267 m a.s.l. on average for the other sites; Table 1) in the Old Crow Range and faces north-east. The remaining sites have no particular aspect.

On average, 33 trees were sampled per site, most of which (~80%) were living. Standard dendrochronology techniques were used to collect and prepare tree-ring samples for ring-width measurement (Speer 2010). Rings were visually cross-dated and measured using a Velmex tree-ring measuring system (precise to 0.001 mm). Two radii per tree were measured in all but a few cases. Cross-dating accuracy was verified with the computer program COFECHA (Holmes 1983). Age-related trends were removed from raw data using conservative negative exponential or negative/zero slope linear curve fits (Fritts et al. 1969). A ‘signal-free’ beta version of the computer program ARSTAN (courtesy of Ed Cook, Lamont-Doherty Earth Observatory Tree-Ring Lab, pers. comm.) was used to calculate standard tree-ring indices according to the ‘signal-free standardisation’ approach described by Melvin and Briffa (2008); mean site chronologies were calculated using the robust bi-weight mean (Cook 1985). Signal-free standardisation was used instead of traditional standardisation because it is ideally suited to avoid ‘trend-distortion,’

an effect that concentrates towards the modern ends of tree-ring series and results in a distorted climate signal in mean tree-ring chronologies (Melvin and Briffa 2008). For some of our sites, trend distortion would be a valid concern if signal-free methods were not used (SI-Figs. 1 and 2).

Temperature-growth analysis

The Old Crow climate record is relatively short (1951-2007) and incomplete. To support a longer comparison with our tree-ring data, we developed a regional composite temperature record from nearby stations including Fairbanks and Fort Yukon in interior Alaska, and Inuvik, Aklavik, and Fort McPherson in the Mackenzie Delta region of Northwest Territories (Fig. 1b). These records compare well with the Old Crow record in terms of trend and interannual variability, and can be used to estimate regional temperatures. Alaskan data were provided by the Alaska Climate Research Center (M. Shulski, 2008, pers. comm.) and Canadian data were obtained from Environment Canada (www.climate.weatheroffice.ec.gc.ca). These records were averaged by region for interior Alaska and the Mackenzie Delta (following normalization), and the regional means were then combined to create a monthly minimum, mean, and maximum temperature record for the larger region centered on Old Crow Flats. The regional composite spans 1930-2007 (pre-1930 data were not used due to limited spatial coverage) and is well correlated with minimum, mean, and maximum temperatures at Old Crow ($r = 0.88, 0.91$, and 0.88 , respectively, for the average month; $p \leq 0.001$).

Growth-year and previous-year May-August temperatures (1930-2007) were compared to each of the 23 site chronologies to determine their dominant response to temperature. Pearson's Product-Moment Correlation Coefficients (r -values) were used to quantify these relations. Sites that shared a common temperature response were averaged into mean regional chronologies to

162 enhance the common signal and then examined for signs of DP as observed in neighbouring
163 regions.

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

Results

Temperature-growth relations in Old Crow Flats

Correlations between our 23 mean site chronologies and the regional temperature record reveal two primary, and distinctly opposite responses to summer temperature (Table 2). In general, all sites are best correlated with June/July temperatures, but roughly half correlate negatively and the other half positively with June/July temperatures (Table 2). The negatively correlated sites exhibit their strongest correlations with maximum July temperatures of the previous year, while the positively correlated sites exhibit their strongest correlations with minimum June temperatures of the growth year. Sites with the negative/positive temperature response are hereafter referred to as ‘Group 1’/‘Group 2’ sites (Table 2). Only one site, DP21, had a ‘mixed’ temperature relation, albeit weak, and could not be differentiated as either Group 1 or Group 2 (Table 2). Since our primary focus is regionally-significant growth responses to temperature, we focus on the Group 1 and Group 2 chronologies that are of greater regional importance.

Group 1/Group 2 regional growth patterns

The distinction between Group 1 and Group 2 sites is clear in terms of correlation with summer temperature (Table 2). In terms of growth index, there are many notable differences and similarities between Groups 1 and 2 (Fig. 3a, b) which we emphasize by comparing the mean Group 1 and Group 2 chronologies (Fig. 3c; mean Group 1/Group 2 chronologies calculated for all years defined by 2 or more site chronologies). The mean Group 1 and Group 2 chronologies are highly coherent over the period 1620-1800, as demonstrated by a strong running correlation coefficient (Fig. 3c; see SI-Fig. 3a for a magnified comparison).

Both groups began the 19th century at their lowest index values on record and maintained similar growth levels until ~1850 before following an upward trend into the 20th century (Fig. 3c). However, Group 1 began its upward trend slightly before Group 2 (as early as ~1825/1875 for Group 1/Group 2) causing a systematic offset between them until the mid-20th century (Fig. 3c; see SI-Fig. 3b for a magnified comparison). The significance of these differences was tested using a running 2-sample t-test (see SI-Note 1). Significant ($p \leq 0.05$) differences between the Group 1 and Group 2 chronologies are found during 63% of years from 1866-1948. However, despite these growth index differences, their respective linear trends were indistinguishable over the period 1866-1948 (+0.043/+0.042 index values per decade for Group 1/Group 2) leading to a strong, stable running correlation (Fig. 3c; mean $r = 0.82$).

Important differences emerge after the 1930s. Group 1 reaches its highest growth values on record in the 1930s (SI-Fig. 3; 1926-1935 mean index = 1.2) and then declines towards low values in the 1980s. Conversely, Group 2 maintains a positive trend throughout the 20th century and reaches record high growth values at present (SI-Fig. 3; 1998-2007 mean index = 2.0). Effectively, this opposing behaviour equates to a ‘growth-trend divergence’ that is evident in the running correlation which drops below the $p \leq 0.05$ significance level in 1953, reaching its lowest point ($r = -0.02$) in 1966 (Fig. 3c). Group 1 growth trends become positive after the 1980s contributing to a small rise in correlation with Group 2. By 1972, correlations become significant again, but remain low relative to pre-1930 correlations (Fig. 3c).

While the Group 1 and Group 2 chronologies are very different in the ‘low-frequency’ domain since the 1930s, the two groups are coherent at higher-frequencies over all periods. To demonstrate this, we compared high-pass filtered mean Group 1 and Group 2 chronologies (40-year cubic smoothing spline; see Cook and Peters 1981) and found that both groups exhibited the

same high-frequency growth variations over the last 400 years, even after the post-1930s growth trend divergence described above (Fig. 3d).

Group 1/Group 2 vs. temperature

As with their constituent site chronologies, the mean Group 1 and Group 2 chronologies are most strongly correlated with previous-year July maximum and growth-year June minimum temperatures, respectively (Table 2). A visual comparison of the mean group chronologies versus their most closely associated temperature index shows no apparent signs of temperature-growth divergence during recent decades of the 20th century (Fig. 4), as observed in other parts of NWNA. From the 1930s to 1980s, Group 1 growth declined as July maximum temperatures increased, but rebounded from the 1980s to present as July temperatures cooled slightly (Fig. 4). Contrary to July maximum temperatures, June minimum temperatures have increased steadily since the 1930s, a trend that is matched by the Group 2 growth response (Fig. 4).

Larger-scale significance of Group 1/Group 2

To determine if the contrasting Group 1 and Group 2 growth patterns are simply a local phenomenon or reflective of larger-scale growth patterns, a number of long white spruce ring-width chronologies from adjacent parts of NWNA (Fig. 5) were obtained from the International Tree Ring Databank (ITRDB; <http://www.ncdc.noaa.gov/paleo/treering.html>) and other sources (Table 3), and compared to the mean Group 1 and Group 2 chronologies. Only chronologies spanning 1700-1975 were considered in order to assess long-term coherence. Also, a number of spatial criteria were considered to determine which sites were used. To avoid sites with a strong maritime climate (e.g., Gulf of Alaska coastline; see L'Heureux et al. 2004, Serreze and Barry

2005), only sites above 65°N were used. The latitudinal range of sites extends eastward from the west coast of Alaska to 115°W. This spatial range coincides with sites near boreal treeline where tree growth is expected to be temperature-limited. The ITRDB has 26 chronologies matching our spatiotemporal criteria (Fig. 5; Table 3). Further, we use 2 ‘sub-population’ chronologies developed from hundreds of white spruce trees at several sites in the Mackenzie Delta (Pisaric et al. 2007; Fig. 5; Table 3). All 28 ‘NWA chronologies’ were processed using signal-free standardisation as outlined in the Methods section.

The NWA chronologies were considered comparable to Group 1 and Group 2 if they met the following criteria: (1) correlates positively and significantly ($p \leq 0.01$) with both Group 1 and Group 2 from 1850-1930; and (2) correlates positively and significantly with only one of Group 1 or Group 2 from 1930-present. Based on these criteria, 14 NWA chronologies are not similar to Group 1 or Group 2 (Table 3; sites 02-03, 05-13, 15, 18, & 25), 11 of which (sites 02-03, & 05-13) are clustered in a small area (~8 km radius) on the Seward Peninsula, Alaska. Of the remaining 14 NWA chronologies, 7 are similar to Group 1: sites 14, 17, 19-21, & 23-24 (Table 4); hereafter, we refer to these chronologies, including the mean Group 1 chronology, as ‘NWA 1 chronologies’ (Fig. 6a). The other 7 NWA chronologies are similar to Group 2: sites 01, 04, 16, 22, & 26-28 (Table 4). These chronologies, plus the mean Group 2 chronology, are now referred to as ‘NWA 2 chronologies’ (Fig. 6b).

A comparison of the mean NWA 1 and NWA 2 chronologies reveals the same general observation seen with the Old Crow Flats tree-ring data. NWA 1 and NWA 2 had a coherent growth pattern from A.D. 1550 until the mid-20th century, after which the mean chronologies diverge from one another (Fig. 6c). This is reflected in the strong running correlation from 1550-1930 and the subsequent decline to non-significant values by 1957 (Fig. 6c). Unlike with Group

1 and Group 2, Nwana 1 and Nwana 2 have little offset from 1866-1948 (Fig. 6c); only 10% of the years over this period are significantly different based on a running 2-sample t-test (SI-Note 1). A second important difference between the Old Crow Flats and the Nwana chronologies is that the timing of mid-20th-century divergence differs slightly. Nwana 1 sites do not peak until ~1950s, which is roughly 2 decades later than the Group 1 sites in Old Crow Flats. Lastly, as we found in Old Crow Flats, high-frequency growth patterns at the Nwana-scale were coherent over all periods (Fig. 6d) suggesting that growth trend divergence is exclusively a low-frequency phenomenon.

Nwana 1/Nwana 2 vs. temperature

Given their similarity to Group 1/Group 2, it seems plausible that Nwana 1/Nwana 2 may also represent negative/positive responses to 20th-century summer temperature. Indeed, this idea is supported by independent climate-growth analyses for several of the Nwana 1/Nwana 2 site chronologies (Szeicz and MacDonald 1994, 1996; D'Arrigo et al. 2004; Pisaric et al. 2007; Visser et al. 2010). At broader scales, this is supported by correlations between the mean Nwana 1/Nwana 2 chronologies and a composite of gridded mean monthly temperatures for the greater Nwana region derived from the CRUTEM3v dataset (Brohan et al. 2006; see SI-Note 3 for details on the composite).

Overall, Nwana 1 sites were most strongly and negatively correlated with previous-year July temperatures over the last century (1900-2003), as with Group 1 in Old Crow Flats (Table 5). However, Nwana 1's temperature-growth relation was not stable over the 20th century as demonstrated by a split-period analysis (Table 5). Nwana 1 sites shared a significant positive relation with growth-year June/July average temperatures in the early-20th century (1900-1950),

and a significant inverse relation with prior-year July temperatures in the late-20th century (1951-2003) (Table 5). This transition from a positive to negative temperature response is effectively illustrated with a plot of Nwana 1 versus June/July temperatures (Fig. 7). However, because of the gradual, low-frequency nature of the change, and of the temperature-growth relation itself, it is difficult to pinpoint the timing of temperature-growth divergence with any precision. Based on a visual inspection of the data, the early-1960s appear to be a reasonable approximation (Fig. 7). Conversely, Nwana 2 sites responded positively to growth-year June/July temperatures during both the early- and late-20th century (Table 5; Fig. 7).

Long-term temperature-growth relations in Nwana

As both Nwana 1 and Nwana 2 had a positive temperature response before the mid-20th century, it seems likely that the Nwana 1 growth pattern represents an anomalous temperature response. Although, this idea lacks long-term verification from Nwana instrumental data which are largely restricted to the 20th century. Alternatively, a simple comparison against the mean of 6 Northern Hemisphere (NH) temperature reconstructions since A.D. 1300 provides independent verification (Fig. 8a, b). Nwana 1 and Nwana 2 track NH temperatures well before the mid-20th century (Fig. 8b) suggesting their pre-divergence growth was a positive function of temperature. We do note that some site chronologies that constitute the mean Nwana chronologies were also used in some of the NH reconstructions; however, the contribution of the shared chronologies to the overall variance of the NH reconstructions is negligible. In other words, virtually all of the coherence between Nwana 1/Nwana 2 and NH temperatures is from independent data. Finally, as was the case with Nwana 1 versus Nwana temperatures (Fig. 7), Nwana 1 failed to track NH temperatures since the mid-20th century (Fig. 8).

Discussion

Implications and timing of growth trend divergence

Here we have provided evidence of widespread growth trend divergence in NWNA. Our Group 1 and Group 2 chronologies showed similar growth patterns prior to the 1930s, but one of two contrasting patterns since. This growth trend divergence (negative/positive trends) is the main distinction between Group 1 and Group 2 since the 1930s. At larger scales, this growth trend divergence was replicated using 14 white spruce chronologies across NWNA, although the timing of divergence was the 1950s. The coherence of these chronologies before divergence implies that they were responding to a similar regional climatic factor; however, their contrasting behaviour since implies that one of the groups diverged from its former response to climate.

Due to the instrumental data limitations in Old Crow Flats we could not directly assess the implied climate-growth response change for Group 1/Group 2 sites. Yet, the NWNA-scale analysis did reveal that NWNA 1 sites transitioned from a positive to negative temperature response in the mid-20th century, while NWNA 2 sites maintained a stable positive temperature response throughout the 20th century. A comparison against reconstructed Northern Hemisphere temperatures confirmed that all sites, including NWNA 1 sites, shared a positive temperature response prior to 20th-century divergence.

As reported elsewhere (Jacoby and D'Arrigo 1995; Briffa et al. 1998; Cook et al. 2004; D'Arrigo et al. 2004; D'Arrigo et al. 2008), our NWNA-scale results largely support the idea that temperature-growth divergence is a mid-20th-century phenomenon. However, there does appear to be some variability in timing. For example, in Old Crow Flats Group 1/Group 2 sites began to trend in opposite directions as early as the 1930s. Briffa et al. (1998) and Pisaric et al. (2007) also found that some regions may have diverged in the 1930s.

On a more cautious note, there is some indication that Group 1 sites began to separate from Group 2 sites in terms of temperature sensitivity during the 19th century given their offset index values thereafter. At the NWNA-scale, this offset was less apparent, but it was evident. Similarly, sub-population (i.e., intra-site dynamics) studies in Alaska have also found evidence of growth pattern divergence during the 19th century (Wilmking et al. 2004). However, in Old Crow Flats, we remain cautious of labelling this offset a true temperature-growth divergence given that Group 1 and Group 2 have virtually identical linear trends over their period of offset. This parallel behaviour suggests Group 1 and Group 2 responded to climate in the same manner over this period, but that they had slight differences in terms of climatic sensitivity. Further, we cannot discount the possibility it is a detrending artefact due to the difficulty in separating age-related trend from two very different externally-forced 20th-century growth patterns, one peaking in the 1930s and another with a strong late-century increase. Regardless of its nature, the offset does not imply a major change in temperature-growth response. By contrast, 20th-century growth trend divergence implies a temperature-growth response reversal for some sites.

Long-term stability of temperature-growth relations

The divergence problem has been the subject of on-going research for nearly 2 decades (D'Arrigo et al. 2008). From a paleoclimatology perspective, it is important to better understand the spatiotemporal extent of the phenomenon, its causes, and the likelihood it has impacted past growth. In turn, such insight would help determine the extent to which affected chronologies can be used to provide robust estimates of past temperature. Current understandings of DP remain tenuous due to the limited number of datasets affected by divergence and the collinearity of potential contributing environmental factors (D'Arrigo et al. 2008). However, a growing body of

evidence suggests that DP may be caused by the anomalously warm temperatures of the 20th century (Jacoby and D'Arrigo 1995; Barber et al. 2000; D'Arrigo et al. 2004; Wilmking et al. 2004; Pisaric et al. 2007) which likely have been unmatched in the last 2000 years (Kaufman et al. 2009).

The idea that DP is unique to the 20th century does have some empirical support. Based on their comparison of divergent northern site chronologies with a number of other temperature sensitive hemispheric chronologies, Cook et al. (2004) concluded that 20th-century DP is unique in the context of the last 1100 years. Similarly, our comparison of the Nwana chronologies to reconstructed NH temperatures suggests that 20th-century divergence is unique in the context of the last 700 years, at least. In fact, the coherence between the Nwana 1/Nwana 2 chronologies and NH temperatures prior to divergence is remarkable considering the spatial-scale differences (i.e., Nwana vs. NH) and the well documented regional-heterogeneity of NH temperatures over past centuries (D'Arrigo et al. 2006; Mann et al. 2009). One notable exception occurs during the late 1400s when the Nwana chronologies indicate a cool period. This period corresponds with the well known Spörer minimum (Stuiver 1961; Bard et al. 2001), one of the longest and most pronounced solar minima of the past millennium.

Large-scale drivers of divergence

Our results support the notion that DP in Nwana is unique to the 20th century suggesting that DP was caused by a large-scale environmental or climatic change that is also unique to the 20th century. Several large-scale factors have been proposed including: temperature-induced drought stress (Jacoby and D'Arrigo 1995; Barber et al. 2000; Lloyd and Fastie 2002; Wilmking and Juday 2005; McGuire et al. 2010), biological temperature thresholds (D'Arrigo et al. 2004;

Wilmking et al. 2004), snow cover changes (Vaganov et al. 1999), UV-B changes (Briffa et al. 2000), and global dimming (D'Arrigo et al. 2008). Depending on the region, DP may have been caused by one, all, or a combination of these factors. However, negative temperature-growth relations in NRNA have been linked to regional moisture gradients (Wilmking and Juday 2005; Lloyd and Bunn 2007) implying that the divergence may be caused by moisture stress. NRNA is already limited in terms of its annual precipitation budget, and the strong 20th-century warming in this region (ACIA 2005) has likely placed an even greater moisture limitation on these forests.

We also introduce another large-scale factor that may have contributed to DP in NRNA, but has been absent in discussions of DP: Pacific derived moisture. Directional warming was one of the most prominent climate system changes following the mid-19th century in high-latitude regions (Overpeck et al. 1997; Kaufman et al. 2009). However, in NRNA, this warming was accompanied by important moisture changes towards interior NRNA (Anderson et al. 2007, 2011) due to a major atmospheric circulation reorganisation linked to the strength of the Aleutian Low (Fisher et al. 2004; Anderson et al. 2005). This transition led to a relatively dry 20th century in many parts of interior NRNA compared to the last 1200 years (Fisher et al. 2004; Anderson et al. 2011). This reorganisation appears to have been widespread as it coincides with several other large climate system changes in Pacific basin (e.g., Thompson et al. 1986; Mann et al. 2000; Hendy et al. 2002). Although we cannot say which large-scale factor ultimately caused some sites to diverge, several moisture related factors may be involved including temperature-induced drought stress and reduced Pacific-derived moisture since the mid-19th century.

The role of small-scale factors

Because DP is found across such a large portion of NWNA, it is clear that DP was caused by a large-scale forcing. However, not all sites were affected, some of which are found within a few kilometres of sites that were affected (e.g., DP30 vs. DP31, or DP23-26 vs. DP27; SI-Fig. 4). Therefore, it is also clear that site-specific factors determine how each site has been impacted by putative large-scale forcing that caused DP. Ecological factors may be particularly important. For example, Wilmking and Juday (2005) reported that sites with lower tree densities tended to have a greater proportion of trees that responded positively to temperature, presumably related to soil moisture competition.

Organic layer thickness may also be important. At a white spruce stand in the Mackenzie Delta, King (2009) found that trees with a positive temperature-growth response were linked to thicker surficial organic layers that maintain cooler active layers and limit direct evaporative moisture loss from the soil. Negative temperature responses were linked to a thinner organic layer and warmer active layer. A larger-scale study of black spruce in western Quebec, Canada, by Drobyshev et al. (2010) also supports the notion that surficial organic layer thickness may lead to contrasting climate-growth responses (see also Nilsson and Wardle 2005; Turetsky et al. 2010 on the importance of the organic layer in boreal ecosystem functioning). The contribution of such site-level (or intra-site) factors to DP has not been examined in detail, but presents an exciting research opportunity to better understand the spatial complexity of tree growth responses in high-latitude boreal ecosystems.

Concluding remarks

The Divergence Problem has been an ongoing issue for dendroclimatologists working in high-latitude regions for the past 16 years. Because of the limited number of datasets affected by DP, progress in characterizing and advancing current understandings of it has been slow. In this study, we made a sizeable contribution to the high-latitude tree-ring network, adding 23 new white spruce chronologies from Old Crow Flats, northern Yukon. Further, we draw comparisons between these sites and 14 other long white spruce chronologies from across NWN and shed new light on potential causes of DP. Our results suggest that white spruce temperature-growth divergence in NWN largely began in the 1950s, and as early as the 1930s in Old Crow Flats. A long-term comparison of NWN chronologies against reconstructed Northern Hemisphere temperatures provides good independent verification of the idea that these chronologies responded positively to temperature since A.D. 1300, and that temperature-growth divergence in NWN is probably restricted to the 20th century.

The large spatial extent of sites that were impacted by DP suggests a large-scale forcing was the cause. As DP occurs during the warmest period of the last 2000 years (Kaufman et al. 2009) it is likely temperature-induced drought stress is involved. A strengthened Aleutian Low since the mid-19th century (Fisher et al. 2004; Anderson et al. 2005), which led to anomalously dry 20th-century conditions in interior NWN (Anderson et al. 2011), may also be a contributing factor. However, considering the proximity of divergent and non-divergent sites in our network, we speculate that site-specific factors (e.g., organic layer thickness; King 2009; Drobyshev et al. 2010) have an overarching role in determining the susceptibility of individual sites to DP.

Acknowledgements

We thank the Vuntut Gwitch'in First Nation for their hospitality and permission to conduct this research in Old Crow Flats. We also thank: D. Maxwell, P. deMontigny, A. Burn, P. Miller, E. Tizya-Tramm, T. Green for field/lab support; G. King, S. St. George for comments on earlier drafts of this paper; E. Cook for a 'signal-free' version of ARSTAN; the contributors of ITRDB data used here; and two anonymous reviewers whose comments helped improve this manuscript considerably. This research was supported by: a NSERC Discovery Grant, NSERC Northern Supplement and Government of Canada IPY grant to M. Pisaric; and a NSERC Graduate Scholarship and Northern Scientific Training Program grant to T. Porter.

References

- ACIA (2005) *Arctic Climate Impact Assessment*. Cambridge University Press, New York, NY, USA, 1042 pp.
- Anderson L, Abbott MB, Finney BP, Burns SJ (2005) Regional atmospheric circulation change in the North Pacific during the Holocene inferred from lacustrine carbonate oxygen isotopes, Yukon Territory, Canada. *Quaternary Research*, **64**, 21-35.
- Anderson L, Abbott MB, Finney BP, Burns SJ (2007) Late Holocene moisture balance variability in the southwest Yukon Territory, Canada. *Quaternary Science Reviews*, **26**, 130-141.
- Anderson L, Finney BP, Shapley MD (2011) Lake carbonate- $\delta^{18}\text{O}$ records from the Yukon Territory, Canada: Little Ice Age moisture variability and patterns. *Quaternary Science Reviews*, **30**, 887-898.
- Barber VA, Juday GP, Finney BP (2000) Reduced growth of Alaskan white spruce in the twentieth century from temperature-induced drought stress. *Nature*, **405**, 668-673.
- Bard E, Raisbeck G, Yiou F, Jouzel J (2000) Solar irradiance during the last 1200 years based on cosmogenic nuclides. *Tellus*, **52B**, 985-992.

505 Briffa KR (2000) Annual climate variability in the Holocene: interpreting the message of ancient
506 trees. *Quaternary Science Reviews*, **19**, 87-105.

507

508 Briffa KR, Bartholin TS, Eckstein D, Jones PD, Karlén W, Schweingruber FH, Zetterberg
509 P (1990) A 1,400-year tree-ring record of summer temperatures in Fennoscandia. *Nature*,
510 **346**, 434-439.

511

512 Briffa KR, Osborn TJ, Schweingruber FH (2004) Large-scale temperature inferences from tree
513 rings: a review. *Global and Planetary Change*, **40**, 11-26.

514

515 Briffa KR, Schweingruber FH, Jones PD, Osborn TJ, Shiyatov SG, Vaganov EA (1998) Reduced
516 sensitivity of recent tree-growth to temperature at high northern latitudes. *Nature*, **391**, 678-682.

517

518 Brohan P, Kennedy JJ, Haris I, Tett SFB, Jones PD (2006) Uncertainty estimates in regional and
519 global observed temperature changes: A new data set from 1850. *Journal of Geophysical*
520 *Research*, **111**, doi:10.1029/2005JD006548.

521

522 Cook ER (1985) *A Time Series Analysis Approach to Tree Ring Standardization*. University of
523 Arizona, Ph.D. dissertation in the School of Renewable Natural Resources, Tuscan, Arizona,
524 USA, 171 pp.

525

526 Cook ER, Esper J, D'Arrigo RD (2004) Extra-tropical Northern Hemisphere land temperature
527 variability over the past 1000 years. *Quaternary Science Reviews*, **23**, 2063-2074.

528

529 Cook ER, Peters K (1981) The smoothing spline: a new approach to standardizing forest interior
 530 tree-ring width series for dendroclimatic studies. *Tree-Ring Bulletin*, **41**, 45-53.

531

532 D'Arrigo R, Kaufmann RK, Davi N, Jacoby GC, Laskowski C, Myneni RB, Cherubini P (2004)
 533 Thresholds for warming-induced growth decline at elevational tree line in the Yukon Territory,
 534 Canada. *Global Biogeochemical Cycles*, **18**, GB3021, doi:10.1029/2004GB002249.

535

536 D'Arrigo R, Wilson R, Jacoby G (2006) On the long-term context for late twentieth century
 537 warming. *Journal of Geophysical Research*, **111**, D03103, doi:10.1029/2005JD006352.

538

539 D'Arrigo R, Wilson R, Liepert B, Cherubini P (2008) On the 'Divergence Problem' in Northern
 540 Forests: A review of the tree-ring evidence and possible causes. *Global and Planetary*
 541 *Change*, **60**, 289-305.

542

543 Drobyshev I, Simard M, Bergeron Y, Hofgaard A (2010) Does soil organic layer thickness affect
 544 climate–growth relationships in the black spruce boreal ecosystem? *Ecosystems*, **13**, 556-574.

545

546 Dyke AS, Andrews JT, Clark PU, England JH, Miller GH, Shaw J, Veillette JJ (2002) The
 547 Laurentide and Innuitian ice sheets during the Last Glacial Maximum. *Quaternary Science*
 548 *Reviews*, **21**, 9-31.

549

Dyke LD (2000) Climate of the Mackenzie River valley. In: *The Physical Environment of the Mackenzie Valley, Northwest Territories: A Baseline for the Assessment of Environmental Change* (eds Dyke LD, Brooks GR), pp. 21-30. Geological Survey of Canada, Ottawa, Ontario.

Esper J, Cook ER, Schweingruber FH (2002) Low-frequency signals in long tree-ring chronologies for reconstructing past temperature variability. *Science*, **295**, 2250-2253.

Fisher DA, Wake C, Kreutz K *et al.* (2004) Stable isotope records from Mount Logan, Eclipse ice cores and nearby Jellybean Lake. Water cycle of the North Pacific over 2000 years and over five vertical kilometers: Sudden shifts and tropical teleconnections. *Géographie Physique et Quaternaire*, **58**, 9033-9048.

Frank D, Esper J, Cook ER (2007) Adjustment for proxy number and coherence in a large-scale temperature reconstruction. *Journal of Geophysical Research*, **34**, L16709, doi:10.1029/2007GL030571.

Fritts HC, Moismann JE, Bortorff CP (1969) A revised computer program for standardizing tree-ring series. *Tree-Ring Bulletin*, **29**, 15-20.

Gostev M, Wiles G, D'Arrigo R, Jacoby G, Khomentovsky P (1996) Early summer temperatures since 1670 A.D. for Central Kamchatka reconstructed based on a Siberian larch tree-ring width chronology. *Canadian Journal of Forest Research*, **26**, 2048-2052.

573 Heginbottom JA, Dubreuil MA, Harker PA (1995) Canada - Permafrost. In: *National Atlas of*
 574 *Canada, 5th Edition. Plate 2.1, scale 1:7,500,000 (MCR 4177)* Natural Resources Canada,
 575 Ottawa, Canada.
 576
 577 Hendy EJ, Gagen MK, Alibert CA, McColloch MT, Lough JM, Isdale PJ (2002) Abrupt
 578 decrease in tropical Pacific sea surface salinity at the end of the Little Ice Age. *Science*, **295**,
 579 1511-1514.
 580
 581 Holmes RL (1983) Computer-assisted quality control in tree-ring dating and measurement. *Tree-*
 582 *Ring Bulletin*, **43**, 69-78.
 583
 584 Hughes MK (2011) Dendroclimatology in High-Resolution Paleoclimatology. In:
 585 *Dendroclimatology: Progress and Prospects* (eds Hughes MK, Swetnam TW, Diaz HF), pp. 17-
 586 34. Springer, New York.
 587
 588 Hughes OL (1972) Surficial geology of northern Yukon Territory and northwestern District of
 589 Mackenzie, Northwest Territories. *Geological Survey of Canada Paper 69-36*, 11.
 590
 591 Hughes OL, Rampton VN (1971) Northern Yukon Territory, and Northwestern District of
 592 Mackenzie. *Geological Survey of Canada, Surficial Geology Map 1319A (1:500,000)*.
 593
 594 Jacoby GC, D'Arrigo RD (1995) Tree ring width and density evidence of climatic and potential
 595 forest change in Alaska. *Global Biogeochemical Cycles*, **9**, 227-234.

596

597 Jacoby GC, Lovelius NV, Shumilov OI, Raspopov OM, Karbainov JM, Frank DC (2000) Long-
598 term temperature trends and tree growth in the Taymir Region of Northern Siberia. *Quaternary*
599 *Research*, **53**, 312-318.

600

601 Jansen E, Overpeck J, Briffa KR *et al.* (2007) Palaeoclimate. In: *Climate Change 2007: The*
602 *Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the*
603 *Intergovernmental Panel on Climate Change* (eds Solomon S, Qin D, Manning M, Chen Z,
604 Marquis M, Averyt KB, Tignor M, Miller HL), Cambridge University Press, Cambridge, United
605 Kingdom and New York, NY, USA.

606

607 Jones PD, Briffa KR, Barnett TP, Tett SFB (1998) High-resolution palaeoclimatic records for the
608 last millennium: interpretation, integration and comparison with General Circulation Model
609 control-run temperatures. *The Holocene*, **8**, 455-471.

610

611 Kaufman DS, Schneider DP, McKay NP *et al.* (2009) Recent warming reverses long-term Arctic
612 cooling. *Science*, **325**, 1236-1239.

613

614 King GM (2009) *Factors influencing the growth of white spruce (Picea glauca) in the Mackenzie*
615 *Delta, NT*. Carleton University, Unpublished M.Sc. thesis in Geography, Ottawa, Canada, 159
616 pp.

617

618 Labrecque S, Lacelle D, Duguay CR, Lauriol B, Hawkings J (2009) Contemporary (1951-2001)
619 evolution of lakes in the Old Crow Basin, northern Yukon, Canada: remote sensing, numerical
620 modelling, and stable isotope analysis. *Arctic*, **62**, 225-238.

621

622 Lauriol B, Duguay CR, Riel A (2002) Response of the Porcupine and Old Crow rivers in
623 northern Yukon, Canada, to Holocene climatic change. *The Holocene*, **12**, 27-34.

624

625 L'Heureux ML, Mann ME, Cook BI, Gleason BE, Vose RS (2004) Atmospheric circulation
626 influences on seasonal precipitation patterns in Alaska during the latter 20th century. *Journal of*
627 *Geophysical Research*, **109**, D06106, doi:10.1029/2003JD003845.

628

629 Lloyd AH, Bunn AG (2007) Responses of the circumpolar boreal forest to 20th century climate
630 variability. *Environmental Research Letters*, **2**, doi:10.1088/1748-9326/2/4/045013.

631

632 Lloyd AH, Fastie CL (2002) Spatial and temporal variability in the growth and climate response
633 of treeline trees in Alaska. *Climatic Change*, **52**, 481-509.

634

635 Mann ME, Bradley RS, Hughes MK (2000) Long-term variability in El-Niño-Southern
636 Oscillation and associated teleconnections. In: *El Niño and the Southern Oscillation* (eds Diaz
637 HF, Markgraf V), pp. 357-412. Cambridge University Press, New York, USA.

638

639 Mann ME, Zhang Z, Rutherford S *et al.* (2009) Global signatures and dynamical origins of the
640 Little Ice Age and Medieval Climate Anomaly. *Science*, **326**, 1256-1260.

641

642 McGuire AD, Ruess RW, Lloyd A, Yarie J, Klein JS, Juday GP (2010) Vulnerability of white

643 spruce tree growth in interior Alaska in response to climate variability: dendrochronological,

644 demographic, and experimental perspectives. *Canadian Journal of Forest Research*, **40**, 1197-

645 1209.

646

647 Melvin TM, Briffa KR (2008) A "signal-free" approach to dendroclimatic standardisation.

648 *Dendrochronologia*, **26**, 71-86.

649

650 Morlan RE (1980) Taphonomy and archaeology in the upper Pleistocene of the northern Yukon

651 Territory: a glimpse of the peopling of the New World. *Archaeological Survey of Canada, Paper*

652 *no. 94*, 398.

653

654 Nilsson M-C, Wardle DA (2005) Understorey vegetation as a forest ecosystem driver: evidence

655 from the northern Swedish boreal forest. *Frontiers in Ecology and the Environment*, **3**, 421-428.

656

657 Ovenden J, Brassard GR (1989) Wetland vegetation near Old Crow, northern Yukon. *Canadian*

658 *Journal of Botany*, **67**, 954-960.

659

660 Overpeck J, Hughen KA, Hardy D *et al.* (1997) Arctic environmental change of the last four

661 centuries. *Science*, **278**, 1251-1256.

662

663 Pisaric MFJ, Carey SK, Kokelj SV, Youngblut D (2007) Anomalous 20th century tree growth,
 664 Mackenzie Delta, Northwest Territories, Canada. *Geophysical Research Letters*, **34**, L05714,
 665 doi:10.1029/2006GL029139.
 666
 667 Serreze MC, Barry RG (2005) Physical characteristics and basic climatic features. In: *The Arctic*
 668 *Climate System* pp. 17-54. Cambridge University Press, Cambridge, UK.
 669
 670 Stuiver M (1961) Variations in radiocarbon concentration and sunspot activity. *Journal of*
 671 *Geophysical Research*, **66**, 273-276.
 672
 673 Speer JH (2010) *Fundamentals of Tree-Ring Research*. University of Arizona Press, Tucson,
 674 Arizona, 252 pp.
 675
 676 Szeicz JM, MacDonald GM (1996) A 930-year ring-width chronology from moisture-sensitive
 677 white spruce (*Picea glauca* Moench) in northwestern Canada. *The Holocene*, **6**, 345-351.
 678
 679 Szeicz JM, MacDonald GM (1995) Dendroclimatic reconstruction of summer temperatures in
 680 northwestern Canada since A.D. 1638 based on age-dependent modelling. *Quaternary*
 681 *Research*, **44**, 257-266.
 682
 683 Szeicz JM, MacDonald GM (1994) Age-dependent tree-ring growth responses of subarctic white
 684 spruce to climate. *Canadian Journal of Forest Research*, **24**, 120-132.
 685

686 Thompson LG, Mosely-Thompson E, Dansgaard W, Grootes PM (1986) The Little Ice Age as
687 recorded in the stratigraphy of the tropical Quelccaya Ice Cap. *Science*, **234**, 361-354.
688

689 Thorson RM, Dixon EJ (1983) Alluvial history of the Porcupine River, Alaska: role of glacial-
690 lake overflow from northwest Canada. *Geological Society of America*, **94**, 576-589.
691

692 Turetsky MR, Mack MC, Hollingsworth TN, Harden JW (2010) The role of mosses in
693 ecosystem succession and function in Alaska's boreal forest. *Canadian Journal of Forest*
694 *Research*, **40**, 1237-1264.
695

696 Vaganov EA, Hughes MK, Kirdyanov AV, Schweingruber FH, Silkin PP (1999) Influence of
697 snowfall and melt timing on tree growth in subarctic Eurasia. *Nature*, **400**, 149-151.
698

699 Visser H, Büntgen U, D'Arrigo R, Petersen AC (2010) Detecting instabilities in tree-ring proxy
700 calibration. *Climate of the Past Discussions*, **6**, 225-255.
701

702 Wahl ER, Ammann CM (2007) Robustness of the Mann, Bradley, Hughes reconstruction of
703 Northern Hemisphere surface temperatures: Examination of criticisms based on the nature and
704 processing of proxy climate evidence. *Climatic Change*, **85**, 33-69.
705

706 Walker DA, Raynolds MK, Daniëls FJA *et al.* (2005) The Circumpolar Arctic vegetation
707 map. *Journal of Vegetation Science*, **16**, 267-282.
708

Wilmking M, Juday GP (2005) Longitudinal variation of radial growth at Alaska's northern treeline – recent changes and possible scenarios for the 21st century. *Global and Planetary Change*, **47**, 282-300.

Wilmking M, Juday GP, Barber VA, Zald HSJ (2004) Recent climate warming forces contrasting growth responses of white spruce at treeline in Alaska through temperature thresholds. *Global Change Biology*, **10**, 1724-1736.

Wilson R, D'Arrigo R, Buckley B *et al.* (2007) A matter of divergence: tracking recent warming at hemispheric scales using tree ring data. *Journal of Geophysical Research*, **112**, D17103, doi:10.1029/2006JD008318.

Youngblut D, Luckman B (2008) Maximum June-July temperatures in the southwest Yukon over the last 300 years reconstructed from tree rings. *Dendrochronologia*, **25**, 153-166.

Table 1. White spruce sites sampled in Old Crow Flats and corresponding tree-ring chronology information.

Site code	Lat. (°N)	Long. (°W)	Elev. (m)	No. series/ trees	First year ≥ 1 series	First year ≥ 4 series	Last year	Mean series length (yrs)
DP21	67.53	139.93	251	67/35	1657	1727	2007	199.7
DP23	67.50	139.96	249	46/24	1711	1716	2007	216.1
DP24	67.51	139.96	249	42/22	1735	1746	2006	208.7
DP25	67.52	139.99	251	43/22	1759	1778	2006	163.7
DP26	67.52	140.02	243	36/20	1728	1739	2006	200.5
DP27	67.50	140.06	243	57/29	1608	1738	2007	150.0
DP30	67.48	140.33	244	71/36	1552	1555	2007	247.2
DP31	67.48	140.26	245	80/41	1617	1620	2007	240.6
JC1	67.95	139.14	286	73/37	1744	1808	2007	157.0
OC2	67.59	139.75	258	67/35	1691	1742	2007	149.7
OC9 _a	67.70	139.81	267	54/27	1618	1620	1846	183.8
OC9 _b	67.70	139.81	267	50/26	1874	1881	2007	109.2
OC10	67.72	139.82	259	61/32	1719	1727	2006	111.7
OC50	68.13	139.93	282	45/25	1668	1768	2007	184.8
OC51	68.07	139.78	272	84/43	1680	1714	2007	186.6
OC52	68.05	139.59	269	59/31	1749	1753	2007	158.9
OC53	67.86	139.79	265	82/44	1648	1649	2007	189.3
OC54	68.22	140.09	292	42/21	1615	1627	2007	140.3
PC17	67.55	139.41	251	94/49	1803	1811	2006	159.2
PC18	67.53	139.32	251	60/33	1686	1711	2006	203.4
SC1	67.76	140.52	647	56/29	1537	1553	2007	214.4
TH1	68.33	140.75	339	97/50	1650	1661	2007	225.4
TM1	68.34	139.72	315	94/48	1631	1648	2007	223.6
TM2	68.17	139.78	305	77/41	1522	1547	2007	217.4

N.B. – OC9_a and OC9_b belong to the same site, but do not overlap because of a stand replacing fire that occurred just prior to circa A.D. 1850; OC9_a represents a population of trees that died before or during the fire and OC9_b represents the post-burn population.

Table 2. Correlations between the 23 Old Crow Flats site chronologies and previous-year June/July maximum temperatures (left) and growth-year June/July minimum temperatures (right) (see SI-Table 1 for May-August correlations); only correlations significant at $p \leq 0.05$ (two-tailed) are presented; sites are classed as Group 1/Group 2 if negatively/positively correlated with June/July temperatures.

	Previous year (t-1) max. temperatures		Growth year (t) min. temperatures	
	June	July	June	July
Mixed				
DP21	-	-0.29	0.31	-
Group 1				
DP27	-0.45	-0.58	-0.28	-0.41
DP30	-0.28	-0.43	-	-
OC9	-0.50	-0.54	-0.42	-0.41
OC10	-0.24	-0.48	-	-
OC50	-0.42	-0.40	-	-
OC52	-0.41	-0.57	-	-0.25
OC53	-0.25	-0.48	-	-
OC54	-0.41	-0.58	-0.31	-0.35
PC18	-0.35	-0.54	-	-0.23
TM1	-	-0.41	-	-
TM2	-0.41	-0.62	-	-
Group 2				
DP23	-	-	0.41	-
DP24	0.35	-	0.54	0.43
DP25	0.35	0.24	0.54	0.48
DP26	0.33	-	0.52	0.38
DP31	-	-	0.48	0.27
JC1	-	-	0.39	-
OC2	-	-	0.44	0.26
OC51	0.24	-	0.57	0.45
PC17	0.38	0.25	0.54	0.41
SC1	0.34	-	0.53	0.43
TH1	0.24	-	0.58	0.42
Regional means				
Group 1	-0.42	-0.61	-	-0.24
Group 2	0.28	-	0.57	0.40

Table 3. List of 300+ year white spruce ring-width chronologies from upper NRNA (65-70°N, 115-170°W) that we compared to the mean Group 1 (G1)/Group 2 (G2) chronologies from Old Crow Flats. Correlations with G1/G2 were calculated for all years of overlapping data for the periods 1850-1930 and 1930-2003; only positive correlations significant at $p \leq 0.01$ (one-tailed) are provided. Site # corresponds to the map of NRNA sites (Figure 5). Mean chronologies for each NRNA site were calculated from raw ring-width data using signal-free standardisation as outlined in the Methods section. All ring-width files were downloaded from ITRDB (accessed Sept. 2009) unless stated otherwise.

Site #	Site name, ITRDB code	Temporal coverage	Correlation with G1/G2			
			1850-1930		1930-2003	
			G1	G2	G1	G2
01	Almond Butter Lower, AK057	1607-2002	0.26	0.32	-	0.58
02	Almond Butter Upper, AK058	1406-2002	-	-	-	-
03	Alpine View, AK059	1542-2002	-	-	-	-
04	Burnt Over, AK060	1621-2002	0.27	0.29	-	0.46
05	Bye Rosanne, AK061	1575-2002	-	-	0.47	-
06	Death Valley, AK062	1358-2002	-	-	-	-
07	Echo Slope, AK063	1590-2002	-	-	-	-
08	Frost Valley, AK064	1611-2002	-	-	0.31	-
09	Gordon's Cat, (AK065	1400-2002	-	-	-	-
10	Hey Bear, AK066	1533-2002	-	-	-	-
11	Hey Bear Upper, AK067	1383-2002	-	-	-	-
12	Mt. Mole, AK068	1550-2002	-	-	-	-
13	Windy Ridge, AK070	1556-2002	-	-	-	-
14	^a Four-Twelve with Revisit, AK031	1515-1990	0.55	0.51	0.63	-
15	^b Kobuk/Noatak, AK046	978-1992	-	-	0.36	0.29
16	^a Arrigetch, AK032	1585-1990	0.41	0.39	-	0.50
17	Sheenjek River and Flats, AK033	1296-1979	0.41	0.37	0.51	-
18	^c Firth River, AK047	1676-2002	0.71	0.71	-	-
19	Spruce Creek, CANA029	1570-1977	0.34	0.26	0.48	-
20	^d Richardson Mountain, CANA121	1547-1992	0.64	0.65	0.59	-
21	^a Twisted Tree Heartrot Hill, CANA157	1459-1999	0.65	0.66	0.69	-
22	^e MDEC Positive Responders	1516-2003	0.77	0.82	-	0.86
23	^e MDEC Negative Responders	1501-2003	0.72	0.73	0.82	-
24	^f Campbell Dolomite Upland, CANA138	1060-1992	0.56	0.45	0.67	-
25	Mackenzie Mountains, CANA156	1509-1984	-	-	0.69	-
26	Franklin Mountains, CANA154	1621-1984	0.39	0.34	-	0.45
27	^d Discovery Ridge, CANA117	1429-1991	0.35	0.26	-	0.63
28	^a Coppermine River, CANA153	1046-2003	0.53	0.59	-	0.29

Data contributors by site #: Church and Fritts (19); D'Arrigo, Mashig, Frank, Wilson, and Jacoby (01-13); Jacoby, D'Arrigo, and Buckley (14, 16-17, 21, 25-26, and 28); King and Graumlich (15); Pisaric (22-23); Szeicz and MacDonald (27); Szeicz, MacDonald, and Lundberg (20 and 24); Wilmking (18).

^aUpdated versions of sites 14, 16, 21, and 28 (not yet available on ITRDB) were provided by R. D'Arrigo (pers. comm.).

^bKobuk/Noatak is a regional (~24,000 km² area) composite developed from hundreds of trees in the Kobuk and Noatak River basins.

^cCross-dating verification using COFECHA indicated that two of the Firth River series (BRFR49 & 88) should be adjusted -1 year. The possibility these series were misaligned was confirmed by M. Wilmking (pers. comm., Sept. 2010). Adjusting these series increased their correlation with the BRFR49/88 master chronologies from 0.03/0.11 to 0.40/0.41.

^dAs per Szeicz and MacDonald (1995), we used only 200+/100+ year series for Discovery Ridge/Richardson Mountain.

^fMDEC (Mackenzie Delta East Channel) represents several sites in the Mackenzie Delta whose individual series were pooled into ‘positive- or negative-responder’ chronologies (Pisaric et al. 2007). The Positive and Negative Responder chronologies used here are modified versions of the originals used by Pisaric et al. (2007); see SI-Note 2 for more details.

^fA subset (11 trees) of samples collected by Szeicz, MacDonald, and Lundberg from Campbell Dolomite Upland was developed into a separate ‘Campbell Dolomite Upland B’ chronology by F. Schweingruber and is available on the ITRDB. Here we only use the Szeicz, MacDonald, and Lundberg version because of its much larger sample depth.

Table 4. NWNA chronologies that meet the following two similarity criteria: (1) correlates positively and significantly ($p \leq 0.01$) with both Group 1 and Group 2 from 1850-1930; and (2) correlates positively and significantly with only one of Group 1 or Group 2 from 1930-present. Chronologies that meet these criteria and are most closely associated Group 1/Group 2 during 1930-2003 are classed as NWNA 1/NWNA 2 chronologies.

NWNA 1 chronologies

Campbell Dolomite Upland
 Four-Twelve with Revisit
 MDEC Negative Responders
 Richardson Mountain
 Sheenjek River and Flats
 Spruce Creek
 Twisted Tree Heartrot Hill

NWNA 2 chronologies

Almond Butter Lower
 Arrigetch
 Burnt Over
 Coppermine River
 Discovery Ridge
 Franklin Mountains
 MDEC Positive Responders

Table 5. Correlations between the mean NWNA 1/NWNA 2 chronologies and regional mean monthly CRUTEM3v temperatures (Brohan et al. 2006; see SI-Note 3); correlations for May-August of the prior and current growth years are provided for three periods: 1900-2003, 1900-1950, and 1951-2003; all correlations are significant at $p \leq 0.05$ (two-tailed); underlined coefficients are significant at $p \leq 0.01$.

		NWNA1			NWNA2		
		1900-2003	1900-1950	1951-2003	1900-2003	1900-1950	1951-2003
Year t-1	May	-	-	-	<u>0.34</u>	-	-
	Jun	-	-	-	<u>0.41</u>	-	<u>0.40</u>
	Jul	<u>-0.36</u>	-	<u>-0.40</u>	<u>0.29</u>	-	-
	Aug	-	-	-	0.21	-	-
Year t	May	-0.20	-	<u>-0.30</u>	0.20	-	-
	Jun	-	<u>0.53</u>	-	<u>0.54</u>	<u>0.47</u>	<u>0.46</u>
	Jul	-	<u>0.39</u>	-	<u>0.54</u>	<u>0.41</u>	<u>0.36</u>
	Aug	-	0.20	-	0.21	-	-

Figure 1 – (a) White spruce sites sampled in Old Crow Flats (site codes indicated, see Table 1). Major lakes and channels are shaded grey. (b) Climate stations at Fairbanks (FAI), Fort Yukon (FTY), Inuvik (INU), Aklavik (AKL), and Fort McPherson (FTM). (c) Large-scale context of Old Crow Flats. Boreal treeline was delineated from the Circumpolar Arctic Vegetation Map dataset (Walker et al. 2005).

Figure 2 – Monthly normals (1971-2000) for minimum, mean, and maximum temperatures (lines), and total precipitation (bars) at Old Crow (www.climate.weatheroffice.ec.gc.ca).

Figure 3 – (a/b) Group 1/Group 2 site chronologies (grey lines); all site chronologies are defined by no less than 4 series (≥ 2 trees). The mean Group 1/Group 2 chronologies (black lines) were calculated using a robust bi-weight mean for all years defined by 2 or more site chronologies. Sample depth curves are indicated above each plot. (c) A comparison of the mean Group 1 (1555-2007) and Group 2 (1620-2007) chronologies. The running 51-year correlation (above plot) indicates coherence between the chronologies (dashed line is $p \leq 0.05$ level). (d) High-pass filtered (40-year cubic smoothing spline with a 50% frequency cut-off; Cook and Peters 1981) comparison of the mean Group 1 and Group 2 chronologies; the high-pass series were smoothed with a 3-year cubic-smoothing spline for ease of comparison.

885 Figure 4 – Comparisons between mean Group 1/Group 2 (black lines) and previous-year July
886 maximum/growth-year June minimum temperatures (grey lines); correlations are significant at p
887 ≤ 0.001 .

888

889 Figure 5 – Locations of all Nwana site chronologies compared to the mean Group 1/Group 2
890 chronologies (site numbers indicated, see Table 3).

891

892 Figure 6 – Same as Figure 3, but for Nwana 1/Nwana 2 chronologies (Table 4; including the
893 mean Group 1/Group 2 chronologies).

894

895 Figure 7 – Comparison between the mean Nwana 1/Nwana 2 (dotted/grey line) chronologies
896 and a regional composite of June/July temperatures for Nwana (solid black line; see SI-Note 3).

897

898 Figure 8 – (a) Northern Hemisphere (NH) temperature reconstructions by Jones et al. (1998),
899 Briffa (2000), Esper et al. (2002), D'Arrigo et al. (2006), Wahl and Ammann (2007), and Wilson
900 et al. (2007) (grey lines); mean reconstruction (black line). The reconstructions were expressed
901 as z-scores relative to the common period of overlap (1750-1980). (b) Comparison of the mean
902 NH temperature reconstruction (thick black line; smoothed with 15-year cubic smoothing spline,
903 Cook and Peters 1981) against the mean Nwana 1/Nwana 2 chronologies (dotted/grey line).
904 The mean Nwana chronologies are defined by a minimum of 2 site chronologies.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

960

961 Fig. 8

Supporting Information

SI-Figure 1. Standardized ring-width indices (grey lines) and mean site chronologies (black lines) for each site in Old Crow Flats based on signal-free standardisation (Melvin and Briffa 2008) using conservative ‘negative exponential’ or ‘negative-to-zero slope linear’ curve fits (Fritts et al. 1969). Sample depth (red lines) indicates the number of series defining the mean chronology. The mean chronologies were calculated with a robust bi-weight mean (Cook 1985); more details on each chronology are provided in Table 1.

Supporting Information

SI-Figure 2. A comparison of each mean site chronology produced using ‘signal-free’ (black lines) and non-signal-free (red lines) methods. Inter-series differences can be considered the result of ‘trend distortion’ (Melvin and Briffa 2008). Due to differences in non-age-related growth (i.e., forced by climate, disturbance, etc.) between sites, trend distortion effects are more pronounced in some chronologies (e.g., JC1, OC9, OC54, SC1, and TH1) than in others (e.g., DP26, OC50, OC52, TM1, and TM2).

Supporting Information

SI-Figure 3. Magnified comparison of the mean Group 1 and Group 2 chronologies during (a) 1600-1800 and (b) 1800-2000. Smoothed chronologies were calculated using a 15-year cubic smoothing spline with a 50% frequency cut-off (Cook and Peters 1981).

SI-Figure 4. Old Crow Flats sites: Group 1 (negative temperature response), Group 2 (positive temperature response), and Mixed (mixed negative/positive temperature response).

Supporting Information

SI-Table 1. Correlations between the 23 Old Crow Flats ‘signal-free’ site chronologies and minimum/maximum temperatures from May-August of the growth year (t) and previous growth year (t - 1); only correlations significant at $p \leq 0.05$ (two-tailed) are presented.

	Minimum temperatures								Maximum temperatures							
	Previous growth year				Growth year				Previous growth year				Growth year			
	May	Jun	Jul	Aug	May	Jun	Jul	Aug	May	Jun	Jul	Aug	May	Jun	Jul	Aug
DP21						0.31					-0.29					
DP23		0.32				0.41										
DP24	0.29	0.45	0.38			0.54	0.43	0.27	0.30	0.35				0.41	0.28	
DP25	0.33	0.46	0.42	0.23	0.25	0.54	0.48	0.31	0.31	0.35	0.24			0.38	0.29	
DP26	0.27	0.46	0.36		0.24	0.52	0.38	0.24	0.24	0.33				0.34		
DP27		-0.38	-0.43		-0.35	-0.28	-0.41	-0.23	-0.27	-0.45	-0.58	-0.40	-0.43	-0.39	-0.41	-0.32
DP30										-0.28	-0.43	-0.33	-0.27		-0.34	-0.34
DP31	0.29			0.27		0.48	0.27		0.28					0.31		
JC1	0.27	0.36				0.57	0.45			0.24				0.34		
OC2	0.24	0.26				0.39										
OC9	-0.33	-0.53	-0.52	-0.35	-0.34	-0.42	-0.41	-0.24	-0.34	-0.50	-0.54	-0.41	-0.36	-0.46	-0.42	-0.34
OC10										-0.24	-0.48	-0.24			-0.27	-0.24
OC50		-0.34	-0.26		-0.26					-0.42	-0.40		-0.31		-0.24	
OC51	0.26					0.44	0.26									
OC52		-0.36	-0.47		-0.36		-0.25	-0.23		-0.41	-0.57	-0.34	-0.41	-0.25	-0.35	-0.34
OC53										-0.25	-0.48	-0.29	-0.25		-0.34	-0.39
OC54	-0.25	-0.42	-0.53	-0.40	-0.36	-0.31	-0.35	-0.30	-0.28	-0.41	-0.58	-0.47	-0.44	-0.40	-0.43	-0.45
PC17	0.30	0.46	0.36		0.23	0.54	0.41		0.32	0.38	0.25			0.42	0.29	
PC18		-0.24	-0.34		-0.24		-0.23			-0.35	-0.54	-0.34	-0.32	-0.24	-0.38	-0.37
SC1	0.36	0.45	0.28	0.27		0.53	0.43		0.33	0.34				0.37	0.25	
TH1	0.35	0.41	0.26	0.27		0.58	0.42		0.26	0.24				0.35		
TM1			-0.38		-0.35						-0.41		-0.37			
TM2		-0.31	-0.42		-0.32					-0.41	-0.62	-0.30	-0.39		-0.35	-0.29

Supporting Information

SI-Note 1. A running 2-sample t-test was used to test the null hypothesis that the Group 1 and 2 site chronologies are derived from the same normal distribution with equal means and variance ($p \leq 0.05$). The null hypothesis was tested for each year that both groups contained four or more site chronologies. The t-test result was calculated using the 'ttest2' function (Statistics Toolbox) in Matlab 7.4.

SI-Note 2. The MDEC negative-responder (neg) and positive-responder (pos) used here are modified versions of the “negative- and positive-responder” chronologies by Pisaric et al. (2007). The main difference is that the modified MDEC neg and MDEC pos chronologies do not contain Campbell Dolomite Upland series (Szeicz and MacDonald 1996). CDU series were excluded from MDEC neg and MDEC pos to ensure our Group 1/Group 2 correlations with CDU, MDEC neg, and MDEC pos would be independent of each other.

SI-Note 3. The mean NWNA 1/NWNA 2 chronologies were compared to a regional average of CRUTEM3v gridded temperatures (Brohan et al. 2006). The regional average includes 22 ($5^\circ \times 5^\circ$) grid cells bounded by $60\text{-}70^\circ\text{N}$ and $170\text{-}115^\circ\text{W}$; only 2 grid cells ($65\text{-}70^\circ\text{N}/160\text{-}155^\circ\text{W}$ and $65\text{-}70^\circ\text{N}/150\text{-}145^\circ\text{W}$) did not contain any data. Temperature data before 1900 were not used as spatial coverage is limited. The number of grid cells with data for the year 1900 is 5; the number of grid cells increases steadily to more than 10 by the early-1920's, more than 15 by the early-1940's, and a high of 20 by 1959.