Canadian Journal of Animal Science
Revue canadienne de science animale

Effects of pork differentiation strategies in Canada on pig performance and carcass characteristics

Journal:	<i>Canadian Journal of Animal Science</i>
Manuscript ID	CJAS-2015-0197.R3
Manuscript Type:	Article
Date Submitted by the Author:	13-Apr-2016
Complete List of Authors:	<p>Zhang, Hongbo; Inner Mongolia Agricultural University, College of Food Science and Engineering; Agriculture and Agri-Food Canada Lacombe Research Centre</p> <p>Aalhus, J.; Lacombe Research Centre, Agriculture and Agri-Food Canada</p> <p>Gariépy, C.; AAFC, Food Research and Development Center</p> <p>Uttaro, Bethany; Lacombe Research Centre, Agriculture and Agri-Food Canada</p> <p>Lopez-Campos, Oscar; Agriculture and Agri-Food Canada, Lacombe Research Centre</p> <p>Prieto, Nuria; Agriculture and Agri-Food Canada Lacombe Research Centre</p> <p>Dugan, M.; Lacombe Research Centre, Agriculture and Agri-Food Canada</p> <p>Jin, Ye; Inner Mongolia University</p> <p>Juarez, M.; Lacombe Research Centre, Agriculture and Agri-Food Canada</p>
Keywords:	Meat Science, Pork, Carcass

SCHOLARONE™
Manuscripts

Effects of pork differentiation strategies in Canada on pig performance and carcass characteristics

Zhang, H.^{1,2,‡}, Aalhus, J.L.², Gariépy, C.³, Uttaro, B.², López, Campos, O.², Prieto, N.⁴, Dugan, M.E.R.², Jin, Y.^{1,‡}, Juárez, M.^{2,*}

¹College of Food Science and Engineering, Inner Mongolia Agricultural University, Hohhot, Inner Mongolia, China 010018

²Lacombe Research Centre, Agriculture and Agri-Food Canada, Lacombe, AB, Canada T4L1W1

³Food Research and Development Centre, Agriculture and Agri-Food Canada, Saint-Hyacinthe, QC, Canada J2S 8E3

⁴Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, AB, Canada T6G 2P5

*Corresponding author: Manuel Juárez. Phone: +1 403 782 7412. Email: manuel.juarez@agr.gc.ca

[‡]First authors

Abstract

Performance and quality traits were measured in carcasses from combinations of genotype, diet supplement, slaughter weight and carcass chilling regime. Iberian-crossed pigs had lower live animal performance than Duroc and Lacombe. From 70 to 115 Kg, Lacombe pigs grew slightly faster than Duroc. Duroc carcasses had a higher lean percentage and heavier ham and picnic primals and lighter loins and bellies, compared to Lacombe and Iberian. Heavier carcasses had lower lean yield, except those from Iberian-crossed pigs, and bigger bellies. Meat from Duroc-crossed pigs was lighter in colour and higher in marbling, with intermediate values for fat hardness. Iberian carcasses displayed dark meat with intermediate marbling and the hardest fat. Meat from Lacombe pigs was the leanest although, like in all three breeds, marbling scores were higher in heavier carcasses. Generally, supplementing with canola and flax decreased and blast-chilling increased fat hardness values. Dietary canola also seemed to affect meat colour traits, but these effects were not consistent among breeds and slaughter weights. The commercial combination of Duroc breed, control diet and 115 Kg slaughter weight showed a balance in terms of performance, carcass and quality traits. Other combinations evaluated in this study showed potential to efficiently produce differentiated pork.

Key words: canola; cut-out; Duroc; flax; growth; Iberian; Lacombe; pork

Abbreviations: ADG: average daily gain; AFI: average feed intake

Introduction

Variability in pork quality is due to a combination of genetic, production and processing factors (Ngapo and Gariépy 2008). While differentiated high quality pork products are commonly marketed in countries such as Spain (e.g. Iberian Pork) or Italy (e.g. Parma Ham), Canadian pork, an established commercial cross-bred of Yorkshire-Landrace female and Duroc boar, is mainly a commodity product, with little differentiation. Such a situation owes to the genetic selection being applied to a limited number of specialized breeds driven by grading systems encouraging higher carcass yield which has successfully led to leaner carcasses, faster growth rate and lower feed conversions in Canada as in many countries (Pommier et al, 2004; Sellier, 2007). Consequences of this strategy of selection are: 1- pig genetic diversity has eroded and 2- the whole meat sensory characteristics and some reproductive and health traits have been undesirably affected (Rauw et al, 1998).

The Canadian pork industry now recognizes the need for differentiation and development of new high quality products, as well as for adaptability and flexibility to successfully compete in priority markets (Young 2011). Genetics is at the basis of differentiation and studies involving non-selected and selected breeds invariably reported superior meat technological and / or sensory traits in non-selected breeds over those from selected ones (Alfonso et al, 2005; Renaudeau and Mourot, 2007; Labroue et al, 2000). Superior meat technological quality traits have also been reported in Iberian compared to Landrace breed (Serra et al, 1998), supporting further the need to increase genetic variability / diversity. Combining more diverse genetics with dietary treatments, slaughter weight and / or chilling regimes would further increase the necessary variability in support of effective differentiation processes and allow for a better identification of the most appropriate source of variation or treatments to efficiently address market needs.

Evaluating the carcass characteristics of pigs is a first step, essential to optimize pork production systems (Schinckel et al., 2008) and even the entire pork marketing chain (Gispert et al., 2007). Pig carcass merit traits usually include composition (yield of lean, fat and bone), percentage of primal cuts, meat pH value, marbling score, and colour (Marcoux et al. 2007; Ngapo and Gariépy 2008; Uttaro and Zawadski 2010). All these traits have been reported to be influenced by factors such as breed (Gispert et al. 2007; Sellier 1976; Straadt et al. 2013), diet (Caine et al. 2007; Dugan et al. 2001; Thacker et al. 2004), slaughter weight (Latorre et al. 2004; Martin et al. 1980) and chilling regime (Jones et al. 1988; Juárez et al. 2009; Springer et al. 2003). Understanding the potential impact of manipulating each factor, as well as their interactions, would allow the Canadian pork industry to develop differentiated value added products for a highly competitive market.

The objective of the present study is to evaluate the effects of using integrated strategies for producing differentiated pork quality in Canada, based on combinations of breed, gender, diet, slaughter weight and chilling regime, on carcass merit traits with economic impact. Duroc, Lacombe and Iberian breeds form the best model for a study on pork quality as they are genetically both far and equidistant from each other in the phylogenetic tree. Flax was chosen as a dietary source of n-3 fatty acids available in Canada (potential differentiation strategy) and canola was chosen as a Canadian source of oleic acid which could be used both to produce Iberian-based products (traditionally rich in oleic acid) or a Canadian heritage pork, using the Lacombe breed and this local product.

Materials and Methods

Experimental design and animal performance

85 The live animal experimental design for this study consisted of a $3 \times 3 \times 2$ factorial
86 (breed combination, diet and slaughter weight, respectively). Commercial dams (42 Large White $\times$
87 Landrace F1; Hypor Inc., Regina, SK, Canada) were inseminated with semen from three
88 different breeds (4 boars per breed): Duroc (Peak Swine Genetics, Leduc, AB, Canada), Lacombe
89 (Peak Swine Genetics Inc.) and Iberian (Semen Cardona, Cardona, Spain). When, after following
90 commercial practices, live weight reached 70 kg, animals were transferred to experimental
91 rooms, sorted into pens of three (same breed combination and gender in each pen) and these pens
92 were then assigned to a slaughter weight group (115 or 135 kg) and a dietary treatment (control,
93 Canola or Flax). Three replicates of the full design (72 pens and 216 pigs per replicate) resulted
94 in a total of 648 pigs in test. Beginning at 70 Kg, all animals were fed a commercial finisher diet
95 (Table 1; Masterfeeds, Winnipeg, MB, Canada) until four weeks before slaughter. Pigs were then
96 fed a the commercial finisher diet (control), a Canola (10% ExtraPro; 50% full fat canola and
97 50% extruded field peas, O&T Farms, Ltd., Regina, SK, Canada), or a Flax (10% LinPro; 50%
98 flaxseed and 50% extruded field peas, O&T Farms) diet for the last three weeks prior to
99 slaughter, with a transitional week (50% commercial and 50% assigned diet) for the Canola and
100 Flax diets (Table 1). Average pen feed intake was manually measured weekly by subtracting
101 refusals once the animals were transferred to the experimental rooms in order to calculate average
102 daily intake (AFI). Individual weights were measured weekly from birth to slaughter in order to
103 calculate average daily gain (ADG). Pigs had *ad libitum* access to feed and water. In all cases,
104 diets were formulated to meet the nutrient requirements of the pigs and validated by experts from
105 Verus Animal Nutrition (Winnipeg, MB, Canada). All the trials were carried out at the Lacombe
106 Research and Development Centre (Agriculture and Agri-Food Canada, Lacombe, AB) in

accordance with guidelines established by the Canadian Council on Animal Care in Science (Canadian Council on Animal Care, 2009).

Carcass evaluation

When average animal weight per pen reached the target weight (115 or 135 Kg), animals were shipped to the Lacombe Research and Development Centre Agriculture and Agri-Food Canada federally inspected abattoir (distance: 1.6 km). Pigs were shipped at 7:00 am, not being fed since the previous day. Pigs were stunned (2.1 Amps for 5 sec), and exsanguinated in a manner simulating commercial conditions. Processing of carcasses included pasteurization (16 nozzles at 12 L/nozzle for 10 s with 86.4 °C water for a total of 192 L/carcass), following scalding and before evisceration, using an on-line stainless steel pasteurizing cabinet (Bryant et al. 2003). Following splitting of carcass, hot side weights were recorded. At approximately 45-minutes *post-mortem*, an estimated lean yield was determined on the left side between the 3rd and 4th last ribs, approximately 7 cm from the midline, using an Anitech PG100 Grading Probe (Anitech Information Systems Inc., Markham, ON, Canada). Initial (45min) pH and temperature were also collected between the 10th and 11th rib on the left and right *longissimus lumborum* (LL) muscles, using a Hanna HI99163 pH meter equipped with a Hanna Smart electrode FC232 (Hanna Instruments, Laval, QC, Canada) and a Mark III, MC4000 (Sumaq Wholesalers, Toronto, ON, Canada), respectively. The right half carcass was blast chilled (-25 °C, 2.5 m s⁻¹ wind speed) for 1 h. Both left and right carcass sides were then placed in a cooler at 2 °C. At 24 h, cold carcass weight, *longissimus* pH and temperature were recorded. Cooler losses were calculated based on hot and cold carcass weights. Marbling scores from both right and left carcass sides were assessed using the National Pork Producers Council scale (NPPC 2000). Fat hardness was measured on the inner layer of subcutaneous fat above the 2nd thoracic vertebra of

the left and right sides using a Rex Durometer (Rex Gauge Company, Buffalo Grove, IL, USA). Left carcass sides were then divided into primal cuts: including picnic, butt, ham, loin and belly. Picnic, butt, ham and loin were then weighed and dissected into fat (subcutaneous, intermuscular and body cavity depots), bone and lean according to the procedures described by Martin et al. (1981). Skinned-trimmed belly and rib weights were also recorded. For statistical analysis, fat, bone and lean values of all primal cuts were combined

A chop (2.5 cm) was cut from the left and right LLmuscle (between 8th and 9th ribs) and allowed to oxygenate for 20 min. Subjective colour of both right and left carcass sides was evaluated using the Japanese (Nakai et al., 1975) and Canadian Pork Quality Standards (CPI, 2013). Objective colour measurements [L^* , chroma and hue; (CIE 1978)] were determined in duplicate using a Minolta CR-300 with Spectra QC-300 Software (Folio Instruments, Kitchener, ON, Canada).

Statistical analyses

Data were analyzed using the Mixed Model computer algorithm of the SAS Institute, Inc. (SAS 2003) including breed combination, diet and slaughter weight group, respectively as fixed factors and replicate as blocking effect. Chilling method was also included as a fixed factor for those traits measured in both sides of the carcass. Actual slaughter weight, nested within slaughter weight group and breed, was used as a covariate. When significance was indicated by the model ($P < 0.05$), group means were determined using the LSMEANS statement. Test of multiple comparisons of LSMEANS were separated using the F-test protected LSD procedure and adjusted according to the Tukey-Kramer method to ensure the significance of individual comparisons at $P \leq 0.05$. In order to avoid outliers, sick animals and those out of the target weight groups were not used in the final analyses. The total number for the whole study after chopping

outliers was 578 animals, with a minimum of 20 animals for any combination of pre-slaughter treatments.

Results

Animal Performance

No interactive effects ($P > 0.05$) between breed and diet were observed for performance traits. While birth weights were not affected by breed ($P > 0.05$), ADG of Iberian pigs, at all growth stages, was significantly ($P < 0.001$) lower than those from Duroc and Lacombe (Figure 1). From birth to penning (70 Kg) and from 115 to 135 Kg, no significant differences in ADG were observed between Duroc and Lacombe sired pigs. However, from 70 to 115 Kg, Lacombe pigs grew slightly faster than Duroc.

Regarding the dietary treatments evaluated in this study, for the last three weeks before slaughter, AFI was slightly lower ($P < 0.001$; Diet effect: data not shown) for the Canola ($3.36 \text{ kg} \cdot \text{kg}^{-1}$) and higher for the Flax diet ($3.57 \text{ kg} \cdot \text{kg}^{-1}$), compared to the control ($3.48 \text{ kg} \cdot \text{kg}^{-1}$). Thus, ADG was also lower ($P = 0.011$; Diet effect: data not shown) for the Canola diet ($0.81 \text{ kg} \cdot \text{d}^{-1}$), compared to the control ($0.85 \text{ kg} \cdot \text{d}^{-1}$) and Flax ($0.86 \text{ kg} \cdot \text{d}^{-1}$) diets.

Carcass Yield

An interactive effect between breed and slaughter weight was observed for carcass weight (Table 2; $P = 0.022$), hot dressing ($P = 0.012$) and estimated lean yield ($P = 0.003$). As expected, carcass weights were higher from heavier animals (135 Kg). However, the heavy carcasses from Lacombe pigs were slightly lighter than those from Duroc and Iberian. The dressing percentage of carcasses from 115 Kg Duroc and 135 Kg Lacombe pigs were lower than those from 135 Kg Duroc and Iberian. The light (115 Kg) Duroc group had the highest estimated lean yield

percentage, followed by its heavy group. The slaughter weight did not affect this trait in Lacombe pigs, while Iberians had the lowest values, especially in heavy carcasses.

The interaction between breed and slaughter weight was only a trend for cooler loss ($P = 0.057$). Iberian carcasses had the lowest values ($P = 0.006$) and were not affected by differences in slaughter weight. Light carcasses from Lacombe and Duroc pigs showed higher cooler losses than heavy ones. The dietary treatments applied during the last three weeks before slaughter had no effect ($P > 0.05$) on any of these traits.

Carcass Cut-out

The percentage of each tissue in carcasses was highly affected by an interaction between breed and slaughter weight (Table 3). As observed for estimated lean yield percentage, due to the effect of this interaction ($P = 0.045$), the actual percentage of lean from manual dissection was highest in carcasses from 115 Kg Duroc pigs, followed by the heavier carcasses from the same breed. Lacombe had intermediate values, and also higher in light carcasses. Iberian carcasses had the lowest percentage of total lean and slaughter weight did not make any difference for this trait. The opposite interactive effect was observed for subcutaneous fat content ($P = 0.030$), with Iberian carcasses having the greatest amounts and light Duroc carcasses the lowest. Diet only affected the lean content of Lacombe pigs (breed $\times$ diet interaction; $P = 0.042$), with higher values in carcasses from animals fed the Flax diet (56.4%) compared to the control diet (54.3%). The highest (breed $\times$ weight interaction; $P = 0.004$) percentage of intermuscular fat was observed in carcasses from 115 Kg Iberian and 135 Kg Iberian and Lacombe pigs (7.69%), followed by 115 Kg Lacombe and 135 Kg Duroc (6.83%). Light (115 Kg) Duroc pigs (6.44%) had the lowest content for this tissue. Carcasses from Duroc pigs fed flax (6.79%) had higher (breed $\times$ diet interaction; $P = 0.016$) intermuscular fat content than those from the same breed-cross fed canola

(6.37%). On the other hand, pigs fed the control diet (7.65%) had higher intermuscular fat content than those fed canola (6.98%) and flax (7.07%). No diet effect ($P > 0.05$) was observed for Iberian pigs. The percentage of body cavity fat in carcasses from 115 Kg pigs fed the control diet (0.78%) was lower (weight $\times$ diet interaction; $P = 0.025$) than in carcasses from the flax diet (0.86%). In the heavy group (135 Kg), the control diet (0.91%) led to higher content of this tissue than the canola diet (0.76%). Light carcasses from Duroc pigs had the highest percentage of bone, while 115 Kg Iberian and 135 Kg Lacombe carcasses had the lowest (breed $\times$ weight interaction; $P = 0.001$).

Loin percentage was only affected by breed, with Duroc showing the lowest percentage compared to Iberian and Lacombe (Table 3; $P = 0.016$). Breed and weight affected ham percentage, with the highest values being observed on Duroc carcasses, followed by Lacombe and finally Iberian ($P < 0.001$) and relatively heavier hams being observed in lighter carcasses ($P < 0.001$). Carcasses from Iberian cross-bred pigs had the highest percentage of belly primal, compared to Lacombe and Duroc ($P < 0.001$). The proportion of the belly in the total carcass weight of 135 Kg pigs (19.2%) was higher than in 115 Kg pigs (18.6 %; $P = 0.036$) and was not affected by the diet. However, flax-fed pigs had greater proportion of belly primal (18.8%) than canola-fed pigs (18.4%). Duroc cross-bred pigs had the highest percentage of picnic, and light Iberians had the lowest (breed $\times$ weight interaction; $P = 0.034$). The percentage of picnic in carcasses from Duroc pigs (9.93%) was the highest and not affected by the dietary treatment (breed $\times$ diet interaction; $P = 0.046$). Canola-fed Iberian and Lacombe pigs and flax-fed Iberian pigs showed the lowest picnic percentage (9.42%). Most combinations of weight, breed and diet had no effect ($P > 0.05$) on the percentage of butt (9.81%). Carcasses from canola-fed 135 Kg Iberian pigs and control-fed 135 Kg Lacombe pigs showed an increase in the percentage of this

primal cut (10.3%). Carcasses from control- and flax-fed 135 Kg Iberian pigs showed lower values (9.5%) than the rest of the population (breed $\times$ diet interaction; $P = 0.001$).

Carcass Quality Traits

Marbling was highest in *longissimus* from Duroc cross-bred pigs, followed by Iberian and then Lacombe (Table 4; $P < 0.001$), and it was always higher in the 135 Kg group ($P = 0.004$). Fat hardness (arbitrary scale from 1-100) was highest (breed $\times$ weight interaction; $P = 0.002$) in carcasses from 115 Kg Iberian pigs (79.1) and lowest in carcasses from 115 Kg Lacombe pigs (72.2). No weight effect was observed for fat hardness in carcasses from Duroc pigs, which had intermediate values (76.1). Both experimental diets (Canola and Flax) decreased fat hardness values when compared with the control diet ($P < 0.05$). As expected, blast chilling resulted on harder fat in all the treatments evaluated. Although diet ($P = 0.016$), chilling regime ($P = 0.027$) and the interaction breed $\times$ weight ($P < 0.001$) had statistical significant effects on pH values 24 h after slaughter, the actual differences were minor.

Both breed and diet had the same effect on carcass lean colour evaluated using either the Japanese ($P < 0.001$ and $P = 0.026$, respectively) or the newly developed Canadian ($P < 0.001$ and $P = 0.041$, respectively) colour scales. Lean from Iberian pigs obtained the highest scores (darker). Lean from pigs fed flax got higher scores than lean from pigs fed the control diet. Objective colour traits (L^* , $P = 0.004$; chroma, $P = 0.042$; and hue, $P = 0.018$) were affected by a three-way interaction between breed, weight and diet (Figure 2). As observed in the Japanese and Canadian colour scores, Iberian carcasses had the darkest meat (lowest L^*), although the canola diet seemed to slightly increase the lightness in heavy Iberian carcasses (Figure 2a). Canola fed 135 Kg Lacombe pigs had darker meat (lower L^*) than those from the same breed and weight group fed the control diet. A very similar interactive effect between breed, weight and diet was

observed for hue (Figure 2c). Iberian pigs fed the control diet had the highest chroma values. The lowest values were observed in carcasses from canola-fed Duroc pigs (Figure 2b). Chroma was also the only colour trait affected by the chilling regime ($P = 0.044$), decreasing with the use of the blast-chilling method (Table 4).

Discussion

When developing differentiated pork and pork products through management strategies, changes in animal growth performance need to be considered in order to assess their viability in a competitive market. Breed composition has been widely reported to impact animal performance (Miao et al. 2004; Sellier 1976), and directly affects production costs. The Duroc breed is commonly used in cross-breeding programs as a terminal sire to improve performance and meat quality in commercial pork production (McGloughlin et al. 1988; Whittemore and Kyriazakis 2008). The results from the present study show high performance values for all studied breed combinations, around $1 \text{ kg} \cdot \text{day}^{-1}$ in the finishing phase, similar to those reported in previous studies, such as Bertol et al. (2013). However, the higher ADG observed in Lacombe crossbred pigs during the finishing phase (70 to 155 kg) would be an advantage for this minority breed in terms of production costs. Fredeen and Stothart (1969) indicated that the Lacombe pig, the only Canadian pig breed, popular in the 1960s for its maternal traits, had higher growth rate compared to Large White. However, no recent information is available in scientific literature regarding the performance of Lacombe crossbred pigs under current commercial conditions. The present results also indicated, as expected, that the growth of Iberian-crossbred pigs was slower, which would account for greater production costs. Serra et al. (1998) reported that Iberian pigs, with a significant lower ADG, needed six additional weeks to reach the 100 kg live weight

compared to Landrace. Purebred Iberian and Iberian×Duroc crossbreeding, in order to enhance performance of this unselected rustic breed, are the typical alternatives for Iberian pork production in Spain (Ibáñez-Escriche et al. 2014). The Iberian breed has been widely reported to produce high-quality (high marbling and dark colour, among other attributes) pork and pork products (Juárez et al. 2009). Thus, although low performance from the Iberian crossbreeding was expected, this alternative was explored due to the potential enhancement of carcass and meat quality.

The results from the dietary treatments also need to be considered from an efficiency point of view. As observed in our study, low levels of dietary flaxseed have been reported to increase AFI with no effect on growth rates (Juárez et al. 2010; Romans et al. 1995), which may be due to an increase in palatability. Higher AFI with no impact on ADG can lead to higher feeding costs. However, this effect on AFI seems to disappear when flaxseed levels increase $\geq 15\%$ (Juárez et al. 2010; Thacker et al. 2004). Including high levels of canola in the diet of pigs has also been reported to decrease AFI (Dugan et al. 2001; Smit et al. 2014; Woyengo et al. 2014), with no effect (Dugan et al. 2001) or a small decrease in ADG (Woyengo et al. 2014). This may be due to negative effects of canola on diet palatability due to glucosinolate levels (Mawson et al. 1993). In canola-fed pigs, the decrease in AFI and ADG may increase the length of the finishing period, with the subsequent increase in production costs.

The increase in slaughter weight for commercial pigs may benefit slaughter plants by increasing the volume of meat harvested per slaughtered pig. From a production point of view, a recent study (CDPQ 2015) concluded that, unless feed prices are significantly lower, it is more profitable to produce lighter pigs (~120 Kg), as this allows an increase in the number of Kg of lean per pig-place/barn/year. According to previous research, increasing slaughter weight results

in higher dressing percentages(García-Macías et al. 1996). Carcasses from Duroc pigs were the only ones to clearly show this effect in our study, and was due to differences in the weight of internal organs, such as the liver and large intestine ($P<0.001$, data not shown).

Hot carcass weight and estimated lean yield are used to determine the commercial value of pig carcasses in Canada(Gazette_du_Canada 1986). Therefore, within a given slaughter weight range, producers benefit from increased lean yield percentages. For Iberian and Lacombe crossbred pigs, estimated lean yield percentage was higher than actual total lean from carcass dissection. For Duroc pigs, the estimated lean yield percentage was lower. These differences were smaller for Duroc and more evident in carcasses from Iberian pigs. This may be due to the fact that the prediction equations were developed using commercial pigs, with a significant influence of the Duroc breed. The high lean and low backfat content of Duroc carcasses is a very positive trait from a production and processing point of view, and leads to higher carcass prices in the form of premiums(Marcoux et al. 2007). At the same time, the levels of lean and fat content observed in Iberian carcasses would lead to great discounts and loss of profitability. Carcasses from Lacombe-crossbred pigs, although not as lean as those from Duroc-crossbred animals, showed acceptable yield percentages. The lack of effects from the dietary treatments on carcass lean yield is consistent with previous research using similar diets (Caine et al. 2007).

Allometric function explains that fat is a late maturing tissue (Fortin et al. 1987) and usually develops faster when animals are closer to maturity. This explains why heavy carcasses from Duroc and Lacombe pigs showed a decrease in lean content and an increase in the relative contribution of fat depots compared to light carcasses from the same breed. At the same time, Fortin et al.(1987)reportedthat higher relative deposition rate for subcutaneous fat was an indication of earlier maturity.The Iberian breed is known for its early maturing fat deposition

(Daza et al. 2006), explaining the high fat content in Iberian carcasses and why no differences were observed between slaughter weights.

As previously mentioned, carcass weight and estimated lean yield are the main parameters determining the payment received by producers in Canada. However, the relative percentage of each primal cut may have a large impact on the final value of the carcass for the processing plant. Loin and belly are the two most profitable cuts in North America (Greenwood and Dunshea 2009). Marcoux et al. (2007) reported that the economic contribution of the ham and the loin to total carcass value tended to decrease in favour of the contribution of the belly, while the economic contribution of other cuts has been fairly stable. Therefore, increasing the relative percentage of loin and belly may be desirable. In general, fatter carcasses show a higher proportion of belly (Eikelenboom et al. 2004), as observed in the current study for carcasses from heavier pigs, with the subsequent decrease in relative contribution of ham to the total carcass weight. Gispert et al. (2007) reported that the big differences among genotypes in the proportion of primal cuts are worthwhile to characterize. Carcasses from Lacombe- and, especially, Iberian-crossbred pigs also had a larger proportion of loin and belly compared to Duroc-crossbred pigs, which had proportionally heavier hams. This is consistent with results previously reported comparing carcasses from Duroc and other commercial breeds (CCSI 2007; Latorre et al. 2003). The lack of effect of the dietary treatments on carcass composition also agrees with results from previous studies (Dugan et al. 2001).

Similar to carcass composition, meat quality is not currently used to economically reward or penalize producers in the Canadian pork industry. However, processors classify pork based on quality characteristics, such as fat consistency, meat colour and marbling. Quality requirements from national, international and niche markets determine the final value of pork carcasses and

primals. Therefore, profitability is directly affected by these traits. Although the impact of marbling scores on pork palatability is not consistent among studies (Ngapo and Gariépy 2008), marbling is one of the main attributes considered by the industry to classify based on quality. The inclusion of flax and canola in the diet may lead to changes in marbling scores (Caine et al. 2007; Dugan et al. 2001). However, these effects have been reported with dietary levels well above those used in this study. Marbling is the last fat depot to develop (Fortin et al. 1983), as observed by the highest scores observed in 135 Kg carcasses. In Spain, the Iberian pig is commonly crossed with Duroc to increase productive traits (Utrilla et al. 2010). Iberian pork has been reported to be darker and have higher marbling than Duroc-sired pigs (Juárez et al. 2009; Straadt et al. 2013). High levels of intramuscular fat reported in purebred Iberian pigs under traditional production systems tend to decrease in hybrids (Juárez et al. 2009). In North America, Duroc is widely used as terminal sire by the pork industry due to its potential to improve meat quality traits, especially intramuscular fat content (Mandell et al. 2006), of white breeds. Our results show the highest marbling scores in meat from Duroc-crossed pigs, confirming its ability to enhance this trait. The lower marbling scores observed in Iberian-crossed pork may be due to the interaction genotype $\times$ slaughter weight. Iberian pigs usually show their potential for intramuscular fat production when raised as purebred and slaughtered at 160-180 Kg (Juárez et al. 2009). Previous studies (Straadt et al. 2013) reported similar intramuscular fat content in meat from the Iberian $\times$ Landrace/Large White and Duroc $\times$ Landrace/Large White crosses, while meat from Iberian $\times$ Duroc hybrids had significantly higher fat content. Meat from Lacombe-crossed pigs was consistently leaner, as expected from this type of breed, selected for maternal and performance traits (Fredeen and Stothart 1969).

Harder fat is desirable for processing of meat products. Softer fat may lead to low fabrication efficiency, oily appearance at retail and reduced product shelf-life (Soladoye et al. 2015). Fatty acid composition has been generally identified as the main factor responsible for fat hardness (Soladoye et al. 2015). Thus, the lower hardness scores observed in carcasses from animals fed the experimental diets (canola and flax) can be explained by the increase in dietary mono- and polyunsaturated fatty acids (Caine et al. 2007; Dugan et al. 2001; Juárez et al. 2010; Miller et al. 1990). Thicker backfat, as observed in heavy carcasses and carcasses from Iberian-crossed pigs, usually leads to higher fat hardness scores, while leaner carcasses have softer fat (Wood and Riley 1982). However, the low hardness scores observed in carcasses from Lacombe-crossed pigs, which contained medium levels of subcutaneous fat, may be linked to different fat composition due to differences in Lacombe's fat metabolism and deposition. Regardless of breed, diet and slaughter weight, blast-chilled carcasses had higher fat hardness scores. Springer et al. (2003) and Jones et al. (1988) also reported higher fat hardness values in blast-chilled compared to conventionally-chilled carcasses. The connective tissue lipid composition, matrix of connective tissue, as well as its water content, influence fat firmness (Wood et al. 2004). Blast-chilling reduces fat crystal size, increasing fat hardness (Fox and McSweeney 2006).

While the differences observed in pH were minimal and would have no practical consequences, the effects of breed, slaughter weight and diet on colour traits were significant and need to be considered. Pork colour is one of the main attributes affecting consumer's purchase decisions at retail (Ngapo and Gariépy 2008). Iberian pork has been widely reported to display a characteristic dark red colour (Juárez et al. 2009; Straadt et al. 2013). Serra et al. (1998) indicated that Iberian meat had the lowest L^* due to a higher percentage of pigments. At the same time, other studies have reported a lighter less intense red colour in meat from Duroc-

crossed pigs (Bertol et al. 2013). These effects were consistent with our findings using both subjective and objective colour measurements. Springer et al. (2003) and Jones et al. (1988) reported that muscle became darker after blast chilling than after control chilling. Minimum chilling effects were observed on colour traits in the present study, maybe due to different temperature and chilling regimes in others studies. Previous studies have shown meat lightness was unrelated to carcass weight (Čandek-Potokar et al. 1998; García-Macías et al. 1996). However, the effect of the dietary treatments on pork colour was inconsistent across the study, with canola leading to changes in L^* , chroma and hue values in certain combinations of breed and slaughter weight only and the highest chroma value being found in 115 Kg Iberian pigs fed the control diet. Miller et al. (1990) and Myer et al. (1992) reported a decrease in subjective colour scores in carcasses from canola-fed pigs, while Bertol et al. (2013) found no differences in objective colour traits between control- and canola-fed pigs. Thus, although the levels of the dietary treatments in this study were low, the effects of feeding pigs with diets containing ingredients with high levels of unsaturated fatty acids on meat quality need to be monitored when attempting to produce differentiated pork products.

In conclusion, pork differentiation can be achieved by combining modifications of pre- and post-slaughter factors. In the present study, the use of Duroc breed, commercial diets and 115 Kg slaughter weight, commonly used by the pork industry, showed a good balance in terms of performance, carcass yield and quality traits. However, the use of different breeds, diets and slaughter weights may result in positive changes in some of these attributes, such as ADG, percentage of valuable cuts or meat colour. The negative impact on other traits needs to be considered in order to assess potential profitability of meat with differentiated attributes. Using

this information, together with market conditions and consumer's preferences, Canadian producers and processors have the opportunity to manipulate pork quality and not be limited to the commodity pork market.

Acknowledgements

Financial support was provided by the Alberta Livestock and Meat Agency Ltd. (2011F035F). The authors thank the Lacombe Research and Development Centre's operational, processing and technical staff for their dedication and expert assistance.

- Alfonso, L., Mouro, J., Insausti, K., Mendizabal, J.A., Arana, A. 2005. Comparative description of growth, fat deposition, carcass and meat quality characteristics of Basque and Large White. *Anim. Res.* **54**: 33-42.
- Bertol, T. M., de Campos, R. M. L. D., Ludke, J. V., Terra, N. N., de Figueiredo, E. A. P., Coldebella, A., dos Santos Filho, J. I., Kawski, V. L. and Lehr, N. M. 2013. Effects of genotype and dietary oil supplementation on performance, carcass traits, pork quality and fatty acid composition of backfat and intramuscular fat. *Meat Science* **93**(3):507-516
- Bryant, J., Brereton, D. A. and Gill, C. O. 2003. Implementation of a validated HACCP system for the control of microbiological contamination of pig carcasses at a small abattoir. *Canadian Veterinary Journal* **44**(1):51-55.
- Caine, W. R., Aalhus, J. L., Dugan, M. E. R., Lien, K. A., Larsen, I. L., Costello, F., McAllister, T. A., Stanford, K. and Sharma, R. 2007. Growth performance, carcass characteristics and pork quality of pigs fed diets containing meal from conventional or glyphosate-tolerant canola. *Canadian Journal of Animal Science* **87**(4):517-526.
- CPI. 2013. Canadian Pork Farm to Table. Canada Pork International. www.canadapork.com
- Canadian Council on Animal Care. 2009. CCAC guidelines on: the care and use of farm animals in research, teaching and testing. Canadian Council on Animal Care, Ottawa, ON.
- Čandek-Potokar, M., Žlender, B. and Bonneau, M. 1998. Effects of breed and slaughter weight on longissimus muscle biochemical traits and sensory quality in pigs. *Animal Research* **47**(1):3-16.
- CCSI. 2007. Effect of breed and sex on growth, carcass and meat quality traits. Report from the Canadian Centre for Swine Improvement.
- CDPQ. 2015. Technical and economic study on heavier market pigs (140 kg). Centre de développement du porc du Québec.
- CIE. 1978. Recommendations on uniform color spaces – color difference equations – psychometric color terms. Commission Internationale de l'Eclairage (publication no. 15, supplement no. 2, pp. 8-12). Paris, France: CIE.

- Daza, A., Mateos, A., Carrasco, C. L., Rey, A., Ovejero, I. and López-Bote, C. J. 2006. Effect of feeding system on the growth and carcass characteristics of Iberian pigs, and the use of ultrasound to estimate yields of joints. *Meat Science* **72**(1):1-8.
- Dugan, M. E. R., Aalhus, J. L., Lien, K. A., Schaefer, A. L. and Kramer, J. K. G. 2001. Effects of feeding different levels of conjugated linoleic acid and total oil to pigs on live animal performance and carcass composition. *Canadian Journal of Animal Science* **81**(4):505-510.
- Eikelenboom, G., Walstra, P., Huiskes, J. H. and Klont, R. E. 2004. Species of Meat Animals | Pigs. Pages 1284-1291 in W. K. Jensen, ed. *Encyclopedia of Meat Sciences*. Elsevier, Oxford.
- Fortin, A., Wood, J. D. and Whelehan, O. P. 1983. Breed and sex effects on the development and proportion of muscle, fat and bone in pigs. *The Journal of Agricultural Science* **108**:39-45.
- Fortin, A., Wood, J. D. and Whelehan, O. P. 1987. Breed and sex effects on the development, distribution of muscle, fat and bone, and the partition of fat in pigs. *Journal of Agricultural Science Cambridge* **108**:141-153.
- Fox, P. F. and McSweeney, P. 2006. *Advanced Dairy Chemistry: Volume 2 Lipids*. Ed. Birkhäuser, NY, USA:277-278.
- Fredeen, H. T. and Stothart, J. G. 1969. Development of a new breed of pigs: The Lacombe. II. Evaluation. *Canadian Journal of Animal Science* **49**(3):247-261.
- García-Macías, J. A., Gispert, M., Oliver, M. A., Diestre, A., Alonso, P., Muñoz-Luna, A., Siggers, K. and Cuthbert-Heavens, D. 1996. The effects of cross, slaughter weight and halothane genotype on leanness and meat and fat quality in pig carcasses. *Animal Science* **63**(3):487-496.
- Gazette_du_Canada. 1986. Hog carcass grading regulation. Pages 1657-1667.
- Gispert, M., Font i Furnols, M., Gil, M., Velarde, A., Diestre, A., Carrión, D., Sosnicki, A. A. and Plastow, G. S. 2007. Relationships between carcass quality parameters and genetic types. *Meat Science* **77**(3):397-404.
- Greenwood, P. L. and Dunshea, F. R. 2009. 2 - Biology and regulation of carcass composition. Pages 19-60 in J. P. Kerry, D. Ledward, eds. *Improving the Sensory and Nutritional Quality of Fresh Meat*. Woodhead Publishing.
- Ibáñez-Escriche, N., Varona, L., Magallón, E. and Noguera, J. L. 2014. Crossbreeding effects on pig growth and carcass traits from two Iberian strains. *Animal : an international journal of animal bioscience* **8**(10):1569-1576.
- Jones, S. D. M., Murray, A. C. and Robertson, W. M. 1988. The effects of spray chilling pork carcasses on the shrinkage and quality of pork. *Canadian Institute of Food Science and Technology Journal* **21**(1):102-105.
- Juárez, M., Clemente, I., Polvillo, O. and Molina, A. 2009. Meat quality of tenderloin from Iberian pigs as affected by breed strain and crossbreeding. *Meat Science* **81**(4):573-579.
- Juárez, M., Dugan, M. E. R., Aldai, N., Aalhus, J. L., Patience, J. F., Zijlstra, R. T. and Beaulieu, A. D. 2010. Feeding co-extruded flaxseed to pigs: Effects of duration and feeding level on growth performance and backfat fatty acid composition of grower-finisher pigs. *Meat Science* **84**(3):578-584.
- Labroue, F., Goumy, S., Gruand, J., Mourot, J., Neelz, V., Legault, C. 2000. Comparaison au large White de quatre races locales porcines françaises pour les performances de croissance, de carcasse et de qualité de viande. *Journées Rech. porcine en France*, **32**: 403-411.
- Latorre, M. A., Lázaro, R., Valencia, D. G., Medel, P. and Mateos, G. G. 2004. The effects of gender and slaughter weight on the growth performance, carcass traits, and meat quality characteristics of heavy pigs. *Journal of Animal Science* **82**(2):526-533.

- 493 Latorre, M. A., Medel, P., Fuentetaja, A., Lázaro, R. and Mateos, G. G. 2003. Effect of gender,
494 terminal sire line and age at slaughter on performance, carcass characteristics and meat
495 quality of heavy pigs. *Animal Science* **77**(1):33-45.
- 496 Mandell, I. B., Campbell, C. P. and de Lange, C. F. 2006. Effects of gender, sire line, and
497 penning environment on growth, carcass characteristics, and aspects of pork meat
498 quality at different locations in the loin. *Canadian Journal of Animal Science* **86**(1):49-61.
- 499 Marcoux, M., Pomar, C., Faucitano, L. and Brodeur, C. 2007. The relationship between different
500 pork carcass lean yield definitions and the market carcass value. *Meat Science* **75**(1):94-
501 102.
- 502 Martin, A., Sather, A., Fredeen, H. and Jolly, R. W. 1980. Alternative market weights for swine.
503 II. Carcass composition and meat quality. *Journal of Animal Science* **50**(4):699-705.
- 504 Martin, A. H., Fredeen, H. T., Weiss, G. M., Fortin, A. and Sim, D. 1981. Yield of trimmed pork
505 product in relation to weight and backfat thickness of the carcass. *Canadian Journal of*
506 *Animal Science* **61**:299-310.
- 507 Mawson, R., Heaney, R. K., Zdunczyk, Z. and Kozłowska, H. 1993. Rapeseed meal-
508 glucosinolates and their antinutritional effects. Part II. Flavour and palatability. *Food /*
509 *Nahrung* **37**(4):336-344.
- 510 McGloughlin, P., Allen, P., Tarrant, P. V., Joseph, R. L., Lynch, P. B. and Hanrahan, T. J. 1988.
511 Growth and carcass quality of crossbred pigs sired by Duroc, Landrace and Large White
512 boars. *Livestock Production Science* **18**(3):275-288.
- 513 Miao, Z. H., Glatz, P. C. and Ru, Y. J. 2004. Review of Production, Husbandry and
514 Sustainability of Free-range Pig Production Systems. *Asian Australas J Anim Sci*
515 **17**(11):1615-1634.
- 516 Miller, M. F., Shackelford, S. D., Hayden, K. D. and Reagan, J. O. 1990. Determination of the
517 alteration in fatty acid profiles, sensory characteristics and carcass traits of swine fed
518 elevated levels of monounsaturated fats in the diet. *Journal of animal science*
519 **68**(6):1624-1631.
- 520 Myer, R. O., Lamkey, J. W., Walker, W. R., Brendemuhl, J. H. and Combs, G. E. 1992.
521 Performance and carcass characteristics of swine when fed diets containing canola oil
522 and added copper to alter the unsaturated:saturated ratio of pork fat. *Journal of animal*
523 *science* **70**(5):1417-1423.
- 524 Nakai, H., F. Saito, T. Ikeda, S. Ando, and A. Komatsu. 1975. Standard models of pork
525 colour. *Bull. Natl. Inst. Anim. Ind. (Chiba)* **29**:69-75.
- 526 Ngapo, T. M. and Gariépy, C. 2008. Factors affecting the eating quality of pork. *Critical Reviews*
527 *in Food Science and Nutrition* **48**(7):599-633.
- 528 NPPC. 2000. Pork composition & quality assessment procedures. National Pork Producers
529 Council, Des Moines, Iowa.
- 530 Pommier, S. A., Murray, Robertson, W., Aalhus, J., Sosnicki, A., Klont, R. 2004. Effect of
531 genetics on meat quality and sensory properties of pork. 50th Congress Meat Sci.
532 Technol. Helsinki, 4pp.
- 533 Rauw, W.M., Kanis, E., Noordhuizen-Stassen, E.N. and Grommers F.J. 1998. Undesirable side
534 effects of selection for high production efficiency in farm animals: A review. *Livestock*
535 *Production Science* **56**:15-33.
- 536 Renaudeau, D. and Mouro, J. 2007. A comparison of carcass and meat quality characteristics of
537 Creole and Large White pigs slaughtered at 90 kg BW. *Meat Sci.* **76**:165-171.
- 538 Romans, J. R., Johnson, R. C., Wulf, D. M., Libal, G. W. and Costello, W. J. 1995. Effects of
539 ground flaxseed in swine diets on pig performance and on physical and sensory
540 characteristics and omega-3 fatty acid content of pork: I. Dietary level of flaxseed.
541 *Journal of Animal Science* **73**(7):1982-1986.
- 542 SAS. 2003. SAS® user's guide: Statistics. SAS for Windows, version 9.1. SAS Institute, Inc,
543 Cary, NC.

- Schinckel, A.P., Mahan, D.C., Wiseman, T.G. and Einstein, M.E. 2008. Impact of alternative energy systems on the estimated feed requirements of pigs with varying lean and fat tissue growth rates when fed corn and soybean meal-based diets. *Prof Anim Sci.* **24**:198–207.
- Sellier, P. 1976. The basis of crossbreeding in pigs; A review. *Livestock Production Science* **3**(3):203-226.
- Sellier, P. 2007. Génétique et qualité de la viande de porc: quelles perspectives? *Viandes Prod. Carnés.* **26**:11-16.
- Serra, X., Gil, F., Pérez-Enciso, M., Oliver, M. A., Vázquez, J. M., Gispert, M., Díaz, I., Moreno, F., Latorre, R. and Noguera, J. L. 1998. A comparison of carcass, meat quality and histochemical characteristics of Iberian (Guadyerbas line) and Landrace pigs. *Livestock Production Science* **56**(3):215-223.
- Smit, M. N., Seneviratne, R. W., Young, M. G., Lanz, G., Zijlstra, R. T. and Beltranena, E. 2014. Feeding increasing inclusions of canola meal with distillers dried grains and solubles to growing-finishing barrows and gilts. *Animal Feed Science and Technology* **189**:107-116.
- Soladoye, P. O., Shand, P. J., Aalhus, J. L., Gariépy, C. and Juárez, M. 2015. Review: Pork belly quality, bacon properties and recent consumer trends. *Canadian Journal of Animal Science* **95**(3):325-340.
- Springer, M. P., Carr, M. A., Ramsey, C. B. and Miller, M. F. 2003. Accelerated chilling of carcasses to improve pork quality. *Journal of Animal Science* **81**(6):1464-1472.
- Straadt, I. K., Aaslyng, M. D. and Bertram, H. C. 2013. Sensory and consumer evaluation of pork loins from crossbreeds between Danish Landrace, Yorkshire, Duroc, Iberian and Mangalitza. *Meat Science* **95**(1):27-35.
- Thacker, P. A., Racz, V. J. and Soita, H. W. 2004. Performance and carcass characteristics of growing-finishing pigs fed barley-based diets supplemented with Linpro (extruded whole flaxseed and peas) or soybean meal. *Canadian Journal of Animal Science* **84**(4):681-688.
- Utrilla, M. C., Soriano, A. and Ruiz, A. G. 2010. Quality attributes of pork loin with different levels of marbling from duroc and iberian cross. *Journal of Food Quality* **33**(6):802-820.
- Uttaro, B. and Zawadski, S. 2010. Prediction of pork belly fatness from the intact primal cut. *Food Control* **21**(10):1394-1401.
- Whittemore, C. T. and Kyriazakis, I. 2008. *Whittemore's Science and Practice of Pig Production*. John Wiley & Sons, Oxford, UK.
- Wood, J. D., Nute, G. R., Richardson, R. I., Whittington, F. M., Southwood, O., Plastow, G., Mansbridge, R., Da Costa, N. and Chang, K. C. 2004. Effects of breed, diet and muscle on fat deposition and eating quality in pigs. *Meat Science* **67**(4):651-667.
- Wood, J. D. and Riley, J. E. 1982. Comparison of boars and castrates for bacon production. 1. Growth data, and carcass and joint composition. *Animal Science* **35**(01):55-63.
- Woyengo, T. A., Yáñez, J., Young, M. G., Lanz, G., Beltranena, E. and Zijlstra, R. T. 2014. Nutritional value of full-fat green canola seed fed to growing-finishing pigs. *Journal of Animal Science* **92**(8):3449-3459.
- Young, M. 2011. Differentiation through Quality: What are the perceptions and attitudes about pork quality in Canada's priority markets? . *Proc. Banff Pork Seminar*, Banff, AB.

Table 1. Ingredient composition and nutrient content of diets

Ingredient	Control	Canola	Flax
Wheat (%)	44.0	27.8	17.1
Barley (%)	37.7	43.7	56.4
Canola meal (%)	15.0	15.0	14.0
ExtraPro (%)	-	10.0	-
LinPro (%)	-	-	10.0
Calcium carbonate (%)	1.20	1.20	0.95
Soyameal 46% (%)	1.00	1.05	-
Salt (%)	0.43	0.42	0.41
Dicalcium phosphate (%)	0.33	0.25	0.31
Lysine (%)	0.14	0.31	0.38
Threonine (%)	-	0.09	0.13
Methionine (%)	-	-	0.09
Vitamin mix (%)	0.10	0.10	0.10
Mineral mix (%)	0.10	0.10	0.10
Ronozyme (%)	0.02	0.02	0.02
Composition	Control	Canola	Flax
ME (Mcal/K)	3.03	3.13	3.12
Crude protein (%)	15.0	15.0	15.0
Crude fat (%)	2.00	3.94	3.88
Crude fibre (%)	5.00	5.72	5.78
Dry matter (%)	87.8	88.5	88.5
Sodium (%)	0.10	0.19	0.19
Calcium (%)	0.70	0.65	0.65
Phosphorus (%)	0.50	0.50	0.50
Vitamin A (KIU/kg)	8.00	8.00	8.00
Vitamin D (KIU/kg)	1.00	1.00	1.00
Vitamin E (IU/kg)	20.0	20.0	20.0
Zinc (mg/kg)	175	175	175
Copper (mg/kg)	26.8	27.4	26.8
Lysine (%)	0.72	0.97	0.95
Methionine (%)	0.26	0.28	0.37
Threonine (%)	0.53	0.68	0.67
Tryptophan (%)	0.18	0.19	0.18

Table 2. Interactive effect of breed and slaughter weight on carcass weight, dressing percentage, cooler loss and estimated lean yield

	Duroc		Iberian		Lacombe		SEM	<i>P</i> value			
	115Kg	135Kg	115Kg	135Kg	115Kg	135Kg		Breed(B)	Weight(W)	Diet	B×W
Commercial hot weight, kg	96.8 ^c	115 ^a	96.8 ^c	115 ^a	96.9 ^c	114 ^b	0.68	0.101	<0.001	0.705	0.022
Hot dressing, %	82.6 ^b	83.2 ^a	82.8 ^{ab}	83.3 ^a	83.0 ^{ab}	82.5 ^b	0.56	0.348	0.240	0.601	0.012
Cooler loss, %	2.38	2.25	2.25	2.24	2.36	2.30	0.02	0.006	0.001	0.232	0.057
Estimated Lean Yield, %	61.6 ^a	60.5 ^b	55.2 ^d	54.4 ^e	59.0 ^c	58.9 ^c	0.34	<0.001	<0.001	0.562	0.003

^{a,b,c,d,e} Different superscripts within the same row indicate significant differences ($P < 0.05$)

Table 3. Interactive effect of breed and slaughter weight on carcass composition

	Duroc		Iberian		Lacombe		SEM	P value						
	115Kg	135Kg	115Kg	135Kg	115Kg	135Kg		Breed(B)	Weight(W)	Diet(D)	B×W	B×D	W×D	B×W×D
Primal Weight, % Total Carcass Weight														
Loin	27.5	27.9	28.4	28.0	28.1	28.1	0.18	0.016	0.980	0.633	0.089	0.609	0.611	0.483
Ham	27.0	26.4	24.2	23.8	26.2	25.8	0.09	<0.001	<0.001	0.192	0.394	0.447	0.533	0.363
Belly	17.6	18.2	20.0	20.6	18.3	18.8	0.11	<0.001	<0.001	0.261	0.944	0.702	0.036	0.622
Picnic	9.96 ^a	9.90 ^a	9.35 ^c	9.67 ^b	9.58 ^b	9.64 ^b	0.07	<0.001	0.061	0.139	0.034	0.046	0.262	0.404
Butt	9.98 ^{ab}	9.92 ^{ab}	9.99 ^{ab}	9.82 ^{ab}	9.83 ^b	10.04 ^a	0.08	0.843	0.938	0.111	0.034	0.000	0.374	0.001
Tissue Weight, % Total Carcass Weight														
Lean	63.5 ^a	62.2 ^b	47.8 ^c	48.0 ^c	57.0 ^c	54.1 ^d	0.60	<0.001	0.004	0.820	0.045	0.042	0.360	0.161
Bone	9.19 ^a	8.77 ^b	8.28 ^c	8.65 ^{bc}	9.02 ^{ab}	8.22 ^c	0.15	0.001	0.017	0.300	0.001	0.311	0.395	0.122
SQ	20.2 ^c	21.5 ^d	35.1 ^a	34.9 ^a	26.3 ^c	29.3 ^b	0.57	<0.001	0.002	0.429	0.030	0.067	0.470	0.143
IM	6.44 ^c	6.81 ^b	7.91 ^a	7.54 ^a	6.86 ^b	7.61 ^a	0.16	<0.001	0.041	0.673	0.004	0.016	0.141	0.148
BC	0.70	0.77	0.94	0.90	0.81	0.80	0.03	<0.001	0.700	0.101	0.205	0.777	0.025	0.192

SQ: subcutaneous fat; IM: inter-muscular fat; BC: body-cavity fat

^{a,b,c,d,e} Different superscripts within the same row indicate significant differences ($P<0.05$)

Table 4. Effects of breed, slaughter weight, diet and chilling method on pig carcass characteristics

	Breed			Weight		Diet			Chill		SEM	<i>P</i> value					
	Duroc	Iberian	Lacombe	115Kg	135Kg	Control	Canola	Flax	Control	Blast		Breed(B)	Weight(W)	Diet(D)	Chill(C)	B×W	B×W×D
Marbling	1.89 ^a	1.69 ^b	1.41 ^c	1.61 ^b	1.71 ^a	1.66	1.70	1.62	1.66	1.66	0.08	<0.001	0.004	0.166	0.972	0.299	0.395
Fat Hardness	76.1 ^b	78.6 ^a	73.9 ^c	75.5 ^b	76.8 ^a	78.8 ^a	74.9 ^b	74.8 ^b	74.2 ^b	78.1 ^a	1.66	<0.001	0.011	<0.001	<0.001	0.002	0.555
pH 24h	5.53 ^a	5.51 ^b	5.50 ^b	5.52	5.51	5.51 ^{ab}	5.52 ^a	5.50 ^b	5.51 ^b	5.52 ^a	0.01	0.569	0.111	0.016	0.027	<0.001	0.388
Jap. Colour	3.62 ^b	4.42 ^a	3.67 ^b	3.90	3.91	3.84 ^b	3.92 ^{ab}	3.95 ^a	3.90	3.91	0.03	<0.001	0.714	0.026	0.797	0.993	0.929
Can. Colour	3.52 ^b	4.36 ^a	3.53 ^b	3.81	3.79	3.73 ^b	3.82 ^{ab}	3.86 ^a	3.84	3.77	0.08	<0.001	0.730	0.041	0.090	0.497	0.516
<i>L</i> *	51.3 ^a	48.5 ^b	51.0 ^a	50.3	50.3	50.4	50.2	50.2	50.4	50.1	0.5	<0.001	0.699	0.334	0.056	0.949	0.004
Chroma	9.32 ^c	10.43 ^a	9.65 ^b	9.79	9.81	9.99 ^a	9.63 ^b	9.78 ^{ab}	9.89 ^a	9.71 ^b	0.27	<0.001	0.757	0.007	0.044	0.003	0.042
Hue	26.1 ^a	22.5 ^c	24.8 ^b	24.5	24.5	24.8	24.2	24.4	24.5	24.5	0.65	<0.001	0.972	0.068	0.870	0.290	0.018

Japanese: Japanese colour score; Canadian: Canadian colour score

^{a,b,c} Different superscripts within the same row and effect indicate significant differences ($P < 0.05$)

Figure 1. Breed effect on average daily gain at different growth stages

^{a,b,c}Different superscripts within each growth stage indicate significant differences ($P<0.05$)

Figure 2. Interactive effect of breed, weight and diet on colour (2a: L^* , 2b: chroma and 2c: hue) of pig carcasses

^{a-g}Different superscripts indicate significant differences ($P < 0.05$)