Field calibration of blowfly-derived DNA against traditional methods for assessing mammal diversity in tropical forests

Journal:	<i>Genome</i>
Manuscript ID	gen-2015-0193.R1
Manuscript Type:	Article
Date Submitted by the Author:	15-Mar-2016
Complete List of Authors:	Lee, Ping-Shin; University of Malaya, Faculty of Science Gan, Han Ming; Monash University Malaysia Clements, Gopalasamy-Reuben ; Universiti Malaysia Terengganu Wilson, John; University of Malaya,
Keyword:	camera traps, cage traps, hair traps, Malaysia, mist nets

Field calibration of blowfly-derived DNA against traditional methods for assessing mammal diversity in tropical forests

Ping-Shin Lee^{1,2}, Han Ming Gan^{3,4}, Gopalasamy Reuben Clements^{3,5,6,7}, John-James Wilson^{1,2*}

¹Museum of Zoology, Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia.

²Ecology and Biodiversity Program, Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia.

³School of Science, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway, 47500 Petaling Jaya, Selangor, Malaysia.

⁴Monash University Malaysia Genomics Facility, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway, 47500 Petaling Jaya, Selangor, Malaysia.

⁵Centre for Tropical Environmental and Sustainability Science, and College of Marine and Environmental Sciences, James Cook University, 4870 Cairns, Queensland, Australia.

⁶Kenyir Research Institute, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Malaysia.

⁷Rimba, 4 Jalan 1/9D, 43650 Bandar Baru Bangi, Selangor, Malaysia.

*Author for correspondence:

Dr. John James Wilson
Institute of Biological Sciences,
Faculty of Science,
University of Malaya,
50603 Kuala Lumpur, Malaysia.
Fax: 03-79674112
Email: wilso04@gmail.com

Abstract

Mammal diversity assessments based on DNA derived from invertebrates have been suggested as alternatives to assessments based on traditional methods; however, no study has field-tested both approaches simultaneously. In Peninsular Malaysia, we calibrated the performance of mammal DNA derived from blowflies (Diptera: Calliphoridae) against traditional methods used to detect species. We first compared five methods (cage trapping, mist netting, hair trapping, scat collection, and blowfly-derived DNA) in a forest reserve with no recent reports of megafauna. Blowfly-derived DNA and mist netting detected the joint highest number of species ($n=6$). Only one species was detected by multiple methods. Compared to the other methods, blowfly-derived DNA detected both volant and non-volant species. In another forest reserve, rich in megafauna, we calibrated blowfly-derived DNA against camera traps. Blowfly-derived DNA detected more species ($n=11$) than camera traps ($n=9$), with only one species detected by both methods. The rarefaction curve indicated that blowfly-derived DNA would continue to detect more species with greater sampling effort. With further calibration, blowfly-derived DNA may join the list of traditional field methods. Areas for further investigation include blowfly feeding and dispersal biology, primer biases, and the assembly of a comprehensive and taxonomically-consistent DNA barcode reference library.

Keywords: camera traps, cage traps, hair traps, Malaysia, mist nets

Introduction

A wide variety of field methods have been used to assess mammal diversity in tropical forests. Frequently used methods include: live trapping such as cage traps (Hanif-Ridzuan et al. 2010; Madinah et al. 2011), mist nets and harp traps (Kingston et al. 2003; Sing et al. 2013), camera traps (Clements 2013; Hedges et al. 2015a), indirect signs such as tracks or scat (Daim 2002), interviews with local communities (Sharma et al. 2005), direct observations by researchers (Syakirah et al. 2000; Jayaraj et al. 2013), and hair traps (Castro-Arellano et al. 2008; Hedges et al. 2015b). Recent additions to the toolbox are secondary sources of mammal DNA, for example, mammal DNA detected from owl-pellet bones (Rocha et al. 2015) and invertebrate gut contents (Calvignac-Spencer et al. 2012; Schnell et al. 2012; Calvignac-Spencer et al. 2013; Lee et al. 2015; Schnell et al. 2015). These methods can provide accurate identification of mammal species, are not stressful to the mammals themselves, require the least ecological and taxonomic expertise, and yet have the potential to detect rare and cryptic species (Calvignac-Spencer et al. 2012; Schnell et al. 2012; Lee et al. 2015).

Blowflies (Diptera: Calliphoridae) have shown promising potential as sources of mammal DNA due to their presence in all habitats (Norris 1965) and broad host preferences as saprophagous and coprophagous generalists (Calvignac-Spencer et al. 2012; Azwandi et al. 2013; Calvignac-Spencer et al. 2013; Lee et al. 2015; Schnell et al. 2015). *Chrysomya bezziana* has been reported feeding on mammal species from the orders Artiodactyla (7 spp.), Carnivora (6 spp.), Diprotodontia (2 spp.), Perissodactyla (4 spp.), Proboscidea (1 spp.), and Primates (1 spp.) at a zoo in Malaysia (Spradbery & Vanniasingham 1980). *Lucilia sericata* is found in Asia, North America, and Europe (James 1947; Picard & Wells 2010), and has been reported to feed on wounds of sheep, cats, a dog, and a horse in Israel (Schnur et al. 2009). Blowflies, such as *Chrysomya megacephala*, were the first and most abundant insects arriving at *Macaca fascicularis*, *Oryctolagus cuniculus*, and *Rattus norvegicus* carcasses in a forest in Peninsular Malaysia (Azwandi et al. 2013). Standardised methods for sampling of blowflies have been developed which take into account the persistence period of mammal DNA in blowfly guts (Lee et al. 2015). Despite the increasing interest in invertebrate-derived mammal DNA for mammal diversity assessments, field calibrations of the performance of invertebrate-derived mammal DNA against traditional methods have yet to be conducted (Schnell et al. 2015).

Each field method listed above differs in terms of the targeted mammal groups and resource costs, with advantages and disadvantages for each method (summarised in Table 1). Recently, a number of studies from the forests of Peninsular Malaysia, home to 222 mammal species (DWNP 2010), have simultaneously tested the effectiveness of different methods in detecting mammal species (Syakirah et al. 2000; Jayaraj et al. 2012; Tingga et al. 2012; Jayaraj et al. 2013). In Gunung Stong State Park, Kelantan, a combination of cage traps, mist nets, and harp traps yielded low capture rates of non-volant small mammals (Jayaraj et al. 2012); cage traps did not detect treeshrews and squirrels (probably due to bait incompatibility and poor positioning), while harp traps captured a low diversity of insectivorous bats. In Taman Negara, Pahang, a combination of cage traps, Sherman traps, mist nets, and harp traps demonstrated the efficiency of harp traps for mammal sampling (Tingga et al. 2012). The harp traps accounted for 84% of the bat species reported and 65% of the total mammal species. In contrast, at Wang Kelian State Park, Perlis, after several years of surveys using a combination of harp traps, cage traps, and direct observations, the inclusion of mist nets in the latest sampling period increased the number of species sampled by 33% (from an average of 24 per survey to 36; Jayaraj et al. 2013).

Here, we evaluate the potential of blowfly-derived DNA for mammal diversity assessments in tropical forests. We conducted two field studies to calibrate the performance of blowfly-derived mammal DNA at generating species inventories and richness estimates against traditional methods in Peninsular Malaysia. We first compared five methods - cage trapping, mist netting, hair trapping, scat collection, and blowfly-derived DNA - in a tropical forest reserve with no recent reports of megafauna. Next, we compared blowfly-derived DNA with the most popular method to assess megafauna diversity - camera traps - in another tropical forest reserve with megafauna.

Materials and Methods

Study sites

Two field calibrations were conducted in two forest reserves in Peninsular Malaysia: 1) Ulu Gombak Forest Reserve (UGFR) in Selangor; and 2) Tembat Forest Reserve (TFR) in Terengganu. UGFR comprises 17,000 ha of logged dipterocarp forest (3°20'N, 101°45'E; Figure 1). UGFR is considered one of the most species-rich localities for Old World bats (Sing et al. 2013) and has been intensively studied since the establishment of Ulu Gombak

Field Studies Centre (University of Malaya) 50 years ago (Medway 1966). We compiled an extensive mammal checklist for UGFR based on previous records (including University of Malaya student theses) of sampled or observed species (Table S1). TFR, comprising logged dipterocarp forest (5°11'N, 102°41'E), is one of 17 ecological linkages recognised in the Malaysia Federal Government's "Central Forest Spine Master Plan for Ecological Linkages" to restore connectivity between four fragmented forest complexes (DTCP & DOF 2012).

Field methods

Our first field survey was conducted at UGFR between 3rd November 2014 and 25th December 2014 for two nights each week (16 nights in total). Two transects of 100 m were established 1500 m apart (Figure 1). The survey incorporated (a) baited cage traps, (b) mist nets, (c) baited hair traps, (d) scat collection, and (e) baited blowfly traps.

- a) Ten wire mesh cage traps (Figure 1) were set per transect at 10 m intervals, monitored and baited with fresh banana daily. Banana is reported as being among the effective bait for cage traps used for small mammal diversity assessments in tropical forests of Malaysia (Bernard 2003; Payne & Francis 2005; Madinah et al. 2011) and is resistant to removal by invertebrates (Bernard 2003). Cage traps were checked daily, and hair samples from any trapped mammals were collected into a 1.5 mL microcentrifuge tube before release.
- b) Ten mist nets (9 m x 4 m x 36 mm mesh size) (Figure 1) were positioned across potential flight paths of bats (trails or streams). Mist nets were set at a single transect from 1900 and were monitored continuously until 2300 or until it rained. A small wing punch was collected from each captured bat into a 1.5 mL ethanol-filled microcentrifuge tube following AMNH (2013). Scissors and forceps were cleaned with alcohol and sterile tissues between bats to avoid cross-contamination. Mist nets were only used for four weeks (two nights per week) in November.
- c) Ten hair traps (PVC pipes covered with reversed duct tape on the inside) (Figure 1) were set per transect at 10 m intervals and monitored daily. Hair traps were baited with banana at the start of the 2-night cycle and collected at the end of the cycle. Any hair samples from traps were collected into a 1.5 mL microcentrifuge tube using sterile forceps.
- d) Any scat (Figure 1) encountered while walking along transects daily was collected into a 1.5 mL microcentrifuge tube.

Table S1. Checklist of mammal species recorded at Ulu Gombak Forest Reserve.

e) Ten blowfly traps baited with rotten fish (see Lee et al. 2015) were set per transect at 10 m intervals and 2 m above the ground. The traps were emptied every 24 h, and collected blowflies (Diptera: Calliphoridae) were frozen at -20°C within 5 h of trap emptying. The guts of all collected blowflies (=315) were then dissected with sterile implements and pooled (five individual blowfly guts per tube) for DNA extraction (63 extractions in total). The DNA extracts were further pooled by week and transect of sampling, resulting in (8 x 2) 16 pooled DNA extracts for bulk PCR.

Our second field survey was conducted at TFR between 11th April 2015 and 27th May 2015 (30 days in total). A 28 km transect was used incorporating (a) baited blowfly traps and (b) camera traps (Figure 1). A longer transect was used, compared to UGFR, in accordance with the longer transects generally used for camera trapping surveys (Clements 2013).

a) Twenty baited blowfly traps (Lee et al. 2015) were set along the transect (Figure 1), 2 m above the ground, 500 m apart. As above, the traps were emptied every 24 h, and collected blowflies were frozen at -20 °C within 5 h of trap emptying. The guts of all collected blowflies (=1,345) were then dissected with sterile implements and pooled (five individual blowfly guts per tube) for DNA extraction (269 extractions in total). The DNA extracts were further pooled by sampling day, resulting in 30 pooled DNA extracts for bulk PCR.

b) Twenty Panthera V4 camera traps with passive infrared motion sensors were deployed along the transect (Figure 1), attached to tree trunks approximately 50 cm above ground level. Camera traps were placed on linear features known to have high detection probabilities for large mammals (e.g. animal trail, ridge, or old logging road; Harmsen et al. 2010). No attractants were employed around the cameras. The cameras were triggered by motion, and only photocaptures that were obtained during the sampling period of the blowfly traps (30 days) were used for analysis and were catalogued using Camera Base version 1.4 software (Tobler 2012).

Species-level identification of blowfly-derived DNA

Our protocol for minimally-invasive collection of mammal DNA samples (hair, wing punches) from mammals trapped in cages or nets has been approved by the University of Malaya Institutional Animal Care and Use Committee (UMIACUC) (Ref. ISB/02/1212013/JJW (R)) and the Department of Wildlife and National Parks, Peninsular Malaysia (Ref. JPHL&TN(IP): 80-4/2 Jld16(24)). DNA was extracted from the collected hair

and bat tissue samples using an alkaline lysis extraction method (100 μ L elution; modified from Ivanova et al. 2012). The samples collected into 1.5 mL microcentrifuge tubes were first cut into smaller fragments using sterile scissors. 35 μ L of alkaline buffer (0.1 N NaOH, 0.3 mM EDTA, pH 13.0) was then added into the tube, followed by centrifugation for 1 min at 110,000 g and incubation at 95 °C for 20 min. 65 μ L of neutralization buffer (0.1 M Tris-HCl, pH 7.0) was then added to the tubes, followed by vigorous vortexing and centrifugation (110,000 g, 1 min). DNA was extracted from scat using a QIAamp DNA stool mini kit (QIAGEN, USA) following the manufacturer's protocol (200 μ L elution). DNA was extracted from blowfly guts using a NucleoSpin Tissue kit (Macherey-Nagel, Germany) (100 μ L elution), following the manufacturer's instructions.

The extracted DNA samples from hair, bat tissue, and scat samples were used for PCR in standard protocols following Wilson (2012) and Wilson et al. (2014). Primers VF1d_t1 / VR1d_t1 (Ivanova et al. 2012) were used for a first pass and RonM / VR1 for a second pass (Ivanova et al. 2012). PCR was performed using EconoTaq PLUS 2X mastermix (Lucigen, USA) and *COI* Fast thermocycling program (see Wilson 2012), with slight modification of the amount of DNA template added (0.5-2 μ L) depending on DNA extraction method. PCR products were then Sanger-sequenced by a local company (MYTACG-Kuala Lumpur, Malaysia) using the reverse PCR primer. Each sequence generated from individual specimens was trimmed of primers and edited using CodonCode Aligner (CondonCode Corp., USA) (following Wilson 2012).

For blowfly-derived DNA, a 205 bp *COI* amplicon was generated using a two-step PCR that utilises universal mammal primers, Uni-Mini-bar F and RonPing R (Lee et al. 2015) with partial Illumina adaptor sequences, TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG and GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG incorporated onto the 5' end of the forward and reverse primers, respectively. EconoTaq PLUS 2X Master Mix (Lucigen, USA) and *COI* Fast thermocycling program (see Wilson 2012) were used to generate amplicons from 1 μ L of DNA extract (five replicate PCRs were performed for each pooled DNA extract, together with a negative control). PCR products were visualised on a 2% agarose gel stained with 1 x GelRed (Biotium, USA), and gels were extracted and purified using NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel, Germany), following the manufacturer's instructions. Of the 16 pooled DNA extracts from UGFR, seven produced

PCR products and were selected for high-throughput (next-generation) DNA sequencing (HTS). Of the 30 pooled DNA extracts from TFR, 17 produced PCR products and were selected for HTS. The purified PCR products were used as templates for a second round of PCR reaction to generate amplicons containing Illumina adaptors and unique dual-index multiplex identifier (MID) tags. Briefly, 6 μ L of the purified PCR product was mixed with 10 μ L of NEBNext® High-Fidelity 2X Master Mix (New England Biolabs, USA) and 2 μ L of each Nextera XT N70X and N50Y dual-index primers (Illumina, San Diego, CA) to make up a total of 20 μ L reaction volume. The thermal cycling conditions were 98°C for 30s; followed by 6 cycles of 98°C for 10 s, 65°C for 30 s, and 72°C for 30 s; and a final extension at 72°C for 5 min. The PCR products were subsequently purified and size-selected using 0.8x volume ratio of AMPure XT® beads (Beckman Coulter Inc., USA). The final amplicons from all samples were quantified using KAPA Library Quantification Kit (KAPA Biosystems, CapeTown, South Africa), normalized, pooled, and then sequenced on the Illumina MiSeq Sequencer (model number: M02133; 2 x 250 bp paired-end read setting) located at the Monash University Malaysia Genomics Facility. The sequencing run produced 5,541,198 paired end reads corresponding to 2,403 megabases data output. The MiSeq outputs (FASTQ) were submitted to the NCBI Sequence Read Archive under project accession number SRP064503. The Illumina reads were demultiplexed and trimmed of adapters onboard the MiSeq using the MiSeq Reporter Software. The ‘raw’ output of paired-end reads was quality filtered using the online PRINSEQ (Schmieder & Edwards 2011) by sequence length (75-350 bp), minimum mean quality score (>25), and GC content (20-50%). Reads with low complexity (threshold using Entropy=80) and characters other than A, C, T, or G were excluded. Redundant reads (exact sequence duplicates, 5’ sequence duplicates, 3’ sequence duplicates, reverse complement exact sequence duplicates, and reverse complement 5’/3’ sequence duplicates) were dereplicated. Reads with mean scores of less than 5 (sliding window size =5, step size =5) or minimum tail length of poly-A/T tails of five at 5’end and five at 3’end were trimmed of low-quality bases at 5’end (1 bp) and 3’end (1 bp). The output data (in FASTA format) was then further filtered and trimmed of primer sequences with quality control and filtering steps performed manually in CodonCode Aligner (CondonCode Corp., USA) following Brandon-Mong et al. (2015). Both consensus and singleton reads surviving the quality control and filtering steps were retained for taxonomic assignment.

The resulting DNA barcodes obtained from all individual specimens collected from cage traps, mist nets, and scat collection, and unique DNA metabarcodes from blowfly-derived

DNA were uploaded to the Barcode of Life Data Systems (BOLD) (Ratnasingham & Hebert 2007) and are available in the public dataset DS-BDNGS. Linnaean species names were assigned to our DNA barcodes (individual specimens collected from cage traps, mist nets, and scat collection) and a representative DNA metabarcode (recovered from blowfly-derived DNA) when they had sequence similarity matches of >98% to DNA barcodes with Linnaean species names (submitted by other BOLD users) in the full database of BOLD (see Wilson et al. 2014). In the case of conflicts, i.e. our DNA (meta)barcodes had sequence similarity matches of >98% to database DNA barcodes with several different Linnaean species names, we assigned the Barcode Index Number (BIN) (Ratnasingham & Hebert 2013) of the matching reference DNA barcodes to our DNA (meta)barcodes without using a species epithet. When our DNA (meta)barcodes had sequence similarity matches of <98% but >94% with reference DNA barcodes, we assigned the genus name of the closest matching reference DNA barcode to our DNA (meta)barcodes (see Zeale et al. 2011). When our DNA (meta)barcodes had sequence similarity matches of <94% but >90% with reference DNA barcodes, we assigned the order name of the closest matching reference DNA barcode to our DNA (meta)barcodes (see Zeale et al. 2011). DNA (meta)barcodes sharing <90% sequence similarity to the closest matching DNA barcode in BOLD were discarded from further analyses but are discussed anecdotally below.

Species identification from camera traps

Identification of species from the photocaptures (Figure 2) was carried out by LPS and GRC based on morphological characters with the aid of Francis (2008). Photocaptures of mammals that could not be identified with certainty due to poor lighting or blurred images were excluded.

Field method calibration

For the methods used at UGFR, the percentage of potentially detectable species in the UGFR checklist (Table S1) detected by each field method was calculated. The expected species richness (using Chao 1) for each field method was computed in EstimateS Version 9.1.0 (Colwell 2006). Chao 1 is suitable for species richness estimation of our sampling sites as it corrects for bias due to sampling effort and species rarity (Chao 1984; Iknayan et al. 2014). Rarefaction curves of expected species richness (using Chao 1) with 95% confidence intervals were generated based on the cumulative sampling days using EstimateS Version

9.1.0 (Colwell 2006; Figure 4). The sampling completeness ratio (observed species richness/expected species richness; Soberón et al. 2000) was calculated for each method.

Results

At UGFR a total of 14 mammal species from the orders Artiodactyla, Chiroptera, Primates, and Rodentia (Figure 3) were detected from a total effort of 1,440 trap days (320 from cage traps; 160 from mist nets; 320 from hair traps; 320 from scat collection; 320 from blowfly traps). Mist nets and blowfly traps contributed the most species-rich samples with six species (including BINs without formal names) detected each, followed by cage traps and scat collection that detected two species each; hair traps detected no species (Figure 3). Blowfly traps recorded species from four orders - Artiodactyla, Chiroptera, Primates, and Rodentia (Figure 3). The other field methods (except hair traps that detected no species) detected only a single order each. Two domesticated mammal species, *Bos* [under BIN - BOLD:AAA2294] and *Felis* [BOLD:AAC2892], and non-mammal taxa including one bird species, *Gallus gallus* [BOLD:AAA3630], a midge species, and a fish species, were also detected in blowfly-derived DNA, but these were excluded from further analyses as our focus was the detection of wild mammal species. Of 20 potentially detectable species of small-bodied rodents and treeshrews recorded in the UGFR checklist, both cage traps and blowfly traps detected two species (10.0%). Of 57 potentially detectable bats recorded in the UGFR checklist, mist nets detected six species (10.5%), and blowfly traps detected two species (3.5%). Of 45 potentially detectable non-volant mammals recorded in the UGFR checklist, blowfly traps detected the highest number of species (n=4; 8.9%). Blowfly traps detected one new record of non-volant mammal for the UGFR checklist - *Trachypithecus obscurus* [BOLD:AAI0540]. Sampling completeness ratios for blowfly traps, cage traps, and scat collection were 1.00, and 0.86 for mist nets. The HTS data from UGFR included reads from five blowfly species - *Chrysomya* [BOLD:AAC4787; BOLD:ACD5557; and BOLD:ACF0516], and *Lucilia* [BOLD:AAA6618; and BOLD:ACQ1337].

At TFR a total of 20 mammal species from the orders Artiodactyla, Carnivora, Cetartiodactyla, Chiroptera, Perissodactyla, Primates, Proboscidea, and Rodentia were detected from a total effort of 1,200 trap days (600 from camera traps; 600 from blowfly traps; see Figure 3). Blowfly traps detected a higher number of species (n=11) than camera traps (n=9), including five orders - Artiodactyla, Carnivora, Chiroptera, Primates, and Rodentia,

while camera traps detected six orders - Artiodactyla, Carnivora, Cetartiodactyla, Perissodactyla, Primates, and Proboscidea. While only one potentially domesticated mammal species was detected - *Bos* [BOLD:AAA2294], its identify could also be that of a wild bovid known from TFR - *Bos gaurus*. Non-mammal taxa including a bird - *Gallus gallus* [BOLD:AAA3630], three fish species, a lizard species, a snake species, another squamate, and a turtle species, were also detected from blowfly-derived DNA but were excluded from further analyses. Additional non-mammal taxa detected, but which had sequence similarity matches of <90% with DNA barcodes in BOLD include an algae (76-80% sequence similarity) and another snake (89% sequence similarity); these were also excluded from further analyses. Sampling completeness ratio of blowfly traps was 0.79, and 1.00 for camera traps (Figure 4). The HTS data from TFR included reads from five blowfly species - *Chrysomya* [BOLD:AAA5667; BOLD:AAB3064; BOLD:ACD5557; and BOLD:ACF0516], and *Lucilia* [BOLD:ACQ1337].

Discussion

Blowfly traps and mist nets detected the highest number of wild mammal species (6 species each) at UGFR. These were followed by cage traps and scat collection, which detected two species each. Our hair traps did not detect any species at UGFR, but the difficulties associated with obtaining mammal DNA samples from hair traps have already been documented in Malaysia (see Hedges et al. 2015b). There was no difference in the number of non-volant small-bodied mammal species detected by blowfly traps and cage traps (2 species each). Only one species was detected by multiple field methods (cage traps, scat collection, and blowfly traps) - *Rattus* [BOLD:AAB2208]. Blowfly traps were less effective than mist nets in detecting bat species (2 versus 6 of 57 in the checklist). The effectiveness of mist nets in capturing bat species, and consequently making a significant contribution to overall mammal species richness estimates, has also been shown in other studies in Peninsular Malaysia (Jayaraj et al. 2012; Jayaraj et al. 2013). The number of species detected by our mist nets is similar to another study at UGFR that detected seven species using ten mist nets and four harp traps over 9 trap days (Sing et al. 2013). There was no species detected by both mist nets and blowfly traps. All species detected from cage traps and scat collection were detected at both transects at UGFR except one species - *Sundamys* [BOLD:AAD5076], only detected from two cage traps at transect 1. For blowfly traps, all species were detected at both transects except two species - *Sus* [BOLD:AAA3445], only detected at transect 1, and

Macroglossus sobrinus [BOLD:AAA9952], only detected at transect 2. At TFR, blowfly traps detected higher species richness than camera traps - 11 species from the blowfly traps versus 9 species (based on visual identification) from camera traps. Blowfly traps detected more mammal species that were volant and arboreal in nature than those detected by camera traps. Only one species was potentially detected by both blowfly traps and camera traps - *Sus* [BOLD:AAA3445] which is probably “*Sus scrofa*” from the camera traps.

The blowfly traps at UGFR detected the highest number of orders (4 orders), in contrast to the traditional methods that detected species from only a single mammal order each. Cage traps can potentially detect small non-volant mammals of several orders but in our study only detected Rodentia, probably due to the placement of cage traps on the ground where they can be avoided by arboreal mammals (see Fontúrbel 2010). Mist nets are restricted in only being able to detect Chiroptera. Scat collection should, theoretically, not be restricted to detecting any specific mammal orders, but detection likely depends on the range size of the individuals of a given species relative to the length of transect patrolled; in our study only Rodentia were detected from scat. The number of orders detected by blowfly traps at UGFR increases (to 5 orders) when the two domestic mammal species were included - *Bos* [BOLD:AAA2294], from Artiodactyla, and *Felis* [BOLD:AAC2892], from Carnivora. The blowfly traps at TFR detected fewer orders compared to the camera traps (5 versus 6 orders). However, the comparison may be less meaningful given the lack of taxonomic equivalency (Bertrand et al. 2006) between the orders of large charismatic megafauna (specifically targeted by the camera traps) and other mammal orders. For example, Perissodactyla and Proboscidea are orders containing only a single extant species in Peninsular Malaysia. Nine vertebrate orders were detected by blowfly traps at TFR when the bird species - *G. gallus* [BOLD:AAA3630], three fish species, a lizard species, a snake species, a squamate, and a turtle species, whose DNA was also detected in the blowfly guts, were included.

The detection of a broad diversity of mammals and other vertebrate orders from blowfly traps is consistent with the study by Calvignac-Spencer et al. (2012), who detected six mammal orders, an avian order, and an amphibian order from blowfly-derived DNA sampled at Tai National Park, Côte d'Ivoire, and Kirindy Forest, Madagascar. Although these results demonstrate the broad range of blowfly hosts, biases in PCR primer binding affinity may lead to sequences of certain species being amplified less efficiently than others (Deagle et al. 2014; Thomsen & Willerslev 2015). This may limit the detections to species with higher primer

binding affinity, resulting in species diversity being underrepresented (Shokralla et al. 2012). Lee et al. (2015) found the primers used in this study exhibited no strong amplification bias across mammal species (from Peninsular Malaysia), but the test only included a limited number of species (41).

One further uncertainty in the use of blowfly-derived DNA for mammal diversity assessments is potential blowfly feeding biases and dispersal distances. Surprisingly the blowfly traps did not detect *Macaca fascicularis*, the long-tailed macaque, at UGFR and TFR despite our frequent observation of numerous individuals at the site throughout the sampling period. Nevertheless, the blowfly traps at UGFR did detect a different primate - *Trachypithecus obscurus* [BOLD:AAI0540], the dusky leaf monkey, which is a near threatened species (IUCN 2015). This species is known to occur in the vicinity of UGFR, but had not been reported from the reserve itself. It is possible, however, that the blowflies fed on *T. obscurus* tissue or faeces outside UGFR before moving into our sampling area. The few studies on blowfly dispersal (from temperate and subtropical regions - Braack & Retief 1986; Smith & Wall 1998; Tsuda et al. 2009) suggested individuals travel 100-2,400 m per day, but there is no data available on blowfly travelling distances in the tropics. It is also not known whether blowflies, as opportunistic feeders (Calvignac-Spencer et al. 2013), are more likely to have fed on carcasses, wounds, or faeces. All field methods recovered a low percentage of their potentially detectable species from the UGFR checklist (2-10%), which may be an artefact of the short sampling period and restricted spatial scale of sampling relative to the total size of forest reserve, rather than feeding or primer biases. However, UGFR has more than 20 years of logging history (Nor Hashim & Ramli 2013), and megafauna recorded in 1961 (Medway 1966), such as *E. maximus*, is highly likely to be extirpated as no records of activities or populations have been published since.

Another current limitation of the blowfly-derived DNA approach is the reliance on public DNA databases to assign detected species formal species names (Collins & Cruickshank 2013). Five out of 14 wild mammal species (36%) detected at UGFR and six out of 20 species (30%) detected at TFR could not be assigned traditional species names. For example, *Rattus* [under BIN - BOLD:AAB2208] detected from blowfly-derived DNA, cage traps, and scat collection likely represents *Rattus* “*sp. R3*” (Pages et al. 2010), a classic “dark taxon” (see Wilson et al. 2014) whereby it has not yet been formally described and is likely to be reported as *R. andamanensis*, *R. argentiventer*, or *R. tanezumi* in studies relying on visual

diagnosis (Robins et al. 2007; Pages et al. 2010; Latinne et al. 2013; Li et al. 2015). The BIN associated with this taxon comprises DNA barcode members under the traditional names *Rattus tanezumi*, *Rattus tiomanicus*, *Rattus rattus*, *Rattus sp.*, and *Rattus sp. abtc47994* in BOLD. The two DNA barcodes named *Rattus rattus* (JF459864 and JF459865 in GenBank) had been flagged earlier as problematic and attributed to misidentification or contamination (Shen et al. 2013). In addition, *Rhinolophus sp. 25Ne* showed no species-level match (>98%) in BOLD despite many bat DNA barcodes from Peninsular Malaysia being present in BOLD (Lim & Wilson 2015). *Sus sp. 4Ba*, detected at TFR, also showed no species-level match (>98%) in BOLD, despite the presence of 218 DNA barcodes of *Sus*, suggesting a cryptic species. We could only assign formal genus names to DNA metabarcodes detected from blowfly-derived DNA, whose closest matches were *Callosciurus notatus* [under BIN - BOLD:AAF8467] (93-94% sequence similarity), and *Paradoxurus hermaphroditus* (KC894743 in GenBank; only two sequences of ≤ 421 bp available in BOLD, so there is no BIN) (95%). The 205 bp DNA mini-barcode region amplified by the primers used in this study was successful in distinguishing 94% of 113 examined mammal species from Peninsular Malaysia (mined from GenBank) in a previous study by Lee et al. (2015), including separating 26 “dark” bat taxa.

The utility of blowfly-derived DNA in identifying threatened mammal species in tropical forests remains questionable. However, this is also true for other field methods (e.g. cage traps, mist nets, and scat collection used in our study) that rely on DNA-based identification for accurate assignment of formal species names (Wilson et al. 2014). As species boundaries are increasingly being refined and recognised on the basis of molecular data, especially for species-rich groups (Francis et al. 2010; Pages et al. 2010), but even for charismatic megafauna (Wilting et al. 2015), field methods that rely solely on visual diagnosis (e.g. camera traps) will have increasingly limited utility. It also remains to be seen whether blowfly-derived DNA can yield nuclear DNA of appropriate quality for individual identification (Calvignac-Spencer et al. 2013). Currently, camera trapping is the most popular approach to monitor populations of mammal species, but this is restricted to species that have individuals with unique markings (e.g. spotted and striped felids; Henschel & Ray 2003; Maffei et al. 2005; Hedges et al. 2015a), or to individuals that can be marked prior to camera trapping (Trolle & Kery 2003; Rowcliffe et al. 2008).

At UGFR the sampling completeness ratio for all field methods exceeded 0.86, while at TFR the sampling completeness ratio of camera traps was 1 and of blowfly traps was 0.79. This suggested that with further sampling, blowfly traps may continue to detect a higher number of species. One reason for the high values for sampling completeness of the traditional field methods, despite low detection of the known species present, could be the limited distribution of traps over representative habitats in the study site, as some mammal species may be restricted to specific habitat types (Geier & Best 1980; Goulart et al. 2009). The dispersal of blowflies could, in theory, overcome the problem of localised trap placement as blowflies will disperse across all habitat types (Norris 1965). This could explain the detection of *T. obscurus* in blowfly traps as noted above. Alternatively, for all the field methods, except blowfly traps and potentially hair traps and scat collection, it is relatively safe to conclude that the species detected was present at the precise location of the trap during a known (short) time interval.

Deciding on the most cost-effective technique to conduct a mammal diversity assessment in tropical forests depends on many factors. For example, blowfly traps are more prone to damage but can be easily replaced at low cost (< \$1 each). During our study, six blowfly traps were lost, as opposed to four cage traps (\$3 each) and two camera traps (\$250 each). Camera traps may be more prone to theft than blowfly traps, and can leave a significant financial dent in the project when that happens (high-end camera traps can cost up to \$1000; Rovero et al. 2013), but are less prone to damage. Nonetheless, there is high initial cost involved in purchasing reagents and equipment for DNA-based detection (DNA extraction, PCR, and next generation sequencing), as well as personnel costs for analysing samples. However, species identification of bats detected from mist nets and harp traps, and rodents detected from cage traps (as exemplified by the case of *Rattus* [BOLD:AAB2208]) with an acceptable degree of accuracy also requires DNA barcoding (Wilson et al. 2014), together with the associated costs. High-throughput-sequencing can now be outsourced to private companies and/or research institutes at increasingly competitive rates (see Brandon-Mong et al. 2015).

Conclusion

We have shown that blowfly-derived DNA is a feasible technique for mammal diversity assessments, especially when considering that a high number of species and orders were detected relative to traditional methods. Blowfly-derived DNA can potentially overcome

ecological and taxonomical challenges associated with traditional methods when detecting threatened and rare or previously unrecorded species. While the sampling completeness ratios obtained from blowfly traps indicates a potential to detect more species with greater sampling effort, the lack of overlap in species detected by different field methods indicates employing multiple field methods may be the fastest way to obtain a representative account of species. Some uncertainties in the use of blowfly-derived DNA remain, such as the impact of blowfly dispersal, biases in primer binding affinity, and lack of a comprehensive and taxonomically-consistent DNA barcode reference library. With further calibration to address these uncertainties, blowfly-derived DNA may join the list of traditional field methods used to assess mammal diversity in tropical forests.

Acknowledgements

This study was supported by grants from the Nagao Natural Environment Foundation Japan to LPS and the Ministry of Higher Education Fundamental Research Grant Scheme (FP042-2014A) to JJW, GRC, and Yong Kien Thai (University of Malaya). GHM is grateful to the Monash University Malaysia Tropical Medicine and Biology Platform for financial and infrastructure support. GRC is grateful to permits from the Economic Planning Unit (Permit no: 3747), Department of Wildlife and National Parks for Project Harimau Selamanya [Rujukan kami: JHPL&TN(P)/80-4/2 Jld. 18(7)], Department of Forestry [Rujukan kami: JH/100Jld. 8(16)], and financial support from the Woodland Park Zoo-Panthera Wildcat Conservation Grant. The collaboration between the University of Malaya and Monash University Malaysia falls under an MoA. Yong Kien Thai provided advice and suggestions on an earlier version of this manuscript. Lee Yoon Hin and Tee Ah Jee provided assistance during fieldwork. Sing Kong Wah, Brandon Mong Guo Jie, and Thary Goh Gazi provided advice and suggestions during laboratory work. Lam Wai Yee, Foon Junn Kitt, and Sri Venkateswara Rao from Project Harimau Selamanya provided advice and technical assistance during camera trapping. This project was presented at the 6th International Barcode of Life Conference as part of an Institutional Links Program (Queen Mary University of London-University of Malaya) supported by the Newton-Ungku Omar Fund.

References

- Abu, N.A.C. 2007. Kajian perbandingan kepelbagaian tikus di 3 kawasan hutan simpan berbeza (Gombak, Kenaboi, Johor). B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- AMNH (American Museum of Natural History). 2013. Wing punch and hair sampling protocols. Available from <http://research.amnh.org/vz/mammalogy/donating-bat-tissue-and-hair-samples-genomic-and-stableisotope-studies/wing-punch-and-hair-sampling>. [accessed 4 March 2015].
- Aziz, A.A. 2010. Ecology and morphology of shrews in PPL, Ulu Gombak. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- Azwandi, A., Nina Keterina, H., Owen, L.C., Nurizzati, M.D., and Omar, B. 2013. Adult carrion arthropod community in a tropical rainforest of Malaysia: analysis on three common forensic entomology animal models. *Trop. Biomed.* **30**(Suppl 3): 481-494.
- Beja Pereira, A., Oliveira, R., Alves, P.C., Schwartz, M.K., and Luikart, G. 2009. Advancing ecological understandings through technological transformations in noninvasive genetics. *Mol. Ecol. Resour.* **9**(5): 1279-1301. doi:10.1111/j.1755-0998.2009.02699.x.
- Bernard, H. 2003. Bait preferences of some small mammal species of North Borneo based on line-trapping with wire-mesh cage traps. *Sabah Parks Nat. J.* **6**: 27-44.
- Bernard, H., Ahmad, A.H., Brodie, J., Giordano, A.J., Lakim, M., Amat, R., Koh, S.P.H., Lee, S.K., Tuuga, A., Malim, P.T., Lim-Hasegawa, D., Yap, S.W., and Sinun, W. 2013. Camera-trapping survey of mammals in and around Imbak Canyon conservation area in Sabah, Malaysian Borneo. *Raffles. Bull. Zool.* **61**(2): 861-870.
- Bertrand, Y., Pleijel, F., and Rouse, G.W. 2006. Taxonomic surrogacy in biodiversity assessments, and the meaning of Linnaean ranks. *Syst. Biodivers.* **4**: 149-159.
- Braack, L.E.O., and Retief, P.F. 1986. Dispersal, density, and habitat preference of the blow-flies *Chrysomya albiceps* (Wd.) and *Chrysomya marginalis* (Wd.) (Diptera: Calliphoridae). *Onderstepoort. J. Vet. Res.* **53**: 13-18.

- Brandon-Mong, G.J., Gan, H.M., Sing, K.W., Lee, P.S., and Wilson, J.J. 2015. DNA metabarcoding of insects and allies: an evaluation of primers and pipelines. *Bull. of Entomol. Res.* **7**: 1-11. doi:10.1017/S0007485315000681.
- Bonvicino, C.R., and Bezerra, A.M.R. 2003. Use of regurgitated pellets of Barn Owl (*Tyto alba*) for inventory small mammals in the Cerrado of central Brazil. *Stud. Neotrop. Fauna E.* **38**: 1-5. doi:10.1076/snfe.38.1.1.14030.
- Buś, M.M., Żmihorski, M., Romanowski, J., Balčiauskienė, L., Cichocki, J., and Balčiauskas, L. 2014. High efficiency protocol of DNA extraction from *Micromys minutes* mandibles from owl pellets: a tool for molecular research of cryptic mammal species. *Acta Theriol.* **59**: 99-109. doi:10.1007/s13364-013-0144-y.
- Calvignac-Spencer, S., Leendertz, F.H., Gilbert, M.T.P., and Schubert, G. 2013. An invertebrate stomach's view on vertebrate ecology. *Bioessays* **35**: 1004-1013. doi:10.1002/bies.201300060.
- Calvignac-Spencer, S., Merkel, K., Kutzner, N., Kühl, H., Boesch, C., Kappeler, P.M., Metzger, S., Schubert, G., and Leendertz, F.H. 2012. Carrion fly-derived DNA as a tool for comprehensive and cost-effective assessment of mammalian biodiversity. *Mol. Ecol.* **22**: 915-924. doi:10.1111/mec.12183.
- Castro-Arellano, I., Madrid-Luna, C., Lacher, T.E., and León-Paniagua, L. 2008. Hair-trap efficacy for detecting mammalian carnivores in the tropics. *J. Wildl. Managemt.*, **72**: 1405-1412. doi:http://dx.doi.org/10.2193/2007-476.
- Catling, P.C., Burt, R.J., and Kooyman, R. 1997. A comparison of techniques used in a survey of the ground-dwelling and arboreal mammals in forests in North-eastern New South Wales. *Wildl. Res.* **24**: 417-432.
- Chao, A. 1984. Nonparametric estimation of the number of classes in a population. *Scand. J. Stat.* **11**(4): 265-270.
- Clements, G.R. 2013. The environmental and social impacts of roads in Southeast Asia. Ph.D. thesis, School of Marine and Tropical Biology, James Cook University, Australia. Available from <http://researchonline.jcu.edu.au/31888>. [accessed 8 July 2015].

- Collins, R.A., and Cruickshank, R.H. 2013. The seven deadly sins of DNA barcoding. *Mol. Ecol. Resour.*, **13**(6): 969-975. doi:10.1111/1755-0998.12046
- Colwell, R.K. 2006. EstimateS: statistical estimation of species richness and shared species from samples. Version 9.1.0. Available from <http://purl.oclc.org/estimates>. [accessed 10 February 2015].
- Daim, M.S. 2002. The care and management of domesticated elephants in Malaysia. *In* Giants on our hands: proceedings of the international workshop on the domesticated Asian elephant, 5-10 Feb 2001. *Edited by* I. Baker and M. Kashio. FAO Regional Office for Asia and the Pacific (RAPA), Bangkok.
- Davison, A., Birks, J.D.S., Brookes, R.C., Braithwaite, T.C., and Messenger, J.E. 2006. On the origin of faeces: morphological versus molecular methods for surveying rare carnivores from their scats. *J. Zool.* **257**(2): 141-143.
- Deagle, B.E., Jarman, S.N., Coissac, E., Pompanon, F., and Taberlet, P. 2014. DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biol. Letters* **10**: 20140562. doi:10.1098/rsbl.2014.0562.
- DTCP, and DOF (Department of Town and Country Planning & Department of Forestry). 2012. Central Forest Spine: summary of master plan for ecological linkages. DTCP, Kuala Lumpur.
- DWNP. 2010. Red list Of Mammals for Peninsular Malaysia. Department of Wildlife and National Parks Peninsular Malaysia, Kuala Lumpur. pp. 1-150.
- Ebert, C., Huckschlag, D., Schulz, H.K., and Hohmann, U. 2010. Can hair traps sample wild boar (*Sus scrofa*) randomly for the purpose of non-invasive population estimation? *Eur. J. Wildl. Res.* **56**(4): 583-590.
- Fontúrbel, F.E. 2010. A methodological approach to assess the small mammal community diversity in the temperate rainforest of Patagonia. *Mamm Biol.* **75**: 294-301.
- Francis, C.M. 2008. A field guide to the mammals of South-East Asia. New Holland Publishers, United Kingdom.
- Francis, C.M., Borisenko, A.V., Ivanova N.V., Eger, J.L., Lim, B.K., Guillen-Servent, A., Kruskop, S.V., Mackie, I., and Hebert, P.D.N. 2010. The roles of DNA barcodes in

- understanding and conservation of mammal diversity in Southeast Asia. PLoS ONE **5**(9): e12575. doi:10.1371/journal.pone.0012575.
- Garden, J.C., McAlpine, C.A., Possingham, H.P., and Jones, D.N. 2007. Using multiple survey methods to detect terrestrial reptiles and mammals: what are the most successful and cost-efficient combinations? *Wildl. Res.*, **34**: 218- 227.
- Geier, A.R., and Best, L.B. 1980. Habitat selection by small mammals of riparian communities: evaluating effects of habitat alterations. *J. Wildl. Managemt.* **44**(1): 16-24.
- Gese, E.M. 2001. Monitoring of terrestrial carnivore populations. USDA National Wildlife Research Center - Staff Publications. Paper 576. Available from http://digitalcommons.unl.edu/icwdm_usdanwrc/576. [accessed 20 August 2015].
- Goulart, F.V.B., Cáceres, N.C., Graipel, M.E., Tortato, M.A., Ghizoni, Jr. IR., and Oliveira-Santos, L.G.R. 2009. Habitat selection by large mammals in a southern Brazilian Atlantic Forest. *Mamm. Biology.* **74**(3): 182-190.
- Hanif-Ridzuan, M.D., Nur-Aida, M.T., Wan-Nurainie, W.I., Zahirunisa, R., Madinah, I., and Abdullah, M.T. 2010. Species diversity, richness and abundance of small mammals in Endau Kluang, Johor, Malaysia. *In* Proceedings of National Biodiversity Seminar 2010.
- Harmsen, B.J., Foster, R.J., Silver, S., Ostro, L., and Doncaster, C.P. 2010. Differential use of trails by forest mammals and the implications for camera-trap studies: a case study from Belize. *Biotropica* **42**: 126-133. doi:10.1111/j.1744-7429.2009.00544.x.
- Hassan, N. 2009. Kepelbagaian order rodentia & scandentia di 3 altitud berbeza. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- Hedges, L., Clements, G.R., Aziz, S.A., Yap, W., Laurance, S., Goosem, M., and Laurance, W.F. 2013 Small carnivore records from a threatened habitat linkage in Terengganu, Peninsular Malaysia. *Small Carnivores Conservation* **49**: 9-14.
- Hedges, L., Lam, W.Y., Campos-Arceiz, A., Mark Rayan, D., Laurance, W.F., Latham, C.J., Saaban, S., and Clements, G.R. 2015a. Melanistic leopards reveal their spots: infrared cameras provide a population-density estimate of leopards in Malaysia. *J. Wildl. Managemt.* **79**: 846-853.

- Hedges, L., Marrant, D.S., Campos-Arceiz, A., and Clements, G.R. 2015b. Feasibility of using scent-baited hair traps to monitor carnivore populations in Peninsular Malaysia. *Trop. Conserv. Sci.* **8**(4): 975-982.
- Heller, K.G., and Volleth, M. 1989. Fledermäuse aus Malaysia. 1. Beobachtungen zur Biologie, Morphologie und Taxonomie. *Senck. Biol.* **69**: 243-276.
- Heller, K.G., and Volleth, M. 1995. Community structure and evolution of insectivorous bats in the Paleotropics and Neotropics. *J. Trop. Ecol.* **11**: 429-442.
- Henschel, P., and Ray, J. 2003. *Leopards in African Rainforests: Survey and Monitoring Techniques*. Wildlife Conservation Society, New York, NY.
- Hill, J.E. 1972. New records of Malayan bats, with taxonomic notes and the description of a new *Pipistrellus*. *Bull. Brit. Mus. Nat. Hist. Zool.* **23**: 23-42.
- Hill, J.E. 1974. New records of bats from South-eastern Asia, with taxonomic notes. *Bull. Brit. Mus. Nat. Hist. Zool.* **27**: 127-138.
- Hoppe-Dominik, B., Kühl, H.S., Radl, G., and Fischer, F. 2011. Long-term monitoring of large rainforest mammals in the biosphere reserve of Tai National Park, Côte d'Ivoire. *Afr. J. Ecol.* **49**: 450-458.
- Iknayan, K.J., Tingley, M.W., Furnas, B. J., and Beissinger, S.R. 2014. Detecting diversity: emerging methods to estimate species diversity. *Trends Ecol. Evol.* **29**: 97-106. doi: 10.1016/j.tree.2013.10.012.
- IUCN (International Union for Conservation of Nature). 2015. IUCN Red List of Threatened Species. Available from <http://www.iucnredlist.org>. [accessed 6 June 2015].
- Ivanova, N.V., Clare, E.L., and Borisenko, A.V. 2012. DNA barcoding in mammals. *Methods Mol. Biol.* **858**: 153-182.
- James, M.T. 1947. The flies that cause myiasis in man (No. 631). Department of Agriculture, United States.
- Jayaraj, V.K., Daud, S.H.M., Azhar, M-I., Sah, S.A.M., Mokhtar, S.I., and Abdullah, M.T. 2013. Diversity and conservation status of mammals in Wang Kelian State Park, Perlis, Malaysia. *Check List* **9**(6): 1439-1448.

- Jayaraj, V.K., Tahir, N.F.D.A., Udin, N.A., Baharin, N.F.K., Ismail, S.K., and Zakaria, S.N.A. 2012. Species diversity of small mammals at Gunung Stong State Park, Kelantan, Malaysia. *J. Threat. Taxa* **4**(6): 2617-2628.
- Jenkins, P.D., and Hill, J.E. 1981. The status of *Hipposideros galeritus* Cantor, 1846 and *Hipposideros cervinus* (Gould, 1854) (Chiroptera: Hipposideridae). *Bull. Brit. Mus. Nat. Hist. Zool.* **27**: 127-138.
- Kingston, T., Francis, C.M., Zubaid, A., and Kunz, T.H. 2003. Species richness in an insectivorous bat assemblage from Malaysia. *J. Trop. Ecol.* **19**: 67-79.
- Kuenzi, A.J., and Morrison, M.L. 1998. Detection of bats by mist-nets and ultrasonic sensors. *Wildl. Soc. Bull.* **26**: 307-311.
- Latinne, A., Waengsothorn, S., Rojanadilok, P., Eiamampai, K., Sribuarod, K., and Michaux, J.R. 2013. Diversity and endemism of Murinae rodents in Thai limestone karsts. *Syst. Biodivers.* **11**: 323-344.
- Lee, P.S., Sing, K.W., and Wilson, J.J. 2015. Reading mammal diversity from flies: the persistence period of amplifiable mammal mtDNA in blowfly guts (*Chrysomya megacephala*) and a new DNA mini-barcode target. *PLoS ONE* **10**(4): e0123871. doi:10.1371/journal.pone.0123871.
- Li, J., Zheng, X., Cai, Y., Zhang, X., Yang, M., and Yue, B. 2015. DNA barcoding of Murinae (Rodentia: Muridae) and Arvicolinae (Rodentia: Cricetidae) distributed in China. *Mol. Ecol. Resour.* **15**(1): 153-167.
- Lim, B.L., Muul, I., and Chai, K.S. 1977. Zoonotic studies of small animals in the canopy transect at Bukit Lanjan Forest Reserve, Selangor, Malaysia. *Malay. Nat. J.* **31**: 127-140.
- Lim, V.C., and Wilson, J.J. 2015. Progress in DNA barcoding the bats of Peninsular Malaysia. *Genome* **58**: 246.
- Madinah, A., Fatimah, A., Mariana, A., and Abdullah, M.T. 2011. Ectoparasites of small mammals in four localities of wildlife reserves in Peninsular Malaysia. *Southeast Asian J. Trop. Med.* **42**(4): 803-813.

- Maffei, L., Noss, A.J., Cuellar, E., and Rumiz, D.I. 2005. Ocelot (*Felis pardalis*) population densities, activity and ranging behaviour in the dry forests of eastern Bolivia: data from camera trapping. *J. Appl. Ecol.* **45**: 1228-1236.
- Medway, L. 1965. Mammals of Borneo. Field keys and annotated checklist. Malaysian Branch of the Royal Asiatic Society, Kuala Lumpur.
- Medway, L. 1966. The Ulu Gombak Field Studies Centre. *Malay. Scient.* **2**: 1-16.
- Medway, L. 1967. A bat-eating, *Megaderma lyra* Geoffroy. *Malay. Nat. J.* **20**: 107-110.
- Medway, L. 1983. The Wild Mammals of Malaya (Peninsular Malaysia and Singapore). 2nd edn. Oxford University Press, Kuala Lumpur.
- Meijaard, E., Mengersen, K., Buchori, D., Nurcahyo, A., Ancrenaz, M., Wich, S., Suci, S., Atmoko, U., Tjiu, A., Prasetyo, D., Nardiyono Hadiprakarsa, Y., Christy, L., Wells, J., Albar, G., and Marshall, A.J. 2011. Why don't we ask ? A complementary method for assessing the status of great apes. *PLoS ONE* **6**(3): e18008. doi:10.1371/journal.pone.0018008.
- Mohd-Azlan, J., Belant, J.L., and Meijaard, E. 2013. Can secondary information inform about population trends of carnivores in Borneo? *Raffles Bull. Zool.* **S28**: 1-8.
- Mukherjee, S., Ashalakshmi, C.N., Home, C., and Ramakrishnan, U. 2010. An evaluation of the PCR-RFLP technique to aid molecular-based monitoring of felids and canids in India. *BMC Res. Notes* **3**: 159. doi:10.1186/1756-0500-3-159.
- Mullins, J., Statham, M., Roche, T., Turner, P., O'Reilly, C. 2009. Remotely plucked hair genotyping: a reliable and non-invasive method for censusing pine marten (*Martes martes*, L. 1758) populations. *Eur. J. Wildl. Res.* **5**(3): 443-453.
- Mumma, M.A., Soulliere, C.E., Mahoney, S.P., and Waits, L.P. 2014. Enhanced understanding of predator-prey relationships using molecular methods to identify predator species, individual and sex. *Mol. Ecol. Resour.* **14**(1): 100-108.
- Nor Hashim, E., and Ramli, R. 2013. Comparative study of understorey birds diversity inhabiting lowland rainforest virgin jungle reserve and regenerated forest. *Sci. World J.* **676507**. doi:10.1155/2013/676507.

- Norris, K.R. 1965. The bionomics of blow flies. *Ann. Rev. Entomol.* **10**: 47-48.
- Pages, M., Chaval, Y., Herbreteau, V., Waengsothorn, S., Cosson, J.F., Hugot, J.P., Morand, S., and Michaux, J. 2010. Revisiting the taxonomy of the Rattini tribe: a phylogeny-based delimitation of species boundaries. *BMC Evol. Biol.* **10**: 184.
- Payne, J., and Francis, C.M. 2005. *A Field Guide to the Mammals of Borneo*. The Sabah Society, Kota Kinabalu.
- Picard, C.J., and Wells, J.D. 2010. The population genetic structure of North American *Lucilia sericata* (Diptera: Calliphoridae), and the utility of genetic assignment methods for reconstruction of postmortem corpse relocation. *Forensic Sci. Int.* **195**(1): 63-67.
- Ratnasingham, S., and Hebert, P.D.N. 2007. Bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Mol. Ecol. Notes* **7**(3): 355-364.
- Roberts, N.J. 2011. Investigation into survey techniques of large mammals: Surveyor competence and camera-trapping vs. transect-sampling. *Biosci. Horiz.* **4**: 40-49.
- Robins, J., Hingston, M., Matisoo-Smith, E., and Ross, H. 2007. Identifying *Rattus* species using mitochondrial DNA. *Mol. Ecol. Notes* **7**: 717-729.
- Rocha, R.G., Justino, J., Leite, Y.L.R., and Costa, L.P. 2015. DNA from owl pellet bones uncovers hidden biodiversity. *Syst. Biodivers.* **13**(4): 403-412. doi: 10.1080/14772000.2015.1044048.
- Rodzuan, S.N. 2006. Diversity of small mammals of Ulu Gombak Forest Reserve and Kenaboi Forest Reserve. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- Rovero, F., Zimmermann, F., Berzi, D., and Meek, P. 2013. "Which camera trap type and how many do I need?" A review of camera features and study designs for a range of wildlife research applications. *Hystrix* **24**: 1-9.
- Rowcliffe, J.M., Field, J., Turvey, S.T. and Carbone, C. 2008. Estimating animal density using camera traps without the need for individual recognition. *J. Appl. Ecol.* **45**: 1228-1236. doi: 10.1111/j.1365-2664.2008.01473.x.
- Sanderson, J.G., and Trolle, M. 2005. Monitoring elusive mammals. *Am. Sci.* **93**: 148-155.

- Schmieder, R. and Edwards, R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* **27**(6): 863-864.
- Schnell, I.B., Sollmann, R., Calvignac-Spencer, S., Siddall, M.E., Yu, D.W., Wilting, A., and Gilbert, M.T.P. 2015. iDNA from terrestrial haematophagous leeches as a wildlife surveying and monitoring tool - prospects, pitfalls and avenues to be developed. *Frontiers Zool.* **12**: 24.
- Schnell, I.B., Thomsen, P.F., Wilkinson, N., Rasmussen, M., Jensen, L.R., Willerslev, E., Bertelsen, M.F., and Gilbert, M.T. 2012. Screening mammal biodiversity using DNA from leeches. *Curr. Biol.* **22**: R262-R263.
- Schnur, H.J., Zivotofsky, D., and Wilamowski, A. 2009. Myiasis in domestic animals in Israel. *Vet. Parasitol.* **161**(3): 352-355.
- Sharma, D.S.K., Siong, B.L.M., Mohd Azlan, J., and Ahmad Zafir, A.W. 2005. Living in harmony: resolving conflicts between people and tigers and FELDA Jerangau Barat, Peninsular Malaysia. WWF-Malaysia, Petaling Jaya, Malaysia.
- Shen, Y.Y., Chen, X., and Murphy, R.W. 2013. Assessing DNA barcoding as a tool for species identification and data quality control. *PLoS ONE* **8**(2): e57125.
- Shokralla, S., Spall, J.L., Gibson, J.F., and Hajibabaei, M. 2012. Next-generation sequencing technologies for environmental DNA research. *Mol. Ecol.* **21**: 1794-1805.
- Silveira, L., Jácomo, A.T.A., and Diniz-Filho, J.A.F. 2003. Camera trap, line transect census and track surveys: a comparative evaluation. *Biol. Cons.* **114**: 351-355.
- Sing, K.W., Syaripuddin, K., and Wilson, J.J. 2013. Changing perspectives on the diversity of bats (Chiroptera) at Ulu Gombak since the establishment of the Field Studies Centre in 1965. *Raffles Bull. Zool.* **S29**: 211-217.
- Sly, G.R. 1975. Second record of the Bronzed Tube-nosed Bat (*Murina aenea*) in Peninsular Malaysia. *Malay. Nat. J.* **28**: 217.
- Smith, K.E., and Wall, R. (1998) Estimates of population density and dispersal in the blowfly *Lucilia sericata* (Diptera: Calliphoridae). *Bull. Entomol. Res.* **88**: 65-73.
doi:10.1017/S0007485300041560.

- Soberón, J.M., Llorente, J.B., Onate, L., 2000. The use of specimen label databases for conservation purposes: an example using Mexican Papilionid and Pierid butterflies. *Biodivers. Conserv.* **9**: 1441-1466.
- Souza, D.P., Asfora, P.H., Lira, T.C., and Astúa, D. 2010. Small mammals in Barn Owl (*Tyto alba* Aves, Strigiformes) pellets from northeastern Brazil, with new records of *Gracilinanus* and *Cryptonanus* (Didelphimorphia, Didelphidae). *Mammal. Biol.* **75**: 370-374.
- Spradbery, J.P., and Vanniasingham, J.A. 1980. Incidence of the screw-worm fly, *Chrysomya bezziana*, at the Zoo Negara, Malaysia. *Malaya Vet. J.* **7**: 28-32.
- Stanton, M.A., and Anderson, J.A. 1998. A survey of ground-dwelling mammals inhabiting forests of the southwestern slopes, New South Wales. *Aust. Zool.* **30**(4): 480.
- Stoner, K.E., and Timm, R.M. 2004. Tropical dry-forest mammals of Palo Verde: Ecology and conservation in a changing landscape. *In* Biodiversity Conservation in Costa Rica: Learning the lessons in a seasonal dry forest. *Edited by* G.W. Frankie, A. Mata, and V.S. Bradleigh. University of California Press, USA.
- Syakirah, S., Zubaid, A., Prentice, C., Lopez, A., Azmin, M.R., and Mohd Yusof, A. 2000. A small mammal survey at Tasek Bera, Pahang, Malaysia's first Ramsar site. *Malay. Nat. J.* **54**(1): 31-41.
- Syaripuddin, K. 2012. Study of diversity of bat at three different elevations using two different techniques. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- Syaripuddin, K., Kumar, A., Sing, K.W., Halim, M.R.A., Nursyereen, M.N., and Wilson, J.J. 2014. Mercury accumulation in bats near hydroelectric reservoirs in Peninsular Malaysia. *Ecotoxicology* **23**(7): 1164-71.
- Teta, P., González-Fischer, C.M., Codesido, M., and Bilenca, D.N. 2010. A contribution from Barn Owl pellets analysis to known micromammalian distributions in Buenos Aires province, Argentina. *Mammalia* **74**: 97-103.
- Thomsen, P.T., and Willerslev, E. 2015. Environmental DNA - an emerging tool in conservation for monitoring past and present biodiversity. *Biol. Conserv.* **183**: 4-18.

- Tidemann, C.R., and Woodside, D.P. 1978. A collapsible bat-trap and a comparison of results obtained with the trap and with mistnets. *Aust. Wildl. Res.* **5**: 355-362.
- Tingga, R.C.T., Anwarali, F.A.K., Mohd Ridwan, A.R., Senawi, J., and Abdullah, M.T. 2012. Small mammals from Kuala Atok-Taman Negara, Pahang. *Sains Malays.* **41**(6): 659–669.
- Tobler, M. 2012. Camera Base. Available from <http://www.atrium-biodiversity.org/tools/camerabase>. [accessed 8 July 2015].
- Torre, I., Arrizabalaga, A., and Flaquer, C. 2004. Three methods for assessing richness and composition of small mammal communities. *J. Mammal.* **85**: 524-530.
- Torre, I., Guixe, D., and Sort, F. 2010. Comparing three live trapping methods for small mammal sampling in cultivated areas of Ne Spain. *Hystrix* **21**(2): 147-155.
- Trolle, M. and Kery, M. 2003. Estimation of ocelot density in the Pantanal using capture–recapture analysis of camera trapping data. *J. Mammal.* **84**: 607-614.
- Tsuda, Y., Hayashi, T., Higa, Y., Hoshino, K., Kasai, S., Tomita, T., Kurahashi, H., and Kobayashi, M. 2009. Dispersal of a blow fly, *Celliphore nigribesis*, in relation to the dissemination of highly pathogenic avian influenza virus. *Jpn. J. Infect. Dis.* **62**: 294-297.
- Valderrama, X., Karesh, W.B., Wildman, D.E., and Melnick, D.J. 1999. Noninvasive methods for collecting fresh hair tissue. *Mol. Ecol.* **8**(10): 1749-1750.
- Vynne, C., Baker, M.R., Breuer, Z.K., and Wasse, S.K. 2012. Factors influencing degradation of DNA and hormones in Maned wolf scat. *Anim. Conserv.* **15**: 184-194. doi:10.1111/j.1469-1795.2011.00503.x
- Wilson, J.J. 2012. DNA barcodes for insects. *In* DNA Barcodes: Methods and Protocols. Edited by W.J. Kress and D.L. Erickson. Springer, New York. pp. 17-46.
- Wilson, J.J., Rougerie, R., Schonfeld, J., Janzen, D.H., Hallwachs, W., Hajibabaei, M., Kitching, I.J., Haxaire, J., and Hebert, P.D.N. 2011. When species matches are unavailable are DNA barcodes correctly assigned to higher taxa? An assessment using sphingid moths. *BMC Ecol.* **11**: 18.

- Wilson, J.J., Sing, K.W., Halim, M.R.A., Ramli, R., Hashim, R., and Sofian-Azirun, M. 2014. Utility of DNA barcoding for rapid and accurate assessment of bat diversity in Malaysia in the absence of formally described species. *Genet. Mol. Res.* **13**(1): 920-925.
- Wilting, A., Courtiol, A., Christiansen, P., Niedballa, J., Scharf, A.K., Orlando, L., Balkenhol N, Hofer, H., Kramer-Schadt, S., Fickel, J., and Kitchener, A.C. 2015. Planning tiger recovery: Understanding intraspecific variation for effective conservation. *Sci. Adv.* **1**(5): e1400175.
- Woods, J.G., Paetkau, D., Lewis, D., McLellan, B.N., Proctor, M., and Strobeck, C. 1999. Genetic tagging of free-ranging black and brown bears. *Wildl. Soc. Bull.* **27**: 616-627.
- Xiao, J.H., Wang, N.X., Li, Y.W., Murphy, R.W., Wan, D.G., Niu, L.M. et al. 2010. Molecular approaches to identify cryptic species and polymorphic species within a complex community of fig wasps. *PLoS ONE* **5**(11): e15067.
doi:10.1371/journal.pone.0015067
- Yenbutra, S., and Felten, H. 1983. A new species of the fruit bat genus *Megaerops* from SE-Asia (Mammalia: Chiroptera: Pteropodidae). *Senck. Biol.* **64**(1-3): 1-11.
- Yusof, M.D.N.N. 2005. Kajian kepelbagaian mamalia kecil di Hutan Simpan Ulu Gombak. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.
- Zeale, M.R.K., Butlin, R.K., Barker, G.L.A., Lees, D.C., and Jones, G. 2011. Taxon-specific PCR for DNA barcoding arthropod prey in bat faeces. *Mol. Ecol. Res.* **11**: 236-244.
- Zulkifli, N.M. 2011. Distribution and abundance of rat communities at different elevations. B.Sc. thesis, Institute of Biological Sciences, University of Malaya, Kuala Lumpur.

Table 1. Comparison of different field methods including targeted mammal groups, advantages, disadvantages and cost.

Field method	Targeted mammal groups	Advantages	Disadvantages	Cost
Cage traps	Rodents, insectivores (Catling et al. 1997)	Operates 24 h without supervision (Catling et al. 1997)	Difficult to standardise (varying dimensions, baited/un-baited and deployment location effects) (Fontúrbel 2010; Torre et al. 2010)	Medium (Garden et al. 2007)
Mist nets	Frugivorous bats (Stoner & Timm 2004)	Readily portable (Kuenzi & Morrison 1998)	Must be monitored constantly, as bats can become easily entangled and must be freed individually (Kuenzi & Morrison 1998)	Low (Kuenzi & Morrison 1998)
Harp traps	Insectivorous bats (Kingston et al. 2003)	Does not require constant monitoring (Tidemann & Woodside 1978)	Bulky and not easy to transport (Tidemann & Woodside 1978)	High (Tidemann & Woodside 1978)
Camera traps	Medium to large-bodied mammals (Bernard et al. 2013)	Effective in detecting species rarely recorded from live traps or direct	May under-represent species with specific habitats and unable to	High (Sanderson & Trolle 2005)

		observations (e.g. Hose's civet, <i>Diplogale hosei</i> ; Bernard et al. 2013)	distinguish closely related species (e.g. muntjac and mouse-deers; Bernard et al. 2013)	
Indirect signs	Medium to large-bodied ground dwelling mammals (Catling et al. 1997).	Effective in detecting species inhabiting open areas (e.g. otters and ungulates; Catling et al. 1997)	Imprecise in species identification (Davison et al. 2006; Mumma et al. 2014); Accuracy and precision are dependent on field conditions and expertise of identifiers (Silveira et al. 2003)	Low (Garden et al. 2007)
Interviews	Medium to large-bodied mammals (Mohd-Azlan et al. 2013)	No equipment required and can provide important collateral data (Mohd-Azlan et al. 2013)	Less reliable due to varying survey design and bias produced by respondents such as different recalling ability of respondents for different periods of time (Meijaard et al. 2011) (e.g. most villagers	Low (Mohd-Azlan et al. 2013)

			interviewed could not recall how many tigers have been killed due to livestock depredation by tigers; Sharma et al. 2005)	
Direct observations	Medium to large-bodied mammals that are readily observable (Gese et al. 2001)	Can confirm species presence directly and effectively detect rare and endangered species (e.g. black-footed ferrets, <i>Mustela nigripes</i> ; Gese et al. 2001)	The detection success relies much on the protocols used and expertise of identifiers (Hoppe-Dominik et al. 2011; Roberts 2011)	Low (Garden et al. 2007)
Hair traps	All ground-dwelling mammals (Stanton & Anderson 1998; Castro-Arellano et al. 2008)	Portable (Castro-Arellano et al. 2008); Reliable in identifying species through genetic analysis (Beja Pereira et al. 2009; Mullins et al. 2009)	Different trap types used could influence the detection success (Castro-Arellano et al. 2008) such as the use of hair-traps with mechanical devices like barb wire or attached glue-like substances that rarely collect high	Low (Castro-Arellano et al. 2008)

			proportions of hairs with roots (Valderrama et al. 1999; Ebert et al. 2010).	
Invertebrate-derived mammal DNA	All mammals (Calvignac-Spencer et al. 2012; Schnell et al. 2012; Calvignac-Spencer et al. 2013; Lee et al. 2015; Schnell et al. 2015)	Can accurately identify rare and cryptic species with least ecological and taxonomic expertise (Calvignac-Spencer et al. 2012; Schnell et al. 2012) (e.g. Annamite striped rabbit was recorded from DNA recovered from leeches in Vietnam; Schnell et al. 2012); Blowflies in particular have good potential as sources of mammal DNA due to their presence in all habitats and broad host preferences	Blowflies must be collected every 24 h as mammal DNA persists at least 24 h in blowfly guts (Lee et al. 2015)	High (Calvignac-Spencer et al. 2012; Calvignac-Spencer et al. 2013; Schnell et al. 2015)

(Norris 1965)				
Owl pellet bones-derived DNA	Small mammals (Bonvicino & Bezerra 2003; Torre et al. 2004; Souza et al. 2010; Teta et al. 2010; Rocha et al. 2015).	Effective in detecting and identifying small mammals including those with elusive behaviours and low densities (Rocha et al. 2015) (e.g. Two undescribed species of <i>Oecomys</i> were recorded from DNA recovered from owl pellet bones of barn owls; Rocha et al. 2015)	The age of pellet remains has an impact on the level of PCR inhibition (Buś et al. 2014); Easily degraded as non-invasive samples in tropical regions due to climatic conditions (e.g. Mukherjee <i>et al.</i> 2010; Vynne et al. 2012)	Low (Rocha et al. 2015)

Figure captions

Figure 1. The locations of field sites A) Ulu Gombak Forest Reserve showing the location of two transects and the experimental design for comparing five different field methods. B) Tembat Forest Reserve showing the locations of twenty trap stations each with a blowfly trap and a camera trap. Map data: Google.

Figure 2. Photographic evidence of nine mammal species detected by camera traps at Tembat Forest Reserve, Terengganu. A) *Elephas maximus* B) *Helarctos malayanus* C) *Macaca fascicularis* D) *Muntiacus muntjak* E) *Panthera pardus* F) *Prionailurus bengalensis* G) *Sus scrofa* H) *Tapirus indicus* I) *Viverra zibertha*. Camera settings: sample rate of passive infrared sensor = 70 times per s; flash brightness = normal (Xenon flash system); LED stealth mode = off; daylight images per trigger = 3 with delay period between images = 1 s; flash images per trigger during night time = 1 with delay period between images = 20 s; image resolution: 3 megapixel.

Figure 3. Species detected by different field methods including species unique to each method and species shared across different methods at A) Ulu Gombak Forest Reserve, and B) Tembat Forest Reserve. The number [in square parentheses] represents the BIN number assigned in BOLD to the species detected.

Figure 4. Rarefaction curves showing expected species richness (using Chao 1 with number of randomisations = 100) and sampling completeness ratios (in brackets) for each field method at A) Ulu Gombak Forest Reserve, and B) Tembat Forest Reserve. Dashed lines showed 95% confidence intervals of expected species richness for each field method at A) Ulu Gombak Forest Reserve: dashed blue lines for cage traps; dashed red lines for mist nets; dashed green lines for scat collection; dashed purple lines for blowfly traps, and B) Tembat Forest Reserve: dashed orange lines for camera traps; dashed black lines for blowfly traps.

Figure 1. The locations of field sites A) Ulu Gombak Forest Reserve showing the location of two transects and the experimental design for comparing five different field methods. B) Tembat Forest Reserve showing the locations of twenty trap stations each with a blowfly trap and a camera trap. Map data: Google.
190x254mm (300 x 300 DPI)

Figure 2. Photographic evidence of nine mammal species detected by camera traps at Tembat Forest Reserve, Terengganu. A) *Elephas maximus* B) *Helarctos malayanus* C) *Macaca fascicularis* D) *Muntiacus muntjak* E) *Panthera pardus* F) *Prionailurus bengalensis* G) *Sus scrofa* H) *Tapirus indicus* I) *Viverra zibertha*. Camera settings: sample rate of passive infrared sensor = 70 times per s; flash brightness = normal (Xenon flash system); LED stealth mode = off; daylight images per trigger = 3 with delay period between images = 1 s; flash images per trigger during night time = 1 with delay period between images = 20 s; image resolution: 3 megapixel. 102x77mm (300 x 300 DPI)

Figure 3. Species detected by different field methods including species unique to each method and species shared across different methods at A) Ulu Gombak Forest Reserve, and B) Tembat Forest Reserve. The number [in square parentheses] represents the BIN number assigned in BOLD to the species detected.

Figure 4. Rarefaction curves showing expected species richness (using Chao 1 with number of randomisations = 100) and sampling completeness ratios (in brackets) for each field method at A) Ulu Gombak Forest Reserve, and B) Tembat Forest Reserve. Dashed lines showed 95% confidence intervals of expected species richness for each field method at A) Ulu Gombak Forest Reserve: dashed blue lines for cage traps; dashed red lines for mist nets; dashed green lines for scat collection; dashed purple lines for blowfly traps, and B) Tembat Forest Reserve: dashed orange lines for camera traps; dashed black lines for blowfly traps.

142x107mm (300 x 300 DPI)

Table S1. Checklist of mammal species recorded at Ulu Gombak Forest Reserve.

	Sources
ORDER: CHIROPTERA	
PTEROPODIDAE	
<i>Balionycteris seimundi</i>	1,10,11,12,13
<i>Chironax melanocephalus</i> GOM01	1, 10, 11, 14
<i>Cynopterus brachyotis</i>	1,10,11,12,13,14
<i>Cynopterus horsfieldi</i>	1,3,10,11,12,13
<i>Cynopterus JLE sp. A</i>	14
<i>Dyacopterus spadiceus</i>	13
<i>Eonycteris spelaea</i>	1,10,11,13
<i>Macroglossus minimus</i>	1
<i>Megaerops ecaudatus</i>	9,11,13,14
<i>Penthetor lucasi</i>	1,10,11
<i>Pteropus vampyrus</i>	1,11
<i>Roussethus amplexicaudatus</i>	10,11,12
EMBALLONURIDAE	
<i>Emballonura monticola</i>	1,3,10,11
<i>Saccolaimus saccolaimus</i>	1,11
<i>Taphozous melanopogon</i>	10,11
NYCTERIDAE	
<i>Nycteris javanica</i> ^A	10,11
<i>Nycteris tragata</i> ^a	13
MEGADERMATIDAE	
<i>Megaderma lyra</i>	2
<i>Megaderma spasma</i>	1,10,11
RHINOLOPHIDAE	
<i>Rhinolophus affinis</i>	3,13
<i>Rhinolophus luctus</i>	1,10,11,13
<i>Rhinolophus refulgens</i>	11
<i>Rhinolophus sedulus</i>	1,3,10,11,13
<i>Rhinolophus stheno</i>	10,11,13
<i>Rhinolophus trifolius</i>	3,10,11,13
HIPPOSIDERIDAE	
<i>Coelops frithii</i>	5,11
<i>Hipposideros atrox</i>	14
<i>Hipposideros bicolor</i>	14
<i>Hipposideros cervinus</i>	8,10,11,13,14
<i>Hipposideros cineraceus</i>	1,3,11
<i>Hipposideros doriae</i>	10,11

<i>Hipposideros galeritus</i>	1
<i>Hipposideros larvatus</i>	1,11,13
<i>Hipposideros masoni</i>	1,3,10,11,13

VESPETILIONIDAE

<i>Arielulus circumdatus</i>	10,11
<i>Glischropus tylopus</i>	10,11,13
<i>Hesperoptenus blanfordi</i>	10,11
<i>Hesperoptenus doriae</i>	4,10,11
<i>Hesperoptenus tomesi</i>	10,11
<i>Kerivoula</i> sp.	1
<i>Miniopterus fuliginosus</i>	10,11
<i>Murina aenea</i>	7,11
<i>Murina peninsularis</i>	11,13
<i>Murina suilla</i>	10,11,13
<i>Myotis federatus</i>	3,10,11
<i>Myotis horsefieldii</i>	11
<i>Myotis muricola</i>	3,10,11
<i>Myotis mystacinus</i>	1
<i>Myotis ridleyi</i>	10,11
<i>Philetor brachypterus</i>	6,10,11,13
<i>Phoniscus atrox</i>	1,3,4,10,11
<i>Pipistrellus stenopterus</i>	1, 11
<i>Scotophilus kuhlii</i>	10,11
<i>Tylonycteris pachypus</i>	1,3,10,11
<i>Tylonycteris robustula</i>	1,10,11,13

MOLOSSIDAE

<i>Chaerephon</i> sp.	1,11
<i>Cheiromeles torquatus</i>	1,11

ORDER: DERMOPTERA**CYNOCEPHALIDAE**

<i>Cynocephalus variegatus</i>	1
--------------------------------	---

ORDER: ARTIODACTYLA**SUIDAE**

<i>Sus scrofa</i>	1
-------------------	---

TRAGULIDAE

<i>Tragulus javanicus</i>	1
---------------------------	---

BOVIDAE

<i>Capricornis sumatraensis</i>	1
---------------------------------	---

CERVIDAE		
<i>Muntiacus muntjak</i>		1
ORDER: INSECTIVORA		
SORICIDAE		
<i>Crocidura malayana</i>		15
<i>Crocidura monticola</i>		15
<i>Suncus malayanus</i>		15
ORDER: PHOLIDOTA		
MANIDAE		
<i>Manis javanica</i>		1
ORDER: PRIMATE		
CERCOPITHECIDAE		
<i>Presbytis melalophos</i>		1
<i>Presbytis obscura</i>		1
<i>Macaca fascicularis</i>		1
<i>Macaca nemestrina</i>		1
<i>Trachypithecus obscurus</i> [BOLD:AAI0540]		20
HYLOBATIDAE		
<i>Hylobates lar</i>		1
<i>Hylobates symphalangus</i>		1
ORDER: PROBOSCIDEA		
ELEPHANTIDAE		
<i>Elephas maximus</i>		1
ORDER: CARNIVORA		
FELIDAE		
<i>Panthera tigris</i>		1
<i>Panthera pardus</i>		1
<i>Felis bengalensis</i>		1
MUSTELIDAE		
<i>Martes flavigula</i>		1
VIVERRIDAE		
<i>Arctitis binturong</i>		1
<i>Arctogalidia trivirgata</i>		1
<i>Paradoxurus hermaphroditus</i>		1
<i>Paguma lawata</i>		1

ORDER: SCANDENTIA**TUPAIIDAE**

<i>Ptilocercus lowii</i>	1
<i>Tupaia glis</i>	1,16,17
<i>Tupaia minor</i>	1

ORDER: RODENTIA**HYSTRICIDAE**

<i>Atherurus macrourus</i>	1
<i>Hystrix brachyura</i>	1

MURIDAE

<i>Chiropodomys gliroides</i>	1,16,17
<i>Leopoldamys sabanus</i>	1,16,17,18
<i>Maxomys inas</i>	1,18,19
<i>Maxomys rajah</i>	16,17,18,19
<i>Maxomys surifer</i>	1,18
<i>Maxomys whiteheadi</i>	1,16,17,18
<i>Niviventer rapit</i>	16,17
<i>Rattus bowersi</i>	1,16,17,18,19
<i>Rattus exulans</i>	1,17,18
<i>Rattus norvegicus</i>	18
<i>Rattus rattus</i>	18,19
<i>Rattus tiomanicus</i>	1,16,17,18,19
<i>Sundamys muelleri</i>	1,16,17,18,19

SCIURIDAE

<i>Callosciurus notatus</i>	16,17
-----------------------------	-------

SPALACIDAE

<i>Rhizomys sumatrensis</i>	1
-----------------------------	---

Species names with same alphabetical superscript have been considered by some researchers to be the same species or synonyms, in such cases, the capital letters are used to denote the valid name.

Sources: 1. Medway 1966; 2. Medway 1967; 3. University of Malaya, Kuala Lumpur collection 1963-1969; 4. Medway 1983; 5. Hill 1972; 6. Hill 1974; 7. Sly 1975; 8. Jenkins & Hill 1981; 9. Yenbutra & Felten 1983; 10. Heller & Volleth 1989; 11. Heller & Volleth 1995; 12. Yusof 2005; 13. Syaripuddin 2012; 14. Sing et al. 2013; 15. Aziz 2010; 16. Hassan 2009. 17. Rodzuan 2006. 18. Zulkifli 2011. 19. Abu 2007; 20. In this study