Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

Responses of invasive silver and bighead carp to a carbon dioxide barrier in outdoor ponds

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2015-0472.R2
Manuscript Type:	Article
Date Submitted by the Author:	07-Jun-2016
Complete List of Authors:	Cupp, Aaron; USGS Upper Midwest Environmental Science Center Erickson, Richard A.; Upper Midwest Environmental Sciences Center Fredricks, Kim; USGS Upper Midwest Environmental Science Center Swyers, Nicholas; USGS Western Fisheries Research Center Hatton, Tyson; USGS Western Fisheries Research Center Amberg, Jon; USGS Upper Midwest Environmental Science Center
Keyword:	INVASIVE SPECIES < Organisms, Asian carp, Carbon Dioxide, Barrier

SCHOLARONE™
Manuscripts

Title: Responses of invasive silver and bighead carp to a carbon dioxide barrier in outdoor ponds

Proposed outlet: Canadian Journal of Fisheries and Aquatic Sciences

Author list: Aaron R. Cupp^{1*}, Richard A. Erickson¹, Kim T. Fredricks¹, Nicholas M. Swyers², Tyson W. Hatton², Jon J. Amberg¹

Author email address: acupp@usgs.gov, rerickson@usgs.gov, kfredricks@usgs.gov, nswyers@usgs.gov, thatton@usgs.gov, jamberg@usgs.gov

Author affiliation:

¹U.S. Geological Survey, Upper Midwest Environmental Sciences Center, 2630 Fanta Reed Rd., La Crosse, WI 54603, USA

²U.S. Geological Survey, Western Fisheries Research Center, 5501A Cook-Underwood Road, Cook, Washington 98605, USA

***Corresponding author:** Aaron R. Cupp, Telephone: (608)781-6266, Fax: (608)783-6066, acupp@usgs.gov, 2630 Fanta Reed Rd., La Crosse, WI 54603, USA

Manuscript is submitted to Canadian Journal of Fisheries and Aquatic Sciences as an original research paper.

Abstract

Resource managers need for effective methods to prevent the movement of silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) from the Mississippi River basin into the Laurentian Great Lakes. In this study, we evaluated dissolved carbon dioxide (CO₂) as a barrier and deterrent to silver (278 ± 30.5 mm) and bighead (212 ± 7.7 mm) carp movement in continuous-flow outdoor ponds. As a barrier, CO₂ significantly reduced upstream movement but was not 100% effective at blocking fish passage. As a deterrent, we observed a significant shift away from areas of high CO₂ relative to normal movement before and after injection. Carbon dioxide concentrations varied across the pond during injection and reached maximum concentrations of 74.5±1.9 mg/L CO₂; 29 532 – 41 393 µatm at the site of injection during three independent trials. We conclude that CO₂ altered silver and bighead carp movement in outdoor ponds and recommend further research to determine barrier effectiveness during field applications.

Keywords

Carbon dioxide; Asian carp; invasive species; barrier

Introduction

Range expansion of the invasive silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) throughout the Mississippi River basin has been documented by resource agencies since the 1970's (Kelly et al. 2011). These prolific fish cause harm to native aquatic ecosystems by competing with native mussels, early life-stage fish, and other filter-feeding fish for food resources from a planktivorous diet overlap (Irons et al. 2007; Sampson et al. 2009; Walleaser et al. 2014). This increased competition has been demonstrated through major shifts in zooplankton communities on the Illinois River where silver and bighead carp are in high abundances (Sass et al. 2014). Recent detections using eDNA and physical captures place the upper range of these invasive carp at tributaries adjacent to the southern Great Lakes (Jerde et al. 2013; Parker et al. 2015). With direct hydrological connection of these tributaries to the Great Lakes, research evaluating potential barriers to prevent further upstream movement has become increasingly important.

Physical barriers are not an option on many navigational rivers due to economic losses from the movement of goods and services. Closure of navigational locks for invasive species control on the Chicago Area Waterway System (CAWS), the shipping channel connecting the Mississippi River basin to the Great Lakes basin, was estimated to cost consumers and commercial traffic \$1.3 billion USD annually (Schwieterman 2010). With obvious economic concerns, resource managers have started exploring non-physical barriers that aim to prevent invasive species movement while concurrently allowing water-body navigation (Noatch and Suski 2012).

Fisheries professionals have described a number of potential non-physical strategies for preventing the spread of invasive carps in North America, but most agree that a general lack of data addressing the effectiveness of these techniques will ultimately delay management actions (Wittman et al. 2014). In response, recent studies have begun to address this gap in published literature. Experiments using acoustic barriers, bubble barriers, hydro-guns, carbon dioxide (CO₂), and electricity have all shown

56 varying levels to efficacy to deter invasive carps (Taylor et al. 2005; Kates et al. 2012; Ruebush et al.
57 2012; Romine et al. 2015). Realizing the need for continued research, the Asian Carp Regional
58 Coordination Committee (ACRCC) prioritized promising technologies (ACRCC MRW 2014) and dissolved
59 CO₂ was identified as a strategy that requires further testing.

60 A review of non-physical barriers by Noatch and Suski (2012) highlights the potential for CO₂ to
61 work as a barrier for invasive fishes. Further, reviews by Ishimatsu et al. (2005) and Perry and Abdallah
62 (2012) describe a suite of negative physiological responses by fish when exposed to hypercapnia. Most
63 fish sense CO₂ through chemoreceptors located on the gills (Ishimatsu et al. 2005; Perry and Abdallah
64 2012) and the potential to exploit CO₂ sensitivity in fish suggests that this technique may also work to
65 deter invasive carp movement (Noatch and Suski 2012). In laboratory tests with silver carp, Kates et al
66 (2012) observed avoidance behavior and irregular movements at 70 – 90 mg/L CO₂. Further, recent tests
67 in a static pond confirmed that silver and bighead carp avoided approximately 60 mg/L CO₂ (Donaldson
68 et al. 2016). With evidence growing, the effectiveness of CO₂ as a barrier to invasive carp movement
69 needs to be tested.

70 The objective for our study was to evaluate silver and bighead carp movement in outdoor ponds
71 in response to a CO₂ barrier over three independent trials. We hypothesized that CO₂ would reduce
72 upstream passage and subsequently deter fish to seek refuge in areas of lower CO₂. Barrier effectiveness
73 was determined by quantifying upstream passage through the injection site. Deterrence effectiveness
74 was determined by quantifying shifts in fish locations relative to changes in CO₂. Using these behavioral
75 metrics, we compared fish movement collected before, during, and after CO₂ injection. Results are
76 focused on the applicability of CO₂ to deter silver and bighead carp passage with implications for
77 reducing the risk transfer between the Mississippi River and Great Lakes basins.

79 **Materials and Methods**

Study Animals – We obtained silver carp ($n = 15$; total length: 278 ± 30.5 mm; wet weight: 229.2 ± 81.9 g; mean $\pm$ standard deviation) and bighead carp ($n = 15$; total length: 212 ± 7.7 mm; wet weight: 101.4 ± 12.3 g; mean $\pm$ standard deviation) from the Missouri River near Columbia, Missouri, USA and a private aquaculture farm in Osage Beach, Missouri, USA . Fish were held at the U.S. Geological Survey (USGS) Upper Midwest Environmental Sciences Center (UMESC) Invasive Species Complex (La Crosse, WI USA) until the time of testing. During the week preceding each trial, five bighead and five silver carp were removed from the 1500-L source tanks and placed in a single 130-L flow-through tank at $12 \pm 1^\circ\text{C}$ where diet was converted from dry feed to an algal feed mixture. Algal mix contained equal parts of 3.6 g/L Chlorella (Yaeyama Shokusan Co. Ltd., Shiraho, Ishigaki Island, Japan) and Spirulina (Stakich, Inc., Royal Oak, MI, USA) and was the same feed administered during the outdoor trials. Approval from UMESC Animal Care and Use Committee (Protocol Number: AEH-12-PPTAC-01), using similar guidelines as described in Guide for the Care and Use of Laboratory Animals (1996), was gained before study initiation.

Pond Setup - Trials were conducted in outdoor concrete ponds at UMESC. Pond dimensions were 10.0 m long x 4.9 m wide x 1.2 m deep with a volume of approximately 58 700 L. Influent water was supplied directly to the pond from two on-site wells and effluent water was discharged from the pond by spilling over at the top of the opposite wall. Setup was similar to a large choice chamber, with the pond divided into two symmetrical halves by a partial concrete block wall and a 3.1 m opening at the effluent end (Figure 1). Each half contained similar water volumes and a continuous supply of inflowing water (300 L/min) down each side. Wood lattice was affixed to all pond walls, including the brick partition, to reduce reflection of acoustic telemetry signals during fish tracking.

Three micro-bubble diffusers (Model: Point Four™ MBD 300, Pentair Aquatic Eco-Systems™, Apopka, FL, USA) were placed at an equal distance across the opening to one half of the pond and were

used to inject CO₂ into the pond. To evaluate changes in fish locations, the pond was divided into two areas (Figure 1). The side where CO₂ was injected was termed “High CO₂ Area” and the opposite side “Low CO₂ Area”. Names for each area also describe relative CO₂ concentrations found during CO₂ injection.

For water chemistry sampling, plastic tubing (4.8 mm ID, Python Products Inc., Milwaukee, WI, USA) was run from six locations near the pond bottom to a peristaltic pump (Model 7553-70, Cole-Parmer®, Vernon Hills, IL, USA) outside of the pond area to prevent the potential influence of sample collection on fish behavior (Figure 1). All tubing was placed before fish were stocked into the pond. In the High CO₂ Area, shade was created near the inflowing water as refuge to entice carp to pass through the area of elevated CO₂. Once-daily, 15 L of the algal feed mixture was injected using a small water pump (Little Giant®, Oklahoma City, OK, USA) under the shaded area and served as an additional attractant for fish to cross the High CO₂ area. Diffusers, shade and feeding were placed on the west half of the pond during Trials 1 and 2. Trial 3 was run as a mirror-image of the first two trials; with diffuser, shade, and feeding moved to the opposite (east) pond half. Study events were identical for all three trials.

Acoustic Telemetry - Fish positions were continuously monitored and recorded using an acoustic telemetry array. One Hydroacoustic Technology Inc. Model 290 acoustic tag receiver (HTI, Seattle, WA, USA) was connected to 16 individual HTI Model 590-series hydrophones. Hydrophones were top-mounted at the water’s surface by suspending each hydrophone from a grid of cable strung above the pond supported with small foam floats. Locations for each hydrophone were manually surveyed from an origin at the bottom of the southwest pond corner. All hydrophones were secured in the same fashion.

Ten modified HTI Model 795LD acoustic tags were used for each trial. To eliminate the need for surgical implantation of acoustic tags, tags were modified to allow for external attachment using a T-bar

whisker (Floy Tag, Seattle, WA, USA) from methods described in Romine et al. (2015). Modifications included a T-bar whisker, foam insulation (approximately 6 mm diameter) and an acoustic tag inserted into heat shrink tubing (BuyHeatShrink.com, Deerfield, FL, USA) with the ceramic end of the acoustic tag and T-bar whisker protruding from the tubing at the same end. Heat was carefully applied to the heat shrink tubing until all components were securely held in place. Modified tags were placed in water and the foam trimmed to achieve near neutral buoyancy. Individual tags were programmed using a HTI Model 490-LP Acoustic Tag Programmer (ATP) with ping rates ranging from 2059 to 2675 ms. Once activated, presence and persistence of pings were verified using the HTI Model 492-B Acoustic Tag Detector before attachment to fish.

During post-processing of acoustic data, valid signal returns were filtered from ambient noise and signal multipath using USGS-developed (public) software, FishCount. Position estimates were then created using USGS-developed (public) software, GeneticFish, from the previous valid detections. Logic filters (i.e., pond borders, distance between detections, velocity, vector angles, and absence of signal) were used to identify and remove any unreasonable position estimates. We determined the accuracy of our telemetry array to be < 0.2 m by comparing known tag locations to post-processed position estimates.

CO₂ Trials - Feed was withheld 48 h before transfer and tagging of fish in the outdoor pond as recommended when sedating fish. All fish were under light sedation using 50 – 100 mg/L AQUI-S®20E (5 – 10 mg/L active ingredient eugenol; AQUI-S New Zealand Ltd., Lower Hutt, New Zealand) during transfer (Cupp et al. *in press*). Modified acoustic tags were disinfected for 15 min using a solution of 3% (v/v) Nolvasan® (active ingredient: chlorohexidine diacetate [2%], Ford Dodge Animal Health, Ford Dodge, IA, USA) and rinsed with deionized water. Five bighead and five silver carp were used during each trial. Fish were removed from the sedative solution and the tag was inserted with a T-bar anchor

tagging gun at a 45° posterior angle into the dorsal musculature approximately 1 cm lateral and posterior to the mid-line of the dorsal fin. This allowed streamline tag protrusion while swimming and ensured that the T-end was securely implanted (Wydoski and Emergy 1983). All fish appeared to recover in <1 min from the light sedation when placed in the pond. However, fish were given until the following day to fully recover from sedation and handling. Water temperature during the stocking of fish was typical of UMESC well water at $12 \pm 1^\circ\text{C}$. During experimentation, recorded pond temperature was $13.0 \pm 3.0^\circ\text{C}$ (Trial 1), $11.3 \pm 0.4^\circ\text{C}$ (Trial 2), and $8.2 \pm 2.5^\circ\text{C}$ (Trial 3) corresponding with air temperatures at the time of testing. Pond water alkalinity measured throughout testing was $137 \pm 1 \text{ mg/L CaCO}_3$.

After recovery, the first trial began and fish positions were continuously recorded for 72 h. All trials were discretized into three sequential 24 h periods: (1) pre-; (2) during; and (3) post- CO_2 injection. No human influence on fish behavior occurred at any time during these 72h trials. Fish positions collected pre- and post- CO_2 were used to determine baseline occupancy and movement of fish throughout the pond in the absence of CO_2 . Study events across all three time periods were identical, except for the date of CO_2 injection. No stimuli (e.g. injection of compressed gas) was added during the pre- or post-injection.

During CO_2 injection, gas was supplied directly from the port on a single 180-L liquid CO_2 tank (Airgas Inc., La Crosse, WI, USA) connected to a manifold of three flow meters (AngelAqua®, Busan, Korea). Diffusers delivered CO_2 at a rate of 3 L/min. Two peristaltic pumps (Model 7553-70, Cole-Parmer®, Vernon Hills, IL, USA) continuously extracted water through the plastic tubing from six locations (Figure 1). Water samples were collected from pumps and analyzed immediately for carbon dioxide by a modified HACH® Method 8205 digital titration method using sodium hydroxide. Briefly, the 100 mL water sample was poured into a 500-mL glass-beaker and placed on a stir plate with a magnetic stir bar. Titrant ($0.3636 \pm 0.00200 \text{ N NaOH}$) was added to the sample using the HACH® Digital Titrator until an endpoint of pH 8.3 (Beckman Coulter Model 410 pH meter, Beckman Coulter Inc., Chaska, MN,

USA) was reached. Partial pressures of CO₂ were calculated from pH, temperature, and alkalinity using USGS CO2Calc (<http://pubs.usgs.gov/of/2010/1280/>). Alkalinity was measured using the pH 4.5 titrimetric method with 0.02 N H₂SO₄ at various times throughout all three trials from the water samples (APHA 1995). Atmospheric carbon dioxide levels were intermittently measured during each trial using a handheld pSense Model AZ-0001 meter (CO2 Meter Inc., Ormond Beach, FL, USA) to ensure human safety.

At the conclusion of the trial, the pond was drained and fish were euthanized using an overdose (200 mg/L) of MS-222. Trials were repeated using the same methods previously described, except for the injection site, shade, and feed being placed on the opposite pond half during Trial 3. All trials began at exactly 07:00 CDT and subsequent transitions between study phases occurred at exactly 24h thereafter. Trials were conducted October 15, 2014 to November 1, 2014.

Statistical Analysis –Exploratory data analysis approaches was used to visualize data (Tukey 1977). Fish locations were visualized using density plots for pre-, during and post-CO₂ injection. Fish location counts were also visualized for the same three time periods. Carbon dioxide and pH trends were visualized through time. Fish crossings per hour were visualized using box-and-whisker plots before and after CO₂ concentrations stabilized. All data visualization was done with the ggplot2 package (Wickham 2009) in R (R Core Team 2015). Confirmatory statistics were used for parameter estimation and to test the research hypotheses. Generalized linear mixed models (GLMM) with Poisson error terms were used for all statistical comparisons because the three exposure tanks created a nested experimental design and individual fish are pseudo-replicates (Bolker 2008). Trial number and fish ID were included as a random effect. Species and time period (pre, during, and post- injection) were included as fixed effects. For the crossing per hour per fish GLMM, trial had an interaction with CO₂ concentration within the

model and was included as a fixed effect. The glmPQL package from the MASS (Venables and Ripley 2002) in R was used for all GLMM's. Statistical significance was declared at $\alpha < 0.05$.

Results

Water Chemistry – CO₂ was stable at ambient levels before and after injection (range: 3.8 - 7.6 mg/L CO₂; 1 776 – 4 688 μ atm). During injection, CO₂ increased and pH decreased (Figures 2-3). Highest CO₂ concentrations (max = 74.5 $\pm$ 1.9 mg/L; 29 532 – 41 393 μ atm) were found at the injection site relative to the rest of the pond. Lowest CO₂ concentrations after 24 h of injection (max: 34.1 $\pm$ 8.6 mg/L; 16 683 – 20 352 μ atm), were found farthest from injection and nearest to the inflowing fresh water. CO₂ stabilized at maximum levels approximately 4 – 6 h after injection began (Figure 2).

Barrier and Deterrence Efficacy– Signal transmission stopped from one tag in Trial 2 (n=9) and two tags from Trial 3 (n=8) early in each trial and those tags were subsequently removed from the dataset. No signals were lost during Trial 1 (n=10). Physical retention of tags by fish was 100% at the completion of all three trials. After applying filters to the raw telemetry data, 679 248 (Trial 1), 777 807 (Trial 2), and 637 665 (Trial 3) valid fish positions were included in analyses. There were no mortalities during any of the trials.

Silver and bighead carp were found significantly more times in the High CO₂ Area during pre- ($t = 8.40$, $p < 0.0001$, $df = 52$) and post-injection ($t = 7.20$, $p < 0.0001$, $df = 52$) compared to when CO₂ was injected (Figure 4). Detections in the High CO₂ Area did not differ between species ($t = -1.329$, $p = 0.19$, $df = 23$). Conversely, silver and bighead carp were detected significantly more times in the Low CO₂ Area during CO₂ injection compared to pre- ($t = -20.63$, $p < 0.0001$, $df = 52$) and post-injection ($t = -20.80$, $p < 0.0001$, $df = 52$; Figure 4). This did not differ by species ($t = -0.444$, $p = 0.66$, $df = 23$). Silver and bighead

223 carp crossed upstream through the injection site significantly fewer times during CO₂ injection
224 compared to pre- ($t = 14.46$, $p < 0.0001$, $df = 52$) and post-injection ($t = 6.76$, $p < 0.0001$, $df = 52$; Figure
225 4). Density plots visualized the change in fish locations for all three trials (Figures 5,6,7). Fish crossed the
226 injection site in both directions and there were no instances where a single fish remained upstream of
227 the CO₂ injection site. Silver carp crossed fewer times than bighead carp ($t = -2.10$, $p = 0.046$, $df = 23$).
228 Carbon dioxide levels began to stabilize at the injection site around 60 mg/L CO₂ (Figure 2). When CO₂
229 levels were > 60 mg/L (approximately $> 24\,000$ μatm), silver and bighead carp passed through the
230 injection site significantly fewer times per hour relative to concentrations ≤ 60 mg/L CO₂ ($t = -9.22$, $p <$
231 0.0001 , $df = 652$; Figure 8). Silver carp also passed through fewer times than bighead carp ($t = -26.75$, $p =$
232 0.016 , $df = 16$). Trial 3 had the lowest number of crosses ($t = -6.48$, $p < 0.0001$, $df = 652$).

233

234 Discussion

235 The potential for CO₂ to function as a non-physical deterrent has been recently described in the
236 literature (Noatch and Suski 2012; Kates et al. 2012), and our study was the first to quantify the
237 effectiveness of a CO₂ barrier to invasive carps movement. We found that that CO₂ injection was
238 effective to reduce the upstream movement of silver and bighead carp, but did not completely stop fish
239 passage. Similarly, we observed a significant shift in fish movement away from areas of high CO₂. Most
240 notably, results were consistent across all three trials, including when CO₂ was injected into the opposite
241 side of the pond during Trial 3, providing evidence that CO₂ modified silver and bighead carp behavior.
242 This study demonstrates the potential use of CO₂ to deter silver and bighead carp movement and results
243 contribute to a growing body of research evaluating the use of CO₂ as a deterrent to invasive fishes.

244 Many fish species have been shown to avoid CO₂ during laboratory experiments. For instance,
245 Clingerman et al. (2007) used 60 – 120 mg/L dissolved CO₂ for non-physical transfer of rainbow trout
246 (*Oncorhynchus mykiss*) between large aquaculture tanks with 100 percent effectiveness. A similar

approach was conducted by Kates et al. (2012), where avoidance responses were documented from silver carp, bluegill (*Lepomis macrochirus*), and largemouth bass (*Micropterus salmoides*) at concentrations ≥ 100 mg/L CO₂. Further, blacknose dace (*Rhinichthys atratulus*) and brook trout (*Salvelinus fontinalis*) exhibited avoidance behavior when exposed to 120 mg/L CO₂ in small laboratory tanks (Ross et al. 2001). When in our outdoor pond, we observed similar avoidance behavior of silver and bighead carp at concentrations around 70 mg/L CO₂ and found that this technique could potentially be used to deter upstream movement.

The sensitivity of fish to elevated CO₂ warrants continued investigation into the use of this non-physical method. However, with any non-physical deterrence strategy, less than 100% deterrence must be an acceptable outcome (Noatch and Suski 2012; Wittman et al. 2014). During periods of hypercapnia, fish undergo physiological detriments such as ion imbalance (Claiborne et al. 2002), increased stress response (Kates et al. 2012), acidosis (Ishimatsu et al. 2005), hyperventilation (Perry and Abdallah 2012), hypoventilation (Kates et al. 2012), loss of sensory function (Nilsson et al. 2012), and changes in protein composition (Dennis et al. 2015). In conjunction with elevated CO₂, a corresponding decrease in pH occurs due to the hydration equilibrium of CO₂ to form bicarbonate (HCO₃⁻) and carbonic acid (H₂CO₃). This decrease in pH may play a role in behavioral avoidance during hypercapnia due to the influence of pH on biochemical processes, however, several studies have shown that fish specifically sense and avoid CO₂ (Ishimatsu et al. 2005; Perry and Gilmour 2006; Perry and Abdallah 2012).

Consideration should be given to the methods used to quantify and units reported of dissolved CO₂, for comparisons with previous literature. Our study quantified CO₂ using methods described in Clingerman et al. (2007), Kates et al. (2012), Dennis et al. (2015), and Donaldson et al. (2016) to allow direct comparisons of the results. Free CO₂ can be easily related with pH, which may also facilitate transition to the field using common water chemistry data loggers (Kates et al. 2012). However, CO₂ gas saturation is dependent on temperature, pressure, alkalinity, biological status and pH. Common units for

carbon dioxide are reported or derived in terms of pressure (e.g. PCO_2 , mmHg, μatm). While there is a general lack of uniformity in units quantifying CO_2 across current research, recognizing units are important for comparison of results.

Results from our study identify many of the consideration with CO_2 as a control tool in fisheries management. For example, the U.S. Army Corps of Engineers operates and maintains an electrical barrier in the CAWS near Romeoville, IL, USA to inhibit further upstream movement of silver and bighead carp into the Great Lakes basin (Moy et al. 2011). While the electric barrier is currently the most important management tool to deter upstream movement of carp in the Great Lakes, there is a need for redundancy and multiple integrated deterrence mechanisms (Noatch and Suski 2012). Additionally, Noatch and Suski (2012) expressed concern that electricity loses effectiveness on smaller fish due to a reduction in total body voltage at smaller body sizes (Reynolds 1996; Dolan and Miranda 2003). The potential for size-related effectiveness with the electric barrier further supports the need for alternative and supplemental deterrence strategies. In our study, CO_2 deterred stock length silver carp and sub-stock length bighead carp (Phelps and Willis 2013). With branchial CO_2 receptors present throughout the life-history of most fish (Perry and Abdallah 2012), the absence of size-related effectiveness suggests the CO_2 could be used to supplement other barriers to deter smaller fish passage.

Carbon dioxide may be useful to supplement other barriers (e.g. lock and dams or electrical barriers). During scheduled usage or maintenance, CO_2 could be beneficial to deter silver and bighead carp until primary barriers are restored. For instance, CO_2 could be applied to reduce fish from entering or passing gates during planned vessel lockage. One important observation during our study was that adequate time must be allowed for CO_2 to accumulate to target concentrations to be most effective at deterring fish (e.g. Figures 5,6,7). We observed 4 -6 h for CO_2 levels to stabilize using a simple diffusion injection system and recommend exploring improved designs to reduce build-up time. Reducing the build-up time would expand the utility of CO_2 under different applications. Further, understanding

operational efficiency of individual CO₂ injection systems under different scenarios will be important during rapid implementation (e.g. unexpected primary barrier failure) and for calculating operational costs. For rapid implementation, other deterrent strategies (e.g. mobile electrical barriers, sound-bubble-strobes, or acoustic noise) may be considered due to their on-and-off style of operation (Taylor et al. 2005; Ruebush et al. 2012; Vetter et al. 2015).

Carbon dioxide was not an absolute barrier to silver and bighead carp in our outdoor ponds. While the success of any barrier is binomial, it is possible that total crossings in our study may have been inflated due to the relatively small spatial size of our test pond. With recorded burst speeds of 1.28 – 1.66 m/s (Hoover et al. 2012), silver and bighead carp in our 10 m pond could have moved upstream through the injection site before sensing elevated CO₂. This hypothesis is further supported by the reduced time that fish spent in the High CO₂ Area. In our study, fish consistently avoided areas of elevated CO₂ when given an option of fresh-water. Further, when concentrations were greater than 60 mg/L CO₂ (~ 24 000 µatm) we also observed the most effective reduction in upstream movement relative to lower concentrations, suggesting a potential dose-response effectiveness (Figure 8). Higher concentrations may be considered for future research. Complementary to the statistical comparisons, visualized distributions of fish in the pond shown in Figures 5,6,7 demonstrate the clear shift in fish locations relative to CO₂ injection. Most notably, similar movement away from the injection site during Trial 3 when CO₂ was moved to the opposite side of the pond confirmed that CO₂ deterred fish. Testing on a larger spatial scale can likely address this shortcoming and application of CO₂ on a larger spatial scale may increase its effectiveness as a deterrent, especially when access to fresh water is made readily available. Locations such as the entrance to approach channels at lock and dams or tributaries on large rivers where fish can choose to stay out in the main channel should be considered for further testing.

Carbon dioxide may provide a safer alternative than other non-physical methods, primarily the electrical barrier in the CAWS. Measurements taken by the U. S. Coast Guard conclude that the

electrical barrier can result in a lethal shock to humans if they fall into the water near the barrier (ADRSC 2011). Carbon dioxide exists naturally in the atmosphere as a byproduct of aerobic respiration and burning of fossil fuels at concentrations of 0.038 - 0.04 percent (NOAA 2015). Acceptable human exposure limits of atmospheric CO₂ established by the Occupational Safety and Health Administration (OSHA) were set at 5 000 ppm as a time weighted average (TWA) for an 8 h exposure and 30 000 ppm for acute exposures (OSHA 29 CFR 1910.1000). Atmospheric CO₂ measured before and during our trials were stable at a range of 433 – 498 ppm and did not increase during periods of CO₂ injection. While these concentrations are well below levels generating safety concerns, adequate monitoring should be incorporated when CO₂ is applied at a larger field-scale.

Acknowledgements

Funding for this work was provided by the U.S. Environmental Protection Agency’s (EPA) Great Lakes Restoration Initiative (Agreement number DW14-92404001). Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References

- Acquisition Directorate Research and Development Center (ADRD). 2011. CSSC Fish barrier simulated rescuer touch point results, operating guidance, and recommendations for rescuer safety. Publication No: CG-D-01-12. Available from <http://www.dtic.mil/dtic/tr/fulltext/u2/a554514.pdf> [Accessed 15 April 2015].
- Asian Carp Regional Coordinating Committee Monitoring and Response Workgroup (ACRCC MRW). 2014. Monitoring and response plan for Asian carp in the upper Illinois River and Chicago Area Waterway System. Available from <http://www.asiancarp.us/documents/MRP2014.pdf> [Accessed 3 January 2015].
- American Public Health Association (APHA). 1995. Standard methods for the examination of water and wastewater. 19th ed. American Public Health Association, American Water Works Association, and Water Pollution Control Federation, Washington, D.C.
- Bolker, B.M., 2008. Ecological models and data in R, 508 edition. ed. Princeton University Press, Princeton, N.J.
- Claiborne, J.B., Edwards, S.L., and Morrison-Shetlar, A.I. 2002. Acid-base regulation in fishes: cellular and molecular mechanisms. J. Exp. Zool. **293**(3): 302-319. DOI: 10.1002/jez.10125.
- Clingerman, J., Bebak, J., Mazik, P.M., and Summerfelt, S.T. 2007. Use of avoidance response by rainbow trout to carbon dioxide for fish self-transfer between tanks. Aquacult. Eng. **37**(3): 234-251. DOI:10.1016/j.aquaeng.2007.07.001.
- Cupp, A.R., Fredricks, K.T., Porcher, S.T., Smerud, J.R., Hartleb, C.F., and Gaikowski, M.P. (*in press*). Survival and behavioral responses of cool and warm water fish sedated with AQUI-S® 20E (10% eugenol) at high loading densities. Aquacult. Res. DOI:10.1111/are.12992.

- 355 Dennis, C.E., Kates, D. F., Noatch, M.R. and Suski C.D. 2015. Molecular responses of fishes to elevated
356 carbon dioxide. *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* **187**: 224-231.
357 DOI:10.1016/j.cbpa.2014.05.013.
- 358 Dolan, C.R., and Miranda, L.E. 2003. Immobilization thresholds of electrofishing relative to fish size.
359 *Trans. Am. Fish. Soc.* **132**(5): 969-976. DOI:10.1577/T02-055.
- 360 Donaldson, M.R., Amberg, J., Adhikari, S., Cupp, A.R., Jensen, N., Romine, J., Wright, A., Gaikowski, M.,
361 and Suski, C.D. 2016. Carbon dioxide as a tool to deter the movement of invasive bigheaded
362 carps. *Trans. Am. Fish. Soc.* **145**: 657-670. DOI:10.1080/00028487.2016.1143397.
- 363 Guide for the care and use of laboratory animals. 1996. National Academy Press, Washington D.C.
- 364 Hoover, J. J., Southern, W., Katzenmeyer, A. W., and Hahn, N. M. 2012. Swimming performance of
365 bighead carp and silver carp: methodology, metrics and management applications. U.S. Army
366 Corps of Engineers, Aquatic Nuisance Species Research Program, Technical Notes Collection
367 ERDC/TN ANSRP-12-3, Vicksburg, Mississippi. Available from
368 <http://el.erdc.usace.army.mil/elpubs/pdf/ansrp12-3.pdf> [Accessed 7 March 2015].
- 369 Irons, K.S., Sass, G.G., McClelland, M.A., and Stafford, J.D. 2007. Reduced condition factor of two native
370 fish species coincident with invasion of non-native Asian carps in the Illinois River, U.S.A. Is this
371 evidence for competition and reduced fitness?. *J. Fish Biol.* **71**: 258-273. DOI: 10.1111/j.1095-
372 8649.2007.01670.x.
- 373 Ishimatsu, A., Hayashi, M., and Lee, K.S. 2005. Physiological effects on fishes in a high-CO₂ world. *J.*
374 *Geophys. Res.* **110**: 1-8. DOI:10.1029/2004JC002564.
- 375 Jerde, C.L., Chadderton, W.L., Mahon, A.R., Renshaw, M.A., Corush, J., Budny, M.L., Mysorekar, S., and
376 Lodge, D.M. 2013. Detection of Asian carp DNA as part of a Great Lakes basin-wide surveillance
377 program. *Can. J. Fish. Aquat. Sci.* **70**: 522-526. DOI.org/10.1139/cjfas-2012-0478.

- 378 Kates, D., Dennis, C., Noatch, M.R., and Suski, C.D. 2012. Responses of native and invasive fishes to
379 carbon dioxide: potential for a nonphysical barrier to fish dispersal. *Can. J. Fish. Aquat. Sci.* **69**:
380 1748-1759. DOI:10.1139/f2012-102.
- 381 Kelly, A.M., Engle, C.R., Armstrong, M.L., Freeze, M., and Mitchell, A.J. 2011. History of introductions and
382 governmental involvement in promoting the use of grass, silver and bighead carps. *In* *Invasive*
383 *Asian carps in North America. Edited by D.C Chapman and M.H. Hoff. American Fisheries Society,*
384 *Bethesda, Maryland. pp. 163-174.*
- 385 Moy, P.B., Polls, I., and Dettmers, J.M. 2011. The Chicago Sanitary and Ship Canal aquatic nuisance
386 species dispersal barrier. *In* *Invasive Asian carps in North America. Edited by D.C Chapman and*
387 *M.H. Hoff. American Fisheries Society, Bethesda, Maryland. pp. 121-137.*
- 388 National Oceanic and Atmospheric Administration (NOAA). 2015. Earth System Research Laboratory –
389 Global Monitoring Division: Global Greenhouse Gas Reference Network. Available from
390 <http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html> [Accessed 7 June 2015].
- 391 Nilsson, G.E., Dixon, D.L., Domenici, P., McCormick, M.I., Sorenson, C., Watson, S., and Munday, P.L.
392 2012. Near-future carbon dioxide levels alter fish behavior by interfering with neurotransmitter
393 function. *Nat. Clim. Chan.* **2**: 201-204. DOI: 10.1038/NCLIMATE1352.
- 394 Noatch, M.R., and Suski, C.D. 2012. Non-physical barriers to deter fish movements. *Env. Rev.* **20**: 1-12.
395 DOI:10.1139/A2012-001.
- 396 Occupational Safety and Health Administration (OSHA). 29 CFR 1910.1000. Available from
397 <http://www.osha.gov/dts/chemicalsampling/data> [Accessed 10 June 2015].
- 398 Parker, A.D., Glover, D.C., Finney, S.T., Rogers, P.B., Stewart, J.G., and Simmonds, R.L. 2015. Direct
399 observations of fish incapacitation rates at a large electrical fish barrier in the Chicago Sanitary
400 and Ship Canal. *J. Gr. Lak. Res.* **41**: 396-404. DOI.org/10.1016/j.jglr.2015.03.004.

- 401 Perry, S.F., and Abdallah, S. 2012. Mechanisms and consequences of carbon dioxide sensing in fish. *Resp.*
402 *Physiol. Neurobiol.* **184**: 309-315. DOI.org/10.1016/j.resp.2012.06.013.
- 403 Perry, S.F., and Gilmour, K.M. 2006. Acid-base balance and CO₂ excretion in fish: unanswered questions
404 and emerging models. *Resp. Physiol. Neurobiol.* **154**: 199-215. DOI:10.1016/j.resp.2006.04.010.
- 405 Phelps, Q.E., and Willis, D.W. 2013. Development of an Asian carp size structure index and application
406 through demonstration. *N. Am. J. Fish. Manag.* **33**: 338-343. DOI:
407 10.1080/02755947.2012.760506.
- 408 R Core Team, 2015. R: A language and environment for statistical computing. R Foundation for
409 Statistical Computing, Vienna, Austria. R Foundation for Statistical Computing, Vienna, Austria.
- 410 Reynolds, J.B. 1996. Electrofishing. *In* Fisheries Techniques 2nd edition. *Edited by* B.R. Murphy and D.W.
411 Willis. American Fisheries Society, Bethesda, Maryland. pp. 221-253.
- 412 Romine, J.G., Jensen, N.R., Parsley, M.J., Gaugush, R.F., Severson, T.J., Hatton, T.W., Adams, R.F., and
413 Gaikowski, M.P. 2015. Response of bighead carp and silver carp to repeated water gun
414 operation in an enclosed shallow pond. *N. Am. J. Fish. Manag.* **35**: 440-453. DOI:
415 10.1080/02755947.2015.1012279.
- 416 Ross, R.M., Krise, W.F., Redell, L.A. and Bennett, R.M. 2001. Effects of dissolved carbon dioxide on the
417 physiology and behavior of fish in artificial streams. *Envir. Tox.* **16**: 84-95.
- 418 Ruebush, B.C., Sass, G.G., Chick, J.H., and Stafford, J.D. 2012. In-situ tests of sound-bubble-strobe light
419 barrier technologies to prevent expansions of Asian carp. *Aquat. Inv.* **7**: 37-48.
420 DOI:10.3391/ai.2012.7.1.005.
- 421 Sampson, S.J., Chick, J.H., and Pegg, M.A. 2009. Diet overlap among two Asian carp and three native
422 fishes in backwater lakes on the Illinois and Mississippi rivers. *Biol. Inv.* **11**:483-496. DOI
423 10.1007/s10530-008-9265-7.

- 424 Sass, G.G., Hinz, C., Erickson, A.C., McClelland, N.N., McClelland, M.A., and Epifanio, J.M. 2014. Invasive
425 bighead and silver carp effects on zooplankton communities in the Illinois River, Illinois, USA. J.
426 Gr. Lak. Res. **40**: 911-921. DOI: 10.1016/j.jglr.2014.08.010.
- 427 Schwieterman, J.P. 2010. An analysis of the economic effects of terminating operations at the Chicago
428 River controlling works and O'Brien Locks on the Chicago Area Waterway System. DePaul
429 University. Available from [http://www.unlockoutjobs.org/wp-](http://www.unlockoutjobs.org/wp-content/themes/unlockourjobs/pdf/DePaul_University_Study.pdf)
430 [content/themes/unlockourjobs/pdf/DePaul_University_Study.pdf](http://www.unlockoutjobs.org/wp-content/themes/unlockourjobs/pdf/DePaul_University_Study.pdf) [Accessed 27 April 2015].
- 431 Taylor, R.M., Pegg, M.A., and Chick, J.H. 2005. Response of bighead carp to a bioacoustics behavioral fish
432 guidance system. Fish. Manag. Ecol. **12**(4): 283-286. DOI: 10.1111/j.1365-2400.2005.00446.x.
- 433 Tukey, J.W., 1977. Exploratory data analysis, 1 edition. ed. Pearson, Reading, Mass.
- 434 Venables, W.N., Ripley, B.D., 2002. Modern applied statistics with S, Statistics and Computing. Springer
435 New York, New York, NY.
- 436 Vetter, B.J., Cupp, A.R., Fredricks, K.T., Gaikowski, M.P., and Mensinger, A.F. 2015. Acoustical
437 deterrents of silver carp. Biol. Inv. **17**: 3383-3392. DOI:10.1007/s10530-015-0964-6.
- 438 Walleser, L.W., Sandheinrich, M.B., Howard, D.R., Gaikowski, M.P., and Amberg, J.J. 2014. Spatial and
439 temporal variation of the gill rakers of gizzard shad and silver carp in three Midwestern rivers. N.
440 Am. J. Fish. Manag. **34**: 875-884. DOI: 10.1080/02755947.2014.920740.
- 441 Wickham, H., 2009. ggplot2: Elegant graphics for data analysis. Springer, New York.
- 442 Wittmann, M.E., Cooke, R.M., Rothlisberger, J.D., and Lodge, D.M. 2014. Using structured expert
443 judgment of assess invasive species prevention: Asian carp and the Mississippi-Great Lakes
444 hydrologic connection. Env. Sci. Tech. **48**: 2150-2156. DOI.org/10.1021/es4043098.
- 445 Wydoski, R., and Emergy, L. 1983. Tagging and marking. *In* Fisheries Techniques. Edited by L.A. Nielson
446 and D.L. Johnson. American Fisheries Society, Bethesda, Maryland. pp. 215-238.

Figure Captions

Figure 1: Outdoor concrete pond setup at the Upper Midwest Environmental Sciences Center in La Crosse, WI during the evaluation of CO₂ as a barrier and deterrent to silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) conducted October 15, 2014 – November 1, 2014. Graphical illustration shows setup during Trials 1 and 2. Trial 3 was a reversed setup, with CO₂ injection moved to the opposite side of the center partition. Black triangles show approximate pH and CO₂ spatial sampling locations within the pond: NW (northwest), MW (middle-west), SW (southwest), NE (northeast), ME (middle-east), and SE (southeast).

Figures 2,3: Carbon dioxide (CO₂) and pH levels in outdoor ponds during the evaluation of CO₂ as a barrier and deterrent to silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) conducted October 15, 2014 – November 1, 2014 at the Upper Midwest Environmental Sciences Center in La Crosse, WI. We observed increases in CO₂ levels (Figure 2) and decreases in pH (Figure 3) throughout the entire pond during the entire 24h that CO₂ was injected into the pond. Injection was located at MW during Trials 1 and 2. During Trial 3, CO₂ was injected at ME. Locations where values were recorded in the pond are defined as: NW (northwest), MW (middle-west), SW (southwest), NE (northeast), ME (middle-east), and SE (southeast). Values describe a single measurement at each location and time.

Figures 4: Occupancy and behavior of silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) in outdoor ponds. Detections in the High CO₂ Area (top) decreased significantly while CO₂ was injected into the pond, while detections the Low CO₂ Area (middle) increased

significantly. Crosses through the CO₂ injection site were significantly reduced during injection (bottom), but CO₂ was not 100% effective to stop upstream fish passage.

Figure 5,6,7: Density plots illustrating the spatial distribution of silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) during the evaluation of carbon dioxide (CO₂) as a barrier and deterrent in outdoor ponds. Independent trials 1 (Figure 5), 2 (Figure 6), and 3 (Figure 7) were discretized into three 24h periods: before (PreCO₂), during (DuringCO₂), and after (PostCO₂) CO₂ injection. The red line represents the approximate location of the CO₂ barrier.

Figure 8: Box-and-whisker plots describing the movement of silver (*Hypophthalmichthys molitrix*) and bighead carp (*H. nobilis*) upstream through a CO₂ injection site in outdoor concrete ponds. The effectiveness of lower concentrations (≤ 60 mg/L CO₂) to impede upstream movement was compared to higher concentrations (> 60 mg/L CO₂). Carp moved upstream significantly fewer times when CO₂ was ≥ 60 mg/L CO₂ ($p < 0.01$) relative to lower concentrations. No differences between species were observed ($p = 0.17$).

Crossing per hour per fish

<https://mc.manuscriptcentral.com/cjfas-pubs>CO₂ concentration