Canadian Journal of Fisheries and Aquatic Sciences
 Journal canadien des sciences halieutiques et aquatiques

Testing for synchrony in recruitment among four Lake Michigan fish species

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2015-0534.R2
Manuscript Type:	Article
Date Submitted by the Author:	28-Jun-2016
Complete List of Authors:	Bunnell, David; U.S. Geological Survey, Höök, Tomas; Purdue University, Troy, Cary D.; Purdue University, Lyles School of Civil Engineering Liu, Wentao; Purdue University, Lyles School of Civil Engineering Madenjian, Charles; U.S. Geological Survey, Adams, Jean; USGS Great Lakes Science Center
Keyword:	RECRUITMENT < General, PHYSICAL LIMNOLOGY < General, FRESHWATER FISHES < General, CLIMATE CHANGE < General, LAKES < Environment/Habitat

SCHOLARONE™
 Manuscripts

Testing for synchrony in recruitment among four Lake Michigan fish species

David B. Bunnell^{1*}, Tomas O. Höök^{2,3}, Cary D. Troy⁴, Wentao Liu⁴,
Charles P. Madenjian¹, and Jean V. Adams¹

¹U.S. Geological Survey, Great Lakes Science Center, 1451 Green Road, Ann Arbor, MI 48105,
cmadenjian@usgs.gov, jvadams@usgs.gov

²Purdue University, Department of Forestry and Natural Resources, 195 Marsteller Street, West
Lafayette IN 47907-2033, thook@purdue.edu

³Illinois-Indiana Sea Grant College Program, University of Illinois, 1101 W. Peabody Drive
374 National Soybean Research Center, MC-635, Urbana, IL 61801

⁴Purdue University, Lyles School of Civil Engineering, 550 Stadium Mall Drive, West Lafayette,
IN 47907-2051, troy@purdue.edu, wentaoliu@purdue.edu

*Corresponding author: dbunnell@usgs.gov, 734-214-9324

Abstract

In the Great Lakes region, multiple fish species display intra-specific spatial synchrony in recruitment success, with inter-annual climate variation hypothesized as the most likely driver. In Lake Michigan, we evaluated whether climatic or other physical variables could also induce spatial synchrony across multiple species, including bloater (*Coregonus hoyi*), rainbow smelt (*Osmerus mordax*), yellow perch (*Perca flavescens*), and alewife (*Alosa pseudoharengus*). The residuals from stock-recruitment relationships revealed yellow perch recruitment to be correlated with recruitment of both rainbow smelt ($r = 0.37$) and alewife ($r = 0.36$). Across all four species, higher than expected recruitment occurred in 5 years between 1978 and 1987 and then switched to lower than expected recruitment in 5 years between 1996 and 2004. Generalized additive models revealed warmer spring and summer water temperatures and lower wind speeds corresponded to higher than expected recruitment for the nearshore-spawning species, and overall variance explained ranged from 14% (yellow perch) to 61% (alewife). For all species but rainbow smelt, higher recruitment also occurred in extremely high or low years of the North Atlantic Oscillation index. Future development of indices that describe the physical Great Lakes environment could improve understanding of how climate can synchronize fish populations within and across species.

Key words: environmental forcing, Moran effect, food-web, climate, invasive species, Pacific Decadal Oscillation

47 Introduction

48   Among fish populations of the same species, spatial synchrony in recruitment and
49 abundance dynamics has been documented in freshwater (Grenouillet et al. 2001; Ruetz et al.
50 2005; Phelps et al. 2008) and marine (Fox et al. 2000; Hollowed et al. 2001; Cheal et al. 2007)
51 ecosystems. Broad-scale climatic variation has been identified as one of the primary drivers of
52 such synchrony (e.g., the “Moran effect”, Moran 1953). Other less commonly identified
53 underlying mechanisms include widespread dispersal of young in years of high reproductive
54 output or trophic interactions such as a mobile predator synchronizing otherwise disparate prey
55 populations (see review by Liebhold et al. 2004).

56   For some marine ecosystems, spatial synchrony has also been documented across
57 multiple fish species. For example, in George’s Bank in the North Atlantic, several groundfish
58 species exhibited common years of exceptional recruitment, and the North Atlantic Oscillation
59 was identified as the most likely forcing factor (Brodziak and O’Brien 2005). In the eastern
60 Atlantic Ocean, landings of herring (*Clupea harengus*), sardine (*Sardina pilchardus*), and
61 sardinella (*Sardinella aurita*) increased with warm phases of the Atlantic Multidecadal
62 Oscillation between 1945 and 2010 (Alheit et al. 2014). In the Gulf of Alaska, the recruitment of
63 five species that rely on cross-shelf transport during the larval phase appeared to be commonly
64 influenced by sea height (Stachura et al. 2014). Hence, climate has been the most commonly
65 invoked mechanism to explain synchrony across different fish species within a given habitat.

66   In the Laurentian Great Lakes region, historical accounts describe “phenomenal” 1943
67 year-classes simultaneously produced for lake whitefish (*Coregonus clupeaformis*), walleye
68 (*Sander vitreus*), and cisco (*Coregonus artedii*) in Green Bay, Lake Michigan, although no
69 mechanism was offered (Hile et al. 1953). More recently, synchrony has been observed among

populations of the same species. For example, bloater (*Coregonus hoyi*) were synchronized across lakes Superior, Michigan, and Huron (Bunnell et al. 2010), cisco were synchronized across Lake Superior and several inland lakes (Myers et al. 2015), and yellow perch (*Perca flavescens*) were synchronized across four Great Lakes and Lake St. Clair (Honsey et al. 2016). Climate was identified as the most parsimonious explanation in these studies. Moreover, several recruitment studies of individual fish populations in the Laurentian Great Lakes have identified inter-annual variation in climate and physical conditions as important contributing predictors of recruitment for a diversity of species, such as alewife (*Alosa pseudoharengus*, e.g., Madenjian et al. 2005), yellow perch (*Perca flavescens*, e.g., Redman et al. 2011), lake whitefish (Brown et al. 1993), and rainbow smelt (*Osmerus mordax*, O'Brien et al. 2014). Although the exact mechanisms by which climate influence recruitment are often unknown and may even be indirect (e.g., thermal conditions influencing zooplankton availability; Winder and Schindler 2004), climate may be more important in the Laurentian Great Lakes than in smaller freshwater systems because density-independent processes are more likely to override density-dependent ones in these large freshwater seas (see Ludsin et al. 2014). We hypothesized that if climate underlies recruitment variation and can cause synchrony among populations of the same species, it also could induce synchrony across different species within a given lake, especially for those species that share habitat during critical life history stages where climatic or physical conditions can exert a common effect. To our knowledge, no previous study in freshwater ecosystems has ever quantitatively documented synchrony across several species.

Herein, we evaluated whether recruitment patterns of alewife, bloater, rainbow smelt, and yellow perch [whose larvae all occupy the upper pelagic waters during the late spring and summer months in Lake Michigan (Table 1)] were synchronous in Lake Michigan from 1973-

2011. Our approach was to first account for the effect of stock size on recruitment success by estimating stock-recruitment relationships and calculating the recruitment residuals for each species in each year. To test whether or not environmental variables can synchronize the recruitment of these four species, we assumed that variation unexplained by stock size could be explained by climatic or other physical variables. We determined whether positive or negative residuals (i.e., higher or lower than expected recruitment for a given stock size) occurred in common years across species, and then whether residual recruitment patterns were similar across species or exhibited temporal autocorrelation. Finally, we sought to determine whether the residual patterns corresponded with time series of climatic or physical variables that could be compiled over the same 39-year time period.

Materials and Methods

Estimating stock-recruit residuals

For all species, we used data collected by the USGS Great Lakes Science Center bottom trawl survey, which has occurred annually each autumn since 1973. Details regarding the survey can be found in Bunnell et al. (2015), but briefly a bottom trawl with a 12-m footrope was towed along the bottom for up to 10 minutes at each of 69 locations organized across 7 distinct ports (Manistique, Frankfort, Ludington, and Saugatuck in Michigan; Waukegan in Illinois; Port Washington and Sturgeon Bay in Wisconsin) at up to 11 depths (ranging 9 – 110 m at approximately 10-m increments) per port. The catch was sorted to species and weighed, and a random subset of alewife, bloater, rainbow smelt, and yellow perch were measured to the nearest mm (total length, TL). A subset of bloater and alewife are also aged with scales and otoliths, respectively (see Bunnell et al. 2006; Madenjian et al. 2005). Appropriate measures for stock

size and recruitment were based on previously published papers when possible (see Table 2). We included data up through the 2011 year-class for all species.

Stock-recruit models were fit in SAS (SAS Institute Inc. 2015) using Procedure NLIN and assuming a log-normal error structure (e^ε). For alewife and rainbow smelt, the Ricker form of stock-recruit models has already been justified and applied in Lake Michigan (Madenjian et al. 2005; Collingsworth et al. 2014; Feiner et al. 2015). For bloater, previous studies (Bunnell et al. 2006; Collingsworth et al. 2014) identified temporal autocorrelation in the recruitment residuals, and removed it by using dynamic linear models (Pole et al. 1994). In this study, however, we were interested in exploring whether any of the temporal autocorrelation corresponded with climatic variables. For yellow perch, there are no previously published stock-recruit models using USGS data. However, recent studies of yellow perch recruitment in Lake Michigan from other surveys have applied the Ricker form (Redman et al. 2011; Forsythe et al. 2012).

To ensure that the Ricker model was the best fit for the USGS data for bloater and yellow perch, we fit both the Beverton-Holt and Ricker forms to our stock (S) and recruit (R) data. For all species, we fit the Ricker form as:

$$\ln(R) = \ln(S) + a - bS + \varepsilon.$$

For only bloater and yellow perch, we fit the Beverton-Holt form of the model as:

$$\ln(R) = \ln(aS) - \ln(b + S) + \varepsilon.$$

For these two latter species, we evaluated which form provided the best fit using Akaike's Information Criterion (corrected for small sample sizes). Residuals were calculated as $\ln(R_{\text{observed}}) - \ln(R_{\text{predicted}})$, from the best (or only) model. To standardize the residuals across species (to prevent species with higher variability from having a disproportionate effect), we

divided the residual by the standard deviation for the $\ln(R_{\text{observed}})$ values for each species. To provide some sense of how well stock size explained recruitment, we used the r^2 from the linear model between $\ln(R)$ and $\ln(R_{\text{predicted}})$.

Recruitment residual patterns

First, we explored whether temporal autocorrelation in recruitment residuals occurred. Walters (1987) described how temporal autocorrelation in residuals can result from stock-recruitment relationships that vary with time either because of changes in the productivity of the stock or long-term environmental changes. Because we were interested in whether climate could influence fish recruitment by altering the stock-recruitment relationship through time, we evaluated whether temporal autocorrelation existed by correlating each time series of residuals against a 1-year time lag using Pearson correlation coefficient. We assumed that autocorrelation in the residuals could indicate periodicity in the time series. Hence for those time series that revealed autocorrelation, we used a power spectral density (PSD) analysis (estimated using Matlab function pmtm; Mathworks 2015) to determine whether periodicity existed in the recruitment residuals. PSD is a non-parametric method that is well-suited for estimating spectra for relatively short time series. Time series of recruitment residuals were detrended, and for all species, spectra were normalized to have the same energy level. Spectral plots were generated where period (year) was on the x-axis and energy ($\text{signal}^2/\text{frequency}$) was on the y-axis. A significant spectral “peak” occurred when the lower confidence interval of that peak rose above the upper confidence interval of the surrounding values.

Second, we sought to determine whether positive or negative deviations from predicted recruitment exhibited common patterns across species. For each of the 37 years for which

recruitment residuals were available for all four species (1995 and 1998 year-classes were missing for at least one species), we determined how the percentage of positive or negative residuals was distributed within each year (e.g., 75% of species one year might have positive residuals and 25% negative residuals). We then counted the number of years that exhibited 100% agreement, and used a randomization test to determine whether that number was greater than what would be expected by chance alone ($\alpha = 0.05$). For this test, we randomly shuffled the time series of residuals for each species independently, and then recorded the number of times 100% agreement occurred. This procedure was repeated 10,000 times.

Third, to evaluate whether recruitment residuals were similar between pairs of species, we calculated the Pearson correlation coefficient between recruitment residual values for each of the six possible species pairs.

Do recruitment residuals correlate with environmental variables?

We compiled several different variables that could describe the climatic or physical environment during potentially critical ecological periods for the egg or larval stage. One criterion for inclusion was that the time series extended back to 1973, which limited our potential variables. First, we obtained annual maximum ice cover data (Ice, Assel et al., 2013; available <http://www.glerl.noaa.gov/data/ice/>), calculated as the percentage of total lake surface area. Ice cover can influence plankton blooms (Vanderploeg et al. 1992; Magnuson et al. 1997) which, in turn, can affect the developmental conditions for fertilized eggs or the feeding environment for fish larvae. Second, we sought to describe the potential availability of nearshore spawning habitat by calculating the mean lake level (LakeLevel) from April – July in Lake Michigan (available <http://www.glerl.noaa.gov/data/dashboard/data/>). Third, we characterized the mean

lake-wide wind speed (Wind) during May – August, the period over which the larvae of the four fish species spend at least part of the time as plankton and are vulnerable to currents. Wind speed was derived from the Great Lakes Monthly Hydrological Dataset (GLMHD), which was estimated from nearshore land stations without overwater corrections (available <http://www.glerl.noaa.gov/data/arc/hydro/mnth-hydro.html>). Fourth, we used lakewide mean surface water temperature (Temp) during May – August to characterize the growth potential for the four fish species. These data were also obtained from the GLMHD and were produced by a one-dimensional large-lake thermodynamics model (Hunter and Croley 1993) and have been used in other fish recruitment studies (e.g., Madenjian et al. 2005; Bunnell et al. 2006). Finally, we used three different regional climate indices that, although distant from the Great Lakes, have been correlated with climate variables, including ice cover and water level, in this region (e.g., Ghanbari and Bravo 2008; Ghanbari et al. 2009; Bai et al. 2012; Rodionov and Assel 2003): the Multivariate ENSO index (also known as El Niño) and the North Atlantic Oscillation (NAO), and the Pacific Decadal Oscillation (PDO). ENSO has also been linked to synchronous zooplankton dynamics in southern Ontario lakes (Rusak et al. 2008). ENSO is based on six observed variables over the tropical Pacific (Wolter and Timlin 1998) and was obtained from NOAA's Earth System Research Laboratory (available <http://www.esrl.noaa.gov/psd/enso/mei/table.html>). NAO is measured as the difference in air pressure between the Low pressure system near Iceland and the High pressure system near the Azores and we obtained the NAO time series from the NOAA Climate Prediction Center (available <http://www.cpc.ncep.noaa.gov/data/teledoc/nao.shtml>). PDO is indexed based on sea surface temperature and sea level pressure observations in the northern Pacific Ocean, and we obtained the PDO index time series from the NOAA National Centers for Environmental

Information (available <https://www.ncdc.noaa.gov/teleconnections/pdo/>). For each regional climate index, we used the average of the winter months (December, January, February) given that most studies have identified these months as influencing the climate for both the winter and subsequent seasons (Hurrell et al. 2003; Bai et al. 2012). As with the fish recruitment variables, we also explored whether temporal autocorrelation and periodicity (if necessary) occurred for all of the climatic and physical variables.

To determine whether the recruitment residuals corresponded with any of these environmental variables, we used a generalized additive model (GAM) owing to its ability to fit non-linear relationships between explanatory and response variables (Wood 2004). GAMs were conducted for each fish species, as well as for a time series of the recruitment residuals that was averaged across the four species. We used R (R Development Core Team 2015; package mcgv) and fit all possible combinations of the seven main variables (Ice, LakeLevel, Wind, Temp, ENSO, NAO, PDO) and a null model with only an intercept. We also included only one *a priori* interaction, Temp and LakeLevel; we hypothesized that the effect of warm water temperature on fish growth and survival would depend on what quantity or quality of nearshore spawning habitat existed. This yielded a total of 160 models for each response variable that were compared using Akaike's Information Criterion corrected for small sample size (AIC_c). We calculated ΔAIC_c for each candidate model i as $AIC_i - AIC_{min}$, where min is the model with the smallest AIC_c for a particular response variable. Those candidate models with ΔAIC values < 2 were considered to have substantial support in explaining variation in the data and were included in a suite of "top-ranked" models (Burnham and Anderson 2002). To determine which variables had the most explanatory power, we calculated the percentage of top-ranked models that included a given explanatory variable.

Results

Estimating stock-recruit residuals

For bloater, the Ricker and Beverton-Holt forms of stock-recruit models were indistinguishable through AIC_c (48.4, 48.2, respectively); we used the Ricker model to calculate residuals. For yellow perch, however, the Beverton-Holt form ($AIC_c = 72.4$) was more parsimonious than the Ricker model ($AIC_c = 76.2$), and thus we used the Beverton-Holt model. For all species, the amount of variation in recruitment explained by stock size alone was relatively low: 5% for alewife, 8% for yellow perch, 19% for rainbow smelt, and 30% for bloater (Fig. 1, Table 3).

Recruitment residual patterns

The residual patterns (Fig. 2) exhibited temporal autocorrelation (with a one-year lag) for both bloater ($r = 0.59$; $P = 0.0001$; Fig. 2c) and yellow perch ($r = 0.43$; $P = 0.001$; Fig. 2d). Spectral analysis for each of these time series indicated the period with the highest peak in energy was about 25-33 years (Fig. S1, S2); this peak was not significant however. Because residuals were not temporally autocorrelated for alewife ($r = 0.11$; $P = 0.51$; Fig. 2a) or rainbow smelt ($r = 0.00$; $P = 0.99$; Fig. 2b), no spectral analyses were conducted for these species.

One-hundred percent agreement in the sign of the recruitment residuals occurred in 11 out of 37 years (Figure 3b), significantly more than expected by chance alone ($P=0.003$). Positive recruitment residuals occurred for all species in five out of 10 years between 1978 and 1987 (1978, 1983, 1984, 1985, and 1987). Then negative recruitment residuals occurred for all

species in five out of eight years between 1996 and 2004 (1996, 1997, 2000, 2001, 2004).

Finally the 2010 year-class had 100% positive recruitment residuals.

Recruitment residuals were positively correlated between yellow perch and rainbow smelt ($r = 0.38$; $P = 0.02$; Fig. 4a) and between yellow perch and alewife ($r = 0.36$; $P = 0.03$; Fig. 4b). For no other pairs of species were recruitment residuals significantly correlated with one another ($P \geq 0.14$). Absence of a positive correlation between bloater and yellow perch (Fig. 4c) was due to the 2005 and 2009 year-classes; removing these two outliers resulted in a correlation coefficient of 0.47 ($P = 0.004$).

Do recruitment residuals correlate with environmental variables?

Several of the climatic or physical time series of explanatory variables (Fig. S3) exhibited temporal autocorrelation: LakeLevel ($r = 0.84$, $P < 0.0001$), Temp ($r = 0.49$, $P = 0.002$), NAO ($r = 0.36$, $P = 0.03$). Spectral analysis indicated that the most likely periods for these environmental variables ranged 8-20 years (Fig. S4), although none of these was significant. Notably, the most likely period for the environmental variables was not markedly different from residual pattern averaged across all four species (i.e., 20 years, Fig. S4). The other environmental variables (Ice, Wind, ENSO, PDO) were not autocorrelated ($P \geq 0.21$) and exhibited more year-to-year variation.

Among the GAMs, the number of top-ranked models (i.e., those with $\Delta AIC < 2$) ranged from 3 for alewife to 21 for yellow perch, and the % deviance explained by AIC_{min} model was relatively low for yellow perch and alewife (14-20%), somewhat higher for average residual time series (35%) and bloater (37%), and highest for alewife (61%, Table 4). Among this suite of top-ranked models for each species, three explanatory variables consistently occurred: water

temperature, wind speed, and ice cover, although ice cover occurred in a relatively low percentage of the models (range = 13-33%, Table 4). Spring-summer water temperature occurred in 71% of top-ranked models, on average, whereas spring-summer wind speed occurred in 66% of them.

Further inspection of the GAM results (see Fig. S5-S9), however, suggests that common environmental variables among the top-ranked models were not necessarily affecting species the same way. For example, higher wind speeds led to lower than expected recruitment for all fish species (and the average time series), with the exception of bloater (Fig. S7), for which the opposite trend was identified. Similarly, warmer water temperatures led to higher than expected recruitment for all species and the average time series, with another exception for bloater, where recruitment residuals became more negative at the warmest water temperatures. Higher lake levels led to higher than expected recruitment for alewife (Fig. S5), yellow perch (Fig. S8), and the average time series (Fig. S9), but lower than expected recruitment for rainbow smelt (Fig. S6). NAO was the only explanatory variable that exhibited a consistent relationship for every GAM where it occurred among the top-ranked models: it was non-linear with the highest recruitment residuals occurring at extreme negative and positive NAO years for alewife, bloater, yellow perch, and the average time series.

Discussion

We documented moderate levels of synchrony across four fish species in Lake Michigan between the 1973 and 2011 year-classes. After accounting for spawning stock size, recruitment residuals of four Lake Michigan fish species were either 100% positive or negative for 11 different year-classes, which was significantly more frequent than would be expected based on

chance alone. Furthermore, residuals were 100% positive over five years between 1978 and 1987 and then 100% negative over five years between 1996 and 2004. Among pairs of species, the recruitment residuals of yellow perch and rainbow smelt, as well as those of yellow perch and alewife, were significantly correlated, but the correlation coefficient was relatively low ($r \leq 0.37$). For bloater and yellow perch, recruitment residuals exhibited multi-year periods of positive and negative recruitment, although no statistically significant periodicity could be identified, in large part because the apparent dominant time scale of the variation (~25 years) was long relative to the record length (39 years). Our attempt to explain variation in recruitment residuals with climatic or physical variables failed to clearly identify at least one common driver for all four species. Although warmer spring and summer water temperatures and lower spring and summer wind speeds were correlated with stronger recruitment for alewife, rainbow smelt, and yellow perch, the opposite patterns were documented for bloater. The NAO index also showed promise as a common driver, as the three fish species for which it was an important explanatory variable all exhibited a similar non-linear pattern with the strongest recruitment occurring in extreme negative and positive NAO years.

The strongest correlations among recruitment residuals were between yellow perch and alewife as well as between yellow perch and rainbow smelt. These three species have more in common from an early life-history perspective than with bloater, as these three species each spawn in shallow nearshore waters, tributaries, adjacent wetlands, or even drowned river mouths of the Great Lakes, whereas bloater spawn in offshore, hypolimnetic waters (see Table 1). The fact that the residuals were not correlated between rainbow smelt and alewife suggests that the phenology of spawning or hatching of embryos may influence the stronger correlations, given that among these three species, rainbow smelt spawn and hatch first while alewife spawn and

hatch last. Given that yellow perch spawns and hatches during an intermediate time, it may not be surprising their residuals correlate with both species. Under this premise, environmental conditions in the nearshore during the months of June and July may be critical to synchronizing the recruitment patterns of yellow perch with both species. Furthermore, the GAMs indicated that all three nearshore-spawning species were similarly affected by water temperature, wind speed, and the NAO index.

Warm spring and summer water temperature led to higher than expected recruitment for three of the four species and the average residual time series. Bloater was the only species to suggest an alternative pattern, as recruitment residuals were lower than expected in years where average water temperatures exceeded 14 °C. Mechanistically, warmer water temperatures could benefit growth rates of larval fish, through increased maximum consumption rates (Post 1990). In turn, faster growth may facilitate higher survival through various mechanisms including outgrowing gape-limited predators (e.g., Miller et al. 1988), decreasing risk of starvation (e.g., Graeb et al. 2004) and overcoming size-dependent overwinter mortality (e.g., Post and Evans 1989; Höök et al. 2007). One additional hypothesis to be evaluated in future studies is that the timing and magnitude of zooplankton production increases during relatively warm years to further improve the growth potential for fish larvae.

Higher wind speeds led to lower than expected recruitment for the three nearshore-spawning species and the average residual time series. We hypothesize that these years could lead to more frequent upwelling events and physical disruption (i.e., stronger currents, water temperature changes) in the dynamic nearshore habitat. Upwellings in the Great Lakes have been documented to both stimulate (Dunstall et al. 1990) and temporarily reduce (Haffner et al. 1984) densities of zooplankton available to first-feeding larvae, and the extreme declines in

temperature associated with an event (e.g., $>6^{\circ}\text{C}$, Haffner et al. 1984; Dunstall et al. 1990; Bocaniov et al. 2012) could reduce survival of incubating eggs. Furthermore, upwelling events in southeastern Lake Michigan were documented to reduce larval alewife densities fivefold, through either mortality from thermal shock or transport out of the nearshore sampling area (Heufelder et al. 1982). Wind direction would be another factor to consider in future work, as this could also affect where their planktonic larvae are transported. Bloater exhibited the opposite pattern of higher than expected recruitment in windier years. Their eggs incubate in the hypolimnion where the impact of high winds could influence dispersal, but it remains unclear how this could improve recruitment.

Higher lake levels were associated with higher than expected recruitment of alewife and yellow perch. This result is consistent with the hypothesis that higher water levels improved the spawning or nursery habitat for these species that spawn in the nearshore, adjacent tributaries, or even drowned river-mouth lakes. Specifically, higher water levels could increase accessibility to high-quality sites that might be unreachable in low-water years (Fracz and Chow-Fraser 2013; Gaeta et al. 2014; O'Brien et al. 2014). At the same time, higher than expected recruitment occurred during low water-level years for the other nearshore spawning species, rainbow smelt. Although water level rarely occurred (i.e., 13% of the time) among the top-ranked models, it is unclear why this inverse pattern was revealed. A recent analysis of rainbow smelt recruitment in Lake Huron revealed spring precipitation and water levels to be positively correlated with strong year-classes (O'Brien et al. 2014), suggesting the relationship we uncovered in Lake Michigan could be spurious within a model that has relatively low overall variation explained (i.e., 20%).

The regional climate index for the north Atlantic Ocean, NAO, was the one explanatory variable that showed a consistent, albeit non-linear, relationship with recruitment residuals for

the four GAMs where it was among the top-ranked models (i.e., all but rainbow smelt). NAO is measured as the difference in air pressure between the Low pressure system near Iceland and the High pressure system near the Azores. Given the location where NAO is measured, it should not be surprising that most of the research linking NAO indices to aquatic biota centers in the north Atlantic (e.g., Alheit and Hagen 1997; Beaugrand et al. 2002) and over inland lakes in Europe (e.g., Straile et al. 2001, 2007). In the Great Lakes region, negative NAO values are associated with colder winter air temperatures, lower snowfall, and longer ice duration (Ghanbari et al. 2009; Bai et al. 2012). Positive phases of NAO are associated with warmer winters (Hurrell et al. 2003). Suski and Ridgway (2007) reported how NAO was positively associated with higher nest survival among nearshore-spawning smallmouth bass (*Micropterus dolomieu*) in Lake Opeongo, Ontario. Because positive NAO winters were also associated with shallower summer thermocline depths and warmer epilimnions (which can lead to a stronger barrier between vertical layers), they speculated that these years produced fewer upwelling events that could disturb nests. Whether or not the winter NAO affects the thermocline and limnology in larger lakes is unknown. In addition, future research will be required to determine how extreme negative NAO years (cold, dry, high ice winters) can also lead to strong recruitment events for these fish species through some second, unknown mechanism.

A dominant regional climate index for the northern Pacific Ocean, PDO, also occurred in the top-ranked models for all GAMs but alewife. PDO generally influences the temperature and precipitation in northwestern North America and the southeastern United States, although warm phases (i.e., positive values) of the PDO are associated with lower winter precipitation in the Great Lakes region (Mantua 2002). Van Cleave et al. (2014) hypothesized that a “regime shift” in Lake Superior ice cover, evaporation, and water temperature was associated with a coincident

shift in the PDO to a cold phase (i.e., negative values). Our model results indicated that warmer phases of the PDO were associated with stronger than expected recruitment for bloater, yellow perch, and rainbow smelt, although rainbow smelt recruitment was also higher than expected for extremely negative PDO values. For both of these regional indices, future research will be required to improve understanding of how they are influencing fish recruitment patterns or whether our results are simply a spurious correlation.

Other climatic explanatory variables could be included in future research. A direct annual index of upwellings (instead of mean wind speed), for example, would have enabled a more explicit understanding of whether this potentially important physical driver influences the early life history dynamics of nearshore-spawning species. Plattner et al. (2006) demonstrated that upwelling frequency in the Great Lakes can be remotely estimated using sea surface temperature data, but the time series was only developed for 1992 through 2000. Another possible addition would be more interaction terms. The lake level $\times$ water temperature interaction that we included was important only for alewife. When we equally divided the alewife data into low and high water years, we found recruitment residuals to slightly increase with water temperature in high water-level years ($F_{1,18} = 4.4$, $P = 0.05$, $r^2 = 0.20$) but not in low water-level years ($F_{1,18} = 2.1$, $P = 0.17$, $r^2 = 0.11$), which was consistent with our hypothesis. Lastly, future efforts could compile explanatory variables at locally or regionally important scales. We summed environmental variables at the lakewide level, because we did not have spatially explicit estimates of fish recruitment within Lake Michigan. Disproportionally high egg or larval survival can occur in discrete locations, however, and unique local environmental conditions that theoretically could have facilitated strong year-classes either may not have been measured or were swamped by more widespread “normal” environmental conditions. For

example, Höök et al. (2008) indicated year classes for which larval alewife survival is poor in the lake are disproportionately supported by tributary-reared alewives.

Previous research evaluating the mechanisms underlying recruitment of these four species in Lake Michigan has indicated that the role of environmental factors such as climate or water level are not necessarily the predominate factors. Alewife recruitment to age 3 was primarily regulated by salmonine predation, although spring and summer water temperatures were also included among the top-ranked models (Madenjian et al. 2005; Collingsworth et al. 2014). For rainbow smelt, the best model allowed the productivity of the stock to vary over the time series, but no environmental variable clearly explained additional recruitment variability (Feiner et al. 2015). Variation in Lake Michigan bloater has been best explained by sex ratio of the adult population, and climate variables offered minimal explanatory power (Bunnell et al. 2006; Collingsworth et al. 2014). The only exception is a recent analysis of yellow perch recruitment in Illinois water of Lake Michigan that revealed spring and summer water temperatures to be just as important as other biotic variables including densities of alewife predators and zooplankton (Redman et al. 2011). Given that biotic factors, such as predation, can also synchronize prey populations (Liebhold et al. 2004), future research evaluating synchrony across multiple fish species could include biomass of piscivores as an explanatory variable.

In conclusion, after accounting for the effect of stock size on recruitment we discerned moderate levels of synchrony among four fish species in Lake Michigan. Only two of six species pairs exhibited positive correlations in their recruitment residuals, and the coefficient of variation (r) never exceeded 0.37. At the same time, for all four species we observed higher than expected recruitment over five years between 1978 and 1987 and lower than expected

recruitment over five years between 1996 and 2004. Decadal-scale oscillations in fish recruitment or harvest dynamics have previously been documented among marine fish populations, and regional climate indices are believed to underlie these dynamics (e.g., Alheit et al. 2014; Brodziak and O'Brien 2005; Stachura et al. 2014). In the Great Lakes region, however, no one regional climate index has been identified by physical scientists. Hence we included indices from other regions of the world as well as other climatic and environmental variables in a GAM to attempt to explain recruitment patterns for these four species, as well as their average residual pattern. Although the NAO and PDO indices offered some explanatory power for some species, the overall variance explained was relatively low. Warm spring and summer water temperatures and low average spring and summer wind speeds were positively associated with higher than expected recruitment for the three nearshore-spawning species. Scientists in the Great Lakes are increasingly seeking to understand how physical factors influence fish recruitment (e.g., Ludsins et al. 2014), above and beyond the well-studied biotic factors that have been documented to be important for some species. Further development of indices that capture the physical dynamics operating in the Great Lakes region should improve understanding of how multiple environmental factors collectively influence and even synchronize the recruitment dynamics of fishes.

Acknowledgements

We acknowledge the hard work of previous scientists, technicians, and vessel crew in maintaining this long-term federal monitoring of Lake Michigan's fish community. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government. We thank Mark Vinson and two anonymous reviewers for constructive

comments that substantially improved this paper. This work was supported, in part, by funding provided by the Great Lakes Fishery Commission's theme on Physical Processes and Fish Recruitment in Large Lakes. This article is Contribution 2070 of the U.S. Geological Survey Great Lakes Science Center and Contribution XXX of Purdue's Climate Change Research Center.

References

- Alheit, J., and Hagen, E. 1997. Long-term climate forcing of European herring and sardine populations. *Fish. Oceanogr.* **6**: 130-139.
- Alheit, J., Licandro, P., Coombs, S., Garcia, A., Giraldez, A., Santamaria, M.T.G., Slotte, A., and Tsikliras, A.C. 2014. Atlantic Multidecadal Oscillation (AMO) modulates dynamics of small pelagic fishes and ecosystem regime shifts in the eastern North and Central Atlantic. *J. Mar. Sys.* **133**: 88-102.
- Assel, R.A., Wang, J., Clites, A.H., and Bai, X. 2013. Analysis of Great Lakes ice cover climatology: Winters 2006-2011, NOAA Technical Memorandum GLERL-157, NOAA Great Lakes Env. Res. Lab., Ann Arbor, MI.
- Auer, N.A. 1982. Identification of larval fishes of the Great Lakes basin with emphasis on the Lake Michigan drainage, Great Lakes Fish. Comm. Spec. Pub. 82-3, Ann Arbor, MI.
- Bai, X., Wang, J., Sellinger, C., Clites, A., and Assel, R. 2012. Interannual variability of Great Lakes ice cover and its relationship to NAO and ENSO. *J. Geophys. Res.* **117**
doi:10.1029/2010JC006932
- Beaugrand, G., Reid, P.C., Ibanez, F., Lindley, J.A., and Edwards, M. 2002. Reorganization of North Atlantic marine copepod biodiversity and climate. *Science* **296**:1692-1694.

- 483 Beletsky, D., Mason, D.M., Schwab, D.J., Rutherford, E.S., Janssen, J., Clapp, D.F., and
484 Dettmers, J.M. 2007. Biophysical model of larval yellow perch advection and settlement
485 in Lake Michigan. *J. Great Lakes Res.* **33**: 842-866.
- 486 Bocianov, S.A., Schiff, S.L., and Smith, R.E.H. 2012. Plankton metabolism and physical forcing
487 in a productive embayment of a large oligotrophic lake: insights from stable isotopes.
488 *Freshwater Biol.* **57**: 481-496.
- 489 Brodziak, J., and O'Brien, L. 2005. Do environmental factors affect recruits per spawner
490 anomalies of New England groundfish? *ICES J. Mar. Sci.* **62**: 1394-1407.
- 491 Brown, R.W., Taylor, W.W., and Assel, R.A. 1993. Factors affecting the recruitment of lake
492 whitefish in two areas of northern Lake Michigan. *J. Great Lakes Res.* **19**: 418-428.
- 493 Bunnell, D.B., Madenjian, C.P., and Croley, T.E. 2006. Long-term trends of bloater recruitment
494 in Lake Michigan: evidence for the effect of sex ratio. *Can. J. Fish. Aquat. Sci.* **63**: 832-
495 844.
- 496 Bunnell, D.B., Adams, J.V., Gorman, O.T., Madenjian, C.P., Riley, S.C., Roseman, E.F., and
497 Schaeffer, J.S. 2010. Population synchrony of a native fish across the upper Great Lakes:
498 evaluating the effects of dispersal and climate. *Oecologia* **162**: 641-651.
- 499 Bunnell, D.B., Madenjian, C.P., Desorcie, T.J., Kostich, M.J., Woelmer, W., and Adams, J.V.
500 2015. Status and trends of preyfish populations in Lake Michigan in 2014. Distributed at
501 the Lake Michigan Committee Meeting of the Great Lakes Fishery Commission during
502 March 2015.
- 503 Bunnell, D.B., Madenjian, C.P., Desorcie, T.J., Kostich, M.J., Woelmer, W., and Adams, J.V.
504 2015. Status and trends of prey fish populations in Lake Michigan in 2014. *In* Compiled

- 505 reports to the Great Lakes Fishery Commission of the annual bottom trawl and acoustics
506 surveys, 2014, Ypsilanti, Mich, 23-27 March 2015. pp. 33-48.
- 507 Burnham, K.P., and Anderson, D.R. 2002. Model selection and multimodel inference: a practical
508 information-theoretic approach, 2nd edition. Springer-Verlag, New York, NY.
- 509 Cheal A.J., Delean, S., Sweatman, H., and Thompson, A.A. 2007. Spatial synchrony in coral
510 reef populations and the influence of climate. *Ecology* **88**:158-169
- 511 Collingsworth, P.D., Bunnell, D.B., Madenjian, C.P., and Riley, S.R. 2014. Comparative
512 recruitment dynamics of alewife and bloater in Lakes Michigan and Huron. *Trans. Am.*
513 *Fish. Soc.* **143**: 294-309.
- 514 Dettmers, J.M., Janssen, J., Pientka, B., Fulford, R.S., and Jude D.J. 2005. Evidence across
515 multiple scales for offshore transport of yellow perch (*Perca flavescens*) larvae in Lake
516 Michigan. *Can. J. Fish. Aquat. Sci.* **62**: 2683–2693.
- 517 Dunstall, T.G., Carter, J.C.H., Monroe, B.P., Haymes, G.T., Weiler, R.R., and Hopkins, G.J.
518 1990. Influence of upwellings, storms, and generating station operation on water
519 chemistry and plankton in the Nanticoke region of Long Point Bay, Lake Erie. *Can. J.*
520 *Fish. Aquat. Sci.* **47**: 14343–1445.
- 521 Feiner, Z.S., Bunnell, D.B., Höök, T.O., Madenjian, C.P., Warner, D.M., and Collingsworth,
522 P.D. 2015. Non-stationary recruitment dynamics of rainbow smelt: the influence of
523 environmental variables and variation in size structure and length-at-maturation. *J. Great*
524 *Lakes Res.* **41**: 246-258.
- 525 Forsythe, P.S., Doll, J.C., and Lauer, T.E. 2012. Abiotic and biotic correlates of yellow perch
526 recruitment to age-2 in southern Lake Michigan, 1984-2007. *Fish. Manage. Ecol.* **19**:
527 389-399.

- 528 Fox, C.J., Planque, B.P., and Darby C.D. 2000. Synchrony in the recruitment time-series of
529 plaice (*Pleuronectes platessa* L) around the United Kingdom and the influence of sea
530 temperature. J. Sea Res. **44**: 159-168.
- 531 Fracz, A., and Chow-Fraser, P. 2013. Impacts of declining water levels on the quantity of fish
532 habitat in coastal wetlands of eastern Georgian Bay, Lake Huron. Hydrobiologia **702**:
533 151-169.
- 534 Gaeta, J.W., Sass, G.G., and Carpenter, S.R. 2014. Drought-driven lake level decline: effects on
535 coarse woody habitat and fishes. Can. J. Fish. Aquat. Sci. **71**: 315-325.
- 536 Ghanbari, R.N., and Bravo, H.R. 2008. Coherence between atmospheric teleconnections, Great
537 Lakes water levels, and regional climate. Adv. Water Res. **31**: 1284-1298.
- 538 Ghanbari, R.N., Bravo, H.R., Magnuson, J.J., Hyzer, W.G., and Benson, B.J. 2009. Coherence
539 between lake ice cover, local climate, and teleconnections (Lake Mendota, Wisconsin).
540 J. Hydrol. **374**: 282-293.
- 541 Graeb, B.D.S., Dettmers, J.M., Wahl, D.H., and Caceres, C.E. 2005. Fish size and prey
542 availability affect growth, survival, prey selection, and foraging behavior of larval yellow
543 perch. Trans. Am. Fish. Soc. **133**: 504-514.
- 544 Grenouillet, G., Hugueny, B., Carrel, G.A., Olivier, J.M., and Pont, D. 2001. Large-scale
545 synchrony and inter-annual variability in roach recruitment in the Rhone River: the
546 relative role of climatic factors and density-dependent processes. Freshwater Biol. **46**: 11-
547 26.
- 548 Haffner, G.D., Yallop, M.L., Hebert, P.D.N., and Griffiths, M. 1984. Ecological significance of
549 upwelling events in Lake Ontario. J. Great Lakes Res. **10**: 28-37.

- 550 Heufelder, G.R., Jude, D.J., and Tesar, F.J. 1982. Effects of upwelling on local abundance and
551 distribution of larval alewife (*Alosa pseudoharengus*) in eastern Lake Michigan. Can. J.
552 Fish. Aquat. Sci. **39**: 1531-1537.
- 553 Hile, R., Lunger G.F., and Buettner, H.J. 1953. Fluctuations in the fisheries of state of Michigan
554 waters of Green Bay. Fish. Bulletin 54: 1-34.
- 555 Hollowed, A.B., Hare, S.R., and Wooster, W.S. 2001. Pacific basin climate variability and
556 patterns of northeast Pacific marine production. Prog. Oceanogr. **49**: 257–282.
- 557 Höök, T.O., Rutherford, E.S., Mason, D.M., and Carter, G.S. 2007. Hatch dates, growth,
558 survival, and overwinter mortality of age-0 alewives in Lake Michigan: implications for
559 habitat-specific recruitment success. Trans. Am. Fish. Soc. **136**: 1298-1312.
- 560 Höök, T.O., Rutherford, E.S., Croley, T.E., Mason, D.M., and Madenjian, C.P. 2008. Annual
561 variation in habitat-specific recruitment success: implications from an individual-based
562 model of Lake Michigan alewife (*Alosa pseudoharengus*). Can. J. Fish. Aquat. Sci. **65**:
563 1402-1412.
- 564 Honsey, A.E., Bunnell, D.B., Troy, C.D., Fielder, D.G., Thomas, M.V., Knight, C.T., Chong,
565 S.C., and Höök, T.O. 2016. Recruitment synchrony of yellow perch (*Perca flavescens*)
566 in the Great Lakes region, 1966-2008. Fish. Res. 181: 214-221.
- 567 Hunter, T.S. and Croley II, T.E. 1993. Great Lakes Monthly Hydrologic Data, NOAA Data
568 Report ERL GLERL, National Technical Information Service, Springfield, VA.
- 569 Hurrell, J.W., Kushnir, Y., Ottersen, G., and Visbeck, M. 2003. An overview of the North
570 Atlantic Oscillation. In: The North Atlantic Oscillation: climatic significance and
571 environmental impact. Geophys. Monogr. **134**:1–35.

- 572 Jude, D.J., Tesar, F.J., Tomlinson, J.C., Miller, T.J., Thurber, N.J., Godun, G.G., and Dorr, III
573 J.A. 1979. Inshore Lake Michigan fish populations near the D. C. Cook Nuclear Power
574 Plant during pre-operational years – 1973-1974, Spec. Rep. 71 of the Great Lakes Res.
575 Div., Univ. Michigan, Ann Arbor, MI.
- 576 Lauer, T.E., Shroyer, S.M., Kilpatrick, J.M., McComish, T.S., and Allen, P.J. 2005. Yellow
577 perch length-fecundity and length-egg size relationships in Indiana waters of Lake
578 Michigan. *North Amer. J. Fish. Manage.* **25**: 791-796.
- 579 Liebhold, A., Koenig, W.D. and Bjørnstad, O.N. 2004. Spatial synchrony in population
580 dynamics. *Ann. Rev. Ecol. Evol. Syst.* **35**:467-490.
- 581 Ludsin, S.A., DeVanna, K.M., and Smith, R.E.H. 2014. Physical-biological coupling and the
582 challenge of understanding fish recruitment in freshwater lakes. *Can. J. Fish. Aquat. Sci.*
583 **71**: 775-794.
- 584 Madenjian C.P., Höök, T.O., Rutherford, E.S., Mason, D.M., Croley, T.E., Szalai, E.B., and
585 Bence, J.R. 2005. Recruitment variability of alewives in Lake Michigan. *Trans. Am.*
586 *Fish. Soc.* **134**: 218-230.
- 587 Magnuson, J.J., Webster, K.E., Assel, R.A., Bowser, C.J., Dillon, P.J., Eaton, J.G., Evans, H.E.,
588 Fee, E.J., Hall, R.I., Mortsch, L.R., Schindler, D.W., and Quinn, F.H. 1997. Potential
589 effects of climate change on aquatic systems: Laurentian Great Lakes and Precambrian
590 Shield region. *Hydrol. Processes* **11**: 825-871.
- 591 MathWorks. 2015. MATLAB software, version R2015b [online]. Available from
592 <http://www.mathworks.com/products/matlab/> [Accessed 19 November 2015].

- 593 Miller, T.J., Crowder, L.B., Rice, J.A., and Marschall, E.A. 1988. Larval size and recruitment
594 mechanisms in fishes: toward a conceptual framework. *Can. J. Fish. Aquat. Sci.* **45**:
595 1657-1670.
- 596 Moran, P.A.P. 1953. The statistical analysis of the Canadian lynx cycle. *Aust. J. Zool.* **1**:291–
597 298.
- 598 Myers, J. T., Yule, D.L., Jones, M.L., Ahrenstorff, T.D., Hrabik, T.R., Claramunt, R.M., Ebener,
599 M.P., and Berglund, E.K. 2015. Spatial synchrony in cisco recruitment. *Fish. Res.* **165**:
600 11-21.
- 601 O'Brien, T.P., Taylor, W.W., Roseman, E.F., Madenjian, C.P., and Riley, S.C. 2014. Ecological
602 factors affect rainbow smelt recruitment in the main basin of Lake Huron, 1976-2010.
603 *Trans. Am. Fish. Soc.* **143**: 784-795.
- 604 Phelps, Q.E., Graeb, B.D.S., and Willis, D.W. 2008. Influence of the Moran effect on
605 spatiotemporal synchrony in common carp recruitment. *Trans. Am. Fish. Soc.* **137**:1701-
606 1708.
- 607 Plattner, S., Mason, D.M., Leshkevich, G.A., Schwab, D.J., and Rutherford, E.S. 2006.
608 Classifying and forecasting coastal upwellings in Lake Michigan using satellite derived
609 temperature images and buoy data. *J. Great Lakes Res.* **32**:63-76.
- 610 Pole, A., West, M., and Harrison, J. 1994. Applied Bayesian forecasting and time series analysis.
611 Chapman and Hall, New York, NY.
- 612 Post, J.R., and Evans, D.O. 1989. Size-dependent overwinter mortality of young-of-the-year
613 yellow perch (*Perca flavescens*) – laboratory, in situ enclosure, and field experiments.
614 *Can. J. Fish. Aquat. Sci.* **46**: 1958-1968.

- 615 Post, J.R. 1990. Metabolic allometry of larval and juvenile yellow perch (*Perca flavescens*): in
616 situ estimates and bioenergetics models. Can. J. Fish. Aquat. Sci. **47**: 554-560.
- 617 R Development Core Team. 2015. R: A language and environment for statistical computing
618 [online]. Available from <http://www.R-project.org> [Accessed 19 November 2015].
- 619 Redman, R.A., Czesny, S.J., and Dettmers, J.M. 2011. Old tales in recent context: current
620 perspective on yellow perch recruitment in Lake Michigan. Trans. Am. Fish. Soc. **140**:
621 1277-1289.
- 622 Rice, J.A., Crowder, L.B., and Holey, M.E. 1987. Exploration of mechanisms regulating larval
623 survival in Lake Michigan bloater: a recruitment analysis based on characteristics of
624 individual larvae. Trans. Am. Fish. Soc. **116**:703-718.
- 625 Rodionov, S., and Assel, R.A. 2003. Winter severity in the Great Lakes: A tale of two
626 oscillations. Clim. Res. **24**: 19–31, doi:10.3354/cr024019
- 627 Ruetz, C.R., Trexler, J.C., Jordan, F., Loftus, W.F., and Perry, S.A. 2005. Population dynamics
628 of wetland fishes: spatio-temporal patterns synchronized by hydrological disturbance. J.
629 Anim. Ecol. **74**: 322-332.
- 630 Rusak, J.A., Yan, N.D., and Somers, K.M. 2008. Regional climatic drivers of synchronous
631 zooplankton dynamics in north-temperate lakes. Can. J. Fish. Aquat. Sci. **65**: 878-889.
- 632 SAS Institute Inc. 2015. SAS/STAT software, version 9.4 [online]. Available from www.sas.com
633 [Accessed 19 November 2015].
- 634 Scott, W.B., and Crossman, E.J. 1973. Freshwater fishes of Canada. Fisheries Research Board
635 of Canada Bulletin 184, Ottawa, Canada.

- 636 Stachura, M.M., Essington, T.E., Mantua, N.J., Hallowed, A.B., Haltuch, M.A., Spencer, P.D.,
637 Branch, T.A., and Doyle, M.J. 2014. Linking northeast Pacific recruitment synchrony to
638 environmental variability. *Fish. Oceanogr.* **23**: 389-408.
- 639 Straile, D., Eckmann, R., Jungling, T., Thomas, G., and Loffloer, H. 2007. Influence of climate
640 variability on whitefish (*Coregonus lavaretus*) year-class strength in a deep, warm
641 monomictic lake. *Oecologia* **151**: 521-529.
- 642 Straile, D., van Nes, E.H., and Hosper, H. 2001. Climatic warming causes regime shifts in lake
643 food webs. *Limnol. Oceanogr.* **46**: 1780-1783.
- 644 Suski, C.D., and Ridgway, M.S. 2007. Climate and body size influence nest survival in a fish
645 with parental care. *J. Anim. Ecol.* **76**: 730-739.
- 646 Tin, H.T., and Jude, D.J. 1983. Distribution and growth of larval rainbow smelt in eastern Lake
647 Michigan, 1978-1981. *Trans. Am. Fish. Soc.* **112**: 517-524.
- 648 Vanderploeg, H.A., Bolsenga, S.J., Fahnenstiel, G.L., Liebig, J.R., and Gardner, W.S. 1992.
649 Plankton ecology in an ice-covered bay of Lake Michigan: utilization of a winter
650 phytoplankton bloom by reproducing copepods. *Hydrobiologia* **243/244**: 175-183.
- 651 Walters, C.J. 1987. Nonstationarity of production relationships in exploited populations. *Can. J.*
652 *Fish. Aquat. Sci.* **44**(Suppl. 2): 156-165.
- 653 Winder, M., and Schindler, D.E. 2004. Climate change uncouples trophic interactions in an
654 aquatic ecosystem. *Ecology* **85**: 2100-2106.
- 655 Wolter, K., and Timlin, M.S. 1998. Measuring the strength of ENSO events - how does 1997/98
656 rank? *Weather* **53**: 315-324
- 657 Wood, S.N. 2004. Stable and efficient multiple smoothing parameter estimation for generalized
658 additive models. *J. Amer. Stat. Assoc.* **99**: 673-686.

Table 1. Early life history characteristics of four species in Lake Michigan for which recruitment residuals were estimated for the 1973 to 2011 year-classes. Information for spawning time and location derive from Scott and Crossman (1973) and Auer (1982).

Species	Bloater	Rainbow smelt	Yellow perch	Alewife
Spawning time	February-March	April-May	May-June	June-August
Spawning location	> 80 m bottom depth	Tributaries, nearshore	Nearshore	Nearshore
When larvae are pelagic	June-August (Rice et al. 1987)	May-August (Jude et al. 1979; Tin and Jude 1983)	June-August (Dettmers et al. 2005; Beletsky et al. 2007)	June-August (Jude et al. 1979; Höök et al. 2007)

Table 2. Indices used to estimate stock size and recruit density for each species, and the accompanying reference if index was same as previously published study.

Species	Stock size index	Recruit index
Alewife	Biomass > 150 mm TL (Collingsworth et al. 2014)	Age-3 biomass (Collingsworth et al. 2014)
Bloater	Population egg production (Collingsworth et al. 2014)	Age-3 density (Bunnell et al. 2006)
Rainbow smelt	Biomass > 90 mm TL (Feiner et al. 2015)	Age-0 density, varying size cut-offs (Feiner et al. 2015)
Yellow perch	Biomass > 170 mm TL ^{&}	Age-0 density, < 100 mm TL

[&]Lauer et al. (2005) indicated 170 mm was minimum size of maturity for females

Draft

Table 3. Summary of stock-recruit model results that were used to estimate residuals. For all species but yellow perch, the Ricker form was used as the stock-recruit model.

Species	a	b	r^2
Alewife	2.63	0.10	0.05
Bloater	-2.18	6.64^{-4}	0.30
Rainbow smelt	4.56	0.16	0.19
Yellow perch	1.34	0.05	0.08

Table 4. Each row summarizes the results of generalized additive models for different response variables (identified in the first column). The second column indicates the number of candidate models (out of a possible 180) that had ΔAIC values less than 2. The third column indicates how much variance was explained in the model with the smallest AIC; the parentheses indicate how many of the potential eight explanatory variables were included in this model. Columns 4-11 indicate the percentage of models for which the given explanatory variables was included among the models with ΔAIC values less than 2. Explanatory variables that occurred in at least 2/3 of the top-ranked models are **bolded**.

Response variable	# models with $\Delta AIC < 2$	% deviance explained in model with AIC_{min}	Ice	Temp	Wind	LakeLevel	NAO	ENSO	PDO	Temp $\times$ LakeLevel
Alewife	3	61 (5)	33	100	100	100	100	0	0	67
Rainbow smelt	8	20 (2)	13	25	88	13	0	13	63	0
Bloater	6	37 (4)	33	50	100	0	67	0	100	0
Yellow perch	21	14 (2)	33	100	29	52	19	19	29	0
Average	14	35 (3)	29	79	14	50	100	7	43	14

Fig. Legends.

Fig. 1. Stock-recruit models developed from USGS bottom trawl survey data (1973-2014) in Lake Michigan and used to estimate recruitment residuals for alewife (a), rainbow smelt (b), bloater (c), and yellow perch (d).

Fig. 2. Time series of standardized recruitment residuals (on the natural-log scale) for alewife (a), rainbow smelt (b), bloater (c), and yellow perch (d).

Fig. 3. Panel (a) plots the standardized recruitment residuals for each species (filled circle = alewife, open circle = rainbow smelt, filled triangle = bloater, open triangle = yellow perch), and the line represents the average residual across the four species. Panel (b) plots the percentage of the four species with positive (dark shading) or negative (light shading) residuals for each year class.

Fig. 4. Scatter plots of recruitment residuals for all possible combinations of species pairs between alewife, bloater, rainbow smelt, and yellow perch. Pearson correlation coefficients shown; associated P-values were ≤ 0.03 for the top two panels and ≥ 0.14 for the bottom four panels.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. S1. Spectral plot for bloater recruitment residuals in Lake Michigan, 1973-2011. Time series of recruitment residuals were detrended (narrow plot above) and taller spectral plots have period on the x-axis and energy (signal²/frequency) on the y-axis, with surrounding 80% confidence intervals. Although the most likely period in the data occurred between 20 and 40 years, this spectral peak was not statistically significant.

Fig. S2. Spectral plot for yellow perch recruitment residuals in Lake Michigan, 1973-2011. Time series of recruitment residuals were detrended (narrow plot above) and taller spectral plots have period on the x-axis and energy (signal²/frequency) on the y-axis, with surrounding 80% confidence intervals. Although the most likely period in the data occurred around 20 years, this spectral peak was not statistically significant.

Fig. S3. Time series of environmental variables used to explain variation in the average recruitment residuals for the 1973-2011 year-classes of four fish species in Lake Michigan. Lake level (panel a) is averaged between April and July. Panel b) is annual maximum ice cover. Water temperature (panel c) and wind speed (panel d) are averaged between May and August. NAO (North Atlantic Oscillation, panel e), ENSO (El Nino, panel f), and PDO (Pacific Decadal Oscillation, panel g) were indexed over December, January, and February for the year in which January and February occurred.

Fig. S4. Spectral analysis for the time series of three explanatory variables that were temporally autocorrelated in Lake Michigan: spring and summer water temperatures (red line), lake level (blue line), and North Atlantic Oscillation (green line). Also plotted is the time series for the recruitment residuals averaged for bloaters, alewife, rainbow smelt, and yellow perch in Lake Michigan (black line) across the 1973-2011 year-classes. The most likely period (x-axis) is indicated by the peak with the highest energy (y-axis) for each time series, but none of these spectral peaks were significant.

Fig. S5. Relationship between the alewife recruitment residuals and the five explanatory variables in the top-ranked generalized additive model. The solid line represents the estimated non-linear smoothed function, dashed lines represent 95% confidence intervals, and the rug plot along the bottom indicates the observed values for each explanatory variable.

Fig. S6. Relationship between rainbow smelt recruitment residuals and three of the top-ranked generalized additive models that illustrate the shape of the relationship between explanatory variables and residuals: a) top-ranked model, b) 3rd-ranked model ($\Delta\text{AIC} = 1.04$) and first that includes water temperature, c) 8th-ranked model ($\Delta\text{AIC} = 1.93$), and first that includes lake level. The solid line represents the estimated non-linear smoothed function, dashed lines represent 95% confidence intervals, and the rug plot along the bottom indicates the observed values for each explanatory variable.

Fig. S7. Relationship between bloater recruitment residuals and three of the top-ranked generalized additive models that illustrate the shape of the relationship between explanatory variables and residuals: a) top-ranked model, b) 3rd-ranked model ($\Delta AIC = 0.45$) and first that includes water temperature. The solid line represents the estimated non-linear smoothed function, dashed lines represent 95% confidence intervals, and the rug plot along the bottom indicates the observed values for each explanatory variable.

Fig. S8. Relationship between yellow perch recruitment residuals and three of the top-ranked generalized additive models that illustrate the shape of the relationship between explanatory variables and residuals: a) top-ranked model, b) 2nd-ranked model ($\Delta\text{AIC} = 0.43$) and first that includes wind speed and lake level, c) 10th-ranked model ($\Delta\text{AIC} = 1.31$), and first that includes NAO. The solid line represents the estimated non-linear smoothed function, dashed lines represent 95% confidence intervals, and the rug plot along the bottom indicates the observed values for each explanatory variable.

Fig. S9. Relationship between average recruitment residuals and three of the top-ranked generalized additive models that illustrate the shape of the relationship between explanatory variables and residuals: a) top-ranked model, b) 2nd-ranked model ($\Delta AIC = 0.06$) and first that includes lake level, c) 9th-ranked model ($\Delta AIC = 1.83$), and first that includes wind speed. The solid line represents the estimated non-linear smoothed function, dashed lines represent 95% confidence intervals, and the rug plot along the bottom indicates the observed values for each explanatory variable.

c)