Canadian Journal of Forest Research
Revue canadienne de recherche forestière

**Thinning effects on foliar elements in eastern hemlock –
implications for managing the spread of the hemlock woolly
adelgid.**

Journal:	<i>Canadian Journal of Forest Research</i>
Manuscript ID	cjfr-2016-0260.R1
Manuscript Type:	Article
Date Submitted by the Author:	19-Sep-2016
Complete List of Authors:	Piatek, Kathryn; Editmyscience.com, Fajvan, Mary Ann; USDA Forest Service Turcotte, Richard; USDA Forest Service,
Keyword:	Eastern hemlock, foliar elements, foliar nutrients, thinning effects, hemlock woolly adelgid

Thinning effects on foliar elements in eastern hemlock – implications for managing
the spread of the hemlock woolly adelgid.

Kathryn B. Piatek^{1*}, Mary Ann Fajvan², and Richard M. Turcotte²

This manuscript is dedicated to the memory of Dan Snider and Tom Elliot.

Corresponding Author:

*Kathryn B. Piatek

Division of Forestry,

West Virginia University

Morgantown, WV 26506

¹Current Address:

Editmyscience.com

Morgantown, WV 26501

E-mail: kbpiatek@gmail.com

² Mary Ann Fajvan

USDA Forest Service,

Northern Research Station

180 Canfield St.

Morgantown, WV 26505

Phone: 304-285-1575

27 E-mail: mfajvan@fs.fed.us

28

29 ³ Richard M. Turcotte

30 USDA Forest Service,

31 State and Private Forestry

32 180 Canfield St.

33 Morgantown, WV 26505

34 Phone: 304-285-1575

35 E-mail: rturcotte@fs.fed.us

36

Draft

Abstract

Stand thinning is being tested as means to limit the impacts of the invasive hemlock woolly adelgid (*Adelges tsugae* Annand) on eastern hemlock (*Tsuga canadensis* (L.) Carriere) (HWA). The efficacy of this strategy may be reduced if thinning increases hemlock foliar nutrients because HWA densities are correlated with foliar concentrations of N, P, K, Ca, and Mn. We determined foliar N, P, K, Ca, and Mn concentrations in 1-year-old and all other (older) needles prior to and for 4 years after thinning in northwestern Pennsylvania stands of eastern hemlock without HWA. Average foliar concentrations in 1-year-old needles were: 1.30-1.80 g N 100g⁻¹, 1300-1700 mg P kg⁻¹, 4200-6300 mg K kg⁻¹, 2500 to 5200 mg Ca kg⁻¹, and 2393 µg Mn g⁻¹. N, P, and K decreased, Ca increased, and Mn first increased, then stabilized. Thinning by itself did not affect the tested foliar nutrients. Treatment*year interaction was significant and evident in temporal trajectories of foliar N and K. However, the differences between thinned and unthinned plots within years averaged only 0.03 g N 100g⁻¹ and 340 mg K kg⁻¹. We concluded that even though thinning changed the temporal trajectories of foliar N and K, the nutritional shifts were minimal, brief, and unlikely to affect the efficacy of thinning in limiting the impacts of HWA.

Keywords:

Eastern hemlock, foliar elements, foliar nutrients, thinning, hemlock woolly adelgid.

Introduction

The hemlock woolly adelgid, (HWA) *Adelges tsugae* Annand is the single greatest threat to the sustainability of eastern and Carolina hemlock (*Tsuga canadensis* and *T. caroliniana*, respectively) in eastern North America (Knauer et al. 2002). HWA affects photosynthesis, carbon allocation, foliar nutrients, and water relations and interacts with environmental variables (i.e. geographic location, drought) to cause eventual tree and stand mortality (Pontius et al. 2008, Domec et al. 2013). Widespread stand mortality due to HWA infestations results in possibly irreversible changes to hemlock ecosystem form and function (Koch et al. 2006, Orwig et al. 2008, Jonas et al. 2012).

Foliar chemistry of eastern hemlock has been highly correlated with HWA herbivory (Pontius et al. 2006, Miller–Pierce et al. 2010). Percent infestation was positively correlated with foliar Ca, K, Mn, N, and P, with N having the strongest relationship (Pontius et al. 2006). Incorporating foliar N, P, Ca, and K levels in stand health models greatly improved the ability to predict hemlock susceptibility to HWA, and suggested that foliar nutrition may determine hemlock susceptibility to HWA. Further, differences in foliar chemistry between eastern hemlock and other, HWA-resistant hemlock species indicated the importance of foliar nutrients to resistance to HWA. Thus, resistant species all had significantly lower N than eastern hemlock (Pontius et al. 2006), suggesting that low foliar N may decrease the nutritive value for the insect.

The effects of HWA on the rate of hemlock-growth decline and mortality have been correlated with tree crown variables such as live-crown ratio (Rentch et al. 2009). Silvicultural

manipulations that reduce tree competition and increase the availability of light, soil nutrients, and water are expected to increase tree growth including crown area, vigor, and the ability to deal with stressors, including insect attack (Waring and Pitman 1985; Knauer et al. 2002; references in Clancy et al. 2004; Fajvan 2008). Therefore, promising biological and chemical control strategies for HWA could be combined with silvicultural treatments (Mayfield et al. 2015).

The effectiveness of silvicultural treatments for improving resource acquisition and tree vigor in hemlock stands is not well known. Thinning alone generally increased foliar N concentrations in conifer stands of different ages at various intervals of time following treatment; mixed outcomes were obtained for foliar P, K, Ca, and Mg (Thiebodou et al. 2000; Jonard et al. 2006; Inagaki et al. 2008). Where assessed, needle expansion over reference, and enhanced leaf area indices and rates of photosynthesis suggested an improvement in the crown condition of residual trees following stand thinning (Thiebodou et al. 2000; Jonard et al. 2006; Navarro et al. 2013).

Responses of hemlock foliar nutrients to thinning are unknown. However, if canopy nutrient levels increase as expected, residual trees may be more attractive to HWA than trees in unthinned stands. Therefore, while thinning may be a potential tool in an integrated strategy to manage HWA, lack of information about thinning effects on hemlock foliar nutrients represents a limitation to an effective use of this practice in hemlock stands. An improved understanding of hemlock foliar nutrient response to thinning will help determine whether silvicultural treatments may be a viable option in an integrated control of HWA.

A long-term project to determine the efficacy of silvicultural thinning in reducing hemlock susceptibility to HWA has been established at the Allegheny National Forest (ANF) in northwestern Pennsylvania in HWA-free, hemlock-hardwood stands containing a minimum of 30 percent hemlock basal area, and at least 5-years prior to an estimated HWA invasion (Fajvan 2008). A supporting short-term objective was to determine whether thinning results in changes in foliar chemistry that may increase susceptibility of these stands to HWA infestation. Specifically in this study, we wanted to determine whether thinning increases macro nutrient levels in the foliage of residual eastern hemlock trees in recently-formed and old foliage. Our hypothesis was that thinning increases macronutrient levels in hemlock foliage. We collected foliage before, and for 4 years after thinning to address our hypothesis.

Methods

Study area

Our study site is located in The Allegheny National Forest in northwestern Pennsylvania, U.S.A., near the town of Warren, 41.64976°N, 79.03913°W. The forest covers about 210,000 ha (517,000 acres) of land, of which 90% is forested. Average elevation is 427 m (1500 ft). Soils in the project area belong to two main soil groups - Cavode silt loams with 0 to 8 % slope, and Cookport very stony silt loams, with 0 to 15% slope <http://websoilsurvey.usda.gov/>; Accessed [11-12-13]. Vegetation consists of second-growth, 70-100 years old Appalachian Mixed Hardwood species, including black cherry (*Prunus serotina*), red maple (*Acer rubrum*), black birch (*Betula lenta*), northern red oak (*Quercus rubra*), and sugar maple (*Acer saccharum*). Hemlock is found either in pure patches or dispersed within the hardwoods.

126
127 ANF has moderate climate, with warm and humid summers. Average temperatures range from
128 23.9 to 26.7° C (75-80° F) in summer to -3.9 to -6.7° C (20-25° F) in winter, and mean annual
129 precipitation ranges from 889 to 1,295 mm (35 to 51 in) (<http://www.fs.usda.gov/> Accessed [2-
130 22-13]).

131
132 ***Experimental design: thinning***
133 Three, 24-ha stands were designated as part of a large silvicultural efficacy study. Stands
134 contained tree species associated with the Allegheny Hardwood forest type (Society of American
135 Foresters Type 28). Stands were overstocked with total basal areas ranging from 39 to 46 m² ha⁻¹
136 ¹. Stands were further distinguished based on the overstory dominance of either black cherry or
137 red oak and red maple. Hemlock content ranged from 11.5 to 16 m² ha⁻¹ within the black cherry-
138 dominated stands, and from 7 to 11.5 m² ha⁻¹ in the oak-red maple – dominated stands (Fig. 1).

139
140 Each stand was divided into 6, approximately 4-ha treatment blocks. Three 4-ha blocks were
141 then randomly selected for thinning, and three for reference. Trees were marked for thinning
142 according to stocking guidelines based on field inventories. Due to the initial high stand
143 densities, a removal of about 35 percent of the basal area was planned (Lancaster 1985) (Fig. 1).
144 In each block, ten, 0.04-ha circular plots were randomly established (approximately 1 per 0.4
145 ha). The center of each plot consisted of a hemlock tree that was targeted for crown release (e.g.
146 had 3-4 surrounding stems marked for removal). A similar procedure was used to establish plots
147 destined to be used as reference (no thinning). The 10 trees in each block were designated as

“subject” trees, for a total of 60 per stand. A commercial thinning was conducted in early to mid-2007. Trees were felled, and non-commercial sized stems were left on site.

Experimental design: foliar nutrients

To ensure adequate representation for foliar analyses, “blocks” were treated as plots, or units, to which treatment was applied (see statistical analyses below), while individual “subject” trees within treatment “blocks” were used for subsampling. In each stand (1, 2, 3), we randomly designated two thinned (out of three) and two reference plots (out of three) for foliar sampling. Thus, foliar sampling was conducted in a total of 6 treatment and in 6 reference plots. Within plots, green hemlock foliage was collected from 6 randomly-chosen subject trees in each treatment, and from 4 trees in each reference plot in 2006 (pre-thinning), and in 2008, 2009, 2010, and 2011 (post-thinning). Thus, 36 trees were sampled across thinned blocks, and 24 trees across reference blocks in each sampling year. More trees were sampled in the treated plots in expectation that more variability in the measured parameters would be introduced by the thinning treatment. We sampled in the first week of April, prior to bud break, and after snowmelt. A height pole was used to collect at least 25-cm long sections from the ends of branches about 10 meters above ground. Care was taken in obtaining branches from east and west sides of tree and upper 1/3 of the crown which was partly-sunlit; however, that was not always possible due to tree and crown size and architecture.

Branches were labeled, placed in trash-size plastic bags, and refrigerated until processing. Branches from both sides of individual trees were composited into one sample per tree. Last-formed twig extensions, identified as light-brown twig endings to the last-formed bud swelling,

were cut with scissors and composited per branch and tree into '1-year-old' foliage class in paper bags. The rest of the branch with older green foliage (any age class) was composited into 'older' age class.

Needles on twigs in paper bags were dried at 65-70°C to a constant weight for up to 3 days. Then foliage was separated from twigs by shaking the bag until needles fell off, and discarding any woody pieces. Subsequently, needles were ground in a dedicated coffee grinder to a fine powder. These samples were submitted for nutrient analysis to Spectrum Analytic (Washington Court, Ohio). Total N, P, K, Ca, and Mn concentrations were analyzed in the foliage. Protocol at Spectrum Analytic includes grinding of sample in a Wiley stainless steel mill to pass a 40 mesh screen and analyzing as follows. For plant N, 6-8 mg of sample is weighed on a Sartorius micro balance, and analyzed for N on an ECS 4010 CHNSO Analyzer (Costech). Quality is controlled with NBS Citrus leaves and an in-house check sample with each batch of plant samples. For other nutrients, a 0.5 gram sample is measured, 10 ml of concentrated HNO₃ acid is added, and the samples are microwave-digested. Subsequently, samples are removed, 10 ml of water is added and the samples mixed. The samples are then analyzed on iCAP 6500 - ICP-OES CID Spectrometer (Thermo Scientific).

Analyses

All response variables were checked for normality (PROC UNIVARIATE, SAS, Inc. v. 9.02), then analyzed. We analyzed element concentrations as dependent variables in a randomized block design using a linear mixed model with fixed terms for the main effects of TREATMENT

and sampling YEAR, and their interaction. A quadratic function was also tested for sampling YEAR, but was found not significant, and subsequently dropped from the model. Needle age (age) was included at the split-plot level. The final form of the model was:

Concentration = treatment, year, treatment*year, age, treatment*age, year* age.

YEAR was the repeated measures factor.

We first examined pre-treatment year 2006 as a covariate, but it was not significant, and data obtained in 2006 were then used in the repeated measures. Because measurement years were not evenly spaced (2006, then 2008, 2009, 2010, and 2011), we considered only those covariance structures appropriate for unequally-spaced repeated measures.

Covariance structures were modeled using the REPEATED statement in the PROC MIXED procedure with SAS software, Version [9] (Littell et al. 1996; Copyright © [2014] SAS Institute Inc.) We tested unstructured, Toeplitz variance components, spatial power, spatial Gaussian, and spatial spherical covariance structures. Smallest AIC and BIC diagnostics were used to assess the most appropriate model fit (Burnham and Anderson, 2002). Unstructured covariance structure had the best fit statistics for all elements, and we used it to model nutrient concentrations as a function of treatment, year, and treatment*year interaction, with needle age as split-plot effect. We used Kenward-Rogers degrees of freedom method. Residuals were tested for homogeneity of variances using the Levene's test. Levene's test indicated that residuals were not homogenous for the thinning treatment or year; however, unstructured covariance was designed for non-homogeneity, therefore, satisfying the requirements.

Results

Effects of thinning on foliar nutrients

Treatment, as a main effect, was not significant for the examined foliar elements (Table 1). Treatment and year interaction was significant for foliar concentrations of N and K (Table 1). Specifically for N concentration, the most evident response was a significant decrease over time, regardless of treatment, with N concentration significantly lower in 2010 and 2011 than in 2006, 2008, or 2009 (Fig. 2). The decrease was subject to annual variability, and N concentration in 2009 was higher than that in 2008, although these differences were statistically significant only within treatment. Temporal trajectories of N concentrations for treated and non-treated plots differed slightly, indicating that treatment minimally modified how plot types responded to “year”; thus, N concentration in the thinning treatment was slightly lower than that in the reference in 2006 and 2011, and higher in 2008, 2009, and 2010 (Fig. 2). These differences between plot types were not statistically significant and averaged 0.03 g N 100g⁻¹ within years.

In both - treated and non-treated plots, K concentration increased from 2006 to 2008, subsequently decreased (2009, 2010, and 2011), and was subject to a high degree of variability, with a significant overall low in 2011 and in the reference in 2010 (Fig. 3). K concentration was consistently higher in the thinned plots than in the reference in all years, but particularly in 2010, three years post-treatment, resulting in a significant treatment * year interaction. However, the differences were not statistically significant within year (Table 1, Fig. 3), even in 2010. These results did not indicate a main effect of treatment.

Effects of time on foliar nutrients

The “year” effect was significant for all foliar element concentrations (Table 1), and it was further modified by interactions. Foliar element concentrations were highly variable among years (Table 2), and the most evident response was a significant decrease in N, P, and K, an increase in Ca, and an initial increase then stabilization of Mn over time, regardless of treatment (Table 2). Decreases were subject to annual variation (the interaction effect), and for example, N concentration in 2009 was higher than that in 2008, and that in 2011 was higher than in 2010.

Effects of needle age on foliar nutrients

Needle age was significant as a main effect, and 1-year-old needles (young) had significantly higher N, P, and K, and lower Ca and Mn concentrations than older needles in most years (Table 2). Needle age also interacted with year (Table 1). Lower concentrations of N, P, and K in older needles in 2006 and 2008 became comparable to those of the younger needles in and after 2009 (Table 2). Foliar Ca was generally higher in older needles in all years except in 2009 when Ca in older needles was comparable to that of younger needles (Table 2). Mn in older needles was uniformly higher than in young needles.

Discussion

Hemlock foliar-nutrient responses to thinning

We anticipated an increase in foliar macronutrient concentrations following treatment in our study because thinning shifts available soil resources - water and mineral nutrients - to the remaining biomass (Misson et al. 2005; Navarro et al. 2013). Thus, increases in foliar N were reported for different coniferous species at various intervals following thinning, and ranged from 0.3 mg N g⁻¹ dry weight in current-year needles of ~57-years-old *Picea abies* (Jonard et al. 2006)

263 to 4.8 mg N g⁻¹ in 20-years-old *Abies balsamea* in well-drained soils (Thiebodou et al. 2000).
264 Positive responses in foliar P, and decreases, increases, and no responses were observed in foliar
265 K, and Ca after thinning in a variety of conifers, on diverse sites, and after different thinning
266 regimes (Wollum and Schubert, 1975; Harrington and Wierman, 1990; Thiebodou et al. 2000;
267 Lopez-Serrano et al. 2005). In our study, however, thinning by itself did not affect foliar nutrient
268 concentrations in residual hemlock trees for 4 consecutive years following treatment.
269
270 Thinning resulted in small and variable changes in the trajectories of foliar elements over time
271 since treatment, as indicated by significant interactions of treatment with time. The main
272 response, however, was a significant decrease in N, P, and K over time. We attribute this
273 decrease to a dilution of foliar nutrient concentration associated with a slowly-expanding crown
274 biomass (the same pool of nutrients in a larger biomass) (Switzer and Nelson, 1972). Canopy
275 expansion in both reference and thinned stands in this study was observed, and was further
276 supported by diameter increases, especially in the thinning treatment (data not shown).
277 Therefore, foliar nutrient outcomes need to be evaluated in light of expanding crowns in both
278 plot types (thinned and reference), and apparent nutrient dilution.
279
280 The non-linear decrease in N concentrations over time (higher in 2009 than in 2008) suggested
281 that the decrease was modified by factors that were related to “year”, and probably driven by
282 climate. Foliar N concentration is determined primarily by soil N supply, derived from microbial
283 mineralization of soil organic matter; therefore, a climatic contribution to this process is highly
284 likely. The interaction effect of treatment with “year” may involve a stimulation of organic
285 matter turnover beyond the possible effects of annual variations in climate. Canopy thinning may

increase soil temperatures due to the opening of the canopy and increased solar irradiation of the forest floor, and can thus stimulate microbial turnover of organic matter and N production (Thibodeau et al. 2000). Further, inputs of labile C from root death and leaching from fresh foliage in the thinned areas are likely to contribute to an increase in the mineralization of organic matter because soil microbial processes are primarily driven by substrate chemistry. Increases in mineralization rates in the thinned plots were confirmed with a related in-situ soil mineralization study (Farcas 2010).

The high annual variability in foliar K concentration was probably due to the high mobility of this element within green tissues (Tripler et al. 2006), its potential ability to leach across membranes (Schreeg et al. 2013), and possibly high influx of K from older tissues (Weatherall et al. 2006). None of these processes have been described for mature eastern hemlock, so their relative importance is uncertain. However, the consistently higher (though not significant) K concentration in the treatment than in the reference plots may be associated with increased overall mobility due to mobilization of resources post treatment.

Young needles are typically associated with higher N, P, and lower Ca and Mn than found in old needles, and that was generally the case in our study. Calcium in particular is a structural component of cell walls, and, together with Mn, is not mobile in plant tissues. Both Ca and Mn accumulate over time in older green tissues (Fink, 1991).

The annual averages of foliar nutrients observed in our study fit well within value ranges found at other sites. Thus, hemlock foliar N exhibited a considerable range of concentrations from a

low of 1.05 g 100g⁻¹ (Aber and Martin, 1999) to a high of 1.92 g 100g⁻¹ (Pontius et al. 2006). Nitrogen levels in our study were in the middle of this wide range in 2011, 4-years post-treatment (at 1.32 g 100g⁻¹), and close to the high values in 2006, the pre-treatment year (at 1.82 g 100g⁻¹). Foliar P ranged from 995 mg kg⁻¹ in New Hampshire (Aber and Martin, 1999) to 4,318 mg kg⁻¹ in HWA-susceptible population (Pontius et al. 2006). Foliar P in our study was at the low end of reported values at an average of 1,500 mg kg⁻¹. Foliar K ranged from a low of 4,400 mg kg⁻¹ in our study in 2006 to a high of 9,433 mg kg⁻¹ in New York (Yorks, 2001). Foliar Ca values in our study in 2006 were below those of the HWA-resistant hemlock species (at 2,844 mg kg⁻¹) and higher than the HWA-susceptible ones (1,989 mg kg⁻¹) (Pontius et al. 2006). Interestingly, eastern hemlock seedlings can apparently maintain significantly higher Mn levels than other eastern conifer species, at 2124 µg g⁻¹, (St. Clair and Lynch, 2005), and this average falls within value ranges observed in our study.

Lack of a strong treatment response in foliar nutrients may be related to the ecophysiology of eastern hemlock. Hemlock is very tolerant of shade and is found throughout all canopy strata. Its growth rate is slow relative to species associates, and at high densities, it can survive in repressed conditions for hundreds of years (Godman and Lancaster 1990). Self-thinning is slow and canopy disturbance is typically required for accelerated growth to occur. However, following crown release, radial growth in hemlock can increase 5-8 % per year, regardless of tree age (Fajvan and Seymour 1993; Marchand and Filion 2014). Tree growth and foliar nutrient levels are generally correlated because both depend on available soil resources (Garrison et al. 2000; Jarvis and Linder 2000).

Foliar-nutrient implications for management of HWA

The observed foliar N, P, K, and Mn values in our study were well below those reported for HWA-susceptible trees (Pontius et al. 2006), indicating that these foliar nutrient levels would not be particularly appealing to HWA. Foliar Ca values in our study in 2006 were low, indicating a possible stand disadvantage for withstanding HWA (Pontius et al. 2006). The fact that foliar Ca levels in our stands appeared to be increasing with time since treatment may be encouraging (Fig. 3). At the same time, by itself, foliar Ca does not determine susceptibility to HWA (Pontius et al. 2006).

Lack of a pronounced response of foliar nutrients to thinning is advantageous for eastern hemlock. Eastern and Carolina hemlock (*Tsuga caroliniana*) stands have been decimated on thousands of acres from Georgia to Connecticut by the invasive insect – the hemlock woolly adelgid (USDA, Forest Service, 2005). Hemlock stands form a unique microclimate, with dense shade and cool temperatures, which comprises a critically-important element of bird and animal habitat; therefore, HWA-caused hemlock decline and loss may precipitate significant impacts on these species (DeGraaf et al. 1992, Yamasaki et al. 2000). Alteration of the physical environment (Ellison et al. 2005), including encroachment of invasive species (Orwig and Foster 1998), associated shifts in hydrologic processes (Daley et al. 2007, Ford and Vose 2007; Hadley et al. 2008), modification in nitrogen (N) export dynamics (Templer and McCann 2010), and changes in carbon pools and dynamics (Albani et al. 2010) have been observed following HWA infestation and hemlock mortality.

Our study was conducted in stands close to the HWA front, but without HWA presence. The insect generally spreads at a rate of about 12 km yr⁻¹ (Evans and Gregoire, 2006) but, until recently, its spread into the Allegheny Plateau of Pennsylvania had been curtailed by cold winters (Stadler et al. 2005, 2006). Several warm winters prior to 2014 have increased the rate of spread, and, currently, hemlock stands in the vicinity of the Allegheny National Forest boundary are infested. Based on published, long-term studies of HWA-infested stands, trees that survive HWA the longest are those with high live-crown ratios and competitive crown positions (Rentch et al. 2009). The goal of silvicultural thinning is to improve these same crown features. Hence improving these crown and stand density features prior to infestation may give residual trees a chance to survive longer. Thinning during an infestation would not be worthwhile because trees are already stressed and declining. Additionally, silvicultural treatments, such as thinning, may be the most economical means of managing this forest type against the deadly insect (USDA, Forest Service, 2005). Thinning generates revenue for the landowner and changes resource allocation within a stand, with more light, moisture, and nutrients accessible to the residual trees. Increased resource levels are associated with higher resistance to stressors, including insect herbivory (Waring and Pitman, 1985).

With the imminent entrance of HWA into these stands, it was critical to understand whether thinning, while producing desirable management and economic outcomes, may also result in increased foliage palatability for the HWA. Our findings suggest that thinning at the level performed in our study may be used in hemlock stands without risking HWA attack due to increased needle nutritional value. Other factors are likely to play a role in whether these stands

376 will be infested or not, among them - proximity to the HWA front and impacts of climate
377 change.

378

379

Draft

Acknowledgments

Funds for this study were provided by the USFS in Morgantown, WV. Initial work was conducted in the Division of Forestry, subsequently in the Biology Department, West Virginia University. Investigators are indebted to Dan Snider and Tom Elliott for collecting foliage for this study. The help of USFS technicians in processing foliage is gratefully acknowledged. The USFS Northern Research Station statistician, John Stanovick, assisted with data analysis.

Draft

References

- Aber, J. D., and Martin, M. 1999. Leaf Chemistry, 1992-1993 (ACCP). [Leaf Chemistry, 1992-1993 (Accelerated Canopy Chemistry Program)]. Data set. Available from <http://www.daac.ornl.gov> Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
- Albani, M., Moorcroft, P.R., Ellison, A.M., Orwig, D.A., and Foster, D.R. 2010. Predicting the impact of hemlock woolly adelgid on carbon dynamics of eastern United States forests. *Can.J.For.Res.* **40**: 119-133.
- Burnham, K.P., and Anderson, D.R. 2002. Model selection and multi-model inference: a practical information-theoretic approach. Second edition. Springer-Verlag, New York, USA. ISBN 978-0-387-22456-5. 351 pp.
- Clancy, K.M., Chen, Z., and Kolb, T.E. 2004. Foliar nutrients and induced susceptibility: genetic mechanisms of Douglas-fir resistance to western spruce budworm defoliation. *Can. J. For. Res.* **34**: 939-949.
- Daley, M.J., Phillips, N.G., Pettijohn, C., and Hadley, J.L. 2007. Water use by eastern hemlock (*Tsuga canadensis*) and Black birch (*Betula lenta*): implications of effects of the hemlock woolly adelgid. *Can. J. For. Res.* **37**: 2031-2040.
- DeGraaf, R.M., Yamasaki, M., Leak, W.B., and Lanier, J.W. 1992. New England wildlife: Management of forested habitats. Gen. Tech. Rep. NE-144. USDA Forest Service, Northeastern Forest Experiment Station. 271 pp.
- Domec, J.C., Rivera, L.N., King, J.S., Peszlen, I., Hain, F., Smith, B., and Frampton, J. 2013. Hemlock woolly adelgid (*Adelges tsugae*) infestation affects water and carbon relations of eastern hemlock (*Tsuga canadensis*) and Carolina hemlock (*Tsuga caroliniana*). *New Phytol.* **199**: 452-463.
- Ellison, A.M., Bank, M.S., Clinton, B.D., Colburn, E.A., Elliott, K., Ford, C.R., Foster, D.R., Kloeppel, B.D., Knoepp, J.D., Lovett, G.M., Mohan, J., Orwig, D.A., Rodenhouse, N.L., Sobczak, W.V., Stinson, K.A., Stone, J.K., Swan, C.M., Thompson, J., Von Holle, B., and Webster, J.R. 2005. Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Front. Ecol. Environ.* **3**(9): 479-486.
- Evans, A.M., and Gregoire, T. G. 2006. A geographically variable model of hemlock woolly adelgid spread. *Biol. Invasions* 12 pp.
- Fajvan, M.A., and Seymour, R.S. 1993. Canopy stratification, age structure, and development of multicohort stands of eastern white pine, eastern hemlock and red spruce. *Can. J. For. Res.* **23**: 1799-1809.
- Fajvan, M.A. 2008. The role of silvicultural thinning in eastern forests threatened by hemlock woolly adelgid (*Adelges tsugae*). In: Deal, R.L. (Tech. Ed.), Integrated restoration of forested ecosystems to achieve multiresource benefits: proceedings of the 2007 national silviculture workshop, Gen. Tech. Rep. PNW-733, USDA Forest Service, Pacific Northwest Research Station, pp. 247-256.
- Farcas, D. 2010. Site characterization of hemlock stands at Allegheny National Forest. M.Sc. thesis, College of Agriculture, Forestry, and Consumer Sciences, West Virginia University. 83 pp.
- Fink, S. 1991. The micromorphological distribution of bound calcium in needles of Norway spruce [*Picea abies* (L.) Karst.]. *New Phytol.* **119**: 33-40.

- Ford, C.R. and Vose, J.M. 2007. *Tsuga canadensis* (L.) Carr. Mortality will impact hydrologic processes in Southern Appalachian forest ecosystems. *Ecological Applications* **17**(4):1156-1167.
- Garrison, M.T., Moore, J.A., Shaw, T.M., and Mika, P.G. 2000. Foliar nutrient and tree growth response of mixed-conifer stands to three fertilization treatments in northeast Oregon and north central Washington. *For. Ecol. Manage.*, **132**: 183–198.
- Godman, R.M., and Lancaster, K. 1990. *Tsuga canadensis* (L.) Carr. Eastern Hemlock Pinaceae. Pine Family. In: Burns, R.M. and B.H. Honkala (eds.), *Silvics of North America*, Vol. 1, Conifers. USDA, Forest Service, Agricultural Handbook 654. Washington, D.C.
- Hadley, J., Kuzeja, P.S., Daley, M.J., Phillips, N.G., Mulcahy, T., and Singh, S. 2008. Water use and carbon exchange of red oak- and eastern hemlock dominated forests in the northeastern USA: implications for ecosystem-level effects of hemlock woolly adelgid. *Tree Physiol.* **28**: 615-627.
- Harrington, C.A., and Wierman, C.A. 1990. Growth and foliar nutrient response to fertilization and precommercial thinning in a coastal western red cedar stand. *Can. J. For. Res.* **20**: 764-773.
- Inagaki, Y., Kuramoto, S., Torii, A., Shinomiya, Y., and Fukata, H. 2008. Effects of thinning on leaf-fall and leaf-litter nitrogen concentration in hinoki cypress (*Chamaecyparis obtusa* Endlicher) plantation stands in Japan. *For. Ecol. Manage.* **255**: 1859-1876.
- Jarvis, P., and Linder, S. 2000. Botany: Constraints to growth of boreal forests. *Nature* **405**: 904-905 doi:10.1038/35016154
- Jonard, M., Misson, M., and Ponette, Q. 2006. Long-term thinning effects on the forest floor and the foliar nutrient status of Norway spruce stands in the Belgian Ardennes. *Can. J. For. Res.* **36**(10): 2684–2695. doi:10.1139/X06-153.
- Jonas, S.Z., Xi, W., Waldron, J.D., and Caulson, R.N. 2012. Impacts of hemlock decline and ecological consideration for hemlock stand restoration following hemlock woolly adelgid outbreaks. *Tree and For. Sci. Biotech.* **6**: 22-26.
- Knauer, K., Linnane, J., Shields, K., and Bridges, R. 2002. An initiative for management of hemlock woolly adelgid. http://na.fs.fed.us/fhp/hwa/pubs/proceedings/2002_proceedings/initiative.pdf
- Koch, F.H., Cheshire, H.M., and Devine, H.A. 2006. Landscape-scale prediction of hemlock woolly adelgid, *Adelges tsugae* Homoptera: Adelgidae, infestation in the southern Appalachian Mountains. *Environm. Entomol.* **35**: 1313-1323.
- Lancaster, K.F. 1985. Managing eastern hemlock a preliminary guide. USDA Forest Service NA-FR-30, 5pp.
- Littell, R.C., Milliken, G.A., Stroup, W.W., and Wolfinger, R.D. 1996. SAS System for Mixed Models. SAS Institute, Inc. Cary, NC, U.S.A. 635 pp.
- López-Serrano, F.R., de las Heras, J., González-Ochoa, A.I., and García-Morote, F.A. 2005. Effects of silvicultural treatments and seasonal patterns on foliar nutrients in young post-fire *Pinus halepensis* forest stands. *For. Ecol. Manage.* **210** (1-3): 321- 336 doi: 10.1016/j.foreco.2005.02.042.
- Marchand, N. and Filion, L. 2014. A dendroecological analysis of eastern hemlock and white pine in relation to logging in La Mauricie National Park (Quebec, Canada). *The Forestry Chronicle* **90**(3): 351-360.
- Mayfield, A.E. III, Reynolds, B.C., Coots, C.I., Havill, N.P., Brownie, C., Tait, A.R., Hanula, J.L., Joseph, S.V. , and Galloway, A.B. 2015. Establishment, hybridization and impact of

- Laricobius predators on insecticide-treated hemlocks: Exploring integrated management of the hemlock woolly adelgid. *For. Ecol. Manage* **335**: 1-10.
- Miller-Pierce, M.R., Orwig, D.A., and Preissler, E. 2010. Effects of hemlock woolly adelgid and elongate hemlock scale on eastern hemlock growth and foliar chemistry. *Environm. Entomol.* **39**(2): 513-519.
- Misson, L., Tang, J., Xu, M., McKay, M., and Goldstein, A. 2005. Influences of recovery from clear-cut, climate variability, and thinning on the carbon balance of a young ponderosa pine plantation. *Ag. For. Meteorol.* **130**:207-222.
- Navarro, F.B., Romero-Freire, A., Del Castillo, T., Foronda, A., Jiménez, M.N., Ripoll, M.A., Sánchez-Miranda, A., Huntsinger, L., and Fernández-Ondoño, E. 2013. Effects of thinning on litterfall were found after years in a *Pinus halepensis* afforestation area at tree and stand levels. *For. Ecol. Manage.* **289**: 354-362.
- Orwig, D.A., and Foster, D.R. 1998. Forest response to the introduced hemlock woolly adelgid in southern New England, U.S.A. *J. Torrey Bot. Soc.* **125**: 60-73.
- Orwig, D.A., Cobb, R.C., D'Amato, A.W., Kizilinski, M.L., and Foster D.R. 2008. Multi-year ecosystem response to hemlock woolly adelgid infestation in southern New England Forests. *Can. J. For. Res.* **38**: 834-43.
- Pontius, J.A., Hallett, R.A., and Jenkins, J.C. 2006. Foliar chemistry linked to infestation and susceptibility to hemlock woolly adelgid (Homoptera: Adelgidae). *Environ. Entomol.* **35**(1): 112-120.
- Pontius, J., Hallett, R., Martin, M., and Plourde, L. 2008. A landscape scale GIS tool to assess eastern hemlock susceptibility to hemlock woolly adelgid infestation Online Encyclopedia of Forest Environmental Threats. Available from <http://www.forestencyclopedia.net/p/p3543>.
- Rentch, J., Fajvan, M.A., Evans, R.A., and Onken, B. 2009. Using dendrochronology to model hemlock woolly adelgid effects on eastern hemlock growth and vulnerability. *Biol. Invasions* **11**: 551-562.
- Schreeg, L.A., Mack, M.C., and Turner, B.L. 2013. Nutrient-specific solubility patterns of leaf litter across 41 lowland tropical woody species. *Ecology* **94**(1): 94-105.
- St. Clair, S.B., and Lynch, J.P. 2005. Element accumulation patterns of deciduous and evergreen tree seedlings on acid soils: implications for sensitivity to manganese toxicity. *Tree Phys.* **25**: 85-92.
- Stadler, B., Muller, T., Orwig, D., and Cobb, R. 2005. Hemlock Woolly Adelgid in New England Forests: canopy impacts transforming ecosystem processes and landscapes. *Ecosystems* **8**: 233-247.
- Stadler, B., Muller, T., and Orwig, D. 2006. The ecology of energy and nutrient fluxes in hemlock forests invaded by hemlock woolly adelgid. *Ecology* **87**(7): 1792-1804.
- Switzer, G.L., and Nelson, L.E. 1972. Nutrient Accumulation and Cycling in Loblolly Pine (*Pinus taeda* L.) Plantation Ecosystems: The First Twenty Years. *Soil Sci. Am. J.* **36**: 143-147.
- Templer, P.H., and McCann, T.M. 2010. Effects of the hemlock woolly adelgid on nitrogen losses from urban and rural northern forest ecosystems. *Ecosystems* **13**:1215-1226. DOI: 10.1007/s10021-010-9382-x.
- Thiebodou, L., Raymond, P., Camiré, C., and Munson, A.D. 2000. Impact of precommercial thinning in balsam fir stands on soil nitrogen dynamics, microbial biomass,

- decomposition, and foliar nutrition. *Can. J. For. Res.* **30**(2): 229–238
doi:10.1139/cjfr-30-2-229.
- Tripler, C.E., Kaushal, S.S., Likens, G.E., and Walter, M.T. 2006. Patterns in potassium dynamics in forest ecosystems. *Ecol Lett.* **9**(4): 451-466.
- U.S.D.A. Forest Service. 2005. Hemlock woolly adelgid (pest alert). NA-PR-09-05. U.S. Department of Agriculture Forest Service, Northeastern Area, State and Private Forestry, Newtown Square, Pennsylvania.
- Waring, R.H., and Pitman, G.B. 1985. Modifying lodgepole pine stands to change susceptibility to mountain pine beetle attack. *Ecology* **66**(3): 889-897.
- Weatherall, A., Proe M.F., Craig, J., Cameron, A.D., and Midwood, A.J. 2006. Internal cycling of nitrogen, potassium and magnesium in young Sitka spruce. *Tree Physiol.* **26**: 673-80.
- Wollum, A.G., and Schubert, G.H. 1975. Effect of thinning on the foliage and forest floor properties of ponderosa pine stands. *Soil Sci. Soc. Am. J.* **39**(5): 968-972.
- Yamasaki, M., DeGraaf, R.M., and Lanier, J.W. 2000. Wildlife habitat associations in eastern hemlock - birds, smaller mammals, and forest carnivores. In: McManus, K, A., Shields, K., S., Souto, D. R., eds. *Proc.: Symposium on sustainable management of hemlock ecosystems in eastern North America*. Gen. Tech. Rep. NE-267. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 135-143.
- Yorks, T.E. 2001. Effects of forest harvest, deer herbivory, and tree mortality on nutrient cycling in the Catskill Mountains of New York. PhD Dissertation, SUNY College of Environmental Science and Forestry, Syracuse, NY.

FIGURE CAPTIONS:

Figure 1. Pre- and Post-harvest diameter distribution by species in the three study stands at the Allegheny National Forest, near Warren, PA. Eastern hemlock (EH) are shown as solid grey cylinders. Species abbreviations are: EH – eastern hemlock (*Tsuga canadensis*), NRO – northern red oak (*Quercus rubra*), BC – black cherry (*Prunus serotina*), BB – black birch (*Betula lenta*), RM – red maple (*Acer rubrum*), AB – American beech (*Fagus sylvatica*).

Figure 2. Treatment * year interaction effect for N concentration. No sampling was conducted the year of treatment (2007; thinning). Means with the same letter are not significantly different from each other (Tukey–Kramer test, $P < 0.01$). N-re and N-th indicate N concentration in reference and in thinned plots, respectively. Poly. (N-re) and Poly. (N-th) show 2nd degree polynomial trendlines for N concentration in reference and in thinned plots, respectively.

Figure 3. Treatment * year interaction effect for K concentration. No sampling was conducted the year of treatment (2007; thinning). Means with the same letter are not significantly different from each other (Tukey–Kramer test, $P < 0.01$). K-re and K-th indicate K concentration in reference and in thinned plots, respectively. Poly. (K-re) and Poly. (K-th) show 2nd degree polynomial trendlines for K concentration in reference and in thinned plots, respectively.

Figure 1.

Figure 2.

Figure 3.

Table 1. Probability values for the effects of the thinning treatment (T), needle age (A), year, and their interactions on element concentrations in eastern hemlock needles at the Allegheny National Forest; df denotes degrees of freedom. Output from proc mixed procedure in SAS.

	df	N	P	K	Ca	Mn
Treatment (T)	1					
Needle Age (A)	1		< 0.0001	< 0.0001	< 0.0001	0.0016
T*A	1					
Year	4	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
T*Year	4	0.0094		0.0224		
A*Year	4	0.0002	<0.0001	<0.0001	<0.0001	<0.0001

Table 2. Concentration of selected nutrients in hemlock (*Tsuga canadensis*) needles at the Allegheny National Forest (Pennsylvania) during 5 years of study. Lsmeans are shown for the significant effect of needle age (1-year-old vs. all older needles) or year alone with corresponding p-values for the effect. Sampling in 2007 omitted due to stand thinning in treatment plots.

Year	N (g 100g ⁻¹)		P (mg kg ⁻¹)		K (mg kg ⁻¹)		Ca (mg kg ⁻¹)		Mn (µg g ⁻¹)	
	1-yr-old	older	1-yr-old	older	1-yr-old	older	1-yr-old	older	1-yr-old	older
2006										
Concentration	1.82	1.75	1720	1390	6000	4700	2500	4300	916	1688
SE	0.04	0.04	60	60	100	100	100	100	95	95
2008										
Concentration	1.55	1.46	1830	1580	6300	5400	3900	4600	3033	4045
SE	0.04	0.04	60	60	200	200	200	200	203	203
2009										
Concentration	1.62	1.64	1440	1470	5200	5200	4000	3900	2622	2617
SE	0.04	0.04	50	50	200	200	200	200	152	152
2010										
Concentration	1.29	1.28	1470	1300	5100	4900	4300	5200	2667	3381
SE	0.04	0.04	90	90	200	200	300	300	215	215
2011										
Concentration	1.32	1.32	1280	1130	4400	4200	4800	5100	2725	3047
SE	0.04	0.04	60	60	100	100	100	100	175	175
p-value (year*age)	0.0002		< 0.0001		< 0.0001		< 0.0001		< 0.0001	