Canadian Journal of Earth Sciences
Revue canadienne des sciences de la Terre

Latest Cretaceous eggshell assemblage from the Willow Creek Formation (upper Maastrichtian-lower Paleocene) of Alberta, Canada reveals higher dinosaur diversity than represented by skeletal remains

Journal:	<i>Canadian Journal of Earth Sciences</i>
Manuscript ID	cjes-2016-0080.R1
Manuscript Type:	Article
Date Submitted by the Author:	01-Aug-2016
Complete List of Authors:	Zelenitsky, Darla; Department of Geoscience, Therrien, François; Royal Tyrrell Museum of Palaeontology, Tanaka, Kohei; University of Calgary, Geoscience Currie, Philip; University of Alberta, Biological Sciences DeBuhr, Christopher; University of Calgary, Geoscience
Keyword:	dinosaurs, diversity, eggshells, Maastrichtian, Willow Creek Formation

SCHOLARONE™
Manuscripts

**Latest Cretaceous eggshell assemblage from the Willow Creek Formation
(upper Maastrichtian-lower Paleocene) of Alberta, Canada reveals higher
dinosaur diversity than represented by skeletal remains**

**Darla K. Zelenitsky^{1*}, François Therrien², Kohei Tanaka¹, Philip J. Currie³, and
Christopher L. DeBuhr¹**

¹ Department of Geoscience, University of Calgary, 2500 University Dr. NW, Calgary, Alberta
T2N 1N4, Canada

² Royal Tyrrell Museum of Palaeontology, Box 7500, Drumheller, Alberta T0J 0Y0, Canada

³ University of Alberta, Department of Biological Sciences, Edmonton, Alberta T6G 2E9,
Canada

E-mail: dkzeleni@ucalgary.ca (D. K. Zelenitsky)

E-mail: francois.therrien@gov.ab.ca (F. Therrien)

E-mail: ktanaka@ucalgary.ca (K. Tanaka)

E-mail: pjcurrie@ualberta.ca (P. Currie)

E-mail: cdebuhr@ucalgary.ca (C. DeBuhr)

*Corresponding author: Darla K. Zelenitsky

Department of Geoscience, University of Calgary, 2500 University Dr. NW, Calgary, Alberta
T2N 1N4 Canada

Tel: +1 (403) 210-6082

D.K. Zelenitsky et al. Latest Cretaceous eggshell assemblage from the Willow Creek Formation (upper Maastrichtian-early Paleocene) of Alberta, Canada reveals higher dinosaur diversity than represented by skeletal remains

Abstract

The Willow Creek Formation (upper Maastrichtian-lower Paleocene) of southwestern Alberta is a poorly fossiliferous formation that preserves a low end-Cretaceous dinosaur diversity compared to most correlative terrestrial deposits in the North American Western Interior. Although only three dinosaur taxa are known from skeletal remains (*Tyrannosaurus rex*, Hadrosauridae indet., and Leptoceratopsidae indet.), study of hundreds of dinosaur eggshells recovered from several sites in the formation reveals the presence of a more diverse dinosaur assemblage. Morphological and histological analyses of the eggshells indicate the presence of at least seven dinosaur ootaxa (*Continuoolithus*, *Montanoolithus*, *Porituberoolithus*, *Prismatoolithus* spp., *Spheroolithus* spp.). These ootaxa are referable to at least two ornithopod and five small theropod species, likely including dromaeosaurids, oviraptorosaurs, and troodontids. When considering the taxonomic affinity of eggshells and skeletal remains, the present study triples the known dinosaur diversity of the Willow Creek Formation, increasing the number of dinosaurs from three to at least nine species. Probable ornithopod eggshells comprise most of the eggshells preserved, although small theropods were likely an important component of the Willow Creek ecosystem as most ootaxa can be ascribed to these dinosaurs. Although fossil bones are rarely found in the Willow Creek Formation, fossil eggshells are common compared to most other dinosaur-bearing formations in Alberta. The caliche-bearing deposits,

indicative of arid to semi-arid conditions, typical of the formation were likely conducive to the preservation of calcareous eggshells.

Key words: dinosaurs, diversity, eggshells, Maastrichtian, Willow Creek Formation.

Draft

Introduction

Discoveries of dinosaur remains in Campanian and Maastrichtian formations of Alberta over the past century have provided evidence for the presence of a great diversity of dinosaurs living in North America near the end of the Cretaceous. Despite the abundance of dinosaur skeletal remains recovered from these formations, egg remains are rare. To date, eggs or eggshells have been described only from the late Campanian Dinosaur Park and Oldman formations in Alberta. In Dinosaur Provincial Park, the fossiliferous upper Oldman Formation and lower Dinosaur Park Formation have produced a small number ($n \sim 150$) of eggshell fragments, recovered primarily from vertebrate microfossil sites (Zelenitsky and Sloboda 2005). In contrast, at the Devil's Coulee locality (200 km south of Dinosaur Provincial Park), the upper Oldman Formation has yielded a considerable number of dinosaur embryos and eggs, as well as countless eggshell fragments (Horner and Currie 1994; Zelenitsky et al. 1996). The stratigraphic position of these localities restricts all known fossil egg remains in Alberta to a narrow temporal interval, ranging between 76.7 – 75 Ma (Horner and Currie 1994; Eberth pers. comm. 2016). Although stratigraphically younger deposits (e.g. Horseshoe Canyon and Scollard formations) leading up to the Cretaceous-Paleogene (K-Pg) boundary are well known for preserving dinosaur bones in Alberta, eggshell has yet to be described from this stratigraphic interval. Similarly elsewhere in North America, egg remains are relatively abundant in Campanian age strata of the Western Interior but are poorly known from Maastrichtian deposits (Jensen 1966; Carpenter and Alf 1994; Bray 1999; Jackson and Varricchio 2016).

Here we report the discovery of hundreds of dinosaur eggshell fragments from several uppermost Maastrichtian sites in the Willow Creek Formation (upper Maastrichtian-lower

Paleocene) of southwestern Alberta, a unit characterized by apparent low dinosaur diversity and few skeletal remains. These eggshell specimens reveal the presence of a considerably greater diversity of dinosaurs in the formation than previously recorded (Brown et al. 2015), particularly with respect to small-bodied theropods. They also significantly expand our understanding of the diversity and distribution of latest Cretaceous egg taxa in the Western Interior of North America.

Geologic setting and fossil localities

The Willow Creek Formation is an eastward-thinning (320–1350 m thick) terrestrial clastic wedge that formed in response to Laramide orogenic pulses during the late Maastrichtian and early Paleocene (Tozer 1956; Mack and Jerzykiewicz 1989; Jerzykiewicz 1985, 1997).

Regionally, it correlates with the Scollard and Coalspur formations of Alberta, the Frenchman and Ravenscrag formations of Saskatchewan, and the Hell Creek, Lance and lower Fort Union formations of Montana, North and South Dakota, and Wyoming (Hamblin 2010). Exposed in the southern foothills of Alberta, the Willow Creek Formation consists primarily of interbedded mudstone and fine-grained, laterally extensive sandstones deposited in an alluvial fan-to-fluvial setting along the eastern margin of the Rocky Mountains. Subdivided into two informal units, with the K-Pg boundary occurring near their contact, the lower (Cretaceous) part of the Willow Creek Formation is characterized by alternating red, green and purple mudstones containing pedogenic caliche nodules and hardpans, whereas the upper (Paleocene) part is dominated by dark grey mudstones and thick sandstones. The differing units record a transition from a semi-arid climate in the lower unit to more humid conditions in upper unit (Jerzykiewicz and Sweet 1988; Jerzykiewicz 1997).

The fossil record of the lower Willow Creek Formation is relatively sparse (Braman and Sweet 1990; Jerzykiewicz 1997; Hamblin 2010). Known dinosaur diversity is low in comparison to contemporaneous end-Cretaceous formations (e.g. Frenchman, Hell Creek, Lance and Scollard formations) from the Western Interior, with only three dinosaur species known, including *Tyrannosaurus rex* and indeterminate species of Hadrosauridae and Leptoceratopsidae (Brown et al. 2015). The low diversity of skeletal remains is partially a consequence of the low number of skeletons represented, but may also be attributable to preservational biases related to the depositional environments (Jerzykiewicz 1997), and perhaps to less intensive field study of the formation.

Despite the paucity of skeletal remains in the Willow Creek Formation, hundreds of fossil eggshell fragments, studied herein, have been recovered at nine individual sites from three general areas in southwestern Alberta (Fig. 1). The eggshells are preserved in overbank deposits, primarily light gray mudstones that contain small carbonate nodules indicative of paleosols, in contrast to skeletal remains that are usually preserved in channel sandstones (Abler 1984; Jerzykiewicz 1997; FT pers. obs.).

Materials and Methods

Approximately 600 eggshell fragments from the Willow Creek Formation were classified into various morphotypes based on macro- and microscopic features observed using a binocular microscope. Several shell fragments of each type were selected for examination with petrographic and scanning electron microscopes (SEM). Individual eggshell fragments were cleaned in an ultrasonic bath prior to examination with an FEI Quanta FEG 250 field emission SEM. Inner and outer surfaces, as well as freshly fractured radial surfaces were examined. Radial

thin sections of each morphotype were produced for examination and photography of their microstructure using a Leica DM 2500P polarizing microscope. Thicknesses of the various eggshell morphotypes, both including and excluding ornamentation as applicable, were measured with calipers.

The eggshells were classified based on their morphological characteristics, according to a parataxonomic scheme for fossil eggs (Zhao 1975; Hirsch 1994). Each eggshell type was assigned to an oogenus and, if possible, an oospecies. Descriptions of the eggshells herein focuses on new features of the ootaxa and comparisons with similar ootaxa found in other formations. Precise locality information (GPS coordinates) for individual eggshell sites is available through collections at the Royal Tyrrell Museum of Palaeontology, Drumheller, Alberta.

Abbreviations

TMP, Royal Tyrrell Museum of Palaeontology, Drumheller, Alberta, Canada; ZEC, Zelenitsky Egg Catalogue, University of Calgary, Calgary, Alberta.

Systematic paleontology

Oofamily Indet.

Oogenus *Continuoolithus* Zelenitsky et al., 1996

Continuoolithus cf. *canadensis* Zelenitsky et al., 1996

LOCALITY: Road cuts in the Whiskey Gap area, Municipal District of Cardston, southwestern Alberta.

MATERIAL: TMP 1987.68.4 and TMP 2010.74.1.

DESCRIPTION AND COMPARISON: *Continuoolithus* is rare in the Willow Creek Formation, with only three fragments referable to this ootaxon. The eggshell ranges from 1.05–1.35 mm, including ornamentation, a slightly thicker range than *C. canadensis* (0.94–1.28 mm) from the Oldman Formation (Zelenitsky et al. 1996). The long axis of the nodes and ridges on the outer surface are oriented in the same general direction, as is typical of elongatoolithid eggs (Zhao 1975). The thickness ratio between the continuous and mammillary layers is 7:1, although it might be slightly lower because the basal part of the mammillae is eroded. The thickness ratio reported for *C. canadensis* ranges from 4:1 (Zelenitsky et al. 1996) to 11:1 (Hirsch and Quinn 1990).

TAXONOMIC AFFINITY: The ootaxon *Continuoolithus* has never been found associated with identifiable embryos (Horner, 1997; Zelenitsky 2000), although the eggshells have two structural layers typical of non-avian theropods. *Continuoolithus* has eggshell characteristics that are similar to those of elongatoolithid eggs (Hirsch and Quinn 1990; Zelenitsky 2000), which have been ascribed to oviraptorosaurs, based on associated embryonic and adult skeletal remains from China and Mongolia (Norell et al. 1994, 1995; Dong and Currie 1995; Clark et al. 1999; Sato et al. 2005; Cheng et al. 2008; Weishampel et al. 2009; Fanti et al. 2012; Wang et al. 2016).

Oogenus *Porituberoolithus* Zelenitsky et al., 1996

Porituberoolithus warnerensis Zelenitsky et al., 1996

(Fig. 2A and 2B)

LOCALITY: Road cuts in the Whiskey Gap area, Municipal District of Cardston, southwestern Alberta; exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: TMP 1987.68.5, TMP 2015.56.27, TMP 2010.74.29, and ZEC-338.

DESCRIPTION AND COMPARISON: Forty-two fragments collected from four sites are referable to *Porituberoolithus*. The eggshells from the various sites are of comparable thickness, ranging between 0.45–0.78 mm without ornamentation, and 0.72–1.12 mm including ornamentation. This thickness range encompasses that of *P. warnerensis* from the Oldman Formation (Zelenitsky et al. 1996), and is slightly thicker than *Porituberoolithus* sp. from the Fruitland Formation (Tanaka et al. 2011). The ornamentation is well developed as it makes up approximately 25–40% of the entire shell thickness. The outer surfaces of most fragments are covered with isolated nodes that are circular, oval, or teardrop-shaped in plan view, and occasionally two nodes join to form short ridges. Pore openings are located at the peaks of the nodes (Fig. 2A). Occasionally, the nodes are slightly flattened and the interior of the node is eroded, which may be artifacts of weathering. The pore canals are straight and tubular. The thickness ratio between the continuous and mammillary layers (excluding ornamentation) is between 1:1 and 2:1 (Fig. 2B).

TAXONOMIC AFFINITY: The oogenus *Porituberoolithus* has yet to be found associated with skeletal remains, although the two-layered microstructure identifies it as non-avian theropod.

Oofamily Prismatoolithidae Hirsch, 1994

Oogenus *Prismatoolithus* Zhao and Li, 1993

Prismatoolithus cf. *levis* Zelenitsky and Hills, 1996

LOCALITY: Exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: ZEC-338.

DESCRIPTION AND COMPARISON: Only one eggshell fragment of this taxon is known from the Willow Creek Formation. The eggshell fragment is thick (1.56 mm), slightly more than the range of thickness reported for *Primatooolithus levis* (0.70–1.28 mm; Zelenitsky and Hills 1996; Varricchio et al. 2002). The outer surface is smooth, but the pore pattern is not visible. Slender shell units are visible in radial view, and the prismatic layer to mammillary layer thickness ratio is about 5:1, similar to *P. levis* (~6:1) (Varricchio et al. 2002).

TAXONOMIC AFFINITY: Eggs of the ootaxon *Prismatoolithus levis* have been ascribed to *Troodon formosus*, based on associated skeletal remains from the Two Medicine Formation of Montana (Horner and Weishampel 1996; Varricchio et al. 1997; Varricchio et al. 2002), an ootaxon that has also been described from Devil's Coulee (Zelenitsky and Hills 1996).

Oofamily Prismatoolithidae Hirsch, 1994

Oogenus *Prismatoolithus* Zhao and Li, 1993

Prismatoolithus sp.

(Fig. 2C and 2D)

LOCALITY: Road cuts in the Whiskey Gap area, Municipal District of Cardston, southwestern Alberta; exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: TMP 1987.68.3, TMP 2010.74.30, and ZEC-427.

DESCRIPTION AND COMPARISON: Twenty-five eggshell fragments collected from three sites are referable to *Prismatoolithus* sp. The eggshell fragments range from 0.31–0.63 mm in thickness and are significantly thinner than the range of thicknesses reported for *Primatooolithus levis* (0.70–1.28 mm; Zelenitsky and Hills, 1996; Varricchio et al., 2002). The outer surface is

smooth, displays relatively large, circular pore openings (Fig. 2C), and lacks the double pores in depressions diagnostic of *Prismatoolithus levis* (Zelenitsky and Hills 1996). Slender shell units are visible in radial view and the thickness ratio between prismatic and mammillary layers is 5:1 (Fig. 2D), similar to *P. levis* (~6:1) (Varricchio et al. 2002).

TAXONOMIC AFFINITY: These eggshells are about half the thickness of *Prismatoolithus levis* and lack the double pore openings on the outer surface. Considering these differences, these eggshells likely represent a different troodontid species, or perhaps another non-avian maniraptoran.

Oofamily Montanoolithidae Zelenitsky and Therrien, 2008

Oogenus *Montanoolithus* Zelenitsky and Therrien, 2008

Montanoolithus cf. *strongorum* Zelenitsky and Therrien, 2008

(Fig. 3A and 3B)

LOCALITY: Exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: ZEC-338-2.

DESCRIPTION AND COMPARISON: One eggshell fragment referable to *Montanoolithus* is known from the Willow Creek assemblage. The eggshell measures 0.58 mm in thickness, and is slightly thinner than *Montanoolithus strongorum* from the upper Oldman and upper Two Medicine formations (0.70-0.85 mm; Zelenitsky and Therrien 2008).

The outer surface is coated with a thin layer of milky translucent calcite, although the underlying shell surface appears to have ridges. The thickness ratio between the continuous and

mammillary layers is approximately 1.5:1, similar to that described for *M. strongorum* (2:1; Zelenitsky and Therrien 2008). The mammillae are closely spaced on the inner surface.

TAXONOMIC AFFINITY: *Montanoolithus* is comparable to eggshells associated with an adult skeleton of the dromaeosaurid *Deinonychus* from the Lower Cretaceous Cloverly Formation of Montana (Grellet-Tinner and Makovicky 2006). A cladistic analysis of a *Montanoolithus* clutch revealed this ootaxon may belong to either a dromaeosaurid or an oviraptorosaur, the former of which is consistent with the *Deinonychus* association (Zelenitsky and Therrien 2008).

Oofamily Spheroolithidae Zhao, 1979

Oogenus *Spheroolithus* Zhao, 1979

Spheroolithus cf. *albertensis* Zelenitsky and Hills, 1997

LOCALITY: Road cuts in the Whiskey Gap area, Municipal District of Cardston, southwestern Alberta; exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: TMP 1987.68.2, TMP 94.146.3, and ZEC-338.

DESCRIPTION AND COMPARISON: Fourteen fragments found at three sites are referable to *Spheroolithus* cf. *albertensis*. The eggshell fragments range from 0.53–1.20 mm in thickness, which is slightly thinner than fragments of *Spheroolithus albertensis* from the Oldman Formation (0.96–1.46 mm; Zelenitsky and Hills 1997). The reticulate nature of the ornamentation and the eggshell microstructure observed in the Willow Creek specimens is comparable to that reported for *S. albertensis* (Zelenitsky and Hills 1997).

Spheroolithus cf. *choteauensis* Jackson and Varricchio, 2008

(Fig. 3A and 3B)

LOCALITY: Exposures at the junction of the Oldman River and Callum Creek, Municipal District of Pincher Creek, southwestern Alberta; road cuts in the Whiskey Gap area, Municipal District of Cardston, southwestern Alberta; exposures in the Todd Creek area, Municipal District of Pincher Creek, southwestern Alberta.

MATERIAL: TMP 1987.68.1, TMP 94.146.1, TMP 95.96.1, TMP 2004.95.1, TMP 2010.74.2, TMP 2015.56.8, ZEC-338, and ZEC-427.

DESCRIPTION AND COMPARISON: Eggshell fragments referable to *Spheroolithus* cf. *choteauensis* are the most abundant ($n \sim 415$), occurring at all sites. Eggshell fragments range from 0.40–0.93 mm in thickness, the lower limit of which is thinner than the 0.66 mm reported by Jackson and Varricchio (2010), which may be due to extreme weathering of some fragments. The outer surfaces of the eggshells are usually coated with thin layers of milky translucent calcite, which obscures most surface features. An anastomosing pattern of ridges is apparent on the outer surface of some fragments (Fig. 3A); however, most are slightly undulatory to smooth, although it is uncertain if these surfaces are weathered. Pore openings are occasionally visible on the external surface of fragments. The shell units are parallel sided or flare slightly toward the outer surface. Spherulitic structure as described by Jackson and Varricchio (2010) is often prominent in the inner 1/3 of the shell units, but also can be seen to extend into the outer layer of horizontal accretion lines (Fig. 3B). The individual mammillae are broad, with large interstices on the inner surface. Basal plate groups or eisospherites are sometimes preserved on the mammillae.

TAXONOMIC AFFINITY: The oogenus *Spheroolithus* has been assigned to hadrosaurs based on its co-occurrence with embryonic remains of the saurolophines *Maiasaura* (Horner and Makela 1982; Hirsch and Quinn 1990; Zelenitsky 2000) and *Saurolophus* (Dewaele et al. 2015). However, it is probable that spheroolithid eggs were produced by a more inclusive clade, such as Ornithopoda, because *Spheroolithus* eggshells are known from the Upper Jurassic Morrison Formation (Zelenitsky et al. 2000), which represents a time period that precedes the appearance of hadrosaurs.

Discussion

Eggshells from the Willow Creek Formation provide insight into the diversity as well as the geographic and stratigraphic distribution of dinosaur egg remains during the latest Cretaceous in the North American Western Interior. In this study, five oogenera (*Continuoolithus*, *Montanoolithus*, *Porituberoolithus*, *Prismatoolithus*, *Spheroolithus*) are recognized in the Willow Creek Formation, ootaxa that are also known from other Maastrichtian and Campanian formations of the Western Interior (Table 1). Interestingly, the oogenera present in the Willow Creek Formation are more similar to Campanian formations of Alberta (i.e., Dinosaur Park and Oldman formations) than to those from contemporaneous but more southerly formations (i.e., Hell Creek and North Horn formations). Their occurrence in the Willow Creek Formation significantly extends the known stratigraphic range of these oogenera from late Campanian through the uppermost Maastrichtian, representing a temporal range of approximately 10 Ma. Given this lengthy time span, each oogenus likely represents more than a single genus/species of dinosaur within a given clade.

The Willow Creek eggshell assemblage, consisting of seven oospecies, can be ascribed to at least two ornithopod and five theropod species. The thickness of the theropod eggshells indicate the animals were of relatively small (<100 kg) body size (Tanaka et al. 2016), and their microstructures reveal they likely belonged to dromaeosaurids, oviraptorosaurs, and troodontids. Although small theropods are currently unreported from the Willow Creek Formation, their presence is not unexpected because these dinosaurs are known from the contemporaneous Frenchman, Hell Creek, Lance, and Scollard formations (Weishampel et al. 2004). Taxonomic overlap between the described ootaxa and known skeletal remains (*T. rex*, hadrosaur, and leptoceratopsid) from the Willow Creek Formation is limited to a single taxon (Ornithopoda). As such, this study increases known dinosaur diversity of the formation from three to at least nine species, tripling the previously known diversity recorded from skeletal remains alone.

The relative abundance of the various Willow Creek ootaxa indicates that ornithopod eggshell is much more common than theropod eggshell in the formation, although theropod eggshells are more diverse. The Willow Creek eggshell assemblage consists primarily of *Spherooolithus* (86%), followed by *Porituberooolithus* (8%), *Prismatoolithus* (5%), *Continuoolithus* (<1%), and *Montanoolithus* (<1%), revealing that most (86%) of the eggshell collected probably belongs to ornithopods. This trend is the opposite of that calculated at other individual eggshell sites where the percentage of theropod eggshell ranges from 87% to 100%, including sites from the Dinosaur Park Formation of Alberta (Zelenitsky and Sloboda 2005), the Fruitland Formation of New Mexico (Tanaka et al. 2011), and the 'lower formation' of the Sasayama Group of Japan (Tanaka et al. 2016). Although ornithopod eggshell is more prevalent as expected, it is evident that small theropods were diverse and thus formed a significant part of the Willow Creek ecosystem, even though they are still unknown from skeletal remains.

Fossil eggshells are relatively abundant in the Willow Creek Formation compared to most other dinosaur-bearing formations in Alberta. This abundance may be related to preservational biases induced by paleoenvironmental conditions, as proposed for the Devil's Coulee site in the upper Oldman Formation (Currie 1988). The arid to semi-arid climate indicated by the presence of pedogenic caliche nodules in both formations (Jerzykiewicz 1997; Currie 1988), would have promoted the preservation of calcium carbonate fossils, potentially yielding a higher abundance of dinosaur egg remains. In contrast, the relatively more humid conditions of the Dinosaur Park Formation (Eberth 2005), which preserves abundant skeletal material including hatchling remains but few eggshells (Tanke and Brett-Surman 2001), produced a taphonomic bias against eggshell preservation (Carpenter 1982; Currie 1988). The existence of paleoenvironmental conditions conducive to the preservation of eggshells, combined with a known abundance of eggshell fragments in the Willow Creek Formation, indicates that future field work will lead to additional discoveries of dinosaur nesting remains.

Acknowledgements

The authors are grateful to Mr. Kevin Aulenback, who took part in the initial discovery of the sites in the Willow Creek Formation in the Whiskey Gap area in 1987 with PJC. The assistance of Mr. Vittorio Ornaghi, who helped DKZ sort and measure eggshell fossils, is also greatly appreciated. All SEM work was done at the Instrumentation Facility for Analytical Electron Microscopy at the University of Calgary. This research was supported by an NSERC Discovery Grant (to DKZ) and by funding provided by the Royal Tyrrell Museum of Palaeontology.

References

- Alber, W.L. 1984. A three-dimensional map of paleontological quarry. *Contributions to Geology, University of Wyoming* 23(1): 9-14.
- Braman, D.R., and Sweet, A.R. 1990. Overview of Campanian to Paleocene stratigraphy, southern Alberta Foothills. In *Field guide to Uppermost Cretaceous Strata in Southern Saskatchewan and Alberta*. Edited by D.R. Braman and A.R. Sweet. Canadian Society of Petroleum Geologists Annual Convention Fieldtrip Guide. pp. 66-70.
- Bray, E.S. 1999. Eggs and eggshell from the Upper Cretaceous North Horn Formation, central Utah. In *Vertebrate Paleontology in Utah*. Edited by D.D. Gillette. Geological Survey Miscellaneous Publication 99-1, Utah. pp. 361-376.
- Brown, C. M., D. C. Evans, M. J. Ryan, and A. P. Russell. 2013. New data on the diversity and abundance of small-bodied ornithopods (Dinosauria: Ornithischia) from the Belly River Group (Campanian) of Alberta. *Journal of Vertebrate Paleontology* 33(3):1–26.
doi:10.1080/02724634.2013.746229.
- Brown, C.M., Ryan, M.J., and Evans, D.C. 2015. A census of Canadian dinosaurs: more than a century of discovery. In *All Animals are Interesting*. Edited by O.R.P. Bininda-Emonds, G.L. Powell, H.A. Jamniczky, A.M. Bauer, and J. Theodor. BIS-Verlag der Carl von Ossietzky Universität Oldenburg. pp. 151-209.
- Carpenter, K. 1982. Baby dinosaurs from the Late Cretaceous Lance and Hell Creek formations and a description of a new species of theropod. *Contributions to Geology, University of Wyoming* 20(2): 123-134.
- Carpenter, K., and Alf, K. 1994. Global distribution of dinosaur eggs, nests, and babies. In *Dinosaur eggs and babies*. Edited by K. Carpenter, K.F. Hirsch, and J.R. Horner. Cambridge University Press, Cambridge, United Kingdom. pp. 15-30.

- Cheng, Y.N., Jui, Q., Wu, X.C., and Shou, H.Y. 2008. Oviraptorosaurian eggs (Dinosauria) with embryonic skeletons discovered for the first time in China. *Acta Geologica Sinica* 82(6): 1089-1094. doi: 10.1111/j.1755-6724.2008.tb00708.x.
- Clark, J.M., Norell, M.A., and Chiappe, L.M. 1999. An oviraptorid skeleton from the Late Cretaceous of Ukhaa Tolgod, Mongolia, preserved in an avianlike brooding position over an oviraptorid nest. *American Museum Novitates* 3265: 1-35.
- Currie, P.J. 1988. The discovery of dinosaur eggs at Devil's Coulee. *Alberta* 1: 3-10.
- Dewaele, L., Tsogtbaatar, K., Barsbold, R., Garcia, G., Stein, K., Escuillie, F., and Godefroit, P. 2015. Perinatal Specimens of *Saurolophus angustirostris* (Dinosauria: Hadrosauridae), from the Upper Cretaceous of Mongolia. *Plos One* 10(10). doi: ARTN e013880610.1371/journal.pone.0138806.
- Dong, Z.M., and Currie, P.J. 1996. On the discovery of an oviraptorid skeleton on a nest of eggs at Bayan Mandahu, Inner Mongolia, People's Republic of China. *Canadian Journal of Earth Sciences* 33(4): 631-636. doi: 10.1139/e96-046.
- Eberth, D.A. 2005. The Geology. In *Dinosaur Provincial Park: a spectacular ancient ecosystem revealed*. Edited by P.J. Currie and E.B. Koppelhus. Indiana University Press, Bloomington. pp. 54-82.
- Fanti, F., Currie, P.J., and Badamgarav, D. 2012. New specimens of *Nemegtomaia* from the Baruungoyot and Nemegt formations (Late Cretaceous) of Mongolia. *Plos One* 7(2): 10.1371/journal.pone.0031330.
- Grellet-Tinner, G., and Makovicky, P. 2006. A possible egg of the dromaeosaur *Deinonychus antirrhopus*: phylogenetic and biological implications. *Canadian Journal of Earth Sciences* 43(6): 705-719. doi: 10.1139/E06-033.

- Hamblin, A.P. 2010. Scollard/Willow Creek/Coalspur formations: summary of literature and concepts. Geological Survey of Canada, Open File 6555: 1-22.
- Hirsch, K.F. 1994. Upper Jurassic eggshells from the Western Interior of North America. In Dinosaur Eggs and Babies. Edited by K. Carpenter, K.F. Hirsch, and J.R. Horner. Cambridge University Press, Cambridge, United Kingdom. pp. 137-150.
- Hirsch, K.F., and Quinn, B. 1990. Eggs and eggshell fragments from the Upper Cretaceous Two Medicine Formation of Montana. *Journal of Vertebrate Paleontology* 10(4): 491-511. doi: 10.1080/02724634.1990.10011832.
- Horner, J.R. 1997. Rare preservation of an incompletely ossified fossil embryo. *Journal of Vertebrate Paleontology* 17(2): 431-434.
- Horner, J.R., and Currie, P.J. 1994. Embryonic and neonatal morphology and ontogeny of a new species of *Hypacrosaurus* (Ornithischia, Lambeosauridae) from Montana and Alberta. In Dinosaur eggs and babies. Edited by K. Carpenter, K.F. Hirsch, and J.R. Horner. Cambridge University Press, Cambridge, United Kingdom. pp. 312-336.
- Horner, J.R., and Makela, R. 1979. Nest of juveniles provides evidence of family structure among dinosaurs. *Nature* 282(5736): 296-298. doi: 10.1038/282296a0.
- Horner, J.R., and Weishampel, D.B. 1996. A comparative embryological study of two ornithischian dinosaurs (correction). *Nature* 383: 103.
- Jackson, F.D., and Varricchio, D.J. 2016. Fossil egg and eggshells from the Upper Cretaceous Hell Creek Formation, Montana. *Journal of Vertebrate Paleontology* e1185432 (15 pages).
- Jensen, J.A. 1966. Dinosaur eggs from the Upper Cretaceous North Horn Formation of Central Utah. *Brigham Young University Geology Studies* 13: 55-67.

- Jerzykiewicz, T. 1985. Stratigraphy of the Saunders Group in the central Alberta Foothills - a progress report. Geological Survey of Canada, Paper 85-1B: 247-258.
- Jerzykiewicz, T. 1997. Stratigraphic framework of the uppermost Cretaceous and Paleocene strata of the Alberta basin. Geological Survey of Canada, Bulletin 510: 1-121.
- Jerzykiewicz, T., and Sweet, A.R. 1988. Sedimentological and palynological evidence of regional climatic changes in the Campanian to Paleocene sediments of the Rocky Mountain Foothills, Canada. *Sedimentary Geology* 59: 29-76.
- Mack, G.H., and Jerzykiewicz, T. 1989. Provenance of post-Wapiabi sandstones and its implications for Campanian to Paleocene tectonic history of the southern Canadian Cordillera. *Canadian Journal of Earth Sciences* 26: 665-676.
- Norell, M.A., Clark, J.M., Chiappe, L.M., and Dashzeveg, D. 1995. A nesting dinosaur. *Nature* 378(6559): 774-776. doi: 10.1038/378774a0.
- Norell, M.A., Clark, J.M., Demberelyin, D., Rhinchen, B., Chiappe, L.M., Davidson, A.R., Mckenna, M.C., Altangerel, P., and Novacek, M.J. 1994. A theropod dinosaur embryo and the affinities of the Flaming Cliffs dinosaur eggs. *Science* 266(5186): 779-782. doi: 10.1126/science.266.5186.779.
- Sato, T., Cheng, Y.N., Wu, X.C., Zelenitsky, D.K., and Hsiao, Y.F. 2005. A pair of shelled eggs inside a female dinosaur. *Science* 308(5720): 375-375. doi: 10.1126/science.1110578.
- Tanaka, K., Zelenitsky, D.K., Saegusa, H., Ikeda, T., DeBuhr, C.L., and Therrien, F. 2016. Dinosaur eggshell assemblage from Japan reveals unknown diversity of small theropods. *Cretaceous Research* 57: 350-363. doi: 10.1016/j.cretres.2015.06.002.

- Tanaka, K., Zelenitsky, D.K., Williamson, T., Weil, A., and Therrien, F. 2011. Fossil eggshells from the Upper Cretaceous (Campanian) Fruitland Formation, New Mexico. *Historical Biology* 23(1): 41-55. doi: 10.1080/08912963.2010.499171.
- Tanke, D.H., and Brett-Surman, M.K. 2001. Evidence of hatchling- and nestling-size hadrosaurs (Reptilia: Ornithischia) from Dinosaur Provincial Park Formation: Campanian), Alberta. In *Mesozoic Vertebrate Life*. Edited by D.H. Tanke and K. Carpenter. Indiana University Press, Bloomington. pp. 206-218.
- Tozer, E.T. 1956. Uppermost Cretaceous and Paleocene non-marine molluscan faunas of western Alberta. Geological Survey of Canada, Memoir 280.
- Varricchio, D.J., Horner, J.R., and Jackson, F.D. 2002. Embryos and eggs for the Cretaceous theropod dinosaur *Troodon formosus*. *Journal of Vertebrate Paleontology* 22(3): 564-576. doi: 10.1671/0272-4634(2002)022[0564:Eaeftc]2.0.Co;2.
- Varricchio, D.J., Jackson, F., Borkowski, J.J., and Horner, J.R. 1997. Nest and egg clutches of the dinosaur *Troodon formosus* and the evolution of avian reproductive traits. *Nature* 385(6613): 247-250. doi: 10.1038/385247a0.
- Wang, S., Zhang, S., Sullivan, C., and Xu, X. 2016. Elongatoolithid eggs containing oviraptorid (Theropoda, Oviraptorosauria) embryos from the Upper Cretaceous of Southern China. *BMC Evolutionary Biology* 16(67): 1-21. doi: 10.1186/s12862-016-0633-0.
- Weishampel, D.B., Barrett, P.M., Coria, R.A., Loeuff, J., Xu, X., Xijin, Z., Sahni, A., Gomani, E.M.P., and Noto, C.R. 2004. Dinosaur Distribution. In *Dinosauria*, 2nd Ed. Edited by D.B. Weishampel, P. Dodson and H. Osmolska. University of California Press, Berkeley and Los Angeles. pp. 517-606.

- Weishampel, D.B., Fastovsky, D.E., Watabe, M., Varricchio, D., Jackson, F., Tsogtbaatar, K., and Barsbold, R. 2008. New oviraptorid embryos from Bugin-Tsav, Nemegt Formation (Upper Cretaceous), Mongolia, with insights into their habitat and growth. *Journal of Vertebrate Paleontology* 28(4): 1110-1119. doi: 10.1671/0272-4634-28.4.1110.
- Welsh, E., and Sankey, J.T. 2008. First dinosaur eggshells from Texas, USA: Aguja Formation (Late Campanian), Big Bend National Park. In *Vertebrate Microfossil Assemblages: Their Role in Paleoecology and Paleobiogeography*. Edited by J.T. Sankey and S. Baszio. Indiana University Press, Bloomington and Indianapolis. pp. 166-177.
- Zelenitsky, D.K. 2000. Dinosaur eggs from Asia and North America. *Paleontological Society of Korea Special Publication* 4: 13-26.
- Zelenitsky, D.K., Carpenter, K., and Currie, P.J. 2000. First record of elongatoolithid theropod eggshell from North America: the Asian oogenus *Macroelongatoolithus* from the Lower Cretaceous of Utah. *Journal of Vertebrate Paleontology* 20(1): 130-138. doi: 10.1671/0272-4634(2000)020[0130:Froete]2.0.Co;2.
- Zelenitsky, D.K., and Hills, L.V. 1996. An egg clutch of *Prismatoolithus levis* oosp. nov. from the Oldman Formation (Upper Cretaceous), Devil's Coulee, southern Alberta. *Canadian Journal of Earth Sciences* 33(8): 1127-1131. doi: 10.1139/E96-085.
- Zelenitsky, D.K., and Hills, L.V. 1997. Normal and pathological eggshells of *Spheroolithus albertensis*, oosp. nov., from the Oldman Formation (Judith River Group, late Campanian), southern Alberta. *Journal of Vertebrate Paleontology* 17(1): 167-171. doi: 10.1080/02724634.1997.10010960.
- Zelenitsky, D.K., Hills, L.V., and Currie, P.J. 1996. Parataxonomic classification of ornithoid eggshell fragments from the Oldman Formation (Judith River Group; Upper Cretaceous),

- southern Alberta. *Canadian Journal of Earth Sciences* 33(12): 1655-1667. doi: 10.1139/E96-126.
- Zelenitsky, D.K., and Sloboda, W. 2005. Eggshells. In *Dinosaur Provincial Park: a spectacular ancient ecosystem revealed*. Edited by P.J. Currie and E.B. Koppelhus. Indiana University Press, Bloomington. pp. 398-404.
- Zelenitsky, D.K., and Therrien, F. 2008. Unique maniraptoran egg clutch from the Upper Cretaceous Two Medicine Formation of Montana reveals theropod nesting behaviour. *Palaeontology* 51: 1253-1259. doi: 10.1111/j.1475-4983.2008.00815.x.
- Zhao, Z. 1979. Progress in the research of dinosaur eggs. In *Mesozoic and Cenozoic Red Beds of South China Selected Papers from the "Cretaceous - Tertiary Workshop, Nanxiong, Guangdong Province"*. Edited by Institute of Vertebrate Palaeontology and Palaeoanthropology and Nanjing Institute of Palaeontology. Science Press, Beijing. pp. 330-340.
- Zhao, Z., and Li, R. 1993. First record of Late Cretaceous hypsilophodontid eggs from Bayan Manduhu, Inner Mongolia. *Vertebrata Palasiatica* 31(2): 77-84.
- Zhao, Z.K. 1975. Microstructures of dinosaurian eggshells of Nanxiong, Guangdong, and the problem in egg classification. *Vertebrata Palasiatica* 13(2): 105-117.

Table 1. Dinosaur oögenera from the Willow Creek also found at other Campanian or Maastrichtian localities in the Western Interior.

Oögenus	Formation	Age	Location	References
<i>Continuoolithus</i> , <i>Montanoolithus</i> , <i>Porituberoolithus</i> , <i>Prismatoolithus</i> , <i>Spheroolithus</i>	Willow Creek	late Maastrichtian	Alberta	this study
<i>Spheroolithus</i>	Hell Creek	Maastrichtian	Montana	Jackson and Varricchio (2016)
<i>Prismatoolithus</i> , <i>Spheroolithus</i> *	North Horn	Maastrichtian	Utah	Bray (1999); *Re-assigned by Jackson and Varricchio (2010)
<i>Continuoolithus</i> , <i>Porituberoolithus</i> , <i>Prismatoolithus</i> , <i>Spheroolithus</i>	Dinosaur Park	late Campanian	Alberta	Zelenitsky and Sloboda (2005)
<i>Continuoolithus</i> , <i>Montanoolithus</i> , <i>Porituberoolithus</i> , <i>Prismatoolithus</i> , <i>Spheroolithus</i>	Oldman	late Campanian	Alberta	Zelenitsky and Hills (1996, 1997); Zelenitsky et al. (1996)
<i>Continuoolithus</i> , <i>Montanoolithus</i> , <i>Prismatoolithus</i> , <i>Spheroolithus</i>	Two Medicine	Campanian	Montana	Hirsch and Quinn (1990); Zelenitsky (2000); Varricchio et al. (2002); Jackson and Varricchio (2010)
<i>Continuoolithus</i> , <i>Porituberoolithus</i> , <i>Prismatoolithus</i>	Fruitland	late Campanian	New Mexico	Tanaka et al. (2011)
<i>Continuoolithus</i> , <i>Porituberoolithus</i>	Aguja	late Campanian	Texas	Welsh and Sankey (2008)

Figure captions

Fig. 1. Map showing three areas where dinosaur eggshells were found in the Willow Creek Formation of southwestern Alberta. Area 1 is exposures at the junction of the Oldman River and Callum Creek. Area 2 is exposures near Todd Creek. Area 3 is exposures from road cuts near Whiskey Gap.

Fig. 2. (A and B) Eggshell of *Porituberoolithus warnerensis*. TMP 1987.68.5 in, (A) outer surface view with pore opening on nodes. Scale bar = 2mm. (B) Radial SEM view with pore canal (arrow). Scale bar = 300 μm . (C and D) Eggshell of *Prismatoolithus* sp. TMP 1987.68.3 in, (C) outer surface view showing circular pores. Scale bar = 1 mm. (D) Radial thin section showing prismatic microstructure. Scale bar = 500 μm .

Fig. 3. Eggshell of *Spheroolithus* cf. *choteauensis*. TMP 2015.56.8 in, (A) outer surface view showing low-relief ornamentation. Scale bar = 1 mm. TMP 1987.68.1 in, (B) Radial thin section showing two-layered microstructure. Scale bar = 500 μm .

Fig. 1. Map showing three areas where dinosaur eggshells were found in the Willow Creek Formation of southwestern Alberta. Area 1 is exposures at the junction of the Oldman River and Callum Creek. Area 2 is exposures near Todd Creek. Area 3 is exposures from road cuts near Whiskey Gap.

181x109mm (300 x 300 DPI)

Fig. 2. (A and B) Eggshell of *Porituberoolithus warnerensis*. TMP 1987.68.5 in, (A) outer surface view with pore opening on nodes. Scale bar = 2mm. (B) Radial SEM view with pore canal (arrow). Scale bar = 300 μ m. (C and D) Eggshell of *Prismaoolithus* sp. TMP 1987.68.3 in, (C) outer surface view showing circular pores. Scale bar = 1 mm. (D) Radial thin section showing prismatic microstructure. Scale bar = 500 μ m.

182x168mm (300 x 300 DPI)

Eggshell of *Spheroolithus* cf. *choteauensis*. TMP 2015.56.8 in, (A) outer surface view showing low-relief ornamentation. Scale bar = 1 mm. TMP 1987.68.1 in, (B) Radial thin section showing two-layered microstructure. Scale bar = 500 μ m.

182x69mm (300 x 300 DPI)

Draft