Canadian Journal of Chemistry
Revue canadienne de chimie

**A practical synthesis and X-ray crystal structure of
tribenzo[*a,d,j*]corannulene**

Journal:	<i>Canadian Journal of Chemistry</i>
Manuscript ID	cjc-2016-0516.R1
Manuscript Type:	Article
Date Submitted by the Author:	15-Oct-2016
Complete List of Authors:	Smith, Natalie; Boston College, Department of Chemistry Scott, Lawrence; Boston College
Keyword:	benzannulation, geodesic polyarene, polycyclic aromatic hydrocarbon

A practical synthesis and X-ray crystal structure of tribenzo[*a,d,j*]corannulene

Natalie J. Smith and Lawrence T. Scott

Abstract: Tribenzo[*a,d,j*]corannulene (**5**) has been synthesized in four linear steps from commercially available starting materials on a half-gram scale in 18% overall yield. An X-ray crystal structure of this triply benzannulated geodesic polyarene confirms the prediction that successive benzannulation of corannulene has a gradual flattening effect on the bowl-shaped core.

Key words: benzannulation, geodesic polyarene, polycyclic aromatic hydrocarbon

Introduction

The pioneering research of Reginald Mitchell on benzannulations of aromatic π -systems¹ inspired us to extend those studies to corannulene (**1**), the benzannulated homologues of which constitute a unique family of curved polycyclic aromatic hydrocarbons (PAHs, **2-8**, Figure 1).

N. J. Smith and L. T. Scott. Merkert Chemistry Center, Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860 USA.

Corresponding author: Lawrence T. Scott (email: Lawrence.scott@bc.edu).
<https://mc06.manuscriptcentral.com/cjc-pubs>

Figure 1. Corannulene (**1**) and all of the possible benzannulated homologues.

Thanks to the work of Siegel *et al.*, the parent corannulene (**1**) is now available in kilogram amounts.² With the lone exception of our gram-scale synthesis of dibenzo[*a,g*]corannulene (**3**),³ however, none of the benzannulated corannulenes has ever been prepared and isolated in more than milligram quantities, and no syntheses of tetrabenzocorannulene (**7**) or pentabenzocorannulene (**8**) have ever been published. Herein, we report the first practical synthesis of tribenzo[*a,d,j*]corannulene (**5**).⁴

The first milligram quantities of the title compound (**5**) were synthesized, isolated, purified, and characterized in 1997 by a talented undergraduate student in this laboratory, Brandon McMahon, as part of his senior thesis project (Figure 2).⁵ The use of flash vacuum pyrolysis (FVP) in the final step, however, limits the scale of this preparation to only tens of milligrams per run.

Figure 2. The first synthesis of tribenzo[*a,d,j*]corannulene (**5**).⁵

Several years later, Mehta *et al.* published a second synthesis of **5** that also employed FVP in the final step (Figure 3).⁶ Since then, no other syntheses of this C₃₂H₁₆ fullerene fragment have been reported.

Figure 3. The second synthesis of tribenzo[*a,d,j*]corannulene (**5**).⁶

Results and Discussion

Capitalizing on the Pd-catalyzed, solution phase, intramolecular arylation strategy introduced by our laboratory for imposing curvature on planar PAHs,³ we have successfully synthesized the title compound (**5**) on a half-gram scale, and the procedure is readily amenable to further scale-up (Figure 4).⁷ In the final step of this synthesis, intermediate **9** is stirred at 150 °C for 3 days in *N,N*-dimethylacetamide (DMAc) containing catalytic amounts of Pd(PCy₃)₂Cl₂ and several equivalents of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). Tribenzo[*a,d,j*]corannulene (**5**) was obtained from this reaction in 56% isolated yield after

recrystallization from chloroform (mp 302-304 °C). Starting from commercially available starting materials, hydrocarbon **5** is obtained by this route in four linear steps in 18% overall yield.

Figure 4. The first practical synthesis of tribenzo[*a,d,j*]corannulene (**5**), requiring no FVP.⁷

Figure 5 shows the steps used to prepare synthetic intermediate **9**.

Figure 5. Synthesis of intermediate **9** from commercially available starting materials: a) oxalyl bromide, AlBr_3 , CS_2 , $-40\text{ }^\circ\text{C}$, 42% b) $\text{Fe}_2(\text{CO})_9$, benzene, 54% c) KOH , MeOH , 82% d) 2,5-norbornadiene, reflux, 99%, or 2-nitroethanol, phthalic anhydride, $140\text{ }^\circ\text{C}$, toluene, sealed tube, 92%.

The synthesis of acephenanthrylene-4,5-dione (**11**) in one step from phenanthrene (**10**) has previously been reported only in the patent literature.⁸ In our hands, however, the described procedure failed to deliver **11** in the reported yield. Thus, we devoted considerable effort to optimizing the direct conversion of **10** to **11**. In the end, double Friedel-Crafts acylation of **10** with oxalyl bromide, using AlBr_3 as the catalyst, at $-40\text{ }^\circ\text{C}$ in carbon

disulfide resulted in a 42% isolated yield of **11**. Unfortunately, additional experimentation never improved this modest yield any further. On the positive side, the reaction is reproducible; it uses inexpensive starting materials and reagents, and it can easily be carried out on a multi-gram scale.

The 1,3-*bis*(2-chlorophenyl)-2-propanone **13** was prepared by treating 2-chlorobenzyl chloride (**12**) with diironnonacarbonyl in hot benzene.⁹ A double aldol condensation between **11** and **13**, catalyzed by potassium hydroxide in methanol, then gave the fused cyclopentadienone **14**, which could be isolated as a dark purple solid in good yield simply by filtration of the reaction mixture.¹⁰

Transformation of the cyclopentadienone ring in **14** into a benzene ring was achieved by Diels-Alder cycloaddition of a “masked acetylene,” followed by chelotropic loss of carbon monoxide.¹¹ Two different “masked acetylenes” were examined. Nitroethylene, generated *in situ* from 2-nitroethanol by the action of phthalic anhydride in toluene at 140 °C in a pressure vessel,¹² adds rapidly to **14**, and the subsequent loss of carbon monoxide and HNO₂ gives **9** in 92% yield. Alternatively, refluxing **14** in neat 2,5-norbornadiene promotes an inverse electron demand Diels-Alder reaction, which then delivers **9** almost quantitatively by loss of carbon monoxide and cyclopentadiene by a retro-Diels-Alder reaction.¹¹

X-ray quality crystals of tribenzo[*a,d,j*]corannulene (**5**) were grown by slow evaporation of a carbon disulfide solution (Figure 6). An examination of the experimental structural parameters confirms the prediction based on theoretical calculations that successive benzannulation of corannulene has a gradual flattening effect on the bowl-shaped core.¹³

Figure 6. X-ray crystal structure of tribenzo[*a,d,j*]corannulene (**5**), ORTEP ellipsoids at 50%. (a) Single molecule, top view. (b) Side view, showing stacking.

Two metrics have come into common use for describing the curvature of bowl-shaped PAHs. One is the atom-by-atom POAV analysis, popularized by R. C. Haddon,¹⁴ and the other is the overall bowl depth.¹⁵ The angle that each of the three bonds radiating from a trigonal carbon atom makes with the π -orbital axis vector (POAV) of that carbon provides a well-defined measure of the pyramidalization of a quasi- sp^2 center. The angle is reported as the deviation of the carbon framework from a 90° angle with the π -orbital of the conjugated system, 90° being characteristic of planar sp^2 centers (Figure 7). For the mono-, di-, and tribenzannulated corannulenes, POAV angles are calculated for each of the unique carbon centers in the hub pentagon of the corannulene core. Table 1 lists the calculated (B3LYP/6-31G*) and experimental (X-ray) POAV angles for each of the hub carbon atoms of corannulenes **1**, **2**, **3**, and **5**, together with the averages of these POAV angles in each compound, for comparison purposes.

Figure 7. Pi-Orbital Axis Vector angle.

The second structural feature of interest is the bowl depth of each molecule. This value is defined as the shortest distance between the average plane of the 5 hub carbon atoms and the average plane created by the 10 carbon atoms on the corannulene rim. This measurement is a good indicator of the curvature of corannulene (**1**), benzo[*a*]corannulene (**2**), and dibenzo[*a,g*]corannulene (**3**); however, it is somewhat more ambiguous for the other members of the benzocorannulene family, owing to a significant “ruffling” of the corannulene core caused by adjacent benzo units. In those compounds, the 10 rim carbons no longer form an even plane, and the measure of curvature observed by this method may be distorted. Still, useful information can be gained about general trends in bowl depth, and the ruffling effect itself can offer interesting structural insights.

Structure	Compound name	Calc. POAV angles at hub ^a	Avg. Calc. POAV angle ^a	Exp. POAV angles at hub	Avg. Exp. POAV angle	Bowl Depth Exp. (Calc.)
1	Corannulene ^b	8.1	8.1	8.6, 8.3, 8.3, 8.4, 8.3	8.4	0.87 Å (0.87 Å)
2	Benzo[<i>a</i>]corannulene ^c	7.7, 7.8, 7.7	7.8	7.5, 8.0, 8.0, 8.2, 7.9	7.9	0.83 Å (0.82 Å)
3	Dibenzo[<i>a,g</i>]- corannulene ^d	7.3, 7.3, 7.4	7.3	7.6, 7.7, 7.9, 7.8, 8.0	7.8	0.83 Å (0.77 Å)
5	Tribenzo[<i>a,d,j</i>]- corannulene ^e	6.8, 6.3, 6.3, 6.7, 6.6	6.5	6.5, 6.3, 6.0, 6.6, 6.5	6.4	0.67 Å (0.68 Å)
7	Tetrabenzo(<i>a,d,g,j</i>)- corannulene	5.8, 5.2, 5.2, 5.5, 5.2	5.4			(0.50 Å)
8	Pentabenzo[<i>a,d,g,j,m</i>]- corannulene	3.8, 3.9, 3.9, 3.3, 4.2	3.8			(0.29 Å)

Table 1. POAV angles and bowl depths of selected benzocorannulenes

(a) Calculated POAV angles (in deg) at hub positions were obtained at the B3LYP/6-31G* level of theory. (b) X-ray: Petrukhina, M. A.; Andreini, K. W.; Mack, J.; Scott, L. T. *J. Org. Chem.* **2005**, *70*, 5713. (c) X-ray: McComas, C. C. B.S. thesis, Boston College, Chestnut Hill, Massachusetts, 1996 (see supplementary material). (d) X-ray: Petrukhina, M. A.; Andreini, K. W.; Tsefrikas, V. M.; Scott, L. T. *Organometallics* **2005**, *24*, 1394. (e) X-ray: this work.

Table 1 demonstrates the good agreement between the calculated and experimentally determined POAV angles for **1**, **2**, and **5**. In contrast, the calculated values for dibenzo[*a,g*]-corannulene (**3**) are significantly lower than those observed in the crystal structure, and the experimental bowl depth for **3** is almost identical to that for benzocorannulene (**2**). Nevertheless, a general trend can still be discerned. Both the POAV angles and the bowl

depths decrease with an increasing number of benzo fusions. The addition of benzo groups to the rim of corannulene does indeed flatten the bowl, and calculations predict a continuation of this trend, with a particularly dramatic effect in the case of pentabenzocorannulene (**8**).

Conclusion

The successful synthesis of tribenzo[*a,d,j*]corannulene (**5**) on a half-gram scale, in four linear steps and 18% overall yield, represents a significant step forward in the study of benzannulated curved PAHs. The key features of the synthesis are the efficient construction of a chlorinated precursor (**9**) and the introduction of curvature in the corannulene core through a pair of palladium-catalyzed, solution-phase cyclization reactions. An X-ray crystal structure of the title compound confirms the predicted trend that sequential benzannulation of corannulene leads to progressive flattening of the bowl.

Experimental

General.

Commercially available chemicals were used as received without further purification unless noted. Dry solvents were purified using a Glass Contour solvent purification system. Silica gel chromatography was carried out using Sorbent Technology or Zeochem silica gel with a porosity of 60 Å and a particle size of 32-63 µm. NMR spectra were recorded in the Boston College Nuclear Magnetic Resonance Center on Varian instruments. NMR shifts are reported in ppm downfield from tetramethylsilane, using the solvent as the standard reference: chloroform-*d*₁ ($\delta_{\text{H}} = 7.26$ ppm, $\delta_{\text{C}} = 77.0$ ppm). High resolution mass spectra were obtained in the Boston College Mass Spectrometry Center on a DART-TOF spectrometer operating in positive ionization mode. X-ray crystallography was performed at the Boston

College X-ray Crystallography Center on a Siemens 3-circle geometry platform diffractometer with a Bruker APEX CCD area detector equipped with a copper source. Melting points are reported uncorrected.

Acephenanthrylene-4,5-dione (11).

In a flame dried, 500 mL round bottom flask, 30.6 g (115 mmol) of aluminum bromide was dissolved in 200 mL of carbon disulfide. The solution was cooled to $-40\text{ }^{\circ}\text{C}$, and a solution comprising 10.0 g (56.1 mmol) of phenanthrene and 5.30 mL (56.5 mmol) of oxalyl bromide in 80 mL of carbon disulfide was added slowly over 15 min. The reaction mixture was stirred at $-40\text{ }^{\circ}\text{C}$ for 3 h before being allowed to warm to room temperature. After being stirred for 1 h more, the mixture was poured carefully into water. It was then thoroughly extracted with dichloromethane. The organic solution was washed three times with water and once with brine, dried over magnesium sulfate, and concentrated to dryness. Column chromatography on silica gel eluted with dichloromethane gave 5.49 g (42%) of acephenanthrylene-4,5-dione (**11**) as a bright yellow solid. **mp** 242-245 $^{\circ}\text{C}$. **^1H NMR** (500 MHz, CDCl_3): δ (ppm): 8.86 (d, $J = 8.1\text{ Hz}$, 1H), 8.69 (d, $J = 8.2\text{ Hz}$, 1H), 8.35 (s, 1H), 8.17-8.13 (m, 2H), 7.95 (dd, $J = 8.2\text{ Hz}$, 7.2 Hz , 1H), 7.90 (ddd, $J = 8.3\text{ Hz}$, 7.0 Hz , 1.3 Hz , 1H), 7.79 (ddd, $J = 8.1\text{ Hz}$, 7.0 Hz , 1.2 Hz , 1H). **^{13}C NMR** (125 MHz, CDCl_3): δ 188.6, 188.2, 142.9, 132.5, 131.8, 131.6, 130.3, 129.3, 129.0, 128.8, 128.7, 128.5, 125.9, 124.8, 123.1, 122.3. **HRMS** (DART-TOF) calcd for $\text{C}_{16}\text{H}_9\text{O}_2$ ($\text{M}+\text{H}$) $^{+}$ 233.0602, found 233.0603. lit.⁸

1,3-Bis(2-chlorophenyl)propan-2-one (13).

In a 500 mL, flame dried, three-neck round bottom flask, 25 g (68.7 mmol) of diironnonacarbonyl was mixed with 200 mL of anhydrous benzene. After the addition of 8.7 mL (69 mmol) of 2-chlorobenzylchloride, the reaction mixture was heated to reflux for 24 h.

The solution was allowed to cool to room temperature and filtered through a plug of silica gel, flushing with dichloromethane. The solvent was removed under reduced pressure, and the crude material was recrystallized from hexanes to yield 5.21 g (18.6 mmol, 54%) of clean product (**11**). $^1\text{H NMR}$ (500 MHz, CDCl_3): δ 7.41-7.36 (m, 2H), 7.25-7.20 (m, 6H), 3.93 (s, 4H). lit.^{3,16}

9,11-Bis(2-chlorophenyl)-10H-cyclopenta[*e*]acephenanthrylen-10-one (**14**).

In a 125 mL Erlenmeyer flask, 1.20 g (4.31 mmol) of 1,3-bis(2-chlorophenyl)propan-2-one (**13**) and 1.00 g (4.31 mmol) of acephenanthrylene-4,5-dione (**11**) were mixed in a solution of 0.405 g (7.22 mmol) of potassium hydroxide in 36 mL of methanol. The reaction mixture was stirred at room temperature for 4 h, during which time a dark precipitate formed. The mixture was left standing overnight without stirring to allow the product to precipitate fully before it was collected by vacuum filtration and dried to yield 1.69 g (3.56 mmol, 82%) of pure product (**14**) as a dark purple solid. mp 205-212 °C (mixture of 2 atropisomers melts in two stages). $^1\text{H NMR}$ (500 MHz, CDCl_3 , for a mixture of diastereomers): δ (ppm): 8.62 (d, J = 8.0 Hz, 1H), 8.57 (d, J = 8.0 Hz, 1H), 7.96 (d, J = 7.9 Hz, 1H), 7.82 (d, J = 5.0 Hz, 1H), 7.76-7.54 (m, 8H), 7.47-7.37 (m, 4H). $^{13}\text{C NMR}$ (125 MHz, CDCl_3 , for a mixture of 2 atropisomers): δ 199.8, 199.5, 156.8, 156.6, 155.3, 155.2, 142.8, 134.2, 134.1, 133.1, 132.1, 132.0, 131.7, 131.6, 131.0, 130.4, 130.11, 130.05, 129.97, 129.6, 129.5, 128.9 (broad), 128.8, 128.5, 127.9, 127.3, 126.6 (broad), 125.0, 124.8, 124.4, 124.2, 124.1, 122.8, 120.0 (broad), 119.0. HRMS (DART-TOF) calcd for $\text{C}_{31}\text{H}_{17}\text{Cl}_2\text{O}$ ($\text{M}+\text{H}$)⁺ 475.0656, found 475.0640.

7,10-Bis(2-chlorophenyl)benzo[*b*]fluoranthene (**9**)

(a) From the reaction of **14** with 2,5-norbornadiene. In a 5 mL round bottom flask, 318 mg (0.669 mmol) of 9,11-bis(2-chlorophenyl)-10H-cyclopenta[*e*]acephenanthrylen-10-

one (**14**) was stirred in 1.0 mL of 2,5-norbornadiene at reflux for 5 days. After allowing the reaction mixture to cool to room temperature, the excess 2,5-norbornadiene was removed under reduced pressure. The product was purified by filtration through a plug of silica gel with dichloromethane. The final purified product (**9**) was collected in greater than 98% yield (316 mg, 0.668 mmol).

(b) From the reaction of 14 with nitroethylene. In a flame dried 250 mL round bottom flask with a reflux condenser, 3.97 g (8.35 mmol) of 9,11-*bis*(2-chlorophenyl)-10H-cyclopenta[*e*]acephenanthrylen-10-one (**14**) and 31.0 g (209 mmol) of phthalic anhydride were mixed in 100 mL of toluene. After the addition of 14.7 mL (209 mmol) of 2-nitroethanol, the reaction mixture was heated to 140 °C and stirred for 26 h. After allowing the reaction mixture to cool to room temperature, it was diluted with dichloromethane. The organic layer was washed three times with 10% aq sodium hydroxide and three times with water before drying over magnesium sulfate and evaporation of the solvent. The crude residue was purified by flash chromatography on silica gel with 3:1 hexanes:dichloromethane. The product collected (**9**) weighed 2.78 g (5.87 mmol, 70%). Further elution with dichloromethane resulted in the recovery of 11% of the starting material. On a small scale, in a sealed tube at 140 °C, the yield by this method is 92%; **CAUTION:** a full equivalent of carbon monoxide gas is released in this reaction, which causes pressure buildup. The sealed tube method is not advised for large scale runs of this reaction. **mp** 130-186 °C (mixture of 2 atropisomers melts in two stages). **¹H NMR** (500 MHz, CDCl₃, for a mixture of 2 atropisomers): δ 8.59 (d, *J* = 8.1 Hz, 1H), 8.41 (d, *J* = 8.2 Hz, 1H), 7.74-7.46 (m, 12H), 7.39-7.35 (m, 2H), 7.18 (s, 1H), 6.98 (d, *J* = 7.3 Hz, 1H). **¹³C NMR** (125 MHz, CDCl₃, for a mixture of 2 atropisomers): δ 139.4, 139.32, 139.29, 138.4, 136.3, 136.21,

136.20, 135.40, 135.38, 134.78, 134.76, 134.24, 134.23, 133.76, 133.75, 133.63, 133.62, 132.3, 131.4, 131.2, 130.5, 129.90, 129.85, 129.82, 129.77, 129.42, 129.40, 129.35, 129.34, 129.27, 129.26, 128.64, 128.63, 128.0, 127.25, 127.24, 127.14, 127.12, 127.10, 127.07, 127.03, 126.5, 124.3, 122.8, 121.8, 121.7. **HRMS** (DART-TOF) calcd for $C_{32}H_{19}Cl_2$ ($M+H$)⁺ 473.0864, found 473.0846.

Tribenzo[*a,d,j*]corannulene (**5**).

In a flame dried 150 mL pressure vessel containing a magnetic stir bar, 1.00 g (2.11 mmol) of 7,10-*bis*(2-chlorophenyl)benzo[*b*]fluoranthene (**9**) and 0.331 g (0.448 mmol) of $Pd(PCy_3)_2Cl_2$ were mixed in 9.0 mL of *N,N*-dimethylacetamide. After the addition of 9.5 mL (63 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene, the pressure vessel was capped and placed in a 150 °C oil bath to stir for 3 days. The reaction mixture was allowed to cool to room temperature before filtering through a plug of neutral alumina and flushing with dichloromethane. As the solution was diluted with dichloromethane from flushing the plug, some of the product precipitated out. This solid was collected by vacuum filtration. The filtrate was washed twice with 10% aq hydrochloric acid, once with water, and once with brine. The solvent was removed under reduced pressure, and the crude material was purified by recrystallization from chloroform. The total amount of clean product (**5**) collected weighed 476 mg (1.19 mmol, 56%). **mp** 302-304 °C. **¹H NMR** (500 MHz, $CDCl_3$): δ 9.36 (dd, *J* = 8.2 Hz, 1.3 Hz, 2H), 8.81 (dd, *J* = 7.8 Hz, 1.7 Hz, 2H), 8.68 (AA' of AA'XX', 2H), 8.38 (d, *J* = 8.6 Hz, 2H), 8.36 (d, *J* = 8.6 Hz, 2H), 7.85-7.78 (m, 4H), 7.76 (XX' of AA'XX', 2H). **¹³C NMR** (125 MHz, $C_2D_2Cl_4$): δ 134.7, 134.5, 133.8, 133.6, 132.8, 132.7, 128.2, 127.9, 127.6, 127.3, 127.2, 126.6, 125.5, 125.1, 124.6, 124.3, 123.9. **HRMS** (DART-TOF) calcd for $C_{32}H_{17}$ ($M+H$)⁺ 401.1330, found 401.1325. X-ray quality crystals were grown by slow

evaporation of carbon disulfide. See the supplementary material or the Cambridge Crystal Data Base for X-ray crystal data (CCDC 856219).

Supplementary material

NMR spectra of compounds **5**, **9**, **11**, **13**, and **14**. X-ray crystal structure CIF files for compounds **2** and **5**. Supplementary material is available with the article through the journal Web site at [0000](#).

Acknowledgements

We thank the U.S. National Science Foundation and Department of Energy for financial support of this work.

References

- ¹ (a) Mitchell, R. H.; Carruthers, R. J. *Tetrahedron Lett.* **1975**, 4331. (b) Mitchell, R. H.; Yan, J. S.-H. *Can. J. Chem.* **1977**, *55*, 3347. (c) Mitchell, R. H.; Yan, J. S.-H. *Tetrahedron Lett.* **1979**, 1289. (d) Mitchell, R. H. *Isr. J. Chem.* **1980**, *20*, 294. (e) Mitchell, R. H.; Mahadevan, R. *Tetrahedron Lett.* **1981**, *22*, 5131. (f) Mitchell, R. H.; Yan, J. S. H.; Dingle, T. W. *J. Am. Chem. Soc.* **1982**, *104*, 2551. (g) Mitchell, R. H.; Chaudhary, M.; Dingle, T. W.; Williams, R. V. *J. Am. Chem. Soc.* **1984**, *106*, 7776. (h) Mitchell, R. H.; Weerawarna, S. A. *Tetrahedron Lett.* **1986**, *27*, 453. (i) Mitchell, R. H.; Iyer, V. S.; Khalifa, N.; Mahadevan, R.; Venugopalan, S.; Weerawarna, S. A.; Zhou, P. *J. Am. Chem. Soc.* **1995**, *117*, 1514. (j) Laali, K. K.; Okazaki, T.; Mitchell, R. H.; Ward, T. R. *J. Org. Chem.* **2001**, *66*, 5329. (k) Laali, K. K.; Okazaki, T.; Mitchell, R. H.; Ayub, K.; Zhang, R.; Robinson, S. G. *J. Org. Chem.* **2008**, *73*, 457.
- ² Butterfield, A. M.; Gilomen, B.; Siegel, J. S. *Org. Proc. Res. Dev.* **2012**, *16*, 664.
- ³ (a) Reisch, H. A.; Bratcher, M. S.; Scott, L. T. *Org. Lett.* **2000**, *2*, 1427. (b) Tsefrikas, V. M. Ph.D. dissertation, Boston College, Chestnut Hill, Massachusetts, 2007. (c) Greene, A. K. Ph.D. dissertation, Boston College, Chestnut Hill, Massachusetts, 2012.
- ⁴ IUPAC name: acephenanthryleno[3,4,5,6-*fg hij*]picene
- ⁵ McMahon, B. J. B.S. thesis, Boston College, Chestnut Hill, Massachusetts, 1997.
- ⁶ Mehta, G.; Srirama Sarma, P. V. V. *Chem. Commun.* **2000**, 19.
- ⁷ Smith, N. J. Ph.D. dissertation, Boston College, Chestnut Hill, Massachusetts, 2011.
- ⁸ Kamatani, J.; Yamada, N.; Saitoh, A. *PCT Int. Appl.* WO 2010071223, June 24, 2010.
- ⁹ Bennett, M. A.; Neumann, H. *Aust. J. Chem.* **1980**, *33*, 1251.

- ¹⁰ (a) Diltthey, W.; ter Horst, I.; Schommer, W. *J. Prakt. Chem.* **1935**, 143, 189. (b) Allen, C. F. H.; Van Allen, J. A. *J. Org. Chem.* **1952**, 17, 845.
- ¹¹ Mackenzie, K. *J. Chem. Soc.* **1960**, 473.
- ¹² Fort, E. H.; Scott, L. T. *Angew. Chem., Int. Ed.* **2010**, 49, 6626.
- ¹³ (a) Dinadayalane, T. C.; Priyakumar, U. D.; Sastry, G. N. *Theochem-J. Mol. Struct.* **2001**, 543, 1. (b) Dinadayalane, T. C.; Sastry, G. N. *J. Org. Chem.* **2002**, 67, 4605. (c) Sastry, G. N. *Theochem-J. Mol. Struct.* **2006**, 771, 141.
- ¹⁴ (a) Haddon, R. C.; Scott, L. T. *Pure Appl. Chem.* **1986**, 58, 137. (b) Haddon, R. C. *J. Am. Chem. Soc.* **1986**, 108, 2837. (c) Haddon, R. C. *Chem. Phys. Lett.* **1986**, 125, 231.
- ¹⁵ See, for example: Seiders, T. J.; Baldrige, K. K.; Grube, G. H.; Siegel, J. S. *J. Am. Chem. Soc.* **2001**, 123, 517.
- ¹⁶ Dolhem, E.; Ocafrain, M.; Nedelec, J. Y.; Troupel, M. *Tetrahedron* **1997**, 53, 17089.

TOC graphic