2012

Vertical and horizontal photobiont transmission within populations of a lichen symbiosis

Post-print/Accepted manuscript

Francesco Dal Grande

Ivo Widmer

Helen H. Wagner

Christoph Scheidegger

DAL GRANDE, F., WIDMER, I., WAGNER, H.H. and SCHEIDEGGER, C. (2012), Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. *Molecular Ecology*, 21: 3159–3172. doi:10.1111/j.1365-294X.2012.05482.x

This is the peer reviewed version of the following article: DAL GRANDE, F., WIDMER, I., WAGNER, H.H. and SCHEIDEGGER, C. (2012), Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. *Molecular Ecology*, 21: 3159–3172, which has been published in final form at [doi:10.1111/j.1365-294X.2012.05482.x](https://doi.org/10.1111/j.1365-294X.2012.05482.x) This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

HOW TO CITE TSPACE ITEMS

Always cite the published version, so the author(s) will receive recognition through services that track citation counts, e.g. Scopus. If you need to cite the page number of the TSpace version (original manuscript or accepted manuscript) because you cannot access the published version, then cite the TSpace version **in addition to** the published version using the permanent URI (handle) found on the record page.

Vertical and horizontal photobiont transmission within populations of a lichen symbiosis

F. Dal Grande^{1,3*}, I. Widmer^{1,4*}, H.H. Wagner², C. Scheidegger¹

¹ Biodiversity and Conservation Biology, WSL Swiss Federal Research Institute, 8903 Birmensdorf, Switzerland

² Department of Ecology and Evolutionary Biology, University of Toronto, L5L 1C6 Mississauga, Canada

³ Present Address: Biodiversity and Climate Research Centre (BiK-F), Senckenberg Gesellschaft fuer Naturforschung, 60325 Frankfurt am Main, Germany

⁴ Present Address: LPED - Laboratory of Population Environment Development, University of Provence, 13331 Marseille Cedex 03, France

*These two authors contributed equally to this work and are considered joint first authors.

Corresponding Author: Dal Grande Francesco

Biodiversity and Climate Research Centre (BiK-F), Senckenberg Gesellschaft fuer

Naturforschung, 60325 Frankfurt am Main, Germany - Tel. +49 (0)69 798 24798 Fax. +49

(0)69 798 24771

e-mail: Francesco.DalGrande@senckenberg.de, francesco.dalgrande@wsl.ch

Keywords: Population Genetics - Empirical, Ecological Genetics, Algae, Fungi, *Lobaria pulmonaria*, Microsatellite

Running head: Dynamics of a lichen symbiosis

Abstract

Lichens are widespread symbioses and play important roles in many terrestrial ecosystems. The genetic structure of lichens is the result of the association between fungal and algal populations constituting the lichen thallus. Using eight fungus- and seven alga-specific highly variable microsatellite markers on within-population spatial genetic data from 62 replicate populations across Europe, North America, Asia and Africa, we investigated the contributions of vertical and horizontal transmission of the photobiont to the genetic structure of the epiphytic lichen *Lobaria pulmonaria*. Based on pairwise comparisons of multi-locus genotypes defined separately for the mycobiont and for the photobiont, we inferred the transmission mode of the photobiont and the relative contribution of somatic mutation and recombination. After constraining the analysis of one symbiont to pairs of individuals with genetically identical symbiotic partners, we found that 77 % of fungal and 70 % of algal pairs were represented by clones. Thus, the predominant dispersal mode was by means of symbiotic vegetative propagules (vertical transmission), which dispersed fungal and algal clones co-dependently over a short distance, thus shaping the spatial genetic structure up to distances of 20 m. Evidence for somatic mutation generating genetic diversity was found in both symbionts, accounting for 30 % of pairwise comparisons in the alga and 15 % in the fungus. While the alga did not show statistically significant evidence of recombination, recombination accounted for 7.7 % of fungal pairs with identical algae. This implies that, even in a mostly vegetatively reproducing species, horizontal transmission plays a role in shaping the symbiotic association, as shown in many coral and other symbioses in nature.

Introduction

Lichens are symbiotic organisms composed of a fungal partner (mycobiont) and a population of algae and/or cyanobacteria (photobiont). Mycobionts express their symbiotic phenotype only in association with compatible photosynthetic partners, and the tight morphological integration and physiological dependence of the symbionts result in a distinct lichen body called thallus (Ahmadjian 1993). In lichens, the mechanism for symbiotic contact and thallus formation in nature is only partially understood. Reproduction and dispersal of lichens is a complex process since both partners have to be present for the successful development of a new lichen thallus (Honegger 1998, 2008; Dobson 2003). A vast majority of lichens have a sexual and asexual life cycle. In the sexual life cycle, fungal spores are released from specialized structures on the thallus (ascomata). Upon germination, fungal spores must obtain a compatible algal or cyanobacterial partner, which may be free-living (Etges & Ott 2001; Sanders & Lücking 2002; Sanders 2005; Handa *et al.* 2007; Hedenås *et al.* 2007; Macedo *et al.* 2009) or obtained through capture from another lichen (Friedl 1987; Ott 1987a,b; Stenroos 1990; Rambold & Triebel 1992; Ott *et al.* 1995; Gaßmann & Ott 2000; Lücking & Grube 2002). In the vegetative life cycle, mycobiont and photobiont are simultaneously dispersed within specialized asexual propagules (e.g., corticated protuberances called isidia or non-corticated clumps called soredia) or through thallus fragmentation.

The genetic structure of a lichen population will be strongly influenced by the manner in which photobionts are dispersed and transmitted to the fungus (Hill 2009). Vertical (or co-dependent) transmission occurs when the photobiont disperses as part of the vegetative propagule of the lichen, thus presumably representing the predominant process in exclusively or nearly exclusively asexual lichen species (Werth & Sork 2010). The vegetative propagules produce physically separate but genetically identical thalli, i.e., thalli with fungal and algal components

genetically identical to the mother thallus (Paulsrud *et al.* 1998; Doering & Piercey-Normore 2009). On the other hand, horizontal (or independent) transmission usually occurs when the fungus reproduces sexually. The sexual life cycle is considered to reshuffle the genetic composition of the lichen, generating new combinations of fungal and algal genotypes (i.e., genetically different thalli). Horizontal transmission may also depend on the dispersal ability of the photobiont. The ability of green-algal photobionts to move is very restricted, as they usually do not disperse (either sexually or asexually) while embedded in the lichen thallus (Sluiman *et al.* 1989; Nash 1996). However, many green-algal photobionts can occur in free-living populations on soil, rocks, or tree stems (Mukhtar *et al.* 1994; Beck *et al.* 1998; Friedl & Büdel 2008), and viable photobiont cells are found in fecal pellets of lichenivorous snails (Meier *et al.* 2002; Boch *et al.* 2011). Moreover, horizontal transmission of algae has been shown in asexual (e.g., Nelsen & Gargas 2008, 2009) or nearly asexual (Piercey-Normore 2006; Wornik & Grube 2010) lichen species.

Studies on the mode of transmission of lichen photobionts in natural populations remain scarce. In particular, the genetic composition of a lichen thallus (i.e., its individual mycobiont and photobiont genotypes) and its fine-scale spatial distribution has never been reliably assessed at the within population scale due to the lack of appropriate genetic markers. Marker resolution becomes in fact critical when studying highly clonal organisms such as lichens, for which multilocus genotypes are the only way to identify genetically distinct individuals (Arnaud-Haond *et al.* 2007).

This work aims to assess the relative contribution of vertical vs. horizontal transmission to the intra-population genetic structure of the mycobiont and of the photobiont of a mainly vegetative lichen species. The model species is the epiphytic lichen *L. pulmonaria*, which is widespread in the Northern Hemisphere (Yoshimura 1971). Recently, microsatellite markers have been developed for

its haploid eukaryotic symbionts (mycobiont: Walser *et al.* 2003; Widmer *et al.* 2010; this study; green algal photobiont: Dal Grande *et al.* 2010). The high mutation rate of microsatellite loci gives them a far greater resolving power than previous, sequence-based studies performed on lichen populations (e.g., Beck *et al.* 2002; Lohtander *et al.* 2003; Printzen *et al.* 2003; Printzen & Ekman 2003; Yahr *et al.* 2004; Lindblom & Ekman 2005, 2007; Selkoe & Toonen 2006; Doering & Piercey-Normore 2009; Lättmann *et al.* 2009; Werth & Sork 2010). *Lobaria pulmonaria* is highly selective towards its green algal photobiont, i.e., it is associated with the coccoid green alga *Dictyochloropsis reticulata* (Tschermak-Woess) Tschermak-Woess throughout its entire distribution range (Dal Grande 2011).

Earlier studies suggested that the predominant dispersal mode in *L. pulmonaria* is by means of vegetative propagules (Zoller *et al.* 1999; Walser 2004; Wagner *et al.* 2005, 2006; Werth *et al.* 2006a,b, 2007), and showed its mycobiont populations to be highly clonal, suggesting a predominance of vertical transmission of the photobiont. However, the mycobiont of *L. pulmonaria* can undertake sexual reproduction, hence the photobiont also needs to be transmitted horizontally. While no evidence of free-living photobiont populations has been found to date (Tschermak-Woess 1978; Dal Grande 2011), the presence of zoospores (motile flagellate asexual cells) indicates that the photobiont has the potential to move locally (i.e., on the same tree) once released from the thallus (Richardson 1999; Friedl & Büdel 2008).

The availability of symbionts may impose limits on the distribution of the other partner, particularly in cases where the association is obligate (Andras *et al.* 2011). Werth *et al.* (2007) demonstrated for the mycobiont of *L. pulmonaria* that gene flow is spatially restricted, resulting in spatial aggregation of fungal clones. Based on the notion that spatial processes, such as reproduction

111 followed by dispersal, leave a characteristic spatial signature (Seabloom *et al.* 2005; Wagner &
112 Fortin 2005), analysis of spatial genetic structure may be used to identify the underlying processes.
113 In particular, vertical transmission of the photobiont is expected to result in short-distance spatial
114 aggregation of fungal and algal clones, while horizontal transmission due to mycobiont sexual
115 reproduction will decouple photobiont-mycobiont pairs at larger distances (Werth & Sork 2010).

116 This paper addresses the following questions: (a) what is the relative contribution of vertical
117 vs. horizontal transmission of the photobiont to the genetic structure of the lichen populations? (b)
118 What is the relative contribution of the micro-evolutionary processes of mutation and recombination
119 to the current fungal and algal intra-population genetic diversity? (c) Are there differences in the
120 within-population spatial genetic structure between mycobiont and photobiont?

121 To address these questions, we introduce an approach that takes advantage of the
122 microsatellite markers for both the fungal and algal partners. This method allows for the reliable
123 identification of clonal thalli (i.e., thalli with identical multilocus genotypes for the fungus and the
124 alga, respectively). Under the assumption that pairs of thalli with identical multilocus genotypes
125 both for the fungus (MLG_F) and for the alga (MLG_A) within a population result from the vegetative
126 co-dispersal of fungal and algal clones, we can infer within-population evolutionary processes (such
127 as mutation and recombination) by restricting analysis for one symbiont to pairs of thalli with
128 identical MLG in the other symbiont. While statistical inference of (spatial) genetic structure within
129 populations is often limited by a lack of independent replicate populations, we illustrate our
130 approach with a data set of 62 range-wide populations that allows robust statistical analysis.

131 This research assesses the way photobionts are transmitted in a predominantly asexual taxon
132 and provides insights into the contribution of the micro-evolutionary processes of mutation and

recombination to the genetic structure of lichen populations.

Materials and Methods

Sample collection and molecular genetic analysis

The goal of our design was to detect the intra-population genetic structure of the fungal and algal symbionts of *L. pulmonaria* among adjacent trees. This design would not detect either the extent of the overall genetic clustering on the same tree or the extent of gene flow among populations. In total, 2229 thalli of *L. pulmonaria* were sampled from 62 populations across Europe, North America, Asia and Africa (Table S1, Supporting Information). The median distance between a population and the nearest neighbor sampled population was 115 km, and all but nine populations were at least 25 km from their nearest neighbor population. For the purpose of our analyses, a population was defined as a stand of trees colonized by *L. pulmonaria*. Across each population, 1 – 3 thalli were randomly taken from an average of 23 nearest neighbor trees (i.e., proceeding from a sampled tree to its nearest unsampled neighbor tree). The maximum distance among the sampled trees within each population typically was < 1500 m except for three populations, with a median maximum distance of 274 m and a minimum of 16 m. Thalli collected on a single tree were separated by about 50 cm and positioned on different sides of the trunk. This sampling design allows for the investigation of microsatellite variation within a population of *L. pulmonaria* (Walser *et al.* 2003; Wagner *et al.* 2005). On average 31 thalli were collected per population, which has been found to be an appropriate number to resolve within-population mycobiont and photobiont genetic structure (Werth 2010).

Eight fungus-specific (LPu03, LPu09, LPu15, LPu23, LPu24, LPu25, LPu28, Walser *et al.*

2003; Widmer *et al.* 2010; MS4, this study) and seven alga-specific microsatellite markers (LPh1 to LPh7; Dal Grande *et al.* 2010) were amplified from total lichen DNA. For primer sequences, including redesigned primers for LPu25, labeling and PCR conditions see Table S2 (Supporting Information). Fragment lengths were determined on a 3730 DNA Analyzer (Applied Biosystems, Foster City, CA), and electropherograms were analyzed with GENEMAPPER 3.7 (Applied Biosystems, Foster City, CA) using LIZ-500 as internal size standard. Multilocus genotypes were defined separately for the fungus (MLG_F, based on eight loci) and for the alga (MLG_A, based on seven loci).

Statistical analyses

Data sets

Recurrent MLGs could either be the result of vegetative reproduction or chance products of sexual reproduction (Arnaud-Haond *et al.* 2007). Therefore, recurrent MLGs were only interpreted as clones if they were unlikely to result from sexual reproduction given the observed allele frequencies in a population. We calculated for each population the probability of observing two sexually produced fungal or algal individuals identical at all eight or seven microsatellite loci, respectively. This method, implemented in the software GENCLONE v2.0 (Arnaud-Haond & Belkhir 2006), is based on the round-robin method proposed by Parks & Werth (1993), which allows for each MLG an estimate of the probability of obtaining the observed number of recurrent MLGs in the data set by sexual reproduction under random mating (P_{sex}). The method thus takes into account relative levels of polymorphism (Supplementary Material, Table S4). We used the P_{sex} to assess the likelihood that identical MLGs were of sexual origin. The significance of P_{sex} was tested at $\alpha = 0.05$

with 1000 simulations. When significant (i.e., $P_{\text{sex}} < 0.05$), we considered recurrent MLGs as true clones. Recurrent MLGs with $P_{\text{sex}} \geq 0.05$ were excluded from analyses (Arnaud-Haond et al. 2007).

To analyze the genetic diversity of the fungal and algal symbionts, each pair of thalli of *L. pulmonaria* was analyzed for the number of microsatellite loci at which they differed in the fungal genotype MLG_F (“*deltaF*”) and in the algal genotype MLG_A (“*deltaA*”) (see Fig. 1 for a graphic representation). All analyses were restricted to pairwise comparisons of thalli within populations. Three subsets A, B and C of the data were used for analysis as defined in Table 1.

Pairwise comparisons within populations are not independent, hence statistical tests cannot rely on parametric tests and true replication requires independent data from multiple study sites. To allow for robust statistical estimation, we pooled data over all 62 populations and derived bootstrap estimates of standard errors in R (R Development Core Team 2008) by leaving out one population at a time.

Relative contribution of vertical vs. horizontal photobiont transmission

Pairs of thalli were scored as resulting from co-dependent dispersal of the symbionts (vertical photobiont transmission) if they had identical MLGs of both the fungus and the alga, i.e., $\text{deltaA} = \text{deltaF} = 0$. We assessed the relative contribution of vertical photobiont transmission to population genetic structure by the proportion of pairs of thalli with identical MLGs for both symbionts among the pairs of thalli in data set B. We derived a bootstrap estimate of this proportion by omitting one population at a time.

Since the sexual life cycle is considered to be the main factor responsible for the independent dispersal of the symbionts (horizontal photobiont transmission), creating new genotypic

combinations of fungi and algae, we further analyzed pairs of thalli with different MLGs for the fungus and/or the alga. We modeled the contribution of somatic mutation and recombination to the observed differences at the microsatellite loci as follows (see Fig. 2):

(i) Empirical null-model of recombination (Fig. 2, top). Recombination may result in pairs of thalli with any number of differing loci. We derived an empirical null model of the distribution of the expected number of loci difference (*deltaA* or *deltaF*) based on observed allele frequencies within each population. We permuted repeat lengths for each microsatellite marker among the thalli sampled from the same population (data set A), separately for the alga and for the fungus. We thus simulated thalli with new MLGs based on the observed allele frequencies within each population under the assumption of random mating within populations, taking into account observed levels of marker polymorphism and clonality in each population. We repeated the simulation 100 times and evaluated for each run the frequency distribution of number of loci differing between each pair of simulated thalli from the same population (*deltaA*, *deltaF*). We pooled the simulated frequencies across the 62 populations for each simulation run and then averaged over all 100 simulation runs. All calculations were performed in R (R Development Core Team 2008). Simulated probabilities for obtaining identical MLGs by recombination were $3.4 * 10^{-4}$ for *deltaF*=0 and $6.4 * 10^{-4}$ for *deltaA*=0 (not shown in Fig. 2).

(ii) Negative exponential distribution model accounting for somatic mutation (Fig. 2, center). Mutations are assumed to occur independently for each locus and for each symbiont. Hence, over many generations and in an otherwise only vegetatively reproducing population, mutation will first lead to difference in a single locus, a subsequent mutation to difference in one additional locus, and so on, following a negative exponential model defined by parameter lambda.

(iii) Model fitting to the fungal and algal data (Fig. 2, bottom). To assess to what degree the observed non-clonal pattern in a symbiont was the result of mutation versus recombination, the parameter lambda of an exponential function was fitted to data set C using the function 'nls' in R (R Development Core Team 2008) and accounting for the null model of recombination. This resulted in estimates for lambda of 0.55 for the fungus and 0.77 for the alga. We then performed a linear regression of the frequency distribution of the number of differing loci as a function of the exponential model (representing mutation) with the fitted parameter lambda and the null model of recombination, with no intercept, and assessed model fit, statistical significance of regression coefficients, and the relative contribution of the exponential model and the empirical null model of recombination to the frequency of pairs per number of differing loci. For each level of *deltaA* or *deltaF*, we assessed the proportional contribution by each component model to the fitted value (e.g., if 100 pairs were predicted, 37 may be predicted by the exponential model and 63 by the empirical null model of recombination). We multiplied these proportions by the observed frequency of each level of *deltaA* or *deltaF* in data set C to estimate the ratio of mutation vs. recombination among the non-clonal component. Bootstrap mean and standard error of this ratio were determined by leaving out one population at a time.

Assuming an average microsatellite mutation rate of 10^{-3} , the expected probability of observing at least one mutation in the alga (with 7 independent loci) is 0.0068, the expected probability of observing at least one mutation in the fungus (with 8 independent loci) is 0.0077. The expected probability of mutation occurring in both symbionts independently at the same time is $(6.8 * 10^{-3}) * (7.7 * 10^{-3}) = 5.2 * 10^{-5}$. The probability that both symbionts show a somatic mutation was thus expected to be less than 1% of the probability for somatic mutation in either symbiont and

considered negligible.

Spatial genetic structure

To assess spatial genetic structure within populations, we determined for data set A the probability of sampling a pair of thalli from the same population belonging to one of the following categories as a function of their distance in space: (i) clonal thalli ($\Delta F = \Delta A = 0$), (ii) fungal clones associated with different algal MLGs ($\Delta F = 0$ and $\Delta A > 0$), (iii) algal clones associated with different fungal MLGs ($\Delta F > 0$ and $\Delta A = 0$) and (iv) different fungal MLGs associated with different algal MLGs ($\Delta F > 0$ and $\Delta A > 0$). The first distance class contained pairs of thalli sampled from the same tree. Distance class boundaries were defined on a logarithmic scale, with the last distance class containing all pairwise comparisons at distances > 500 m.

Results

After exclusion of recurrent MLGs that were not assessed as true clones (P_{sex} values > 0.05 , thalli), had incomplete genotype assessment (55 thalli) or missing spatial coordinates (5 thalli), the data set consisted of 1960 thalli. We found 1051 MLGs for the haploid fungus and 1025 MLGs for the haploid alga, with a total of 1256 MLGs based on all 15 markers from both symbionts (Table S1, Supporting Information: numbers of different MLGs per population; Table S4, Supporting Information: allele frequency distribution per population at eight fungal and seven algal loci). Multiple fungal or algal genotypes within the same thallus were not found in any of the populations.

Our analyses were based on 36,218 pairwise comparisons within populations pooled over 62 populations (Table 1). The relative frequency distribution of the number of loci differences (ΔA ,

deltaF; calculated over all pairs within populations (data set A) was similar for both symbionts, with the highest frequency of thalli differing by four loci each for the fungus and for the alga (Fig. 3A). In both symbionts, we found a high frequency of identical pairs of thalli (2977 pairs with $\Delta A=0$; 3285 pairs with $\Delta F=0$; Fig. 3A).

Vertical transmission of the photobiont

2294 pairs had identical MLGs both for the alga and the fungus ($\Delta A=\Delta F=0$). When the analysis was restricted per symbiont to those pairs of thalli displaying an identical MLG in the other symbiont (data set B), both the alga and the fungus displayed a high degree of clonality, with $\Delta A=0$ and $\Delta F=0$ as the predominant classes (Fig. 3B). The proportion of clonal comparisons was higher for the fungus (77.06 % $\pm$ 0.72 %) than for the alga (69.85 % $\pm$ 0.60 %).

Identifying micro-evolutionary processes of mutation and recombination

After the exclusion of recurrent genotypes within each population and constraining by clonality in the other symbiont ($\Delta F=0$ for algal MLGs and vice versa, data set C), we found 215 algal and 269 fungal pairs of MLGs that differed from each other in at least one locus (Table 1, Fig. 3C). For both symbionts, the largest proportion of these pairs differed at only one locus (ΔA or $\Delta F=1$). The alga showed a strongly skewed distribution of the number of loci differences as expected under a negative exponential model resulting from mutation (Fig. 3C, left). In the fungus, the distribution was bimodal, suggesting the presence of an additional process (Fig. 3C, right).

In fungal sexual reproduction, each ascoma (i.e., reproductive structure of the fungus) may either form meiotic fungal spores with the same MLGs or spores with different MLGs. Both spore

types may form new associations with either the same or a different algal MLG. The empirical null models of recombination based on the observed allele frequencies estimated that under random mating within each population, 0.033 percent of fungal recombinations and 0.065 percent of algal recombinations would result in the same MLG as the mother thallus. The secondary peak in the distribution of ΔF given $\Delta A=0$ was proportional to the frequency distribution expected from the empirical null-model of recombination (Fig. 3C, right). The combination of an exponential model representing mutation and the empirical null model of recombination explained the distribution of the number of loci differences for each pair of fungi with identical algae well, explaining a total of 96.5% of the variation for the fungus, whereas for the alga, the exponential model alone explained 98.5 % of the variation (Table S3, Supporting Information). Taking into account the above estimate of 77.06 % (± 0.72 %) clonality in the fungus, the fitted models resulted in an overall estimate of 15.21 % (± 0.25 %) of pairwise comparisons of fungal MLG being affected by mutation and 7.73 % (± 0.25 %) being affected by recombination. For the alga with 69.85 % clonality (± 0.60 %), mutation thus accounted for 30.15 %.

Spatial genetic structure

Clonality depended strongly on distance (Fig. 4). There was a marked decrease in the frequency of pairs of thalli with identical fungal and algal MLGs (vertically transmitted photobionts, $\Delta F=\Delta A=0$) within the first 20 m, compensated by an increase in the frequency of pairs that differed both in the alga and in the fungus ($\Delta F>0$ and $\Delta A>0$).

The relative frequency of distance classes for each type of pairs showed significant differences between the two symbionts (Fig. 5). For the alga, pairs with differences at 1 or more loci

($\Delta A > 0$, $\Delta F = 0$) decreased in number over short distances, similarly to the distribution of clonal thalli. Fungal MLG pairs differing at least at one locus ($\Delta F > 0$, $\Delta A = 0$) showed a different spatial pattern similar to that of pairs differing in both symbionts ($\Delta F > 0$, $\Delta A > 0$).

Discussion

Prevalence of vertical transmission of the photobiont

Based on previous evidence that the fungus reproduces mainly clonally (Walser 2004; Wagner *et al.* 2005; Werth *et al.* 2006b, 2007), we expected the photobiont of *L. pulmonaria* to disperse primarily vertically within vegetative propagules. Vegetative reproduction will recreate the MLG of the mother thallus unless there is mutation in at least one symbiont. This should result in a dominating component of pairs of thalli displaying identical MLGs for both symbionts. Indeed, when the analysis was restricted per symbiont to those pairs of thalli displaying an identical MLG in the other symbiont (data set B), we found that the predominant class of MLG comparisons was composed of pairs of thalli having identical MLGs for both symbionts. The probability of creating identical MLGs through sexual reproduction was small enough to be neglected (as tested here with P_{sex} for each MLG, and further supported by the empirical null model of recombination, which estimated the overall probability, combined for all MLGs within a population, at less than 0.1 % for either symbiont), therefore we interpreted recurring MLGs as clones, i.e., resulting from vegetative reproduction. Thus, based on microsatellite fingerprinting of both lichen symbionts, we demonstrated that the photobiont of *L. pulmonaria* is mostly vertically transmitted.

Vegetative dispersal with symbiotic propagules ensures the continuity of a successful combination of MLGs of the two partners across generations (Margulis 1993; Yahr *et al.* 2004,

2006; Reeve & Hölldobler 2007; Zilber-Rosenberg & Rosenberg 2008). It has been suggested that vegetative dispersal in lichens has the advantage of producing large numbers of locally adapted propagules that can readily exploit and colonize the local environment (Ott 1987b; Sanders & Lücking 2002; Walser 2004). A predominance of vertical transmission of photobionts has also been confirmed in other multicellular symbiotic systems for which genetic uniformity is favored by selection for cooperative traits (Gilbert *et al.* 2009). For instance, in some symbiotic systems, such as maternally inherited endosymbionts (Saffo 1992; Moran & Baumann 1994; Huigens *et al.* 2000), grass endophytes (Clay 1990; Saikkonen *et al.* 2002), corals (Coates & Jackson 1987) or sea anemones (Geller & Walton 2001), strong population structure and shared phylogenetic history of symbionts are expected because of the vertical transmission of symbionts (Brem & Leuchtman 2003).

Contribution of mutation and recombination to within-population genetic structure of lichen symbionts

The high variability of the microsatellite markers used in this study, combined with the approach constraining the analyses of one symbiont to pairs of individuals with genetically identical symbiotic partners and the availability of data from 62 replicate populations, provided robust evidence for patterns of mutation and recombination in *L. pulmonaria* symbionts. The alga showed a clear signal of mutation as indicated by the exponential distribution of the number of loci differences (Fig. 3C, left). Considering that no statistically significant signal of recombination was found, our results indicate that the photobiont *D. reticulata* may reproduce strictly asexually. In lichen photobionts, the production of zoospores (motile, flagellate spores) is considered a means to escape from the thallus,

close to which they can form colonies (Slocum *et al.* 1980; Scheidegger 1985). The occurrence of free-living colonies is known for the green-algal genus *Trebouxia* (Tschermak- Woess 1978; Bubrick *et al.* 1985; Mukhtar *et al.* 1994), and their zoospores are known to frequently undergo sexual fusion in fresh cultures (Ahmadjian 1959). Despite extensive investigation, the production of zoospores was never observed in the photobiont of *L. pulmonaria* (Skaloud 2008).

It is remarkable that the alga, with one marker less than the fungus, exhibited a comparable level of genetic diversity within populations to the fungus (Fig. 3A). With no evidence for recombination and having shown that the alga is mostly co-transmitted vertically with the mycobiont, this may be the result of faster mutation rates in the algal microsatellites combined with a greater number of generations in the photobiont. An alternative explanation involves the introduction of new alleles into the populations through the horizontal transmission (symbiont capture) from other photobiont populations found in lichen species associated with *D. reticulata* (genera *Lobaria* and *Sticta*; Dal Grande 2011). The evidence of mutation obtained in our study concurs with the hypothesis that mutation is the key process creating genetic diversity in clonal organisms (Higgs & Woodcock 1995; Tomiuk *et al.* 1998; Butlin 2002; Vogler *et al.* 2006; Ally *et al.* 2008; Mock *et al.* 2008).

The sporadic presence of fruiting bodies in *L. pulmonaria* indicates that the mycobiont can undertake sexual reproduction by forming ascospores. Sexual reproduction involves the process of relichenization, i.e., the formation of a new thallus once fungal spores found a suitable alga (horizontal transmission). Our results show that recombination significantly contributes to the fungal genetic structure (7.73 %, ± 0.25 % of pairwise comparisons of fungal MLG; Table S3). Hence, despite the predominance of vertical transmission, horizontal transmission plays a non-negligible

375 role in shaping the genetic composition of the lichen population.

376 Our approach, however, does not allow to distinguish the effect of horizontal transmission of
377 the photobiont related to fungal sexual reproduction from the process of horizontal movement of
378 photobiont from nearby vegetative propagules, which may affect the interpretation of our results
379 through reshuffling of the genetic composition of lichen thalli independently from fungal sexual
380 reproduction. Previous studies have shown that, even where both partners are co-dispersed in
381 specialized propagules, de-differentiation (separation of algal and fungal partners) allows vertically
382 transmitted algae to be replaced by others available in the environment, or even to be captured from
383 other nearby lichen species (Friedl 1987; Ott *et al.* 1995; Ohmura *et al.* 2006; Wornik & Grube
384 2010; Dal Grande 2011). The frequency of such algal substitutions in nature is unknown (Piercey-
385 Normore & DePriest 2001), but this strategy may provide a mechanism for optimizing symbiotic
386 composition in a local environment (Friedl 1987; Ott 1987b; Ohmura *et al.* 2006; Yahr *et al.* 2006).

387 The way horizontal algal movement and relichenization take place remains elusive, and these
388 processes deserve further attention. They may well be key evolutionary processes in lichen
389 communities, allowing the formation of photobiont-mediated guilds among unrelated lichen-forming
390 fungi (Beck *et al.* 2002; Rikkinen *et al.* 2002; Rikkinen 2003). In other symbioses, evidence for
391 horizontal symbiont transmission has been reported, for instance in certain corals and their
392 symbiotic dinoflagellates (Rowan 1998; Loh *et al.* 2001), in insects and their endosymbiotic bacteria
393 (Huigens *et al.* 2000; Sirviö & Pamilo 2010), or in fungus-gardening ants or termites and fungal
394 cultivars (Aanen *et al.* 2002; Mikheyev *et al.* 2007). Our results showed that the lichen symbiosis is
395 formed by a strictly asexual partner (alga) and by a fungal partner that conserved the sexual pathway
396 together with the formation of asexual diaspores co-dispersing both partners. Sexual propagules are

considered important for long-distance dispersal of the mycobiont (Walser 2004; Seymour *et al.* 2005; Cassie & Piercey-Normore 2008; Scheidegger & Werth 2009) and increase the number of genotypes in local populations, thus potentially enhancing adaptation (Maynard Smith 1986; Samadi *et al.* 1999; Rice & Chippindale 2001; Blaha *et al.* 2006; Foucaud *et al.* 2006).

The symbiotic relationship is obligatory for the fungal partner in *L. pulmonaria* to complete its life-cycle (Ott 1987b; Ingold & Hudson 1993; Honegger 2001). Yet, little is known about how often and under what conditions sexual reproduction and relichenization occur in natural habitats. So far, no corresponding estimates from molecular data were available (Honegger 2001; Dobson 2003). Our study suggests that independent dispersal of the symbionts does occur in natural populations of *L. pulmonaria* and that it has a considerable impact on the genetic diversity of lichen populations.

Spatial genetic structure

We analyzed the spatial genetic structure of lichen populations to infer about dispersal processes related to horizontal and vertical transmission of the photobiont. While statistical analysis of spatial patterns is often limited by the lack of replicate study areas, the availability of comparable spatial genetic data from 62 replicate populations allowed robust statistical analysis based on bootstrap estimates.

Here we showed that the fungal and algal clonal components had a large impact on the small-scale spatial genetic structure of the lichen association, and the signal of clonality markedly decreased within a distance of about 20 m (Fig. 4). Our results indicate that vegetative propagules play a dominant role to disperse genetically identical symbionts of *L. pulmonaria* over short spatial distances within populations. They are thus a means of rapid lichen spread at the local scale

(Hawksworth & Hill 1984; Heinken 1999; Dettki *et al.* 2000; Sillett *et al.* 2000). The restricted dispersal can be explained by these propagules' larger size compared to fungal ascospores, since the larger the propagule the shorter the distance they can be carried by wind, water or animals (Heinken 1999; Walser 2004; Werth *et al.* 2006b; Scheidegger & Werth 2009).

The differences in the reproductive modes between the two symbionts of *L. pulmonaria* described above were clearly reflected in their spatial genetic structure. We expected that the symbionts mainly spread within the vegetative propagules of the lichen, and thus would present similar spatial structures. The alga, which only showed a signal of mutation, confirmed this assumption by displaying almost an identical spatial pattern as the clones (with $\Delta A = \Delta F = 0$; Fig. 5). The fungus, which showed signals of both mutation and recombination, exhibited a different spatial genetic structure suggesting dispersal over larger distances.

Conclusions

This paper presents a novel approach to analyze relatively recent, within population micro-evolutionary processes from the population genetic structure of the lichen *L. pulmonaria*. We provided robust evidence for the predominance of vertical transmission of the photobiont at the intra-population level in a mainly vegetative species.

As a caveat, we recall some key assumptions to our analysis. First, we assume that long-distance migration leads to the introduction of new genotypes, i.e., that it is unlikely that the same MLG would originate independently in two populations and migrate from one to the other, thus reflecting larger-scale processes within our data set B. Based on observed allele frequencies within populations, we estimated the probability of independent origin of clones as <0.1 % each for the

fungus and the alga. The probability of independent origin in different populations and subsequent immigration is expected to be much lower still. Second, we assume that each mutation leads to a new allele, such as expected under an infinite-alleles mutation model. Multiple identical but independent mutations within the same population, as might be expected under a step-wise mutation model, would lead to underestimation of the relative contribution of mutation due to the exclusion of recurrent MLGs in data set C. Third, we assumed independent mutation in both symbionts at the same time to be negligible, and we estimated its probability as <1% of the probability of mutation in either symbiont. Concurrent mutation in both symbionts would reduce the relative size of data set B but should not otherwise bias results.

We inferred the different processes shaping the genetic structure of the symbionts, highlighting that, even in a species with rare sexual reproduction such as *L. pulmonaria*, fungal recombination is a process shaping the genetic structure between the two lichen symbionts. The possibility of sexual reproduction is important to population genetics. Considering the low germination rate in some lichen species, it may seem unlikely that their ascospores would ever develop into a lichen thallus. However, even if only a few out of the thousands of ascospores produced in one ascoma find the proper photobiont to reconstitute the symbiosis, as long as the new thallus multiplies and disperses through vegetative propagules, this may suffice to alter lichen population genetic structure (Honegger & Zippler 2007).

Acknowledgments

This research was founded by the Swiss National Science Foundation (projects 31003A-105830 and 31003A-127346 to C Scheidegger) and an NSERC Discovery Grant (to HH Wagner). We are

grateful to James B. Anderson, Rolf Holderegger and four anonymous reviewers for valuable suggestions on the manuscript, Ariel Bergamini for assistance with statistical analyses, Heather Cole, Carolina Cornejo and Silke Werth for comments of the manuscript, Vladimir Mikryukov, Christine Keller and other collaborators (listed in Table S1, Supporting Information) for field collection of samples. Authors acknowledge the Genetic Diversity Center at ETH Zürich, Switzerland (CCED-GDC) for technical assistance.

Literature cited

- Aanen DK, Eggleton P, Rouland-Lefevre C, Guldberg-Froslev T, Rosendahl S, Boomsma JJ (2002) The evolution of fungus-growing termites and their mutualistic fungal symbionts. *Proceedings of the National Academy of Sciences USA*, **99**, 14887-14892.
- Ahmadjian V (1959) Experimental observations on the algal genus *Trebouxia* de Puymaly. *Svensk Botanisk Tidskrift*, **53**, 71-80.
- Ahmadjian V (1993) The lichen symbiosis. John Wiley, New York, USA.
- Ally D, Ritland K, Otto SP (2008) Can clone size serve as a proxy for clone age? An exploration using microsatellite divergence in *Populus tremuloides*. *Molecular Ecology*, **17**, 4897-4911.
- Andras JP, Kirk NL, Harvell CD (2011) Range-wide population genetic structure of *Symbiodinium* associated with the Caribbean Sea fan coral, *Gorgonia ventalina*. *Molecular Ecology*, **20**, 2525-2542.
- Arnaud-Haond S, Belkhir K (2006) GENCLONE: a computer program to analyse genotypic data, test for clonality and describe spatial clonal organization. *Molecular Ecology Notes*, **7**, 15-17.
- Arnaud-Haond S, Duarte CM, Alberto F, Serrao EA (2007) Standardizing methods to address

- 485 clonality in population studies. *Molecular Ecology*, **16**, 5115-5139.
- 486 Beck A, Friedl T, Rambold G (1998) Selectivity of photobiont choice in a defined lichen
487 community: inferences from cultural and molecular studies. *New Phytologist*, **139**, 709-720.
- 488 Beck A, Kasalicky T, Rambold G (2002) Myco-photobiontal selection in a Mediterranean
489 cryptogam community with *Fulgensia fulgida*. *New Phytologist*, **153**, 317-326.
- 490 Blaha J, Baloch E, Grube M (2006) High photobiont diversity associated with the euryoecious
491 lichen-forming ascomycete *Lecanora rupicola* (Lecanoraceae, Ascomycota). *Biological Journal*
492 *of the Linnean Society*, **88**, 283-293.
- 493 Boch S, Prati D, Werth S, Rüetschi J, Fischer M (2011) Lichen endozoochory by snails. *PloS ONE*
494 **6(4)**, e18770. doi:10.1371/journal.pone.0018770.
- 495 Brem D, Leuchtmann A (2003) Molecular evidence for host-adapted races of the fungal endophyte
496 *Epichloe bromicola* after presumed host-shifts. *Evolution*, **57**, 37-51.
- 497 Brown BE, Dunne RP, Goodson MS, Douglas AE (2000) Marine ecology: bleaching patterns in reef
498 corals. *Nature*, **404**, 142-143.
- 499 Bubrick P, Frensdorff A, Galun M (1985) Selectivity in the lichen symbiosis. In: *Lichen Physiology*
500 *and Cell Biology* (ed. Plenum), pp. 319-334. Brown, D.H., New York.
- 501 Butlin R (2002) The costs and benefits of sex: new insights from old asexual lineages. *Nature*, **3**,
502 311-317.
- 503 Cassie D, Piercey-Normore MD (2008) Dispersal in a sterile lichen-forming fungus, *Thamnia*
504 *subuliformis* (Ascomycotina, Icmadophilaceae). *Canadian Journal of Botany*, **86**, 751-762.
- 505 Clay K (1990) Fungal endophytes of grasses. *Annual Review of Ecology and Systematics*, **21**, 275–
506 297.

- 507 Clay K (1988) Fungal endophytes of grasses: a defensive mutualism between plants and fungi.
508 *Ecology*, **69**, 10-16.
- 509 Coates AG, Jackson JBC (1987) Clonal growth, algal symbiosis, and reef formation by corals.
510 *Paleobiology*, **13**, 363-378.
- 511 Dal Grande F (2011) Phylogeny and co-phylogeography of a photobiont-mediated guild in the
512 lichen family Lobariaceae. PhD Thesis, University of Bern, Bern, Switzerland.
- 513 Dal Grande F, Widmer I, Beck A, Scheidegger C (2010) Microsatellite markers for *Dictyochloropsis*
514 *reticulata* (Trebouxiophyceae), the symbiotic alga of the lichen *Lobaria pulmonaria* (L.).
515 *Conservation Genetics*, **11**, 1147-1149.
- 516 Dettki H, Klintberg P, Esseen PA (2000) Are epiphytic lichens in young forests limited by local
517 dispersal? *Ecoscience*, **7**, 317-325.
- 518 Dobson F (2003) Getting a liking for lichens. *Biologist*, **50**, 263-267.
- 519 Doering M, Piercey-Normore MD (2009) Genetically divergent algae shape an epiphytic lichen
520 community on Jack Pine in Manitoba. *Lichenologist*, **41**, 69-80.
- 521 Etges S, Ott S (2001) Lichen mycobionts transplanted into the natural habitat. *Symbiosis*, **30**, 191–
522 206.
- 523 Foucaud J, Jourdan H, Le Breton J, Loiseau A, Konghouleux D, Estoup A (2006) Rare sexual
524 reproduction events in the clonal reproduction system of introduced populations of the little fire
525 ant. *Evolution*, **60**, 1646–1657.
- 526 Friedl T (1987) Thallus development and phycobionts of the parasitic lichen *Diploschistes*
527 *muscorum*. *Lichenologist*, **19**, 183–191.
- 528 Friedl T, Büdel B (2008) Photobionts. In: Lichen Biology. 2nd ed. (ed. Nash TH III), pp. 9-26.

- Cambridge University Press, Cambridge UK.
- Gaßmann A, Ott S (2000) Growth strategy and the gradual symbiotic interactions of the lichen *Ochrolechia frigida*. *Plant Biology*, **2**, 368–378.
- Geller JB, Walton ED (2001) Breaking up and getting together: evolution of symbiosis and cloning by fission in sea anemones (genus *Anthopleura*). *Evolution*, **55**, 1781–1794.
- Gilbert OM, Queller DC, Strassmann JE (2009) Discovery of a large clonal patch of a social amoeba: implications for social evolution. *Molecular Ecology*, **18**, 1273–1281.
- Handa S, Ohmura Y, Nakano T, Nakahara-Tsubota M (2007) Airborne green microalgae (Chlorophyta) in snowfall. *Hikobia*, **15**, 109–120. (in Japanese)
- Hawksworth DL, Hill DJ (1984) The lichen-forming fungi. pp. Glasgow & London: Blackie.
- Hedenås H, Blomberg P, Ericson L (2007) Significance of old aspen (*Populus tremula*) trees for the occurrence of lichen photobionts. *Biological Conservation*, **135**, 380–387.
- Heinken T (1999) Dispersal patterns of terricolous lichens by thallus fragments. *Lichenologist*, **31**, 603–612.
- Higgs P, Woodcock G (1995) The accumulation of mutations in asexual populations and the structure of genealogical trees in the presence of selection. *Journal of Mathematical Biology*, **33**, 677–702.
- Hill DJ (2009) Asymmetric co-evolution in the lichen symbiosis caused by a limited capacity for adaptation in the photobiont. *Botanical Review*, **75**, 326–338.
- Honegger R (1998) The lichen symbiosis-what is so spectacular about it? *Lichenologist*, **30**, 193–212.
- Honegger R (2001) The symbiotic phenotype of lichen-forming ascomycetes. In: *The Mycota IX*,

- 551 *Fungal Associations* (ed. Hock B), pp. 165-188. Springer, Berlin.
- 552 Honegger R (2008) Morphogenesis. In: *Lichen Biology*. 2nd ed. (ed. Nash TH III), pp. 69-93.
- 553 Cambridge University Press, Cambridge UK.
- 554 Honegger R, Zippler U (2007) Mating systems in representatives of Parmeliaceae, Ramalinaceae
- 555 and Physciaceae (Lecanoromycetes, lichen-forming ascomycetes). *Mycological Research*, **111**,
- 556 424-432.
- 557 Huigens ME, Luck RF, Klaassen RH, Masas MF, Timmermans MJ, Stouthamer R (2000) Infectious
- 558 parthenogenesis. *Nature*, **405**, 178–179.
- 559 Ingold CT, Hudson HJ (1993) *The Biology of Fungi*. 6th ed. 224 pp. Chapman and Hall, London.
- 560 Lättman H, Lindblom L, Mattsson J-E, Milberg P, Skage M, Ekman S (2009) Estimating the
- 561 dispersal capacity of the rare lichen *Cliostomum corrugatum*. *Biological Conservation*, **142**,
- 562 1870–1878.
- 563 Lindblom L, Ekman S (2005) Molecular evidence supports the distinction between *Xanthoria*
- 564 *parietina* and *X. aureola* (Teloschistaceae, lichenized Ascomycota). *Mycological Research*, **109**,
- 565 187-199.
- 566 Lindblom L, Ekman S (2007) New evidence corroborates population differentiation in *Xanthoria*
- 567 *parietina*. *Lichenologist*, **39**, 259-271.
- 568 Loh WKW, Loi T, Carter D, Hoegh-Guldberg O (2001) Genetic variability of the symbiotic
- 569 dinoflagellates from the wide ranging coral species *Seriatopora hystrix* and *Acropora*
- 570 *longicyathus* in the Indo-West Pacific. *Marine Ecology Progress Series*, **222**, 97-107.
- 571 Lohtander K, Oksanen I, Rikkinen J (2003) Genetic diversity of green algal and cyanobacterial
- 572 photobionts in *Nephroma* (Peltigerales). *Lichenologist*, **35**, 325-339.

- 573 Lücking R, Grube M (2002) Facultative parasitism and reproductive strategies in *Chroodiscus*
574 (Ascomycota, Ostropales). *Stapfia*, **80**, 267-292.
- 575 Macedo MF, Miller AZ, Dionisio A, Saiz-Jimenez C (2009) Biodiversity of cyanobacteria and green
576 algae on monuments in the Mediterranean basin. *Microbiology*, **155**, 3476-3490.
- 577 Margulis L (1993) Symbiosis in cell evolution: microbial communities in the Archean and
578 Proterozoic eons. 2nd ed. 452 pp. Freeman WH and Co, New York.
- 579 Maynard Smith J (1986) Evolution: contemplating life without sex. *Nature*, **324**, 300-301.
- 580 Meier FA, Scherrer S, Honegger R (2002) Faecal pellets of lichenivorous mites contain viable cells
581 of the lichen-forming ascomycete *Xanthoria parietina* and its green algal photobiont, *Trebouxia*
582 *arboricola*. *Biological Journal of the Linnean Society*, **76**, 259-268.
- 583 Mikheyev AS, Mueller UG, Boomsma JJ (2007) Population genetic signatures of diffuse co-
584 evolution between leaf-cutting ants and their cultivar fungi. *Molecular Ecology*, **16**, 209-216.
- 585 Mock KE, Rowe CA, Hooten MB, Dewoody J, Hipkins VD (2008) Clonal dynamics in western
586 North American aspen (*Populus tremuloides*). *Molecular Ecology*, **17**, 4827-4844.
- 587 Moran N, Baumann P (1994) Phylogenetics of cytoplasmically inherited microorganisms of
588 arthropods. *Trends in Ecology and Evolution*, **9**, 15-20.
- 589 Mukhtar A, Garty J, Galun M (1994) Does the lichen alga *Trebouxia* occur free-living in nature:
590 further immunological evidence. *Symbiosis*, **17**, 247-253.
- 591 Nash TH III (1996) Nitrogen, its metabolism and potential contribution to ecosystems. In: Nash TH
592 III (ed) *Lichen biology*, pp 121-135. Cambridge University Press, Cambridge UK.
- 593 Nelsen MP, Gargas A (2008) Dissociation and horizontal transmission of codispersing lichen
594 symbionts in the genus *Lepraria* (Lecanorales: Stereocaulaceae). *New Phytologist*, **177**, 264-275.

- 595 Nelsen MP, Gargas A (2009) Symbiont flexibility in *Thamnolia vermicularis* (Pertusariales:
596 Icmadophilaceae). *Bryologist*, **112**, 404-417.
- 597 Ohmura Y, Kawachi M, Kasai F, Watanabe MM, Takeshita S (2006) Genetic combinations of
598 symbionts in a vegetatively reproducing lichen, *Parmotrema tinctorum*, based on ITS rDNA
599 sequences. *Bryologist*, **109**, 43-59.
- 600 Ott S (1987a) Sexual reproduction and developmental adaptations in *Xanthoria parietina*. *Nordic*
601 *Journal of Botany*, **7**, 219-228.
- 602 Ott S (1987b) Reproductive strategies in lichens. *Bibliotheca Lichenologica*, **25**, 81-93.
- 603 Ott S, Meier T, Jahns HM (1995) Development, regeneration, and parasitic interactions between the
604 lichens *Fulgensia bracteata* and *Toninia caeruleonigricans*. *Canadian Journal of Botany*, **73**,
605 S595-S602.
- 606 Parks JC, Werth CR (1993) A study of spatial features of clones in a population of bracken fern
607 *Pteridium aquilinum* (L) Kuhn. *American Journal of Botany*, **80**, 537-544.
- 608 Paulsrud P, Rikkinen J, Lindblad P (1998) Cyanobiont specificity in some *Nostoc*-containing lichens
609 and in a *Peltigera aphthosa* photosymbiodeme. *New Phytologist*, **139**, 517-524.
- 610 Piercey-Normore MD (2006) The lichen-forming ascomycete *Evernia mesomorpha* associates with
611 multiple genotypes of *Trebouxia jamesii*. *New Phytologist*, **169**, 331-344.
- 612 Piercey-Normore MD and DePriest PT (2001) Algal switching among lichen symbioses. *American*
613 *Journal of Botany*, **88**, 1490-1498.
- 614 Printzen C, Ekman S (2003) Local population subdivision in the lichen *Cladonia subcervicornis* as
615 revealed by mitochondrial cytochrome oxidase subunit 1 intron sequences. *Mycologia*, **95**, 399-
616 406.

- 617 Printzen C, Ekman S, Tønsberg T (2003) Phylogeography of *Cavernularia hultenii*: Evidence of
618 slow genetic drift in a widely disjunct lichen. *Molecular Ecology*, **12**, 1473-1486.
- 619 R Development Core Team (2008) R: A language and environment for statistical computing. R
620 Foundation for statistical computing, Vienna, Austria. <http://www.R-project.org>.
- 621 Rambold G, Triebel D (1992) The inter-Lecanoralean associations. *Bibliotheca Lichenologica*, **48**,
622 1-201.
- 623 Reeve HK, Hölldobler B (2007) The emergence of a superorganism through intergroup competition.
624 *Proceedings of the National Academy of Sciences USA*, **104**, 9736-9740.
- 625 Rice WR, Chippindale AK (2001) Sexual recombination and the power of natural selection. *Science*,
626 **294**, 555-559.
- 627 Richardson DHS (1999) War in the world of lichens: parasitism and symbiosis as exemplified by
628 lichens and lichenicolous fungi. *Mycological Research*, **103**, 641-650.
- 629 Rikkinen J (2003) Ecological and evolutionary role of photobiont-mediated guilds in lichens.
630 *Symbiosis*, **34**, 99-110.
- 631 Rikkinen J, Oksanen I, Lohtander K (2002) Lichen guilds share related cyanobacterial symbionts.
632 *Science*, **297**, 357-357.
- 633 Rowan R (1998) Diversity and ecology of Zooxanthellae on coral reefs. *Journal of Phycology*, **34**,
634 407-417.
- 635 Saffo MB (1992) Invertebrates in endosymbiotic associations. *Annals of Zoology*, **32**, 557-565.
- 636 Saikkonen K, Ion D, Gyllenberg M (2002) The persistence of vertically transmitted fungi in grass
637 metapopulations. *Proceedings of the Royal Society of London, Series B*, **269**, 1397-1403.
- 638 Samadi S, Mavarez J, Pointier JP, Delay B, Jarne P (1999) Microsatellite and morphological

- 639 analysis of population structure in the parthenogenetic freshwater snail *Melanoides tuberculata*:
640 insights into the creation of clonal variability. *Molecular Ecology*, **8**, 1141-1153.
- 641 Sanders WB (2005) Observing microscopic phases of lichen life cycles on transparent substrata
642 placed in situ. *Lichenologist*, **37**, 373-382.
- 643 Sanders WB, Lücking R (2002) Reproductive strategies, relichenization and thallus development
644 observed in situ in leaf-dwelling lichen communities. *New Phytologist*, **155**, 425-435.
- 645 Scheidegger C (1985) Systematische Studien zur Krustenflechte *Anzina carneonivea* (Trapeliaceae,
646 Lecanorales). *Nova Hedwigia* **41**, 191-218.
- 647 Scheidegger C, Werth S (2009) Conservation strategies for lichens: insights from population
648 biology. *Fungal Biology Reviews*, **23**, 55-66.
- 649 Seabloom EW, Bjørnstad ON, Bolker BM, Reichman OJ (2005) The spatial signature of
650 environmental heterogeneity, dispersal, and competition in successional grasslands.
651 *Ecological Monographs*, **75**, 199-214.
- 652 Selkoe KA, Toonen RJ (2006) Microsatellites for ecologists: a practical guide to using and
653 evaluating microsatellite markers. *Ecology Letters*, **9**, 615-629.
- 654 Seymour FA, Crittenden PD, Dyer PS (2005) Sex in the extremes: lichen-forming fungi. *Mycologist*,
655 **19**, 51-58.
- 656 Sillett SC, McCune B, Peck JE, Rambo TR, Ruchty A (2000) Dispersal limitations of epiphytic
657 lichens result in species dependent on old-growth forests. *Ecological Applications*, **10**, 789-799.
- 658 Sirviö A, Pamilo P (2010) Multiple endosymbionts in populations of the ant *Formica cinerea*. *BMC*
659 *Evolutionary Biology*, **10**, 335.
- 660 Skaloud P (2008) Polyphasic approaches in the taxonomy of green aerophytic algae. PhD Thesis,

- 661 University of Prague. Prague, Czech Republic.
- 662 Slocum RD, Ahmadjian V, Hildreth KC (1980) Zoosporogenesis in *Trebouxia gelatinosa*:
663 ultrastructure potential for zoospore release and implications for the lichen association.
664 *Lichenologist*, **12**, 173-187.
- 665 Sluiman HJ, Kouwets FAC, Blommers PCJ (1989) Classification and definition of cytokinetic
666 patterns in green algae: sporulation versus (vegetative) cell division. *Arch. Protistenk*, **137**, 277-
667 90.
- 668 Stenroos S (1990) *Cladonia luteoalba* - an enigmatic *Cladonia*. *Karstenia*, **30**, 27-32.
- 669 Tschermak-Woess E (1978) Über die Phycobionten der Sektion *Cystophora* von *Chaenotheca*,
670 insbesondere *Dictyochloropsis splendida* und *Trebouxia simplex*, spec. nova. *Plant Systematics*
671 *and Evolution*, **129**, 185-208.
- 672 Tomiuk J, Guldbrandtsen B, Loeschcke V (1998) Population differentiation through mutation and
673 drift - a comparison of genetic identity markers. *Genetica*, **102/103**, 545-558.
- 674 Vogler AJ, Keys C, Nemoto Y, Colman RE, Jay Z, Keim P (2006) Effect of repeat copy number on
675 variable-number tandem repeat mutations in *Escherichia coli* O157:H7. *Journal of Bacteriology*,
676 **188**, 4253-4263.
- 677 Wagner HH, Fortin MJ (2005) Spatial analysis of landscapes: concepts and statistics. *Ecology* **86**,
678 1975-1987.
- 679 Wagner HH, Holderegger R, Werth S, Gugerli F, Hoebee SE, Scheidegger C (2005) Variogram
680 analysis of the spatial genetic structure of continuous populations using multilocus microsatellite
681 data. *Genetics*, **169**, 1739-1752.
- 682 Wagner HH, Werth S, Kalwij JM, Bolli JC, Scheidegger C (2006) Modelling forest recolonization

by an epiphytic lichen using a landscape genetic approach. *Landscape Ecology*, **21**, 849-865.

Walser JC, Sperisen C, Soliva M, Scheidegger C (2003) Fungus-specific microsatellite primers of lichens: application for the assessment of genetic variation on different spatial scales in *Lobaria pulmonaria*. *Fungal Genetics and Biology*, **40**, 72-82.

Walser, JC (2004) Molecular evidence for limited dispersal of vegetative propagules in the epiphytic lichen *Lobaria pulmonaria*. *American Journal of Botany*, **91**, 1273-1276.

Werth S (2010) Optimal sample sizes and allelic diversity in studies of the genetic variability of mycobiont and photobiont populations. *Lichenologist*, **43**, 73-81.

Werth S, Gugerli F, Holderegger R, Wagner HH, Csencsics D, Scheidegger C (2007) Landscape-level gene flow in *Lobaria pulmonaria*, an epiphytic lichen. *Molecular Ecology*, **16**, 2807-2815.

Werth S, Sork VL (2010) Identity and genetic structure of the photobiont of the epiphytic lichen *Ramalina menziesii* on three oak species in southern California. *American Journal of Botany*, **97**, 821-830.

Werth S, Wagner HH, Gugerli F, Holderegger R, Csencsics D, Kalwij JM, Scheidegger C. (2006)a. Quantifying dispersal and establishment limitation in a population of an epiphytic lichen. *Ecology*, **87**, 2037-2046.

Werth S, Wagner HH, Holderegger R, Kalwij JM, Scheidegger C. (2006)b. Effect of disturbances on the genetic diversity of an old-forest associated lichen. *Molecular Ecology*, **15**, 911-921.

Widmer I, Dal Grande F, Cornejo C, Scheidegger C (2010) Highly variable microsatellite markers for the fungal and algal symbionts of the lichen *Lobaria pulmonaria* and challenges in developing biont-specific molecular markers for fungal associations. *Fungal Biology* **114**, 538-544.

Wornik S, Grube M (2010) Joint dispersal does not imply maintenance of partnerships in lichen

symbioses. *Microbial Ecology*, **59**, 150-157.

Yahr R, Vilgalys R, DePriest PT (2004) Strong fungal specificity and selectivity for algal symbionts in Florida scrub *Cladonia* lichens. *Molecular Ecology*, **13**, 3367-3378.

Yahr R, Vilgalys R, DePriest PT (2006) Geographic variation in algal partners of *Cladonia subtenuis* (Cladoniaceae) highlights the dynamic nature of a lichen symbiosis. *New Phytologist*, **172**, 377-391.

Yoshimura, I (1971) The genus *Lobaria* of Eastern Asia. *Journal of the Hattori Botanical Laboratory*, **34**, 231-331.

Zilber-Rosenberg I, Rosenberg E (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiology Reviews*, **32**, 723-735.

Zoller S, Lutzoni F, Scheidegger C (1999) Genetic variation within and among populations of the threatened lichen *Lobaria pulmonaria* in Switzerland and implications for its conservation. *Molecular Ecology*, **8**, 2049-2059.

Figure legends

Fig. 1 Graphic representation of the pairwise analysis used in this study to infer the mode of photobiont transmission and to analyze within-population micro-evolutionary processes of the fungal and algal symbionts of *L. pulmonaria*. Each circle represents a single thallus of *L. pulmonaria*. Each slice of the circle represents a microsatellite locus (eight for the fungal symbiont, seven for the algal symbiont). Black (for fungal loci) or dark grey (for algal loci) slices represent differences at the given microsatellite locus between Lichen A and Lichen B (top) or C (bottom). The number of loci differing between a pair of thalli is defined for each symbiont as *deltaA* (alga) or

deltaF (fungus).

Fig. 2 Models used to analyze the contribution of mutation and recombination to within-population genetic structure of lichen symbionts in *L. pulmonaria*. Each barplot shows the relative frequency of pairs of thalli differing in *deltaA* (alga, left) or *deltaF* (fungus, right) loci, as expected for each symbiont under the empirical null model of recombination (top) or an exponential model of somatic mutation (center). The bottom barplots show the observed relative frequency of pairs in data set C. Empty circles indicate for each level of *deltaA* or *deltaF* the relative frequency predicted by the fitted model combining the empirical null model of recombination and the exponential model of mutation.

Fig. 3 Barplot of the relative frequency ($\pm$ bootstrap standard errors) of pairs of thalli (within populations pooled over all populations) differing by 0-7 loci for the alga (*deltaA*) and 0-8 loci for the fungus (*deltaF*) for all pairwise comparisons within each population and pooled over all populations (data set A, top) and for pairwise comparisons in one symbiont constrained to identical pairs of multilocus genotypes for the other symbiont (data set B, bottom).

Fig. 4 Spatial distribution of the following categories of thalli: vegetative propagules ($\delta F = \delta A = 0$; black bars, "00"), fungal clones associated with different algal multilocus genotypes ($\delta F = 0$ and $\delta A > 0$; white bars, "01"), algal clones associated with different fungal multilocus genotypes ($\delta F > 0$ and $\delta A = 0$; dark grey, "10"), and different fungal multilocus genotypes associated with different algal multilocus genotypes ($\delta F > 0$ and $\delta A > 0$; light grey,

"11"). The first distance class contained pairs of thalli sampled from the same tree. The last distance class contained all sample comparisons at distances >500 m.

Fig. 5 Relative frequency of distance classes for each group of multilocus genotype pairs defined in Fig. 2. Distance classes: 0-10m (first bar from bottom; darkest bars); 10-20m (second bar from bottom); 20-50m (third bar from bottom); 50-100m (fourth bar from bottom); 100-200m (fifth bar from bottom), >500m (last bar from bottom; faintest bars). The first distance class of pairs of thalli sampled from the same tree was not included in the analysis. The total number of multilocus genotype pairs per group is given on top of the corresponding bar.

Supporting Information

Table S1 Information on sampled populations of *Lobaria pulmonaria*.

Table S2 Microsatellite analysis: (a) primer sequences (Walser *et al.* 2003, 2004; this study; Dal Grande *et al.* 2010), labeling, primer concentrations and (b, c) PCR conditions for genetic analyses of *Lobaria pulmonaria*.

Table S3 Model fitting and residuals of observed frequencies of deltaF vs. fitted frequencies (combined exponential and binomial fitting).

Table S4 Allele frequency distribution at eight fungal (LPu03, LPu09, LPu15, LPu23, LPu24, LPu25, LPu28, Walser *et al.* 2003; Widmer *et al.* 2010; MS4, this study) and seven algal (LPh1 to LPh7; Dal Grande *et al.* 2010) loci per population. Each line is one repeat length (allele) and each number represents the absolute frequency of that allele in the particular population.

Author Information Box

This work was part of Francesco Dal Grande and Ivo Widmer's PhD research on the evolutionary history and co-phylogeography of a lichen symbiosis. Helene H. Wagner focuses on spatial analysis and modelling of dispersal and inter-specific interactions applied in meta-community dynamics and landscape genetics. Christoph Scheidegger's research interests cover the biodiversity evaluation, population genetics and conservation biology of lichens and plants.

Table 1 Data sets used in this study.

Data set	Definition	Number of Pairs	Analysis
A	All pairs from within the same population (same for both symbionts)	All: 36,218 pairs from 62 populations	Quantification of intra-population genetic diversity and spatial genetic structure.
B	Fungus: all pairs of data set A with $\Delta A=0$ Alga: all pairs of data set A with $\Delta F=0$	Fungus: 2977 pairs from 62 populations Alga: 3285 pairs from 62 populations	Restriction for each symbiont to pairs with identical MLG in the other symbiont to partial out larger-scale evolutionary processes when assessing the relative contribution of vertical transmission to population genetic structure.
C	Data set B without recurrent MLGs within the same population	Fungus: 269 pairs from 38 populations Alga: 215 pairs from 50 populations	Exclusion of recurrent MLGs within populations to identify signals of within-population mutation and recombination and to assess the relative contribution of horizontal transmission. This avoids potential underestimation of horizontal transmission in genetically uniform or depauperate populations.

Empirical null model of recombination

Exponential model of somatic mutation

Observed distribution (data set C) and fitted combined model

All pairs of thalli (data set A)

Pairs of thalli with identical MLG in other symbiont (data set B)