Impacts of eutrophication and oil spills on the Gulf of Finland herring stock

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0108.R3
Manuscript Type:	Article
Date Submitted by the Author:	30-Nov-2016
Complete List of Authors:	Rahikainen, Mika; University of Helsinki, Department of Environmental Sciences Hoviniemi, Kirsi-Maaria; Oulun Yliopisto Teknillinen Tiedekunta, Water Resources and Environmental Engineering Laboratory Mäntyniemi, Samu; University of Helsinki, Department of Environmental Sciences Vanhatalo, Jarno ; University of Helsinki, Department of Mathematics and Statistics and Department of Biosciences Helle, Inari; University of Helsinki Lehtiniemi, Maiju; Finnish Environment Institute (SYKE), Pönni, Jukka; Luonnonvarakeskus Kuikka, Sakari; University of Helsinki
Keyword:	EUTROPHICATION < General, oil spill, RECRUITMENT < General, herring, BAYESIAN STATISTICS < General

SCHOLARONE™
Manuscripts

1   Impacts of eutrophication and oil spills 2   on the Gulf of Finland herring stock

---

3   Mika Rahikainen^{1,*}, Kirsi-Maaria Hoviniemi², Samu Mäntyniemi¹, Jarno Vanhatalo³, Inari Helle¹,
4   Maiju Lehtiniemi⁴, Jukka Pönni⁵, Sakari Kuikka¹

5   ¹University of Helsinki, Department of Environmental Sciences, Fisheries and Environmental
6   Management Unit, Viikinkaari 2a, 00014 University of Helsinki, Finland

7   ²University of Oulu, Faculty of Technology, Water Resources and Environmental Engineering
8   Laboratory, P.O. Box 4300, 90014 University of Oulu, Finland

9   ³University of Helsinki, Department of Mathematics and Statistics and Department of Biosciences,
10   Viikinkaari 2a, 00014 University of Helsinki, Finland

11   ⁴Finnish Environment Institute, Marine Research Centre, Mechelininkatu 34a, 00260 Helsinki,
12   Finland

13   ⁵Natural Resources Institute Finland, Viikinkaari 4, 00790 Helsinki, Finland

14   * email mika.rahikainen@helsinki.fi; tel. +358 294140804

15

Abstract

The Baltic Sea is one of the world's most stressed sea areas. Major threats to the ecosystem include eutrophication and oil spills. The progression of anthropogenic nutrient enrichment is lengthy and gradual while oil spills cause rapid changes in the system, with varying impact time. We quantify the impact of eutrophication and the key ecological covariates on the population dynamics of the major pelagic fish stock, the Baltic herring, in the Gulf of Finland. The full life-cycle of herring is represented with a probabilistic state-space model. Moreover, we analyse the impact of the oil spill from M/T Antonio Gramsci, in 1987, on herring survival. The results confirm impact of the spill on the early life-stage survival: the observed high frequency of malformed herring larvae in surveys signaled elevated mortality of the year-class. The optimal July-August chlorophyll α concentration for herring reproduction is approximately 5 $\mu\text{g/l}$. This level is currently exceeded suggesting recruitment impairment due to eutrophication. The herring stock was also recruitment overfished. Analysis suggests deceleration of herring growth as salinity descends below 6 psu.

Keywords: eutrophication, oil spill, recruitment, mortality, herring, Baltic Sea, Bayesian modeling

Running title: Impacts of eutrophication and oil exposure on herring

33 **1. Introduction**

34

35 Aquatic ecosystems are affected by multiple stressors (Halpern et al. 2008). The global key stressors
36 include fishing and aquaculture (Naylor et al. 2000), nutrient enrichment (Cloern 2001, Smith 2003),
37 a wide range of toxic contaminants including residues of accidental and intentional oil spills (Hassler
38 2011), and climate change (Rabalais et al. 2009). There is abundant qualitative knowledge about
39 how eutrophication impacts ecosystems in general (Grall and Chauvaud 2002) but little quantitative
40 evidence for how it effects the population dynamics of commercially important marine fish stocks. In
41 particular, there is currently a critical knowledge gap on how eutrophication and climate variables
42 separately and interactively impact the dynamics of marine ecosystems (Kotta et al. 2009). Notably,
43 it is not possible to infer impacts of these stressors in isolation because of their interaction,
44 therefore novel modelling techniques are called for.

45 The Baltic Sea, one of the largest brackish water areas in the world, provides a data-rich area
46 influenced by several stressors, one of the most dominant being eutrophication (HELCOM 2010).
47 Since the mid-1900s, the Baltic Sea has changed from an oligotrophic clear-water system into a
48 eutrophic environment (Fleming-Lehtinen and Laamanen 2012). Earlier literature suggests that a
49 moderate increase in nutrient input enhances primary and secondary production, including fish,
50 whereas further nutrient input will reduce fish production and inflict a change in the fish taxa (Colby
51 et al. 1972, Hartmann and Nümann 1977, Persson et al. 1991, Caddy 1993). The Gulf of Finland (GoF)
52 is one of the most stressed sea areas of the Baltic Sea (Andersen et al. 2011, Korpinen et al. 2012),
53 and on the global scale (Jackson 2001, Carstensen et al. 2014). Hence, we anticipate the current level
54 of eutrophication may have exceeded the initial positive effect on fish production of commercially
55 harvested species.

56 The Baltic Sea is also one of the most intensively trafficked areas in the world. Both the number and
57 size of the ships, especially oil tankers, have been growing during the last years, and this trend is

58 expected to continue (Brunila and Storgård 2012). A major oil spill could have wide-spread and long-
59 lasting impacts on the ecosystem, as exposure to oil can lead to the immediate death of organisms,
60 or decrease their fitness via various sub-lethal effects (NRC 2003). In winter 1987, the tanker Antonio
61 Gramsci ran aground off the Finnish coast, spilling 570-650 tonnes of crude oil. In spring 1987,
62 floating oil and oiled shores were observed between Helsinki and Kotka (Hirvi 1990a, 1990b; Fig. 1).
63 Although abnormally formed herring (*Clupea harengus membras*) larvae were abundant near the
64 grounding site after the oil spill (Urho 1991), understanding of the influence on the herring
65 population has been absent so far.

66 The Baltic Sea pelagic fish biomass is dominated by herring and sprat (*Sprattus sprattus*). Herring is
67 one of the key species due to its high abundance and role as a consumer in the pelagic food web
68 (Flinkman et al. 1998, Kornilovs et al. 2001), and as forage for cod (*Gadus morhua*; Sparholt 1994).
69 Sprat competes for food with herring (Lindegren et al. 2011), and herring growth is considerably
70 lower at high sprat densities than at low sprat levels (Rönkkönen et al. 2004, Casini et al. 2010).
71 Herring and sprat overlap in the GoF and they are fished as a mixture. Sprat has been the choke
72 species for the Finnish herring fishery in many years.

73 Baltic herring spawn in coastal areas typically in May-June at 8–12 °C on hard bottom vegetation
74 avoiding sites covered by soft sediments. Their spawning beds are restricted in number. They usually
75 reach to about 8 m in depth, depending on the water quality (Aneer 1989, Rajasilta et al. 1989; 1993,
76 Kääriä et al. 1997). The water temperature also has a major effect on the fundamental biotic
77 processes of fish (Pauly 1980, Pepin 1991). The critical period hypothesis states that the bottleneck
78 of reproduction is in the larval phase, and the transition phase of yolk-sack larvae to external feeding
79 is of special importance. The temperature fluctuates markedly from year to year in the Baltic Sea,
80 potentially masking the influence of other factors on these processes. These fluctuations obviously
81 call for considering the impact of temperature on the survival, recruitment and growth of fish.

82 Many Baltic Sea species live at the margin of their salinity tolerance and are sensitive to changes in
83 salinity. The Baltic Sea zooplankton assemblage is a mixture of marine, brackish and freshwater
84 species. Salinity exhibits temporal fluctuations, which in turn influence zooplankton assemblage
85 (Viitasalo et al. 1995). Abundance of neritic copepods correlates positively with salinity, while that of
86 freshwater cladocerans is negative (Vuorinen et al. 1998). Neritic copepods have a larger body size
87 and higher energy content than cladocerans, which strongly influences the feeding success and
88 growth of herring (Flinkman et al. 1998).

89 Rönnerberg and Bonsdorff (2004) address that the Baltic Sea is not an uniform water mass and,
90 therefore, regional ecological assessments in relation to basin-wide eutrophication are required. We
91 use a probabilistic state-space population model to quantify the impact of eutrophication, sea
92 surface temperature (SST), salinity, and abundance of sprat and cod stock on GoF herring stock
93 dynamics. The model is a further development of Mäntyniemi et al. (2013) to account for the
94 interactions of model parameters with environment, competition and predation. With an additional
95 Gaussian Process models we evaluate uncertainties of most of the covariates and use them in the
96 population model. We also analyse whether herring stock data provides detectable signals about the
97 impacts of a historical oil spill. We apply Bayesian theory to fuse the rich information content in the
98 data with prior knowledge on herring stock biology (Kuparinen et al. 2012). The informative prior
99 denotes knowledge in addition to the data and model structure (Romakkaniemi (Ed) 2015). This
100 allows us to separate the effects of otherwise confounding sources.

101 **2. Material and methods**

102

2.1 Data

Fishery data

GoF herring are fished by the Finnish, Estonian and Russian fleets. We used fisheries data from these countries compiled by the ICES Baltic Fisheries Assessment Working Group (WGBFAS) for 1980-2006.

The data set consisted of age-specific commercial catches, mean weight-at-age, and maturation. Weight-at-age was computed from length-based observations which originate from random sampling for 1980-1997 and a quarterly disaggregated age-length-key sampling strategy from 1998 onwards (J. Pönni, unpubl.). Annual length-weight relationships were computed but we did not account for uncertainty in them. However, we accounted for the observation error of the total catches by treating the historical catch data as uncertain observations from the true catches.

Although there is evidence that the GoF herring constitutes a separate stock it is since 1991 assessed in aggregation with the Central Baltic herring (ICES 2007). Nevertheless, herring exhibits homing behaviour and there are clear genetic differences between areas (Jørgensen et al. 2005) and between ecotypes in the same area (Bekkevold et al. 2016) supporting reproductive isolation in the Baltic Sea. We use the GoF herring data solely and assume it as a unit stock. This improves the chance to detect a signal in the recruitment compared to an analysis for the entire Central Baltic herring stock. Also WGBFAS has assessed the GoF herring stock as a discrete component until 1990. Even later there have been attempts to assess smaller areas (ICES 2001).

Juntunen et al. (2012) estimated the herring biomass in the GoF in 2004 based on an acoustic survey. According to their results the 95% credible interval of the biomass was [2000, 103000] tonnes, the posterior mean 24611 tonnes and variance 7.09×10^7 . The posterior distribution from their study was used to construct a log-Gaussian approximation for the likelihood function with respect to the total biomass in 2004 in the population dynamics model (see Kuikka et al., 2014 for similar treatment of results from earlier studies). Here, the posterior mean and variance of Juntunen et al. (2012) are

analogous to the observed data value and observation error variance, respectively, in traditional likelihoods.

Annual point estimates of cod catches and sprat abundance were adopted from ICES working group reports (ICES 2008, 2009). For sprat, we used the annual sprat biomass in the entire Baltic Sea (ICES 2009) to describe influence on herring in the GoF. The Baltic cod stock has experienced marked fluctuations in abundance and distribution during the recent decades (ICES 2009) and, the total Baltic cod biomass poorly reflects its prevalence in the GoF. Hence, we apply the annual total catches of cod in the GoF (ICES, 2008 Table 2.1.2) as a proxy for cod abundance. There are no national differences or changes in management of the cod fishery which could cause the observed temporal landing pattern in the GoF. This fact supports the use of catch data.

Environmental data

We linked three environmental covariates to herring demography: salinity, SST and chlorophyll- α concentration. The chlorophyll- α was used to index the eutrophication. We used the COMBINE¹ archive to retrieve the salinity and chlorophyll data. Swedish Meteorological and Hydrological Institute (SMHI) provided the temperature data.

The surface salinity has been measured once in a year from the 0-1 metre depth from two regular sampling sites in 1980-2006 (Fig. 1). The data comprised one annual average from these sites. The SST (depth 0-1 m) data was a time series of daily measurements from two stationary sampling sites. The data included 7642 temperature measurements from 12.10.1974 until 16.5.2007. There were several missing measurements in the time series. The data on chlorophyll- α concentration comprised measurements on the Finnish coast of the GoF from 1974 to 2010. The number and the locations of measurements varied by year (Fig. 1 lower panel).

¹ http://www.helcom.fi/groups/monas/CombineManual/PartA/en_GB/main/

2.2 Estimating the environmental explanatory variables

The fundamental problem in many environmental data sets is the fact that sampling sites vary by month and year, or temporal sampling frequency is inconsistent over the years at fixed sampling sites. Therefore, comparison among years is misleading unless the random deviations in the sampling programme are accounted for. A valid modelling approach is necessary to interpolate the environmental covariates in areas and times with no observations and smooth out the effects of missing data points.

Our general approach is summarised below and treated more extensively in the Appendix. The observation, $y(t, s)$, SST e.g., at time t and location s , was assumed to be a noisy realisation of an underlying true value of an environmental covariate $f(t, s)$ so that

$$y(t, s) = f(t, s) + \varepsilon_i \quad (1)$$

where $\varepsilon_i \sim N(0, \sigma^2)$. We gave $f(t, s)$ a Gaussian process (GP) prior (Rasmussen and Williams, 2006, Gelfand et al. 2010) and solved its posterior distribution by conditioning on the measurement data (see Fig. 2 and Appendix). GP is a stochastic process which defines probability distribution over functions. A GP prior is defined by a mean and a covariance function which describe the properties, such as the smoothness and magnitude, of the spatio-temporal variation in the environmental variable. The covariance describes the spatio-temporal association between observations and can be represented as a function that describes the correlation between pairs of points with distance in time and space.

We indexed the eutrophication by the July-August average chlorophyll- α concentration on the Finnish coast of the GoF, where herring spawning and nursery areas were expected to be located (Urho and Hildén 1990). In practice, we calculated the posterior distribution for the July-August average chlorophyll- α concentration for the sea area within 10 km off the Finnish shoreline (Fig. 1).

Then we calculated the posterior distribution for the average concentration over this zone for each year from 1974 to 2010. Similarly, we calculated the posterior distribution for the average SST at the two sampling sites for each day from 1980 to 2006, after which the SST indices were formed by calculating the posterior distribution for the average SST in April-June (spawning season) and August-October (growth season) each year. The salinity index was the posterior distribution for the average salinity at the two measurement stations at the measurement day. We, thus, moved beyond using point estimates as explanatory variables and admit uncertainty about the environmental variables by using their posterior distributions in the population dynamics model (Fig. 2).

2.3 Herring stock dynamic model

We built a hierarchical Bayesian model combining the age- and length-structure of the GoF herring population. The core model is described in detail by Mäntyniemi et al. (2013) and in this section we summarise its essential components. The model covers the full life-cycle and also features size-dependency in parameters such as natural mortality, fecundity and fishery selection. This improves the realism of the model, and also allows explicit examination of the population dynamics.

The fishing mortality of each age group each year was assumed to depend on the availability of fish in the main fishing areas, the selectivity of the gears, and fishing effort. Stock abundance and structure were estimated based on the catch-at-age and weight-at-age data and annual growth and mortality rates. The abundance of the year class 1 was estimated based on the stock-recruitment model. Individual growth followed the von Bertalanffy curve, and was modelled as mean length-at-age for each year separately. Priors for the growth curve were derived from FishBase².

The fecundity of fish depends on the age and size structure of the population, and on the quality of the parents (Marshall et al. 2006, Lambert 2008). For example, major changes in average body size

---

² <http://www.fishbase.org/>

have been observed in the GoF herring stock (Rönkkönen et al. 2004), and the spawning stock biomass might give biased information for the real spawning potential. Therefore, the stock-recruitment function in the model was defined on the basis of spawned eggs, allowing the explicit consideration of changes in the demography of the stock (Mäntyniemi et al. 2013). The fecundity of female herring was assumed to increase as a function of weight. The functional form for eggs produced per unit mass of female, and priors for its parameters, were based on empirical studies conducted in the northern coast of the GoF (Parmanne and Kuittinen 1991). Recruitment was assumed to follow either the Beverton-Holt or Ricker stock-recruitment curve (Quinn and Deriso 1999).

We expanded Mäntyniemi et al.'s (2013) model to include information about chlorophyll- α , SST, salinity and abundance of sprat and cod stock as explanatory variables to account for their impact on herring recruitment, growth rate and natural mortality rate (Fig. 3). Moreover, we included the possible additional mortality caused by the Antonio Gramsci oil spill in 1987. In the following sections we summarise the changes compared to Mäntyniemi et al.'s (2013) model.

Impact of spilled oil on herring mortality

We modeled the impact of the tanker Antonio Gramsci oil spill on the GoF herring stock as additional mortality during the year of accident. Mortality was modelled to impact offspring and the adult component at different ratios. The former refers 1987 year-class and the latter to 1986 and older. Oil-induced immediate mortality was assumed to influence the spawning stock size and, hence, the total amount of eggs spawned (S) in 1987 was

$$S_{1987} = N_{1987} \times (1 - OilmorA) \times r \times \sum_{A=1}^8 e_A \quad (2)$$

where N_{1987} is the number of herring in the population during the spawning season 1987 without oil spill-induced mortality, $OilMorA$ is the oil induced mortality in ages 1 and older, r is the sex ratio (1:1 at all ages), e_A is the age-structured and length-based fecundity in the population. Recruitment was a stochastic variable (Mäntyniemi et al., 2013) and depended on the number of spawned eggs, the

underlying stock-recruitment relationship and oil-induced mortality (*OilMorE*) among the offspring. The expected recruitment in 1987, that is the number of age 1 herring at the beginning of 1988, is then

$$E[R_{1987}] = f(S) \times (1 - OilMorE) \tag{3}$$

where $f(S)$ denotes either the Beverton-Holt or Ricker stock-recruitment relationship, as parameterised in Mäntyniemi et al. (2013). The prior odds for Beverton-Holt and Ricker stock-recruitment relationships was one. We calculated the posterior probabilities for these alternative hypotheses and marginalized over them using Bayesian model averaging (see section 2.4).

Priors for the oil induced additional mortalities (Table 1) were adopted from Lecklin et al. (2011, and unpublished data, T. Lecklin). They conditioned immediate mortality on proportion of oiled coast line, oil type and season of an accident. We applied the mortality estimates developed for pelagic fishes and conditioned them on the actual attributes of the Antonio Gramsci accident (Hirvi 1990a, 1990b). The tanker ran aground in February 1987, spilling oil in the ice-covered GoF. The oil divided into a number of separate patches with limited movement in March. Observation of the drift trajectory was incomplete due to the ice cover. In May, the spilled oil drifted ashore at various places on the northern coast of the GoF, the furthest distance between two separate oiled locations being about 120 km. Applying the discretisation used by Lecklin et al. (2011), this corresponds to 7-15% of the whole GoF coastline.

Similarly to the treatment of the stock-recruitment functions, our model included two hypotheses for oil-induced mortality; either oil induced additional mortality, as we have defined by the above model structure and the priors, or the Antonio Gramsci oil spill did not cause any additional mortality in the GoF herring population. In the latter case $OilMorA=OilmorE=0$. We calculated the posterior distribution for both hypotheses as described in section 2.4.

Herring recruitment

The effect of eutrophication on herring stock dynamics was described by a dome-shaped function yielding maximum recruitment at intermediate chlorophyll- α level. The influence of annual SST in April-June on the year-class abundance was described by the non-linear relationship between recruitment and SST. The expected recruitment $E(R_y)$ was

$$E(R_y) = f(S_{y-1}) \times e^{-\left(\frac{Chl_{y-1} - OptChl}{sd}\right)^2 + b \times (\log(SST_{y-1}^{AJ}) - \overline{\log(SST^{AJ})})} \quad (4)$$

where Chl_y is the chlorophyll- α concentration index, $OptChl$ is a parameter for optimal chlorophyll- α concentration, parameter sd defines the width of the parabola, SST_{y-1}^{AJ} is the April-June SST index in year $y-1$ and $\overline{\log(SST^{AJ})}$ is the average of log April-June SST indices over years 1980-2006. Extracting $\log(SST_{y-1}^{AJ}) - \overline{\log(SST^{AJ})}$ centres the fluctuations in the SST indices to an average value of zero.

The idea of model (4) is that $f(S)$ represents the expected recruitment in a situation where the chlorophyll concentration is at the optimal value and $\log(SST)$ is at the mean of the past time series. Deviation of the chlorophyll concentration from the optimal value reduces the expected recruitment compared to $f(S)$. When $\log(SST)$ is higher than average, the recruitment is scaled up from $f(S)$ and down when $\log(SST)$ is lower than average implying positive relationship between recruitment and temperature (Cardinale et al. 2009).

The applied formulation implies that salinity did not have a direct impact on the recruitment of herring (Fig. 3). Instead, it influenced the growth rate of herring which, in turn, affects natural mortality, which varies by length. Hence, salinity had an indirect impact on survival and population abundance.

The parameters of the dome-shaped function for chlorophyll- α concentration were given reasonably informative priors. The prior mean for optimal chl- α concentration, regarding herring reproduction,

was adopted from the estimated level in the Baltic Sea at the beginning of the 1970s (Fleming-Lehtinen et al. 2008). We assume this concentration was near the optimal chl- α level.

Growth of herring

Growth is typically expressed as the mean somatic growth of individuals in a population, relating size to age i.e. weight-at-age (WAA). We modelled the growth in mean length-at-age with von Bertalanffy curve. The relationship between annual growth and the environmental covariates was modelled with linear model for the expected log von Bertalanffy growth parameter, K_y , so that

$$E(\log(K_y)) = (cLS \times \log(\text{Sprat}_y) + ck \times \log(SST_y^{AO}) + \tau_y) - 2. \quad (5)$$

Here, cLS is the regression weight for the log of the Baltic Sea sprat stock total biomass, Sprat_y , ck is the regression weight for the log of the August-October SST index, SST_y^{AO} , and τ_y is a function describing the influence of salinity on $E(\log(K_y))$ (see equation (6) below). The Baltic Sea sprat stock's total biomass was assumed to have been estimated without error. The annual biomass estimates were retrieved from the ICES (2009) stock assessment report. However, as presented in Section 2.2, we took into account the uncertainty about the SST and salinity covariates. SST in August-October was used, as Parmanne (1990) argues that the majority of the annual growth takes place in late autumn. We gave informative priors for cLS and ck so that prior probability was given for negative relationship between sprat stock biomass and herring growth rate, and positive relationship between SST and growth (Brunel & Dickey-Collas 2010) (Table 2). We described the effect of salinity on herring growth rate with logit function

$$\text{logit}(\tau_y) = \varphi + \omega \times \text{Sal}_y \quad (6)$$

$$\varphi = -(L_{50} \times \omega) \quad (7)$$

$$\omega = \frac{2 \times \log 3}{SR} \quad (8)$$

where L_{50} is the salinity where τ_y equals 0.5, and SR is a steepness parameter defining how fast the logistic curve increases (see Table 2 for priors). This function reflects the hypothesis of a ratcheted change in the zooplankton community at a threshold salinity level. According to expert judgement, the abundance of large, neritic zooplankton species (*Pseudocalanus* and *Temora spp.*) decreases markedly when salinity declines below 5.5‰ (unpublished monitoring data by Finnish Environment Institute). In equation 5 the term (-2) is a scaling factor which ensures that the prior median of $E(\log(K_y))$ corresponds to 0.2, which is the ballpark figure for K in FishBase³ for the GoF herring population.

Impact of cod abundance on natural mortality

Natural mortality of herring was assumed to decrease as individual weight increases (Beyer 1989). We included an additive mortality component Me_y which depended on the abundance of herring's main predator, cod, to the natural mortality in the model of Mäntyniemi et al. (2013). In our model, cod abundance affected herring's natural mortality rate as an additional harvesting fleet referring to attack rate-type interaction between cod and herring

$$M_{y,a} = MM_{y,a} + Me_y \quad (9)$$

$$\log(MM_{y,a}) = G \times \log(w_{y,a}) \quad (10)$$

$$\log(Me_y) \sim N(muLMe_y, Te) \quad (11)$$

$$muLMe_y = \log(cM) + \log(Cod_y) - \frac{0.5}{Te} \quad (12)$$

where $MM_{y,a}$ denotes the size-dependent natural mortality of Mäntyniemi et al., (2013), $w_{y,a}$ is the mean weight in stock at age a and year y , Te describes the amount of unexplained variation among

³

<http://www.fishbase.org/PopDyn/PopGrowthList.php?ID=24&GenusName=Clupea&SpeciesName=harengus&fc=43>

years, cM is a constant, Cod_y is the observed catch, and G describes the relationship between weight and natural mortality (Gislason et al. 2010). The applied priors (Table 2) yield realisations around value 0.1 for Me , which we anticipate to be a reasonable figure. The cod-induced natural mortality is independent of herring size.

2.4 Computation

The computation for the joint posterior distribution for the population dynamics model parameters was conducted using Markov chain Monte Carlo (MCMC, e.g. Gilks et al. 1996) methods. The simulation was implemented using JAGS (Plummer 2012). The Bayesian model averaging over competing model structures was also implemented within a single MCMC run by adopting the approach introduced by Carlin and Chib (1995). The model run consisted of three independent MCMC chains. Each MCMC chain was run on a separate processor core. Generation of 20,000,000 samples took about 7 days. Only every 2,000th sample was retained to remove auto-correlation. The first 2,000 samples were discarded as the burn-in period. Neither visual inspection of the chains nor Gelman-Rubin diagnostic plots implied any non-convergence issues. The remaining 6,000 samples from each chain were merged to form a sample of 18,000 that was used for posterior inference.

The inference for the GP models for the environmental covariates (Section 2.2) was conducted by optimising the covariance function parameters in their maximum a posterior (MAP) estimate, after which we were able to solve the posterior distribution for the environmental covariates, given the covariance function parameters, analytically using the standard GP posterior predictive equations (Rasmussen and Williams 2006). The MAP estimate for the covariance function parameters is faster to calculate than full MCMC and, with large data sets, the predictive performance of GP is insensitive to the exact value of the covariance function parameters (Vanhatalo et al. 2010). All GP computations were conducted with Matlab using the GPstuff toolbox (Vanhatalo et al. 2013).

3. Results

The applied Bayesian approach updates knowledge about environmental forcing on the herring stock. We first describe how eutrophication impacts recruitment of herring, secondly portray how the Antonio Gramsci oil spill did influence mortality in two life-stage based sub-stocks in the GoF, and thirdly indicate the effect of salinity, sea surface temperature, and cod and sprat abundance on the herring stock dynamics. We also describe how Bayesian model averaging informs us about the credibility of the alternative hypothesis concerning oil-induced mortality (no additional mortality versus additional mortality) and the stock-recruitment relationship (Beverton-Holt versus Ricker function).

The dome-shaped effect of chlorophyll- α on recruitment has a maximum of 4.7 $\mu\text{g/l}$ (Fig. 4). Chlorophyll levels below and above this optimum will yield decreased recruitment and, thus, in the long run, decreasing stock biomass and sustainable catch, unless compensated with increased growth rate. The posterior mean of the chlorophyll- α ranged from 5.4 to 11.2 in the period 1980-2006 (Fig. 2). These levels are clearly above the optimal chlorophyll- α concentration (Table 2). Moreover, the chlorophyll- α concentration increased towards the end of the studied period. In the current state of eutrophication, every spawning herring produces only about 40% of the offspring it could produce if the GoF was in the optimal condition with respect to eutrophication.

Among offspring, the oil induced mortality was higher a posteriori than in our prior. This is indicated by the shift of the posterior distribution towards the right (Fig. 5). The median value is 17% but the uncertainty is large (Table 2). The two modes in the posterior distribution for offspring mortality are a product of prior probability and likelihood, which, in turn, is determined by the model structure and data. Oil spill-induced mortality appears to have been lower than anticipated among herring of age two years and older (Fig. 5). Mortality caused by the oil spill is about 10% in the older age groups. Uncertainty is lower compared to the offspring mortality.

Uncertainty about the appropriate model structure influences the conclusions about oil-induced mortality and herring recruitment dynamics. The hypothesis that the oil spill caused elevated mortality among herring is 1.2 times more probable than the hypothesis that there was no extra mortality in 1987 (hypothesis credibilities 54% and 46% for “oil induced mortality” and “no oil induced mortality”, respectively). According to the posterior probabilities of alternative models, the Ricker form stock-recruitment relationship is 3.2 times more probable than the Beverton-Holt (hypothesis credibilities 76% and 24% for Ricker and Beverton-Holt, respectively. Fig. 6) function. Tree-parameter stock-recruitment models (Maunder and Deriso 2011) can represent both Ricker and Beverton-Holt functions and their intermediates and might be worthy of further investigation.

Posterior distributions of all model parameters are weighted averages of the posterior distributions under alternative hypothesis related to oil induced mortality and stock-recruitment function, where the posterior probabilities of these hypotheses are used as weights. Whatever the underlying functional relationship between spawning stock size and the following recruitment, the scatterplot is noisy (Fig. 6). Importantly, however, the probabilistic model is capable of providing information about the model and the parameter uncertainty which can be highly relevant in practical fisheries management. A manager might seek for a harvest control rule to cope with the uncertain impact on eutrophication on herring recruitment, coupled with uncertainty about the future nutrient loads. Using the Bayesian decision analysis approach it would be possible to choose from a set of alternative harvest controls rules the one which maximizes the expected utility.

The model suggests that April-June SST influences recruitment (Fig. 7). The center 95% credible interval of the parameter b is between 0 and 1.5 (Table 2). Values below 1 indicate a concave relationship between SST and recruitment, while values above 1 point to a convex relationship. The median is about 0.7. The majority of the posterior probability thus supports a concave relationship between SST and recruitment and with 97,5% probability SST has influenced herring recruitment (Table 2 and Fig. 7). The median posterior value of b implies that $\pm 1^\circ\text{C}$ deviation of the average SST

has about $\pm 14\%$ impact on recruitment so that the coldest spring, in 1997, would have reduced recruitment by 22% from the recruitment at an average SST condition which is comparable to the estimated impact of the Antonio Gramsci oil spill.

The salinity threshold leading to slower growth rate of herring - hypothetically through decreased abundance of energy rich, large neritic zooplankton - appears to be lower than anticipated a priori. The posterior estimate for L_{50} is about 4.7 psu while the applied prior was 5.5 psu (Fig. 4). There are two asymptotes and a threshold in the function. Growth rate is intermediate between these asymptotes: growth rate starts degrading as salinity drops below 6 psu, and increases as salinity grows above 4 psu. Although the salinity-growth function reaches zero at low salinities, this does not imply cessation of growth but in the model parameterization higher salinities boost growth rate beyond the base level determined by herring length.

The data did not update our prior related to the influence of sprat abundance or late summer SST on the herring growth rate (Table 2). The lack of new information is indicated by the similarity between the prior and posterior probabilities of the parameters cLS and ck .

Posterior estimates for natural mortality $M_{y,a}$ at age 1, 5 and 8 are shown in Fig. 8. Natural mortality decreases as a function of age but there is no clear time-dependent trend in the size-based component $MM_{y,a}$. The natural mortality estimates for age 1 from the model are reasonably similar to the estimates by the ICES stock assessment working group (WGBFAS) (ICES 2016) for the early 1980s. The estimates by ICES decrease markedly after this period. Predation obviously has had an influence on herring mortality in the early 1980s when the Baltic cod stock peaked (ICES 2008). The additional natural mortality induced by cod, Me_y , decreases in synchrony with the cod abundance. The two components of mortality mix with each other to some extent. As a result, uncertainty in both of these natural mortality components is highest in the 1980s. However, use of information about fluctuations in the cod stock serves in the estimation of natural mortality. If information about cod stock and concomitant predation were omitted, their effect would have been mostly explained

by the random effects. The natural mortality estimates are only slightly higher for ages 2-8 than those applied by the WGBFAS (ICES 2016).

Our results are also of interest in conventional fish stock assessment. These include annual estimates of stock biomass and reproduction, and fishing mortality (Fig. 9). The Gulf of Finland herring spawning stock size peaked in the mid-1980s but since then has had a monotonic downward trend. The continuous decline of the total biomass commenced about 10 years later. Fishing mortality of the important age-groups displays an increase since the 1990s. The highest instantaneous fishing mortality rates are at a level of 0.5-1 in 1996-2003. Herring growth, in terms as length has varied among years without a distinctive pattern. Although the K_y is an annual estimate, we characterised the posterior estimate pooled over the study period (Table 2).

5. Discussion

The discussion section starts with the study's main conclusions and then deliberates about the mechanisms by which eutrophication may impact herring populations. Then, we elucidate the dynamics of the Gulf of Finland's herring fishery in the context of a changing environment, and deal with the issue of the Baltic herring stock structure. Next, we make conclusions about herring growth rate based on the model, before focusing on the impacts of the oil spill. Finally, we discuss challenges and solutions for pragmatic environmental impact evaluation.

We have applied a probabilistic modelling framework to infer the influence of eutrophication, a major oil spill and a number of environmental covariates on the population dynamics of the Gulf of Finland herring. Changes in the abiotic conditions inflict recruitment fluctuations, while eutrophication forces a recruitment trend, which is negative for the GoF. Our results indicate also a positive relationship between recruitment and April-June sea surface temperature. The Antonio Gramsci oil spill increased the mortality of herring, especially at the early life-stage. The analysis

suggests, however, that the current eutrophication status has more adverse implications on herring population production than this oil spill has had. The mechanisms causing changes in population dynamics via environmental fluctuations and trends may be direct or indirect. There is an apparent need to understand these actual mechanisms, because what happens at the level of the individual will have consequences for population response to natural and anthropogenic pressure (Hutchings and Kuparinen 2014). Our analysis is not helpful in this sense, as we pooled all of the underlying processes into the estimated chlorophyll index. It is assumed to capture the many complex responses of the ecosystem to eutrophication that influence the recruitment of herring. They are mainly processes inducing reduced quality and area of spawning and nursery habitats: increased sedimentation impairing herring spawning beds and increased biomass of filamentous algae restraining habitat-forming submerged plants, and increased turbidity reducing the depth of occurrence of plants which herring prefer as spawning substrate (Aneer and Nellbring 1982, Rajasilta et al. 1989, Kääriä et al. 1997). It has been also suggested that filamentous algae may increase herring egg mortality via low oxygen levels (Aneer 1985) or toxic exudates (Aneer 1987). Furthermore, there may exist a weak link between herring recruitment and coastal hypoxia. We suggest a weak link because spawning beds do not reach the depths (Rajasilta et al. 1989) where hypoxia mostly occurs. Moreover, herring spawning time does not match with the peak months of hypoxia in August and September (Conley et al. 2011).

In the GoF, eutrophication has been an important driver of benthic hypoxia, particularly during the last 50 years, although its elemental basis is the strong halocline causing permanent water-column stratification (Laine et al. 2007, Zillen et al. 2008). As the extent of anoxic bottoms has increased in the coastal zone of GoF, the area suitable for the reproduction of herring and for supplying benthic fauna has decreased. The latter potentially affects adversely the feeding conditions of the post-juvenile herring. Urho et al. (2003) observed a strong negative correlation between total primary production and abundance of herring offspring in three adjacent bays in the GoF. As herring larvae and juveniles remain in in-shore and sheltered areas (Urho and Hildén 1990), the amount and quality

461 of the semi-pelagic coastal nursery areas may also impact year-class abundance as well as the
462 availability of dietary resources, i.e. zooplankton.

463 Herring fishery in the GoF has markedly declined since the late 1980s (Fig. 10, Stephenson et al. 2001,
464 Rahikainen 2005), but has been faintly recovering since the mid-2000 (Natural Resources Institute
465 Finland, unpubl. statistics). The annihilation of the fishery was driven by changes in the markets and
466 fluctuations in biological factors. Stephenson et al. (2001) proposed that the decline of large herring
467 was a consequence of reduced weight-at-age. Our analysis suggests that stock biomass also declined
468 markedly at the same time under heavy fishing pressure (Fig. 9). The excessive fishing mortality rate
469 influenced the size distribution of the stock by reducing the fraction of large herring in the
470 population. As a consequence, the Finnish trawl fleet withdrew from the GoF due to insufficient bulk
471 of large herring of suitable size for filleting. The decline of the abundance was enhanced by
472 progressive eutrophication (Fig. 2) causing impaired reproduction of herring (Fig. 9). Also anecdotal
473 information suggests the loss of occupation of traditional herring spawning areas in the 1980s which
474 was reflected in collapsing herring trap net effort in the Gulf of Finland.

475 The highest fishing mortality estimates concern the period 1996-2003 (Fig. 9), when they were well
476 above the level which is regarded sustainable for the Central Baltic herring stock component
477 including the Baltic Proper and the GoF (ICES Advice 2012). The estimates of the most recent
478 spawning stock abundance and recruitment are located nearest to the origin in the scatterplot (Fig.
479 6). This indicates both stock decline and recruitment impairment, likely incurring from a combination
480 of overfishing and eutrophication. The sustainable catch is decreasing as eutrophication progresses.
481 The fisheries sector could benefit from policies which reduce excessive nutrient load from terrestrial
482 sources. In practise, the control from these types of environmental and agricultural policies is weak
483 and the society's ability to manage the herring population in the GoF is only effective in the
484 conventional fishery management context (Rahikainen et al. 2014).

The importance of stock structure information for effective fisheries management is widely accepted, but defining discrete stocks is problematic (Stephenson 1999). The complex but unresolved herring stock structure is an assessment and management concern in the Baltic also (Rahikainen and Stephenson 2004, Reiss et al. 2009, Gröhsler et al. 2013). However, significant genetic differentiation is found within the Baltic Sea (Jørgensen et al. 2005). The analysis by Jørgensen et al. (2005) indicates a strong structuring effect of the salinity gradients on herring populations. The main barrier for gene flow is identified between the GoF and the Baltic Proper, yet ICES WGBFAS (2008) has been assessing the GoF component in aggregation with the large Central Baltic stock. The management issue is that components within a large assessment area may have diverse dynamics, which cannot be evaluated under a pooled assessment structure, and the weak subcomponents are at risk of being depleted without notice. On the other hand, uncertainty about the migrations among the areas decreases confidence in the disaggregated assessment approach. There are no quantitative estimates available about the Gulf of Finland herring migrating out of the gulf and being fished in the Baltic Proper, nor are there estimates about the Central Baltic stock migrating in the GoF and being fished there. Due to lack of data, the impact of the stock mixing on the model estimates would be completely driven by the priors. Our prior is that these migrations are equal. Therefore, accounting for the stock mixing would not have influenced the conclusion drawn from the model.

Salinity, sprat abundance and the July-August SST have an effect on herring growth rates. The applied model and data updated most of the knowledge about the effect of salinity (Fig. 4) which has been proposed to be linked to growth by inducing changes in the zooplankton community. However, the data was uninformative about the effect of SST on growth. This lack of new information is indicated by the factually identical prior and posterior distributions for parameter ck (Table 2). This, nevertheless, specifies a positive effect on growth rate. The data include the information that sprat stock abundance negatively affects herring growth. However, the location and shape of the posterior distribution for cLS has changed only little compared to the prior (Table 2).

Our analysis of the oil spill effect indicates additional mortality in the GoF herring stock in 1987. Oil-induced mortality was nearly 20% among the offspring, but in the older age-groups the posterior median of additional mortality was about 10% (Table 2, Fig. 4). The centred posterior 95% credible interval is 0-57% for offspring and 0-22% for older ages (Table 2). This result conforms with earlier reports which denote that the larval phase is the most sensitive life stage for oil exposure (Reinharz and Michel 1996). According to the post-incidence hazard monitoring programme in the GoF, herring larvae with abnormal posterior notochord and short-bodied larvae were frequent near the grounding site of the Antonio Gramsci in 1987 and 1988, suggesting elevated mortality among herring offspring (Urho 1991). Unfortunately, oil had arrived to the control area after the first samplings and only compromised data were available about the baseline frequency of malformed larvae (Urho 1991). So far it has not been possible to make conclusions about the impact of this accident on the herring stock dynamics. Our results now support the hypothesis that the high frequency of abnormal herring larvae signalled the elevated mortality of the 1987 year-class. Oil may have remained on the spawning grounds in subsequent years but we modelled only the immediate oil induced mortality on herring.

The negative influence of the Antonio Gramsci oil spill on herring recruitment is at the same level as the effect of very cold weather during the spawning and larval phase. The coldest April-June average SST in 1980-2006 was 3.7 °C, which reduces recruitment by 0.012-41% with 95% probability as compared with an average SST (Fig. 7). The amount of spilled oil was 570 t in the Antonio Gramsci case. A realistic worst-case scenario in the GoF is assumed to be a collision of an oil tanker with another vessel, which could result in an oil spill of approximately 30,000 t of crude oil (Hietala and Lampela 2007). An accident of this magnitude would no doubt exceed the effect of sea temperature fluctuations, with a severe and long-lasting effect on herring recruitment, and other fish and marine biota.

All of the estimates resulting from the model that includes oil induced mortality are conditional on the assumption that oil affects herring mortality (Fig. 3). Since the model with oil effect has larger posterior probability and since the prior probabilities of these effects have clearly been updated in the posterior, we can conclude that there is evidence that the Antonio Gramsci oil spill had a negative impact on GoF herring reproduction.

Spilled oil will sediment on the bottom over the course of the weathering process (Reed et al. 1999). Our analysis of the oil induced mortality focuses on herring, which is a benthic spawner (Rajasilta et al. 1989; 1993), making the spawners, roe, and hatched larvae vulnerable to exposure to sedimented oil. Marine fishes with pelagic eggs, including cod and sprat, spawn in the Baltic's deep basins (MacKenzie et al. 2000, Nissling et al. 2003). Cod and sprat spawning and nursery areas are outside the GoF, so their offspring are unlikely to have been impacted by the Antonio Gramsci accident.

There are typical challenges in verifying the impact of accidents on ecosystem functioning. Underwood (1997) proposes that in the classical statistical framework, data should be gathered and analysed ideally in the BACI-setting (before-after-control-impact). Unfortunately, these conditions, which factually require experimental design, are rarely achieved in actual environmental impact evaluations. In many instances, monitoring starts only after the accident making conclusions on the status before the accident vague. The value of pre- versus post-incident surveys in quantifying oil-related injuries to fish will depend on natural variability in the measured parameters and degree of injury caused by the incident. Overall, testing hypotheses is usually complex due to large natural variability in space and time (Hilborn 1996).

Bayesian model averaging provides an alternative framework to discriminate amongst different explanations (models) by contrasting the data with our prior understanding of the ecosystem. The model structure and the number of unknown parameters depend primarily on biological knowledge about the species and environmental forcing, and on the knowledge about the survey design and behaviour of the fishing fleets. Our modelling approach intends to describe the system as

realistically as possible and makes inferences about latent (unobserved) parameters. Here, we use accumulated knowledge in the scientific literature about toxic effects of oil and the effects of several biotic and abiotic factors shaping herring stock dynamics. It is necessary to take them into account to reach a credible understanding of the impacts of eutrophication and the oil spill on the herring stock. Otherwise, their potential for confounding leverage would be high.

It is noteworthy that we analyzed both prior and post-incident conventional fish stock assessment data, 3 decades worth of, to identify the impact of the oil spill on the GoF herring stock. We expect that a shorter data period would hamper the ability to indicate the influence of the oil spill. Our findings suggest short term data would not allow detection of the impacts of an accident on the ecosystem. Costly environmental monitoring no doubt provides an incentive to keep the monitoring period short. In such cases, there will be a higher probability that actual negative impacts are not identified at all, or at least convincing scientific evidence will not be obtained. This will naturally act in favour of the polluter, leading to lower compensation costs or rehabilitation effort. The problem will emerge especially if uncertainty is ignored during the impact assessment. The Bayesian approach can be used to create a learning chain from one accident to the next in order to effectively reduce the uncertainty about the environmental impacts of anthropogenic disasters.

Acknowledgements

This study was supported by the project “Integrated Bayesian risk analysis of ecosystem management in the Gulf of Finland” (IBAM), funded by the European Community’s Seventh Framework Programme under grant agreement no 217246 made with the joint Baltic Sea research and development programme BONUS, and the Academy of Finland. JV has additionally been funded by the Academy of Finland (grant 266349). We thank two anonymous reviewers for their constructive comments which improved the manuscript.

6. References

- Andersen, J.H., Axe, P., Backer, H., Carstensen, J., Claussen, U., Fleming-Lehtinen, V., Järvinen, M., Kaartokallio, H., Knuuttila, S., Korpinen, S., Kubiliute, A., Laamanen, M., Lysiak-Pastuszek, E., Martin, G., Murray, C., Møhlenberg, F., Nausch, G., Norkko, A. and Villnäs, A. 2011. Getting the measure of eutrophication in the Baltic Sea: towards improved assessment principles and methods. *Biogeochemistry* 106: 137-156.
- Aneer, G. 1985. Some speculations about the Baltic Herring (*Clupea harengus membras*) in connection with the eutrophication of the Baltic Sea. *Can. J. Fish. Aquat. Sci.* 42 S1: 83-90.
- Aneer, G. 1987. High natural mortality of Baltic herring (*Clupea harengus*) eggs caused by algal exudates? *Mar. Biol.* 94: 163-169.
- Aneer, G. and Nellbring, S. 1982. A SCUBA-diving investigation of Baltic herring (*Clupea harengus membras* L.) spawning grounds in the Askö-Landsort area, northern Baltic proper. *J. Fish Biol.* 21: 433-442.
- Aneer, G. 1989. Herring (*Clupea harengus* L.) spawning and spawning ground characteristics in the Baltic Sea. *Fish Res* 8: 169–195.
- Bekkevold, D., Gross, R., Arula, T., Helyar, S.J. and Ojaveer, H. 2016. Outlier Loci Detect Intraspecific Biodiversity amongst Spring and Autumn Spawning Herring across Local Scales. *PLoS ONE* 11(4): e0148499. doi: 10.1371/journal.pone.0148499
- Beyer, J.E. 1989. Recruitment stability and survival: simple size-specific theory with examples from the early life dynamics of marine fish. *Dana* 7: 47–147.
- Blaxter, J.H.S. and Hempel, G. 1963. The influence of egg size on herring larvae. *J. Cons. int. Explor. Mer.* 28: 211-240.

- 606 Brunel, T. and Dickey-Collas, M. 2010. Effects of temperature and population density on von
607 Bertalanffy growth parameters in Atlantic herring: a macro-ecological analysis. Mar. Ecol. Prog.
608 Ser. 405: 15-28.
- 609 Brunila, O.-P. and Storgård, J. 2012. Oil transportation in the Gulf of Finland in 2020 and 2030.
610 Publications from the Centre for Maritime Studies University of Turku, A 61.
- 611 Caddy, J.F. 1993. Toward a comparative evaluation of human impacts on fishery ecosystems of
612 enclosed and semi-enclosed seas. Rev. Fish. Sci. 1: 57–95.
- 613 Cardinale, M., Möllmann, C., Bartolino, V., Casini, M., Kornilovs, G., Raid, T., Margonski, P., Grzyb, A.,
614 Raitaniemi, J., Gröhsler, T. and Flinkman, J. 2009. Effect of environmental variability and spawner
615 characteristics on the recruitment of Baltic herring *Clupea harengus* populations. Mar. Ecol. Prog.
616 Ser. 388: 221–234.
- 617 Carlin, B.P. and Chib, S. 1995. Bayesian model choice via Markov chain Monte-Carlo methods. J. R.
618 Stat. Soc. B 57: 473–484.
- 619 Carstensen, J., Andersen, J.H., Gustafsson, B.G. and Conley, D.J. 2014. Deoxygenation of the Baltic
620 Sea during the last century. Proceedings of the National Academy of Sciences 111: 5628-5633.
- 621 Casini, M., Bartolino, V., Molinero, J.C. and Kornilovs, G. 2010. Linking fisheries, trophic interactions
622 and climate: threshold dynamics drive herring *Clupea harengus* growth in the central Baltic Sea.
623 Mar. Ecol. Prog. Ser. 413: 241–252.
- 624 Cloern, J.E. 2001. Our evolving conceptual model of the coastal eutrophication problem. Mar. Ecol.
625 Prog. Ser. 210: 235-265.
- 626 Colby, P.J., Spangler, G.R., Hurley, D.A. and McCombie, A.M. 1972. Effects of eutrophication on
627 salmonid communities of oligotrophic lakes. J. Fish. Res. Bd. Can. 29: 975-983.

- 628 Conley, D.J., Carstensen, J., Aigars, J., Axe, P., Bonsdorff, E., Eremina, T., Haahti, B.-M., Humborg, C.,
629 Jonsson, P., Kotta, J., Lännegren, C., Larsson, U., Maximov, A., Rodriguez Medina, M., Lysiak-
630 Pastuszak, E., Remeikaite-Nikien, N., Walve, J., Wilhelms, S. and Zillén, L. 2011. Hypoxia is
631 increasing in the coastal zone of the Baltic Sea. *Environmental Science & Technology* 45: 6777-
632 6783.
- 633 Fleming-Lehtinen, V. and Laamanen, M. 2012. Long-term changes in Secchi depth and the role of
634 phytoplankton in explaining light attenuation in the Baltic Sea. *Estuarine, Coastal and Shelf*
635 *Science* 102–103: 1–10.
- 636 Fleming-Lehtinen, V., Laamanen, M., Kuosa, H., Haahti, H. and Olsonen, R. 2008. Long-term
637 Development of Inorganic Nutrients and Chlorophyll a in the Open Northern Baltic Sea. *Ambio* 37:
638 86-92.
- 639 Flinkman, J., Aro, E., Vuorinen, I. and Viitasalo, M. 1998. Changes in northern Baltic zooplankton and
640 herring nutrition from 1980s to 1990s: top-down and bottom-up processes at work. *Mar. Ecol.*
641 *Progr. Ser.* 165: 127-136.
- 642 Gelfand, A., Diggle, P.J., Fuentes, M. and Guttorp, P. 2010. *Handbook of Spatial Statistics*. CRC press
- 643 Gilks, W.R., Richardson, S. and Spiegelhalter, D.J. 1996. *Markov chain Monte Carlo in practice*.
644 Chapman & Hall, London.
- 645 Gislason, H., Daan, N., Rice, J.C. and Pope, J.G. 2010. Size, growth, temperature and the natural
646 mortality of marine fish. *Fish Fisheries* 11: 149–158. doi:10.1111/j.1467-2979.2009.00350.x.
- 647 Grall, J. and Chauvaud, L. 2002. Marine eutrophication and benthos: the need for new approaches
648 and concepts. *Global Change Biol.* 8: 813-830.

- 649 Gröhsler, T., Oeberst, R., Schaber, M., Larson, N. and Kornilovs, G. 2013. Discrimination of western
650 Baltic spring-spawning and central Baltic herring (*Clupea harengus* L.) based on growth vs. natural
651 tag information. ICES J. Mar. Sci. 70: 1108-1117.
- 652 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey,
653 K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig,
654 E.R., Spalding, M., Steneck, R. and Watson, R. 2008. A global map of human impact on marine
655 ecosystems. Science 319: 948-952.
- 656 Hartmann, J. and Nümann, W. 1977. Percids of Lake Constance, a lake undergoing eutrophication. J.
657 Fish. Res. Bd. Can. 34: 1670-1677.
- 658 Hassler, B. 2011. Accidental Versus Operational Oil Spills from Shipping in the Baltic Sea: Risk
659 Governance and Management Strategies. AMBIO 40: 170–178.
- 660 HELCOM 2010. Ecosystem Health of the Baltic Sea 2003–2007: HELCOM Initial Holistic Assessment.
661 Balt. Sea Environ. Proc. No. 122. 63 pp.
- 662 Hietala, M. and Lampela, K. (Editors) 2007. Oil Pollution Preparedness on the Open Sea – Final
663 Report of the Working Group, Finnish Environment Institute, Helsinki.
- 664 Hilborn, R. 1996. Detecting population impacts from oil spills: a comparison of methodologies. In
665 Proceedings of the Exxon Valdez oil spill symposium. Edited by Rice, S.D., Spies, R.B., Wolfe, D.A.
666 and Wright, B.A. American Fisheries Society Symposium 18. Bethesda, Maryland, USA, p 639–644.
- 667 Hirvi, J.P. 1990a. The Oil Spill in the Gulf of Finland in 1987. Proceedings IAHR 1990, 10th
668 International Symposium on Ice, Espoo, Finland.
- 669 Hirvi, J.P. 1990b. Summary of the environmental effects of the Antonio Gramsci oil spill in the Gulf of
670 Finland in 1987. Proceedings of the Thirteenth Arctic and Marine Oilspill Program Technical

- 671 Seminar, June 6-8-. 1990, Edmonton, Alberta. Ministry of Supply and Services Canada. Catalogue
672 No. En 40-11/5-1990. ISBN 0-662-57535-0.
- 673 Hutchings, J.A. and Kuparinen, A. 2014. Ghosts of fisheries-induced depletions: do they haunt us still?
674 ICES J. Mar. Sci. 71: 1467-1473.
- 675 ICES 2001. Report of the Baltic Fisheries Assessment Working Group. ICES Document, CM
676 2001/ACFM:18. 546 pp.
- 677 ICES 2007. Report of the Baltic Fisheries Assessment Working Group (WGBFAS). ICES CM
678 2007/ACFM:15. 735 pp.
- 679 ICES 2008. Report of the Baltic Fisheries Assessment Working Group (WGBFAS). ICES CM
680 2008\ACOM:06. 692 pp.
- 681 ICES 2009. Report of the Baltic Fisheries Assessment Working Group. ICES CM 2009/ACOM:07. 626
682 pp.
- 683 ICES 2016. Report of the Baltic Fisheries Assessment Working Group. ICES CM 2016/ACOM:11. 584
684 pp.
- 685 ICES Advice 2012. Report of the ICES Advisory Committee 2012. Book 8, Baltic Sea. 155 pp.
- 686 Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H.,
687 Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi,
688 J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J. and Warner, R.R. 2001. Historical overfishing and
689 the recent collapse of coastal ecosystems. Science 293: 629-637.
- 690 Juntunen, T., Vanhatalo, J., Peltonen, H., Mäntyniemi, S. 2012. Bayesian spatial multispecies
691 modeling to assess pelagic fish stocks from acoustic and trawl survey data. ICES J. Mar. Sci. 69:
692 95-104.

- 693 Jørgensen, H.B.H., Hansen, M.M., Bekkevold, D., Ruzzante, D.E. and Loeschcke, V. 2005. Marine
694 landscapes and population genetic structure of herring (*Clupea harengus* L.) in the Baltic Sea.
695 Molecular Ecology 14: 3219–3234.
- 696 Kornilovs, G., Sidrevics, L. and Dippner, J.W. 2001. Fish and zooplankton interaction in the Central
697 Baltic Sea. ICES J. Mar. Sci. 58: 579-588.
- 698 Korpinen, S., Meski, L., Andersen, J.H. and Laamanen, M. 2012. Human pressures and their potential
699 impact on the Baltic Sea ecosystem. Ecological Indicators 15: 105-114.
- 700 Kotta, J., Kotta, I., Simm, M. and Põllupüü, M. 2009. Separate and interactive effects of
701 eutrophication and climate variables on the ecosystem elements of the Gulf of Riga. Estuarine,
702 Coastal and Shelf Science 84: 509-518.
- 703 Kuikka, S., Vanhatalo, J., Pulkkinen, H., Mäntyniemi, S. and Corander, J. 2014. Experiences in
704 Bayesian Inference in Baltic Salmon Management. Statistical Science 29:42-49.
- 705 Kuparinen, A., Mäntyniemi, S., Hutchings, J. and Kuikka, S. 2012. Increasing biological realism of
706 fisheries stock assessment: towards hierarchical Bayesian methods. Environmental Reviews 20:
707 135-151.
- 708 Kääriä, J., Rajasilta, M., Kurkilahti, M. and Soikkeli, M. 1997. Spawning bed selection by the Baltic
709 herring (*Clupea harengus membras*) in the Archipelago of SW Finland. ICES J. Mar. Sci. 54: 917–
710 923.
- 711 Laine, A.O., Andersinm A.-B., Leiniöm S. and Zuur, A.F. 2007. Stratification-induced hypoxia as a
712 structuring factor of macrozoobenthos in the open Gulf of Finland (Baltic Sea). J. Sea Res. 57: 65-
713 77.
- 714 Lambert, Y. 2008. Why should we closely monitor fecundity in marine fish populations? J. Northwest
715 Atl. Fish. Sc. 41: 93–106.

- 716 Lecklin, T., Ryömä, R. and Kuikka, S. 2011. A Bayesian network for analyzing biological acute and
717 long-term impacts of an oil spill in the Gulf of Finland. Mar. Poll. Bull. 62: 2822–2835.
- 718 Lindegren, M., Östman, Ö. and Gårdmark, A. 2011. Interacting trophic forcing and the population
719 dynamics of herring. Ecology 92: 1407-1413.
- 720 MacKenzie, B.R., Hinrichsen, H.H., Plikshs, M., Wieland, K. and Zezera, A.S. 2000. Quantifying
721 environmental heterogeneity: habitat size necessary for successful development of cod *Gadus*
722 *morhua* eggs in the Baltic Sea. Mar. Ecol. Prog. Ser., 193. 143-156.
- 723 Marshall, C.T., Needle, C.L., Thorsen, A., Kjesbu, O.S. and Yaragina, N.A. 2006. Systematic bias in
724 estimates of reproductive potential of an Atlantic cod (*Gadus morhua*) stock: implications for
725 stock recruit theory and management. Can. J. Fish. Aquat. Sci. 63:980-994.
- 726 Maunder, M. N. and Deriso, R. B. 2011. A state – space multistage life cycle model to evaluate
727 population impacts in the presence of density dependence : illustrated with application to delta
728 smelt (*Hyposmesus transpacificus*). Can. J. Fish. Aquat. Sci. 68: 1285–1306.
- 729 Mäntyniemi, S., Uusitalo, L., Peltonen, H., Haapasaari, P. and Kuikka, S. 2013. Integrated age-
730 structured length-based stock assessment model with uncertain process variances, structural
731 uncertainty and environmental covariates: case of Central Baltic herring. Can. J. Fish. Aquat. Sci.
732 70: 1317-1326.
- 733 National Research Council (NRC) 2003. Oil in the Sea III: Inputs, Fates and Effects. National Research
734 Council of the National Academies. The National Academies Press, Washington, D.C. 265 p
- 735 Naylor, R.L., Goldburg, R.J., Primavera, J.H., Kautsky, N., Beveridge, M.C.M., Clay, J., Folke, C.,
736 Lubchenco, J., Mooney, H. and Troell, M. 2000. Effect of aquaculture on world fish supplies.
737 Nature 405: 1017-1024.

- 738 Nissling, A., Müller, A. and Hinrichsen, H.-H. 2003. Specific gravity and vertical distribution of sprat
739 eggs in the Baltic Sea. J. Fish Biol. 63: 280–299.
- 740 Parmanne, R. 1990. Growth, morphological variation and migrations of herring (*Clupea harengus* L.)
741 in the northern Baltic Sea. Finn. Fish. Res. 10: 1–48.
- 742 Parmanne, R. and Kuittinen, E. 1991. Comparison of the fecundity of herring (*Clupea harengus* L.) in
743 various areas of the northern Baltic Sea. Finn. Fish. Res. 12: 157-164.
- 744 Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean
745 environmental temperature in 175 fish stocks. Journal du Conseil 39: 175-192.
- 746 Pepin, P. 1991. Effect of temperature and size on development, mortality, and survival rates of the
747 pelagic early life history stages of marine fish. Can. J. Fish. Aquat. Sci. 48: 503-518.
- 748 Persson, L., Diehl, S., Johansson, L., Andersson, G. and Hamrin, S.F. 1991. Shifts in fish communities
749 along the productivity gradient of temperate lakes - patterns and the importance of size-
750 structured interactions. J. Fish Biol. 38: 281-293.
- 751 Plummer, M. 2012. JAGS version 3.2.0 user manual.
- 752 Quinn, T.J. and Deriso, R.B. 1999. Quantitative fish dynamics. Oxford University Press, New York. 542
753 pp.
- 754 Rabalais, N.N., Turner, R.E., Díaz, R.J. and Justić, D. 2009. Global change and eutrophication of
755 coastal waters. ICES J. Mar. Sci. 66: 1528–1537.
- 756 Rahikainen, M. 2005. Evaluation and management of the Finnish herring fishery. Academic
757 dissertation. University of Helsinki. 70 pp.
758 <http://ethesis.helsinki.fi/julkaisut/bio/bioja/vk/rahikainen/>

- 759 Rahikainen, M., Helle, I., Haapasaari, P., Oinonen, S., Kuikka, S., Vanhatalo, J., Mäntyniemi, S. and
760 Hoviniemi, K.-M. 2014. Toward Integrative Management Advice of Water Quality, Oil Spills, and
761 Fishery in the Gulf of Finland: A Bayesian Approach. *AMBIO* 43: 115-123.
- 762 Rahikainen, M. and Stephenson, R.L. 2004. Consequences of growth variation in northern Baltic
763 herring for assessment and management. *ICES J. Mar. Sci.* 61: 339-351.
- 764 Rajasilta, M., Eklund, J., Hänninen, J., Kurkilahti, M., Kääriä, J., Rannikko, P. and Soikkeli, M. 1993.
765 Spawning of herring (*Clupea harengus membras* L.) in the Archipelago Sea. *ICES J. Mar. Sci.* 50:
766 233-246.
- 767 Rajasilta, M., Eklund, J., Kääriä, J. and Ranta-aho, K. 1989. The deposition and mortality of the eggs of
768 the Baltic herring *Clupea harengus membras* L., on different substrates in the south-west
769 archipelago of Finland. *J. Fish Biol.* 34: 417-427.
- 770 Rasmussen, C.E. and Williams, C.K.I. 2006. Gaussian Processes for Machine Learning. The MIT Press:
771 Cambridge MA.
- 772 Reed, M., Johansen, Ø., Brandvik, P. J., Daling, P., Lewis, A., Fiocco, R., Mackay, D. and Prentki, R.
773 1999. Oil Spill Modeling towards the Close of the 20th Century: Overview of the State of the Art.
774 *Spill Science & Technology Bulletin*, 5: 3-16.
- 775 Reinharz, E. and Michel, J. 1996. Preassessment Phase Guidance Document for Natural Resources
776 Damage Assessment under the Oil Pollution Act of 1990. Prepared for the Damage Assessment
777 and Restoration Program, National Oceanic and Atmospheric Administration, 1305 East-West
778 Highway, SSMC #4, Silver Spring, Maryland. 47 pp. + appendices. (NOAA's Damage assessment,
779 remediation & restoration program. Adverse effects of oil.)

- 780 Reiss, H., Hoarau, G., Dickey-Collas, M. and Wolff, W.J. 2009. Genetic population structure of marine
781 fish: mismatch between biological and fisheries management units. *Fish and Fisheries* 10: 361–
782 395.
- 783 Rönkkönen, S., Ojaveer, E., Raid, T. and Viitasalo, M. 2004. Long-term changes in Baltic herring
784 (*Clupea harengus membras*) growth in the Gulf of Finland. *Can. J. Fish. Aquat. Sci.* 61: 219-229.
- 785 Romakkaniemi, A. *Editor* 2015. Best practices for the provision of prior information for Bayesian
786 stock assessment. ICES Cooperative Research Report No. 328: 93 pp.
- 787 Rönnerberg, C. and Bonsdorff, E. 2004. Baltic Sea eutrophication: area-specific ecological
788 consequences. *Hydrobiologia* 176: 227-241.
- 789 Smith, V.H. 2003. Eutrophication of freshwater and coastal marine ecosystems. A global problem.
790 *Environ. Sci. & Pollut. Res.* 10: 126-139.
- 791 Sparholt, H. 1994. Fish species interactions in the Baltic Sea. *Dana* 10: 131-162.
- 792 Stephenson, R.L. 1999. Stock complexity in fisheries management: a perspective of emerging issues
793 related to population sub-units. *Fish. Res.* 43: 247-249.
- 794 Stephenson, R., Peltonen, H., Kuikka, S., Pönni, J., Rahikainen, M., Aro, E. and Setälä, J. 2001. Linking
795 biological and industrial aspects of the Finnish commercial herring fishery in the northern Baltic
796 Sea. *In* Herring: Expectations for a new millennium. *Edited by* Funk, F., Blackburn, J., Hay, D., Paul,
797 A.J., Stephenson, R., Toresen, R. and Witherell, D. University of Alaska Sea Grant, AK-SG-01-04,
798 Fairbanks, p 741-760.
- 799 Underwood, A.J. 1997. Experiments in ecology. Cambridge University Press. 504 pp.
- 800 Urho, L. 1991. Impact of an oil spill on herring stock. - International Herring Symposium, Anchorage,
801 Alaska USA, October 23-25, 1990. Lowell Wakefield Fisheries Symposium (9th). Alaska Sea Grant
802 Report 91-01. Proceedings of the International Herring Symposium, p. 569-582

- 803 Urho, L. and Hildén, M. 1990. Distribution patterns of Baltic herring larvae, *Clupea harengus* L., in the
804 coastal waters off Helsinki, Finland. J. Plankton Res. 12: 41-54.
- 805 Urho, L., Kjellman, J. and Pelkonen, T. 2003. Eutrophication and herring reproduction success in the
806 northern Baltic Sea. ICES Mar. Sci. Symp. 219: 430-432.
- 807 Vanhatalo, J., Pietiläinen, V. and Vehtari, A. 2010. Approximate inference for disease mapping with
808 sparse Gaussian processes. Statistics in Medicine 29: 1580-1607.
- 809 Vanhatalo, J., Riihimäki, J., Hartikainen, J., Jylänki, P., Tolvanen, V. and Vehtari, A. 2013. GPstuff :
810 Bayesian Modeling with Gaussian Processes. Journal of Machine Learning Research 14: 1175–
811 1179.
- 812 Viitasalo, M., Vuorinen, I. and Saesmaa, S. 1995. Mesozooplankton dynamics in the northern Baltic
813 Sea: implications of variations in hydrography and climate. J. Plankton Res. 17: 1857-1878.
- 814 Vuorinen, I., Hänninen, J., Viitasalo, M., Helminen, U. and Kuosa, H. 1998. Proportion of copepod
815 biomass declines with decreasing salinity in the Baltic Sea. ICES J. Mar. Sci. 55: 767–774.
- 816 Zillén, L., Conley, D.J., Andrén, T., Andrén, E., Björck, S. 2008. Past occurrences of hypoxia in the
817 Baltic Sea and the role of climate variability, environmental change and human impact. Earth-Sci.
818 Rev. 91: 77-92.

Table 1. Prior probabilities of additional mortality in two population sub-groups of herring conditioned on M/T Antonio Gramsci accident (oil type: heavy oil, proportion of oiled coast line: 7-15%, season: spring, species: pelagic fish). Mortality denotes the fraction of population killed immediately by oil in 1987.

mortality	developmental stage	
	offspring	older ages
< 20%	0.75	0.9
20-50%	0.23	0.1
50-80%	0.02	0
>80%	0	0

819

820

Draft

Table 2. The prior and posterior distributions for the parameters of the herring stock dynamics model. The indicated posterior location parameter is mean for variables with Gaussian prior, otherwise we report posterior median. We use standard deviation as the shape parameter.

prior distribution	interpretation of the parameter	affecting	posterior mean/median	posterior 95% PI
parameters related to environmental covariates				
$\text{optchl} \sim N(3, 1)$	optimum chl- α concentration, $\mu\text{g/l}$	recruitment	4.73	[3.1, 6.4]
$\text{sd} \sim N(6, 0.58)$	shape parameter for the width of the chl- α parabola	recruitment	6.56	[5.5, 7.6]
$b \sim N(1, 0.71)$	influence parameter for April- June SST	recruitment	0.76	[0.0003, 1.519]
OilMorE, Table 1.	additional mortality caused by the oil spill	offspring survival	0.17	[6.04e-06, 0.569]
OilMorA, Table 1.	additional mortality caused by the oil spill	adult survival	0.09	[8.94e-06, 0.217]
$\text{cLS} \sim N(-0.25, 0.03)$	influence parameter of sprat abundance	growth	-0.24	[-0.29, - 0.19]
$\text{ck} \sim N(0.25, 0.03)$	influence parameter of August- October SST	growth	0.25	[0.19, 0.31]
τ_y , Eq. 10-14	influence function of salinity	growth	See fig. 4.	
$L_{50} \sim N(5.5, 0.55)$	salinity where 50% of herring have an increased growth rate	growth	5.1	[4.3, 5.9]
$\text{SR} \sim N(0.5, 0.1)$	a steepness parameter defining the increase rate of the logistic	growth	0.49	[0.29, 0.69]

	curve				
Me, Eq. 17-19	cod induced natural mortality	mortality	See fig. 8.		
$cM \sim LN(\ln(1 \cdot 10^{-5}), \ln(10))$	component independent of cod abundance and herring size	mortality	$4 \cdot 10^{-6}$	$[1 \cdot 10^{-6}, 8 \cdot 10^{-6}]$	
stock parameters					
$G \sim LN(\log(0.25), \sqrt{\log(0.1^2+1)})$	the relationship between size and M	natural mortality	0.30	$[0.25, 0.35]$	
$L_{inf} \sim N(25, 3)$	the mean maximum length	growth	28.36	$[24.6, 32.4]$	
K_y , Eq. 5	annual von Bertalanffy growth rate	growth	0.12	$[0.0, 0.38]$	

821

822

Draft

Figure captions

Fig. 1. Top panel: the Baltic Sea showing sampling sites in the Gulf of Finland for salinity (triangles) and sea surface temperature (stars). Bottom panel: predicted chlorophyll- α concentration ($\mu\text{g/l}$) in the GoF in July-August 2004 (see Section 2.3). Black dots represent locations of all sampling sites from 1974 to 2010. The annual chlorophyll- α index for the herring population model was calculated for the 10 km zone in the northern coast of GoF.

Fig. 2. The posterior of sea surface temperature (April-June (A) and August-October (B)), chl- α (C), and salinity (D). The solid line shows the posterior mean and the shaded area 95% central posterior credible interval. These posteriors were used as environmental priors in the herring stock dynamics model.

Fig. 3. The modeled linkages between the environment and the herring stock dynamics in the GoF. The ovals with grey edge feature observations - dashed edge indicates the covariates described by a probability distribution. The ovals with black edge are latent variables inferred by the model. Indirect causal relationship is illustrated with a dashed arrow. Oil spill is modeled to cause additional mortality from other reason than fishing in 1987. Acoustic index is available for 2004. The figure is not a directed acyclic graph (DAG) despite reminiscence of Bayesian net.

Fig. 4. Influence of the GoF chl- α concentration on herring recruitment (A) and the logistic relationship between salinity and herring growth rate (B). The blue lines are 100 random realizations of the prior distribution and the red ones realizations of the posterior distribution. The horizontal bar indicates the range of the posterior medians for chl- α concentration and salinity (see Fig. 2).

Fig. 5. The prior (bars) and posterior estimates (line) of the oil induced mortality in herring offspring (A) and older ages groups (B) in 1987. A posterior distribution is proportional to the product of a

likelihood and a prior. Hence, two humps in the posterior in the subplot A mean that the (model induced) marginal likelihood of the oil induced mortality ratio parameter is an increasing function of that parameter.

Fig. 6. The GoF herring stock-recruitment relationship. The red dots are annual posterior medians of spawned eggs and concomitant recruits. Black dots denote samples of the realized simulations. Blue lines are the estimated relationships. The downward bending lines are realizations of the Ricker model.

Fig. 7. Effect of the April-June sea surface temperature (SST) on the herring recruitment. The mean SST in 1980-2006 was 5.3 °C, range [3.7, 6.8]. The posterior estimates of parameter b are indicated in table 2. The dotted, solid, and dashed lines represent 2.5%, median, and 97.5% of the posterior estimates of b , respectively. The lines intersect at the mean SST as the effect at the mean SST was adjusted to value 1 (Eq. 4).

Fig. 8. Natural mortality rate caused by two components: size dependent mortality M at ages 1 (A), 5 (C), and 8 (D), and additional mortality caused by cod predation Me (B). Solid line illustrates posterior median and dashed lines 95% credible interval. Note the different scale for cod induced natural mortality.

Fig. 9. The posterior median and 95% credible interval for herring total biomass (A), eggs spawned (describing spawning stock size) (B), fishing mortality rate at ages 3-7 (C), and number of recruits (D) at the beginning of their second year.

Fig. 10. The posterior median (solid line) and 95% credible interval (dashed lines) for herring catch in the GoF in 1980-2006. The dots indicate observed catches.

Fig. 1. Top panel: the Baltic Sea showing sampling sites in the Gulf of Finland for salinity (triangles) and sea surface temperature (stars). Bottom panel: predicted chlorophyll- α concentration ($\mu\text{g/l}$) in the GoF in July-August 2004 (see Section 2.3). Black dots represent locations of all sampling sites from 1974 to 2010. The annual chlorophyll- α index for the herring population model was calculated for the 10 km zone in the northern coast of GoF.

Fig. 2. The posterior of sea surface temperature (April-June (A) and August-October (B)), chl- α (C), and salinity (D). The solid line shows the posterior mean and the shaded area 95% central posterior credible interval. These posteriors were used as environmental priors in the herring stock dynamics model.

Fig. 3. The modeled linkages between the environment and the herring stock dynamics in the GoF. The ovals with grey edge feature observations - dashed edge indicates the covariates described by a probability distribution. The ovals with black edge are latent variables inferred by the model. Indirect causal relationship is illustrated with a dashed arrow. Oil spill is modeled to cause additional mortality from other reason than fishing in 1987. Acoustic index is available for 2004. The figure is not a directed acyclic graph (DAG) despite reminiscence of Bayesian net.

Fig. 4. Influence of the GoF chl- α concentration on herring recruitment (A) and the logistic relationship between salinity and herring growth rate (B). The blue lines are 100 random realizations of the prior distribution and the red ones realizations of the posterior distribution. The horizontal bar indicates the range of the posterior medians for chl- α concentration and salinity (see Fig. 2).

Fig. 5. The prior (bars) and posterior estimates (line) of the oil induced mortality in herring offspring (A) and older ages groups (B) in 1987. A posterior distribution is proportional to the product of a likelihood and a prior. Hence, two humps in the posterior in the subplot A mean that the (model induced) marginal likelihood of the oil induced mortality ratio parameter is an increasing function of that parameter.

Fig. 6. The GoF herring stock-recruitment relationship. The red dots are annual posterior medians of spawned eggs and concomitant recruits. Black dots denote samples of the realized simulations. Blue lines are the estimated relationships. The downward bending lines are realizations of the Ricker model.

Fig. 7. Effect of the April-June sea surface temperature (SST) on the herring recruitment. The mean SST in 1980-2006 was 5.3 °C, range [3.7, 6.8]. The posterior estimates of parameter b are indicated in table 2. The dotted, solid, and dashed lines represent 2.5%, median, and 97.5% of the posterior estimates of b , respectively. The lines intersect at the mean SST as the effect at the mean SST was adjusted to value 1 (Eq. 4).

Fig. 8. Natural mortality rate caused by two components: size dependent mortality M at ages 1 (A), 5 (C), and 8 (D), and additional mortality caused by cod predation M_e (B). Solid line illustrates posterior median and dashed lines 95% credible interval. Note the different scale for cod induced natural mortality.

Fig. 9. The posterior median and 95% credible interval for herring total biomass (A), eggs spawned (describing spawning stock size) (B), fishing mortality rate at ages 3-7 (C), and number of recruits (D) at the beginning of their second year.

Fig. 10. The posterior median (solid line) and 95% credible interval (dashed lines) for herring catch in the GoF in 1980-2006. The dots indicate observed catches.

Appendix

Temporal model for temperature and salinity data

The models for temperature and salinity are identical so we concentrate only on the former. The measured temperature $y(t)$, on day t , is assumed to be a noisy realisation of an underlying true temperature

$$y(t) = f(t) + \varepsilon_i$$

where $\varepsilon_i \sim N(0, \sigma^2)$. We are interested in the underlying function $f(t)$ which is given a Gaussian process (GP) prior (Rasmussen and Williams, 2006, Gelfand *et.al.*, 2010). Giving a GP prior for the function implies that any set of function values $\mathbf{f} = (f(t_1), \dots, f(t_n))$ has a multivariate normal distribution

$$\mathbf{f} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

where $\boldsymbol{\mu}$ is the mean and $\boldsymbol{\Sigma}$ the covariance. Here we will use a zero mean GP since we centre the observations to have zero mean before modelling. The covariance matrix $\boldsymbol{\Sigma}$ incorporates the temporal association between different observations and can be represented as a function that describes the correlation between pairs of points with distance in time. We will assume that the covariance function is a sum of three components: a periodic, a long-term and a short-term component. The periodic covariance function is constructed following Rasmussen and Williams (2006, page 118)

$$k_p(t_i, t_j) = \sigma_p^2 \exp\left(-\frac{2\sin^2(\pi(t_i - t_j)/T)}{l_p}\right)$$

where T is the period (here 365 days), σ_p^2 the magnitude governing the overall variability of the function and l_p the length-scale parameter which governs the smoothness of the periodic function. The length-scale and magnitude are given a half Student-t prior which works as a weakly informative prior (Vanhatalo *et al.* 2010). The long and short term trends are modelled with a squared exponential covariance function

$$k_{se}(t_i, t_j) = \sigma_{se}^2 \exp\left(-\frac{(t_i - t_j)^2}{l_{se}^2}\right)$$

where σ_{se}^2 is the magnitude and l_{se}^2 is the length-scale parameter. In the squared exponential covariance function the length-scale governs how fast the correlation between time points decreases as their timely difference increases. We give a uniform prior for the length-scale of the long-term trend and initialise it in high value, whereas the short-term component is given a half Student-t prior, with mode 0, scale 3 and 4 degrees of freedom, and initialised to a small value. The total prior covariance can then be written as.

$$\Sigma(t_i, t_j | \theta) = k_p(t_i, t_j | \theta) + k_{se}^{long}(t_i, t_j | \theta) + k_{se}^{short}(t_i, t_j | \theta)$$

where θ denotes a vector that collects all the parameters in the covariance function $\Sigma(t_i, t_j | \theta)$ and the noise variance σ^2 .

The inference for the GP models was conducted as follows. The posterior distribution of the covariance function parameters was approximated by a maximum a posterior (MAP) estimate

$$\hat{\theta} = \arg \max_{\theta} p(\mathbf{y} | \theta) p(\theta),$$

where $p(\theta)$ is the prior distribution for the covariance parameters, $p(\mathbf{y} | \theta) = p(\mathbf{y} | \mathbf{f}, \theta) p(\mathbf{f} | \theta) d\mathbf{f}$ is the marginal likelihood of the covariance parameters and $p(\mathbf{f} | \theta) = N(\mathbf{f} | \mu, \Sigma)$ is the prior for the latent function values. After solving the MAP estimate for the covariance function parameters the (conditional) posterior distribution of function values $\tilde{\mathbf{f}} = [f(\tilde{t}_1), \dots, f(\tilde{t}_N)]$ at time points $\tilde{t}_1, \dots, \tilde{t}_N$ is a multivariate Gaussian distribution (Rasmussen and Williams, 2006)

$$\tilde{\mathbf{f}} | \mathbf{y}, \theta \sim N(\mathbf{K}_{\tilde{\mathbf{f}}, \mathbf{f}} (\Sigma + \sigma^2 \mathbf{I})^{-1} \mathbf{y}, \mathbf{K}_{\tilde{\mathbf{f}}} - \mathbf{K}_{\tilde{\mathbf{f}}, \mathbf{f}} (\Sigma + \sigma^2 \mathbf{I})^{-1} \mathbf{K}_{\tilde{\mathbf{f}}, \mathbf{f}}^T),$$

where $\mathbf{K}_{\tilde{\mathbf{f}}, \mathbf{f}}$ is the covariance matrix between $\tilde{\mathbf{f}}$ and $\mathbf{f}$, and $\mathbf{K}_{\tilde{\mathbf{f}}}$ is the covariance matrix of $\tilde{\mathbf{f}}$.

In order to evaluate the posterior distribution of the average temperature over time period from May to June, A , we need to integrate $f(t)$ over that time period, $\bar{f}(A) = 1/|A| \int_A f(t) dt$. This results in a classical

change of support problem (Gelfand et al. 2010), since our model was constructed for time points and we are now interested in time intervals. The covariance between $f(t)$ and $\bar{f}(A)$ is

$$\Sigma(t, A) = \int_A \Sigma(t, u) du.$$

Thus we can predict directly for the time intervals using the standard GP formalism with this integrated covariance function. In practice the integral cannot be solved analytically for which reason we discretized the time interval A to daily resolution and approximated the integral with a finite sum over values of $f(t)$ in A . We used Monte Carlo method to approximate the posterior distribution for this sum by sampling 1000 discrete sets of $f(t)|\mathbf{y}, \theta$ values from their joint posterior.

Spatio-temporal model for chlorophyll-a

For the chlorophyll- α concentrations, the covariance structure was appended with a spatial component. The observed average July-August chlorophyll- α concentration, $y(s, t)$, at location s in year t , is assumed to be a noisy realisation of an underlying true average July-August chlorophyll-a concentration

$$y(s, t) = f(s, t) + \varepsilon_i$$

where $\varepsilon_i \sim N(0, \sigma^2/n_i^2)$ is the observation error. Since the number of averaged measurements, n_i , per observation varies we scaled the variance of individual measurements accordingly. $f(s, t)$ is given a GP prior with zero mean and separable covariance function

$$k([s, t], [s', t']) = k_s(\|s - s'\|)k_t(\|t - t'\|) = \sigma_e^2 \exp(-\|s - s'\|) \exp(-\|t - t'\|),$$

where $\|\cdot\|$ is a scaled Euclidean distance, e.g., $\|s - s'\| = \sqrt{\sum_{i=1}^2 (s_i - s'_i)^2 / l_i^2}$, and l_i is the length-scale along the co-ordinate axis s_i .

In order to evaluate the average chlorophyll- α concentration for the sea area within 10 km off the Finnish shoreline, S , we need to integrate $f(s, t)$ over that region, $\bar{f}(S, t) = 1/|S| \int_S f(s, t) ds$. We used a Monte

Carlo approximation for this so that we predicted 1000 sets of $f(s, t)$ values on a lattice grid spanning the GoF, after which we took an averages over the grid cells in that region.

Draft