Canadian Journal of Forest Research
Revue canadienne de recherche forestière

Comparative growth trends of five northern hardwood and montane tree species reveal divergent trajectories and response to climate

Journal:	<i>Canadian Journal of Forest Research</i>
Manuscript ID	cjfr-2016-0308.R2
Manuscript Type:	Article
Date Submitted by the Author:	06-Feb-2017
Complete List of Authors:	Kosiba, Alexandra; University of Vermont, Rubenstein School of Environment and Natural Resources Schaberg, Paul; USDA Forest Service Rayback, Shelly; University of Vermont, Geography Hawley, Gary; University of Vermont, Rubenstein School of Environment and Natural Resources
Keyword:	red spruce, sugar maple, yellow birch, red maple, balsam fir
Is the invited manuscript for consideration in a Special Issue? :	

SCHOLARONE™
Manuscripts

Comparative growth trends of five northern hardwood and montane tree species reveal divergent trajectories and response to climate

Alexandra M. Kosiba^{a*}, Paul G. Schaberg^b, Shelly A. Rayback^c, and Gary J. Hawley^a

^a The University of Vermont, Rubenstein School of Environment and Natural Resources, Burlington, VT 05405, USA, p: +1 (802) 656-2911, f: +1 (802) 656-8683, akosiba@uvm.edu, ghawley@uvm.edu

^b Forest Service, U.S. Department of Agriculture, Northern Research Station, Burlington, VT 05405, USA, p: +1 (802) 656-1715, f: +1 (802) 656-8683, pschaberg@fs.fed.us

^c The University of Vermont, Department of Geography, Burlington, VT 05405, USA, p: +1 (802) 656-3019, f: +1 (802) 656-3042, srayback@uvm.edu

* Corresponding author. *E-mail address:* akosiba@uvm.edu (A.M. Kosiba).

Abstract: In the northeastern US, tree declines associated with acid deposition-induced calcium depletion have been documented, notably for red spruce (*Picea rubens* Sarg.) and sugar maple (*Acer saccharum* Marsh.). There is conflicting evidence if co-occurring tree species have capitalized on these declines or suffered similar growth reductions; and, how growth has fluctuated relative to environmental variables. We examined five species along three elevational transects on Mt. Mansfield, Vermont: sugar maple, red spruce, red maple (*Acer rubrum* L.), yellow birch (*Betula alleghaniensis*, Britton), and balsam fir (*Abies balsamea*, [L.] Mill.). We found baseline differences in growth. Red maple and yellow birch had the highest growth, sugar maple and red spruce intermediate, and balsam fir the lowest. While some of year-to-year declines were associated with specific stress events, protracted patterns, such as recent increases in red spruce and red maple growth, were correlated with increased temperature and cooling degree days (heat index). For most species and elevations, there was a positive association between temperature and growth, but a negative association with growth the following year. Based on our comparisons, for some species growth at Mt. Mansfield aligns with regional trends and suggests that patterns assessed here may be indicative of the broader region.

Key words: *red spruce*, *sugar maple*, *yellow birch*, *red maple*, *balsam fir*

36 Introduction

37 Regional forests in the northeastern United States (US) reflect temporally and
38 spatially complex land use change in the 19th and 20th centuries that included
39 deforestation, agricultural expansion, and subsequent reforestation (Whitney 1994).
40 Resulting second growth forests were then subjected to novel anthropogenic stress in the
41 20th century – notably acid deposition-induced nutrient perturbations that altered forest
42 health and productivity (DeHayes et al. 1999; Greaver et al. 2012). In particular, acidic
43 deposition and resulting calcium (Ca) depletion have been associated with health and
44 productivity declines for red spruce (*Picea rubens* Sarg.) (DeHayes et al. 1999) and sugar
45 maple (*Acer saccharum* Marsh.) (Schaberg et al. 2006) – two dominant and economically
46 important tree species in the northeastern US. Added to the stress of pollutants and
47 decades of soil cation depletion, anthropogenic emissions of carbon dioxide (CO₂) and
48 other greenhouse gases have induced changes in the global climate. Greenhouse gas
49 accumulations have led to an increase in the annual mean temperatures in the
50 northeastern US of 0.09°C decade⁻¹ (1895-2011), resulting in a lengthening of the freeze-
51 free period and functional growing season (Kunkel et al. 2013). While precipitation has
52 not significantly changed for winter, spring, or summer, fall and annual precipitation
53 totals have increased 0.61 and 0.99 cm decade⁻¹, respectively (Kunkel et al. 2013).
54 Continued changes in temperature and precipitation, alone and in combination with other
55 stressors, could push trees beyond the environmental thresholds to which they are
56 adapted and result in reduced competitive capacity, growth, and survival for some species
57 compared to others.

Whereas the declines of several tree species associated with acid deposition are well documented in the region, research suggests that some of these chronically stressed species are experiencing contemporary growth increases (Kosiba et al. 2013) and possible range expansion (Beckage et al. 2008; Foster and D'Amato 2015). Both deviations are hypothesized to be associated with changes in environmental factors, such as elevated atmospheric CO₂ concentrations, fluctuating atmospheric pollution, and rising temperature. Elevated CO₂ has been proposed to cause a fertilization effect in some trees (Salzer et al. 2009; Soule and Knapp 2006), but other findings have not supported this (Bader et al. 2013; Girardin et al. 2016; Körner et al. 2005). While acid deposition initially caused declines in some tree species, the enforcement of the Clean Air Act and subsequent amendments has reduced sulfur (S) and, less dramatically, nitrogen (N) deposition (Burns et al. 2011). Removal of this source of damage could allow for a recovery for chronically stressed species and, further, N deposition can promote growth, particularly in N-limited ecosystems (Aber et al. 1998). Rising temperatures could also extend the functional growing season, especially for temperate conifers that retain foliage year-round (Kosiba et al. 2013).

While anthropogenic stress continues to mount within northeastern forests, there is limited evidence of clear trends in forest response; yet, large alterations in tree growth rates could have profound effects on the ecology of these forests, including changes to carbon (C) pools, biogeochemical and hydrological cycles, and species composition. These fluctuations could alter important ecosystem services, including ones with direct economic impacts, such as wood products and recreation.

Here, we present growth patterns and trends determined using xylem annual increment measurements for five tree species that characterize the northern hardwood and montane spruce-fir forests: sugar maple, red spruce, red maple (*Acer rubrum* L.), yellow birch (*Betula alleghaniensis*, Britton), and balsam fir (*Abies balsamea*, [L.] Mill.) growing along elevational transects on Vermont's tallest mountain: Mt. Mansfield. We hypothesized that trees would have different growth rates dependent on species, elevation, and year, with some species displaying muted growth in recent decades. We expected that sugar maple, a species that has experienced geographically broad decline since the 1980s (Horsley et al. 2002), and yellow birch, a species with limited reports of decline (van Doorn et al. 2011), would have reduced recent growth. Conversely, we predicted that red spruce at Mt. Mansfield would show recent increases in growth synchronous with a regional growth pattern of unknown cause (Kosiba et al., 2013). We expected that sympatric balsam fir may have experienced competitive reductions in growth concurrent with increases in red spruce growth. Although red maple stocking and growth in the understory has increased regionally (Abrams 1998), growth trends for dominant and co-dominant red maple are unresolved.

By associating the nature and timing of observed growth relative to local weather and pollution data, we assessed how putative environmental drivers may have influenced species growth and productivity over time. Specifically, we anticipated that deposition of S and N would be negatively associated with red spruce and sugar maple growth, particularly in the 1970s-90s when deposition was the greatest. We hypothesized that atmospheric CO₂ concentration, summer temperature, and growing season length would be positively associated with growth and that temperature relationships would strengthen

with increasing elevation. We expected that increased fall and spring temperatures would be beneficial to evergreen conifers, as they can photosynthesize outside the traditional growing season when sympatric deciduous trees are leafless. We also predicted that metrics of precipitation would have a variable association with tree growth at Mt. Mansfield considering that these trees persisted through severe droughts in the 1930s and 60s (Dupigny-Giroux 2002) as well as a current pluvial (Pederson et al. 2013).

Methods

Site description

Mt. Mansfield State Forest, part of the northern Appalachian mountain chain, located in north-central VT (Underhill, VT, USA; 44.5439° N, 72.8143° W), is a 18 000 hectare parcel comprised of multiple forest types across an altitudinal range of approximately 1 000 m, with a summit at 1 339 m. A northern hardwood forest, dominated by sugar and red maple, American beech (*Fagus grandifolia* Ehrh.), and yellow birch, extends to ~800 m, where it merges in a transition zone with a montane spruce-fir forest, primarily composed of balsam fir and red spruce. Red spruce, a temperate conifer, is frequently found intermixed in the northern hardwood ecotone. The dynamics of the forests are controlled by localized wind events and individual tree mortality; however, historical timber harvest occurred sporadically in the early 20th century through parts of the Mt. Mansfield State Forest (Cogbill 1996). Soils are primarily stony podzols with considerable areas of rock outcrops, particularly approaching the summit (Soil Survey Staff 2015).

The climate of Mt. Mansfield State Forest is continental, encompassing a large temperature range (Fig. S1) and is affected by its proximal location to Lake Champlain, a 126 910 hectare lake 24 km west of the study site, which moderates temperature and increases snow fall amounts. The annual mean temperature is 4.2°C with an average of 154.6 cm of precipitation deposited uniformly throughout the year (Fig. S1) (PRISM Climate Group 2004). January is the coldest month and July is the warmest (-10.5°C and 17.6°C average temperature 1925-2012 for study location, respectively) (PRISM Climate Group 2004). Continuous snow cover is the winter norm, persisting over 5 months at higher elevations. Humidity and water availability increase with elevation.

Plot selection

Elevational transects were set up in three of the four watersheds on Mt. Mansfield (Brown's River, Stevensville Brook, and Ranch Brook Watersheds). No transect was established in the fourth watershed because this area has experienced anthropogenic disturbance associated with a commercial ski area. Along each transect, three plots were selected – one within each of the following elevational zones: low (450-650 m a.s.l.), mid (750-850 m) and high (900-1000 m) (n plots = 9, Fig. 1), which align with northern hardwoods, transition, and montane spruce-fir ecotones. No obvious stand mortality or substantial recent disturbance was evident in any of the plots.

Plots contained 10-14 dominant or co-dominant trees of each of the target tree species equally distributed around plot center to avoid differing competition pressures among trees. Trees with obvious bole or crown damage or those growing in anomalous conditions were not selected. We sampled red maple, sugar maple, and red spruce at low

elevation; sugar maple, yellow birch, and red spruce at mid elevation; and red spruce and balsam fir at high elevation (Fig. 1). Due to differential species densities across the landscape and different number of species sampled per plot, plots were of variable radius (approx. 20-35 m).

We selected species if they were the dominant components of the montane spruce-fir (red spruce and balsam fir) or northern hardwood (sugar maple and yellow birch) forests, or if the species has experienced increased dominance in this latter forest type (red maple; Abrams 1998). A dominant component of the northern hardwood forest, American beech, was not assessed because its growth dynamics have been altered following widespread damage from the beech bark disease complex (Gavin and Peart 1993).

Dendrochronology

We increment cored selected trees ($n = 256$) following standard dendrochronological techniques (Stokes and Smiley 1968) in the fall of 2012. We collected two 5 mm increment cores per tree at stem DBH (diameter at breast height, 1.37 m above ground level) at 180° and perpendicular to the slope. Cores were air-dried, sanded with progressively finer grit sandpaper (ranging from 100-1500 grit dependent on species) and visually crossdated using the list method (Yamaguchi 1991). We microscopically measured rings to 0.001 mm resolution using a Velmex sliding stage unit with MeasureJ2X software (VoorTech Consulting, Holderness, NH) and used the computer program COFECHA to detect and correct for potential crossdating errors in ring series (Holmes 1983). Individual cores were discarded if they were poorly correlated

with the master chronology (i.e., below Pearson critical correlation level of 0.328 [99% CL]). For descriptive purposes, tree age (at breast height) was calculated using the maximum number of rings per tree if pith was evident or was estimated per core using a pith indicator (Speer 2010) if pith was not reached. For all trees, we were able to estimate age from at least one core.

For the climate analyses, we detrended, standardized, and prewhitened all raw ring width series and computed biweight robust mean chronologies per species per elevation. Descriptive statistics (sample sizes, $\bar{R}$, EPS [expressed population signal] and SNR [signal:noise]) of resulting chronologies were calculated using a running 30-year window with 15-year overlap using Spearman correlation coefficients (Table S1) and used to select the best detrending and standardization technique. For all series chronologies, a cubic smoothing spline (CSS) (dplR package in R), using a frequency response cutoff of 0.5 was used. If this spline was a poor fit, a more conservative horizontal line was fit to the data. Traditionally, chronologies are truncated at the year when the EPS value falls below 0.85 (Speer 2010), but this would preclude the use of a portion of both the red maple and sugar maple chronologies at low elevations due to their young age and comparatively small sample size. Because results here were solely used to investigate growth-climate or –deposition relationships rather than for climatic reconstructions, we reduced the threshold to 0.80, allowing for the use of a common period spanning from 1925-2012. However, lower EPS values indicate that these trees have a weaker stand-wide signal and may be more influenced by micro-site factors, which could mask a cohesive stand-wide signal. Summary statistics and data on the nine plots and resulting chronologies can be found summarized by species and elevational

zone in Table S2. Comparisons of growth among species and elevations was done with Kruskal-Wallis rank sums test followed by the Wilcoxon method for pairwise comparisons, as data did not meet the assumptions of equal variance ($P \leq 0.05$).

We also averaged raw ring widths by tree and converted measurements into basal area increment (BAI, $\text{cm}^2 \text{ year}^{-1}$) assuming a circular outline of stem cross sections (Cook and Kairiukstis 1990) and accounting for bark thickness. BAI is considered a more meaningful indicator of tree growth from a physiological standpoint because it provides an indication of annual stemwood production while accounting for the effects of stem geometry on radial growth associated with tree-maturation (Hornbeck and Smith 1985; LeBlanc 1992; West 1980). We then computed biweight robust mean BAI chronologies per species per elevation using the dplR package (Bunn et al. 2015) for R (Version 3.1.1) (R Development Core Team 2015) to moderate the effect of large BAI values and/or outliers on the mean chronology.

Growth associated with climate, deposition

Climate data (maximum and minimum monthly temperature [$T_{\min}$, $T_{\max}$] and total monthly precipitation [P]) from 1925-2012 were obtained from the PRISM Climate Group (PRISM Climate Group 2004). To reduce the number of variables assessed and limit the occurrence of type II errors, we computed seasonal mean (T_{mean}), maximum, and minimum temperatures (Winter: previous December–February, Spring: March–May, Summer: June–August, Fall: September–November), annual water year (previous [p] October to current September) mean temperature, and seasonal and water year precipitation totals. Using monthly data, we also calculated the average temperature for

217 an extended growing season (May–August and June–September) (Fig. S1).

218 Pollutant S and N deposition data by water year were obtained from the Hubbard
219 Brook Experimental Forest (HBEF; Watershed 1), Thorton, NH, USA, spanning 1965-
220 2010 (Likens 2010) (Fig. S1). As this dataset covers more years than the one from Mt.
221 Mansfield (1984-2012) (National Atmospheric Deposition Program 2014) and the
222 datasets were highly correlated ($R = 0.94$, $P < 0.0001$), we combined them by adjusting
223 HBEF data via regression analysis and then added this to the Mt. Mansfield dataset for
224 continuous coverage from 1965-2012. We gathered data of annual atmospheric CO₂
225 concentration from Mauna Loa Observatory, HI, USA (NOAA Earth System Research
226 Laboratory Global Monitoring Division 2013) and from Law Dome, Antarctica ice core
227 data (World Data Center for Paleoclimatology and NOAA Paleoclimatology Program);
228 combined for a CO₂ dataset spanning 1925-2012. We also collected the following data:
229 Vermont Palmer Drought Severity Index (PDSI) (National Drought Mitigation Center
230 2014), and three measures of degree days: growing degree days (GDD, measured as
231 cumulative degrees $> 5^{\circ}\text{C}$ [41°F]), cooling degree days (CDD, $> 18.3^{\circ}\text{C}$ [65°F]) and
232 heating degree days (HDD, $< 18.3^{\circ}\text{C}$ [65°F])) for Burlington International Airport, VT
233 (NOAA National Weather Service 2014), and monthly Standardized Precipitation-
234 Evapotranspiration Index (SPEI, 1-month) for VT, a multi-scalar drought and
235 temperature index that includes the effects of evapotranspiration (The National Drought
236 Mitigation Center 2012). These datasets were chosen based on proximity or length of
237 record; these datasets covered 1945-2012 (Fig. S1).

238 Growth relationships with climate, deposition, and other data were assessed using
239 *treeclim* (Zang and Biondi 2015) for R. *Treeclim* uses time-dependent bootstrapped

resampling (1000 iterations) to test for linear correlations between the residual ring width data and each subvector of the climate matrix (Zang and Biondi 2015). To investigate the dominant drivers of tree growth, we first examined stationary correlations over the common period (or a limited subset for climate/deposition datasets spanning fewer years), including previous year's climate and deposition on current year's growth. Second, to assess the temporal stability of climate-growth relationships, we used the significant variables from the first analysis to evaluate relationships with growth per temporal quartile. Third, we used principal component analysis to examine common modes of growth among species per elevation. Using the first principle component (PC) per elevation, we again examined stationary correlations with our climate and deposition variables. For all analyses, Spearman correlation coefficients were computed with a 99% CI to reduce the number of input variables, type II errors, and covariation among variables.

Lastly, we used the significant climate and pollution deposition variables from the first analysis to model growth per species and elevation. All variables were first standardized ($\mu = 0$, $\sigma = 1$). If multicollinearity was evident among variables, we used forward stepwise linear regression (using AIC criteria) to reduce selected variables. For all species and elevation pairs, we then created linear models to best explain growth; non-significant variables were removed and the model was reassessed. Using the residuals of these models, we then regressed CO₂ data to examine if, after accounting for the dominant climate effects, there was a detectable CO₂ affect (Girardin et al. 2016).

Comparisons with other chronologies and datasets

We used other crossdated (± 0 yrs.) tree ring chronologies collected by the authors or affiliates (Engel et al. 2016; Hansen 2015; Weverka 2012) to compare growth trends on Mt. Mansfield to other locations in the region using Pearson Product-Moment Correlation. All chronologies were collected following similar methods described above and converted into BAI. Due to age limits of the chronologies from the HBEF, we constrained analyses to 1950-2012 for sugar maple and yellow birch (Hansen 2015).

Results

Absolute growth

We found significant and consistent growth differences among the species ($P < 0.001$) (Fig. 2A). Yellow birch and red maple had the largest mean growth (mean BAI $\pm$ SD: 15.2 ± 5.1 and 13.0 ± 4.3 cm², respectively). Sugar maple and red spruce were intermediate in growth (7.1 ± 2.0 and 7.7 ± 2.9 cm², respectively) and balsam fir had the lowest growth (4.7 ± 2.4 cm²). These patterns were fairly stable over the quartiles of the chronology, excluding the first quartile (1925-1946) and the most recent growth (2002-2012).

Growth was consistently higher at low and mid elevations compared with high elevation (Fig. 2B) ($P < 0.001$). For the chronology overall and for the central two quartiles (1947-1968, 1969-1990), growth was greater at mid elevations than low elevation. For all three elevations, overall growth increases in the most recent decade were driven by increases in two species: red spruce and red maple.

For the two species sampled at multiple elevations (sugar maple and red spruce), species-specific growth across elevations showed differing patterns (Fig. 3). Sugar maple

at low elevation had significantly higher mean growth than at mid elevation (8.1 ± 2.6 and $6.5 \pm 1.7 \text{ cm}^2$, respectively; $P < 0.01$), juxtaposing the general patterns that mid elevation plots had higher mean BAI overall. For both elevations, sugar maple maximum growth was only slightly higher than the mean and occurred in the third quartile: 1985-95 for low elevation and 1970-90 for mid elevation trees. Following this peak, both chronologies exhibited slight declines in growth. Mid elevation red spruce had the highest average growth ($13.0 \pm 4.3 \text{ cm}^2$), with low elevation intermediate ($7.8 \pm 3.1 \text{ cm}^2$), and high elevation the lowest ($4.1 \pm 3.2 \text{ cm}^2$) (Fig. 3) ($P < 0.01$). For red spruce at all elevations, maximum growth occurred in 2012 (year of sampling), and was considerably higher than overall chronology means. At both mid and high elevations, four out of five years of maximum red spruce growth occurred in the years immediately prior to sampling (2009-12).

Growth trends

Species-to-species comparisons per elevation (Fig. 3) reveal that peak growth varied, but for all species, the highest growth years were in the second half of the chronology (e.g., 1960s to 2012). At low elevation, both red spruce and red maple displayed recent increases in growth, while sugar maple did not. For all three species at low elevation, peak growth roughly coincided; although for both red maple and red spruce, maximum growth persisted longer and was more recent than for sugar maple. At mid elevation, highest growth for yellow birch and sugar maple corresponded (1960s-80s), yet red spruce peak growth occurred when these neighboring species were showing

decreases. At high elevation, balsam fir experienced maximum growth at the end of the 1980s through the early 2000s.

The recent growth surge of red spruce was not seen in the other species at mid elevation. While this increase was most pronounced for low and mid elevation red spruce, red spruce growing at high elevation, where growth over the length of the chronology was more stable and constrained compared to lower elevation red spruce, also exhibit this upward trend. At all elevations, red spruce exhibited periods of depressed growth beginning in the mid-twentieth century that align with suspected or known winter injury events (Johnson et al. 1986; Lazarus et al. 2004).

Climate and deposition relationships with growth

Examining climate-growth relationships using the principal components (PCs) derived from chronologies per elevation, we saw that overall, all three elevations display a positive correlation between growth and growing season temperature ($P < 0.01$, Table 1). At low elevation, Summer T_{mean} was best associated with growth ($\rho = 0.38$), along with NH_4^+ deposition ($\rho = 0.35$). Both mid and high elevations showed positive correlations with CDD and growing season temperatures (e.g., Spring, Summer T_{mean}). However, non-growing season temperatures and PDSI were important to high elevation trees only. A model of growth at both mid and high elevations identified CDD as the best predictor ($R^2_{\text{adj}} = 0.13$, $P = 0.001$ and $R^2_{\text{adj}} = 0.10$, $P = 0.04$, respectively). At low elevation, the growth model included a positive effect of May-Aug T_{mean} ($R^2_{\text{adj}} = 0.08$, $P = 0.005$). Atmospheric CO_2 concentration was not significantly associated with growth

for any of the species or elevations using either growth, PCs by elevation, or residuals (not shown).

At high elevation, balsam fir growth was positively correlated with temperature ($P < 0.01$; Table 2). The strongest positive correlations occurred with previous year's (p) Fall $T_{\min}$ ($\rho = 0.38$) and Spring $T_{\min}$ ($\rho = 0.24$), while there was a negative association with pCDD ($\rho = -0.37$) and pGDD ($\rho = -0.31$) (Table 2). Although significant overall ($P < 0.01$), the strength of the relationships between balsam fir growth and other variables declined in the middle of the chronology (1969-1990). The best model to predict balsam fir growth ($R^2_{\text{adj}} = 0.19$, $P = 0.0008$) included a negative effect of pCDD ($R^2_{\text{adj}} = 0.13$, $P = 0.002$) and a positive effect of Spring $T_{\min}$ ($R^2_{\text{adj}} = 0.05$, $P = 0.04$).

Red spruce exhibited the strongest positive relationships ($P < 0.01$) with temperature (CDD, GDD, and water year T_{mean}), including a strong correlation with non-growing season temperature (Table 2). Indeed, water year T_{mean} was positively associated with red spruce growth at all elevations and while not always significant, displayed a consistently positive correlation with growth across quartiles. Like the other species, pCDD was strongly, negatively correlated to red spruce growth, and this pattern persisted through time ($P < 0.01$). At both mid and high elevations, there was a positive association between growth and pWinter T_{max} ($\rho = 0.27$ and 0.23 , respectively). This relationship was negative in the third quartile, 1969-1990. Interestingly, pNO₃ deposition was negatively associated with growth of red spruce at low elevation ($P < 0.01$, $\rho = -0.37$), which strengthened through the quartiles of the chronology.

For red spruce at low elevation, we found that the negative effects of both HDD ($R^2_{\text{adj}} = 0.11$, $P < 0.0001$) and pCDD ($R^2_{\text{adj}} = 0.15$, $P < 0.0001$) were the best predictors of

353 growth in a linear model ($R^2_{\text{adj}} = 0.40$, $P > 0.0001$). At mid elevation, the best model (R^2_{adj}
 354 $= 0.31$, $P < 0.0001$) included a positive effect of water year T_{mean} ($R^2_{\text{adj}} = 0.13$, $P = 0.0006$)
 355 and, like low elevation, a negative effect of pCDD ($R^2_{\text{adj}} = 0.12$, $P = 0.0002$). At high
 356 elevation, the best model included a negative effect of both pCDD ($R^2_{\text{adj}} = 0.09$, $P =$
 357 0.0001) and HDD ($R^2_{\text{adj}} = 0.11$, $P = 0.018$), and a positive effect of GDD ($R^2_{\text{adj}} = 0.19$, P
 358 $= 0.017$) (overall model $R^2_{\text{adj}} = 0.35$, $P < 0.0001$).

359 Measures of degree-days and temperature were also important for red maple at
 360 low elevation ($P < 0.01$) (Table 2). Growing season temperature (i.e., Summer T_{mean} ,
 361 June-September T_{mean} , and CDD) was positively correlated with red maple growth ($P <$
 362 0.01 , $\rho = 0.30$, 0.26 , and 0.39 , respectively), while preceding year's heat accumulation
 363 (pCDD) was negatively associated ($\rho = -0.34$). For red maple, the best growth model
 364 ($R^2_{\text{adj}} = 0.20$, $P = 0.0004$) included a negative effect of pCDD ($R^2_{\text{adj}} = 0.11$, $P = 0.002$)
 365 and a positive effect of Summer T_{mean} ($R^2_{\text{adj}} = 0.08$, $P = 0.006$).

366 At low elevation, sugar maple growth was not correlated ($P > 0.01$) with the
 367 selected variables (Table 2); therefore, we were unable to create a model of growth. For
 368 sugar maple at mid elevation, Summer T_{mean} was positively associated with growth ($P <$
 369 0.01 , $\rho = 0.26$). However, this relationship was not consistent through the quartiles. Like
 370 many of the other species in this study, growth for these trees also exhibited a negative
 371 correlation ($P < 0.01$) with pSummer T_{mean} ($\rho = -0.25$) and pSummer T_{max} ($\rho = -0.27$), as
 372 well as with pCDD ($\rho = -0.38$). To model growth, we were limited by the number of and
 373 high collinearity of input variables. Both pSummer T_{mean} and pSummer T_{max} explained a
 374 similar amount of growth variation when fit separately ($R^2_{\text{adj}} = 0.06$, $P = 0.01$ for both).

We did not find any negative correlations between yellow birch growth and the variables selected (Table 2). Yellow birch, however, did display positive associations with summer T_{mean} ($P < 0.01$, $\rho = 0.29$) and these trends were consistent through the quartiles of the chronology. However, when we modeled growth using Summer T_{mean} , only a small amount of growth variation was explained ($R^2_{\text{adj}} = 0.07$, $P = 0.008$)

Comparisons with other chronologies

We compared red spruce growth from this study with red spruce from a wide range of plots in VT, New Hampshire, and Massachusetts (n trees = 452) collected in 2010, 2011, and 2012 (Engel et al. 2016; Kosiba et al. 2013; Weverka 2012). The chronologies were significantly correlated ($R = 0.92$, $P < 0.0001$; Fig. 4A). We also compared sugar maple and yellow birch from this study to those species from the HBEF collected in 2012 (Hansen 2015). For sugar maple (n trees HBEF = 163; Fig. 4B) the chronologies were significantly correlated ($R = 0.57$, $P < 0.0001$), but for yellow birch (n trees HBEF = 170; Fig. 4C) they were not ($R = -0.19$, $P = 0.13$).

Discussion

Patterns of growth

Significant and consistent growth differences among the species highlight their divergent silvics (Burns and Honkala 1990). Yellow birch and red maple, which are intermediate in shade tolerance with moderately high growth potentials (Burns and Honkala 1990), had the greatest mean growth. Sugar maple and red spruce, which are both shade tolerant with the ability to display higher growth when a mature component of

the canopy (Burns and Honkala 1990), were intermediate in growth, while balsam fir, highly shade tolerant and with the potential for constrained growth with elevated stocking (Burns and Honkala 1990), displayed the lowest growth. Species-to-species comparisons per elevation reveal that peak growth varied over time, but for all species, the highest growth years occurred in the second half of the chronology.

Lower growth and more muted disparities among the species in the first quartile of the chronology may signify that competition during stand maturation was a strong constraint on growth. Historical anthropogenic land clearing has left a legacy on the forests of Mt. Mansfield evident in the ages of sample trees (Table S1). In the absence of a substantial natural disturbance, it is likely that selective logging has resulted in the age divergence among the species and elevations. Cogbill (1996) outlined several logging events in the Stevensville Brook Watershed on the western flank of Mt. Mansfield, and it is suspected that widespread harvesting occurred in the other watersheds as well. Assessment of stand dynamics (data not shown) confirmed that release events were stochastic and not consistent within or among plots.

Some of the year-to-year variations in growth (Fig. 3) can be attributed to known biotic and abiotic stressors. For example, specific low growth years for sugar maple are likely associated with the 1993 infestation of pear thrips (*Taeniothrips inconsequens* [Uzel]) and a late spring frost in 2010 (Hufkens et al. 2012; Vermont Agency of Natural Resources Department of Forest Parks & Recreation 2010). While this late frost event could have affected sugar maple at low elevation, because no coincident downturn in growth was evident at mid elevation (Fig. 3), we suspect that mid elevation trees were unaffected due to a later timing of bud break relative to frost exposure. Additionally, a

severe ice storm in 1998 that broke the limbs of many hardwood species (Rhoads et al. 2002), was reported for many mid and high elevation sites (Vermont Agency of Natural Resources Department of Forest Parks & Recreation 2013). Areas of high ice loading align with our mid elevation zone, and when followed by a severe drought in 1999, may have caused the growth declines for yellow birch and sugar maple evident in 1999 through 2000 (Fig. 3). Lastly, declines in growth apparent for red spruce at all elevations (Fig. 3) correspond with documented winter injury events (Johnson et al. 1986), including in 2003, which was a severe event across the region (Lazarus et al. 2004).

At high elevation, balsam fir experienced maximum growth at the end of the 1980s through the early 2000s (Fig. 3), which parallels the documented decline of red spruce (Lazarus et al. 2004) and suggests that balsam fir was able to capitalize on the resulting foliar losses and growth declines of sympatric red spruce. Balsam fir is not affected by winter injury due to its extreme cold tolerance (DeHayes et al. 2001). Since 2003, a substantial winter injury event has not occurred in the region (Vermont Agency of Natural Resources Department of Forest Parks & Recreation 2013), which may in part, explain the increase in red spruce growth relative to balsam fir. However, despite some differences over time, at high elevation plots, both red spruce and balsam fir had relatively low growth due to limitations in growing season length and lower annual temperature.

We detected growth reductions for sugar maple at both low and mid elevations that are temporally consistent with regional declines attributed to acid deposition-induced Ca depletion combined with other stress exposures (Schaberg et al. 2006). In this study, growth declines were more dramatic for sugar maple at mid elevation than at low

elevation. As acid deposition is more severe with increasing elevation (Johnson and Siccama 1983), mid elevation sugar maple may have suffered from Ca depletion more extensively and earlier than trees at lower elevation. Indeed, other studies have indicated that sugar maple at mid and high elevations display greater signs of physiological stress from Ca depletion than those at low elevation (Minocha et al. 2010). Although sugar maple growth measurements reported here are low compared to the other sampled species, they are not atypical when compared to other dendrochronological studies in eastern North America (Bishop et al. 2015; Duchesne et al. 2003; Hansen 2015; Long et al. 2009).

Red spruce on Mt. Mansfield have experienced a recent growth increase consistent with other locations (Fig. 2, 3) (Kosiba et al. 2013). This increase is especially pronounced at mid elevations, which has been previously demonstrated (Kosiba et al. 2013). However, the other species at mid elevations did not display an analogous increase in growth, discounting stand dynamics for this rapid change.

Regardless of species-specific growth, mid elevation plots displayed significantly higher growth than low and high elevations plots. This could signify that mid elevation sites provide more advantageous conditions for tree growth than either lower or higher elevation (e.g., for red spruce), or it could be confounded by the fact that the fastest growing species for all quartiles (yellow birch) was only sampled at mid elevations.

Red maple also exhibited increased growth recently. However, since this study surveyed a relatively small number of red maple, and comparably few studies have examined red maple growth in the region, it is difficult to assess the extent of this growth surge at this time (see Silva et al. 2010; Zhang et al. 2015). Additionally, because red

maple was only sampled at low elevation, we do not know if elevation influences this apparent growth increase. Limited reports on red maple have revealed that its abundance increased dramatically in the 20th century (Siccama 1971), due in part to its purported niche as a “super-generalist” and large ecological amplitude, which permitted competitive growth advantages at a range of site conditions (Abrams 1998).

Relationships with climate and deposition

As expected, growth associations among species and elevations had differing relationships to local climate and deposition. Two general growth associations were evident: (1) a positive correlation with higher temperature in the year of growth (particularly for red maple and red spruce, and to a lesser extent, balsam fir, yellow birch, and sugar maple) and (2) a negative correlation with excessive heat (CDD: cumulative temperatures > 18.3°C) during the previous year (for all species except yellow birch). Positive relationships between temperature and growth imply that low temperature restricted growth. Negative relationships between temperature and growth the following year suggest a legacy effect of higher temperature, conceivably through limitations of C or other compulsory elements. For example, higher temperature could increase contemporary growth if the tree consumes C for immediate needs, such as increased maintenance respiration, growth, or reproductive buds, rather than to store C to support growth the next year (Rennenberg et al. 2006). Similarly, temperature-induced increases in growth could result in the sequestration of other nutrients (e.g., N, Ca) that temporarily become limiting and indirectly suppress growth the following year (Rennenberg et al. 2006).

Using CDD in growth-climate relationships provides a readily available index of integrated heat exposure that was consistently associated with growth for trees at Mt. Mansfield. Measures of growing season temperatures were important for both mid and low elevations, while water year temperature and pFall $T_{\min}$ were important at high elevations. Trees at both high and mid elevations displayed a positive growth correlation with CDD. By quartile, it is clear that climate-growth relationships were not steady through time (Table 2). For some species and variables, associations reversed for one or more quartiles, complicating our ability to predict tree responses to these variables.

Non-growing season temperature was important for the two conifers in this study. For both balsam fir and red spruce, mild autumn temperatures can delay cold hardening, which allows the foliage to remain photosynthetically active for a prolonged period. Additionally, because red spruce is a temperate conifer that can photosynthesize in fall, winter, and spring, provided that temperatures are moderate (Schaberg and DeHayes 2000), it has been hypothesized that the notable and recent increase in growth could be due to climatic warming that has lengthened the functional growing season for red spruce (Kosiba et al. 2013).

At mid elevations, there was a positive association between red spruce growth and previous Winter $T_{\max}$, suggesting that warmer winter temperatures could provide increased growth opportunities for this species. This finding has been reported by others (Cook et al. 1987; McLaughlin et al. 1987). Interestingly, this relationship was negative in the third quartile, 1969-1990, a period of peak acid deposition (Greaver et al. 2012) and documented red spruce decline (Hornbeck et al. 1987; Johnson et al. 1988). When warm winter temperatures cause thaws, red spruce can photosynthesize while sympatric

species remain dormant; however, when cold temperatures return, freezing damage can ensue. This process is exacerbated by acid deposition, which reduces the cold tolerance of red spruce foliage (Schaberg and DeHayes 2000).

Red spruce growth at the Mt. Mansfield study plots corresponds well with red spruce growth region-wide, implying that growth trends in this study are not a byproduct of local stand dynamics and, further, that climate- and deposition- growth relationships presented here may be representative of a region-wide phenomenon. Positive associations with measures of growing season length and temperature (e.g., water year T_{mean} , Jun-Sept T_{mean} , GDD, CDD) allude to continued favorable growth conditions and potential range expansion (e.g., Foster and D'Amato 2015) for red spruce into the future. This conclusion contrasts with projected species range maps that depict red spruce range constriction in the future (Iverson et al. 2008). Interestingly, the period of most divergent growth between Mt. Mansfield and regional red spruce was from roughly 1980 through 2008. This period was interspersed with a series of winter injury events with varied intensities and legacies across the region (Lazarus et al. 2004, Kosiba et al. 2013). The higher than average growth at Mt. Mansfield relative to the regional chronology may indicate that red spruce on Mt. Mansfield experienced less winter injury than the regional norm. Likewise, the recent synchronous increase in growth evident in the chronologies indicates that this rebound is a regional phenomenon, similar to the ubiquitous decline in growth following the 2003 winter injury event.

In contrast with the other species, sugar maple growth at low elevation was not correlated with the selected variables, suggesting that they may be responding to micro-site parameters not considered in this analysis (e.g., nutrient availability, stand dynamics,

local soil moisture, etc.). For example, PDSI is a metric for the whole state of VT that does not consider site based soil and hydrological differences that may help explain tree growth. Others have also shown similar weak climate-growth relationships for sugar maple in the region (e.g., Bishop et al. 2015). Sugar maple growth at mid elevations exhibited a negative correlation with pSummer T_{mean} and T_{max} , indicating that elevated summer temperature due to climate change (Kunkle et al. 2013) could limit sugar maple growth further in the future.

Sugar maple at HBEF (Hansen 2015) had higher growth than those at Mt. Mansfield, particularly during the first half of the chronology. After 1982, the chronologies are more analogous. Sugar maple at Mt. Mansfield displayed a subdued decline in growth in the 1980s, but otherwise maintained consistent, low growth. It is unclear why sugar maple at HBEF had higher growth than those at Mt. Mansfield for the first part of the chronology, though higher acidic loading in the west of the region (e.g., Ollinger et al. 1993) could have stressed sugar maple trees there first. However, the near-steady growth of sugar maple at Mt. Mansfield since 1950 (Fig. 2) seems inconsistent with the possibility of early stress from acidic inputs. Another possibility is that Mt. Mansfield is a more marginal site for sugar maple growth, so baseline growth there was consistently lower.

Growth of Mt. Mansfield yellow birch did not correspond well to yellow birch growing at HBEF (Hansen 2015). One possible explanation for this could be the lower sample size for yellow birch ($n = 34$ trees) relative to sugar maple ($n = 63$ trees) at Mt. Mansfield, which could reduce the accuracy of yearly growth mean estimates and

increase the variation around those means. These results demonstrate the importance of having high replication when looking at region-wide patterns in tree cores.

Surprisingly, annual S and N deposition only related to poor growth for red spruce at low elevation, but not at mid and high elevations where documented declines have been tied to acid deposition (DeHayes 1992; Schaberg et al. 2006). While growth of sugar maple and red spruce, aside from low elevation, did display negative correlations with S deposition, these relationships were not significant. One possible reason for this outcome is the absence of long-term S and N deposition data. Datasets in this region only span from 1965-2012, which excludes over a decade of acid deposition inputs. In addition, detrending growth data with a flexible spline removes high frequency variation and biological trends while preserving decadal and longer trends. Nevertheless, a smoothing spline is one of the best options for reducing the effects of tree competition and age-related growth trends. Lastly, relating growth to annual deposition data may overlook more complex impacts of pollutant loading, such as long-term cumulative Ca depletion. For example, research has shown that red spruce growth was related to the historic accumulation of acidic inputs relative to the capacity of the site to buffer these inputs (Engle et al. 2016).

Atmospheric CO₂ concentration trends were not associated with growth (or residuals of growth-climate models after accounting for the effects of climate) for any of the species or elevations. Our findings align with others who have shown no significant effect of elevated CO₂ concentration on stem growth (e.g., Bader et al. 2013; Girardian et al. 2016; Körner et al. 2005). While it is currently debated if elevated CO₂ concentrations will provoke changes in tree growth, there is evidence that CO₂ effects are muted or

nonexistent on sites where other environmental factors, including soil fertility (particularly N-limitation) (Oren et al. 2001), temperature, and water (Körner et al. 2005), are more limiting to stem growth than C (Körner 2003; Silva et al. 2013). Based on the strong associations with growth of trees in this study and measures of heat that we reported here, we hypothesize that temperature is more limiting to these trees than C availability.

Conclusions

We presented a comparison of growth trends for five key tree species. While this is a case study, by comparing the chronologies developed at this site to others in the region, we show that at least for red spruce, there is strong alignment to the broader region and propose that patterns evident here may be indicative of the region's forested ecosystems. Comparisons of the chronologies (Fig. 2) highlight that the five species often experienced changes in growth that were either species- or elevation-specific. There was a high degree of year-to-year variability in growth, likely due to local abiotic and biotic factors. For some species, the repeated stress of abiotic and biotic factors in succession may help explain prolonged growth declines.

For most species and elevations, there was a positive association between higher temperatures and growth during that same year, suggesting that contemporary warming has improved the competitive status of many trees. However, there are two noted exceptions to this trend. The first is that excessive heat (cumulative temperatures > 18.3°C) the previous year was broadly associated with lower growth the following year. Further investigation is needed to understand the complex interplay of elevated

604 temperature and net growth in successive years and over time. The second exception is
605 that sugar maple trees at low elevation showed no positive association with increased
606 warming. This may provide field-based evidence of the sensitivity of sugar maple to the
607 warming that has been projected by climate change range models (e.g., Iverson et al.
608 2008).

609 Declines of red spruce have also been well documented, but the recent and
610 surprising growth surge had not been investigated. Here we show that it occurred across
611 elevations and watersheds and that perhaps another species, red maple, has had a
612 synchronous increase in growth. Both species show positive correlations with growing
613 season temperatures, including CDD. Indeed, CDD was consistently associated with
614 increased growth for most of species and elevations overall. This finding highlights the
615 potential value of CDD as an integrated temperature index with relevance to tree growth
616 in a warming world. While CDD was often positively associated with growth, pCDD
617 often had a negative relationship. As CDD increases in the future, its influence on tree
618 growth could be mixed. Results of this analysis suggest that some tree species may be
619 responding favorably to changing environmental conditions, while others are either
620 declining or appear stable in growth. Though this specific study covered a small spatial
621 scale, it opens avenues for future work to examine more fully some of the patterns that
622 emerged here. Indeed, there have been relatively few studies that have examined growth
623 trends and responses to the environment among species in the region; yet, this is of
624 interest as the climate changes and the future state of forest health and productivity
625 remains uncertain.

627 **Acknowledgments**

628 We value the helpful manuscript edits from Drs. Andrea Lini (University of
629 Vermont, UVM) E. Carol Adair (UVM), Neil Pederson (Harvard Forest), and two
630 anonymous reviewers. We greatly appreciate the assistance in the field and laboratory by
631 Ben Engel, Dan Hale, Josh Halman, Kindle Loomis, Allyson Makuch, Jill Spies, and
632 Sam Wallace. Thanks to Chris Hansen, Ben Engel, and Aiko Weverka for sharing data.
633 Thanks are also extended to Jim Duncan and Carl Waite (Vermont Monitoring
634 Cooperative) for their assistance and approval of field collection. We also appreciate the
635 cooperation of the Vermont Department of Forest, Parks, and Recreation for providing
636 timely permission for field sampling. This manuscript is a contribution of the Hubbard
637 Brook Ecosystem Study. Hubbard Brook is part of the Long-Term Ecological Research
638 (LTER) network, which is supported by the National Science Foundation. The Hubbard
639 Brook Experimental Forest is operated and maintained by the USDA Forest Service,
640 Newtown Square, PA. This research was supported, in part, by the Forest Service
641 Northern Research Station and by grants from the Vermont Monitoring Cooperative and
642 USDA McIntire-Stennis Forest Research Program.

643

References

- Aber, J., McDowell, W., Nadelhoffer, K., Alison, M., Berntson, G., Kamakea, M.,
McNulty, S., Currie, W., Rustad, L., and Fernandez, I. 1998. Nitrogen saturation
in temperate forest ecosystems. *BioSci* **48**(11): 921-934.
- Abrams, M.D. 1998. The red maple paradox. *BioSc* **48**(5): 355-364.

- Bader, M.K.F., Leuzinger, S., Keel, S.G., Siegwolf, R.T.W., Hagedorn, F., Schleppi, P., and Körner, C. 2013. Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project. *J Ecol* **101**(6): 1509-1519. doi: 10.1111/1365-2745.12149.
- Beckage, B., Osborne, B., Gavin, D.G., Pucko, C., Siccama, T., and Perkins, T. 2008. A rapid upward shift of a forest ecotone during 40 years of warming in the Green Mountains of Vermont. *P Natl Acad Sci* **105**(11): 4197-4202. doi: 10.1073/pnas.0708921105.
- Bishop, D.A., Beier, C.M., Pederson, N., Lawrence, G.B., Stella, J.C., and Sullivan, T.J. 2015. Regional growth decline of sugar maple (*Acer saccharum*) and its potential causes. *Ecosphere* **6**(10): art179. doi: 10.1890/ES15-00260.1.
- Bunn, A., Korpela, M., Biondi, F., Campelo, F., Merian, P., Qeadan, F., and Zang, C. 2015. dplR: Dendrochronology Program Library in R. R package version 1.6.3. <http://r-forge.r-project.org/projects/dplr/>.
- Burns, D., Lynch, J., Cosby, B., Fenn, M., and Baron, J. 2011. National acid precipitation assessment program report to Congress 2011: an integrated assessment. U.S. Environmental Protection Agency, Washington, DC, USA.
- Burns, R., and Honkala, B. 1990. Silvics of North America: 1. Conifers; 2. Hardwoods Agriculture Handbook 654. US Department of Agriculture, Forest Service, Washington, DC.
- Cogbill, C.V. 1996. Historical ecology of the upper Stevensville Brook watershed, Underhill, Vermont. For Vermont Department of Forest, Parks and Recreation and Vermont Monitoring Cooperative.

- Cook, E.R., Johnson, A.H., and Blasing, T.J. 1987. Forest decline: modeling the effect of climate in tree rings. *Tree Phys* **3**(1): 27-40. doi: 10.1093/treephys/3.1.27.
- Cook, E.R., and Kairiukstis, L.A. (eds). 1990. *Methods of Dendrochronology: Applications in the Environmental Sciences*. Kluwer Academic Publishers, Dordrecht.
- DeHayes, D.H. 1992. Winter injury and developmental cold tolerance of red spruce. *In* *The Ecology and Decline of Red Spruce in the Eastern United States. Edited by C. Eagar and M.B. Adams*. Springer-Verlag New York Inc., New York, NY, USA. pp. 295-337.
- DeHayes, D.H., Schaberg, P.G., Hawley, G.J., and Strimbeck, G.R. 1999. Acid rain impacts on calcium nutrition and forest health. *BioSci* **49**(10): 789-800.
- DeHayes, D.H., Schaberg, P.G., and Strimbeck, G.R. 2001. Red spruce cold hardiness and freezing injury susceptibility. *In* *Conifer Cold Hardiness. Edited by F. Bigras*. Kluwer Academic Publishers, Dordrecht, the Netherlands. pp. 495-529.
- Duchesne, L., Ouimet, R., and Morneau, C. 2003. Assessment of sugar maple health based on basal area growth pattern. *Can J Forest Res* **33**(11): 2074-2080. doi: 10.1139/x03-141.
- Dupigny-Giroux, L.-A. 2002. Climate variability and socioeconomic consequences of Vermont's natural hazards: a historical perspective. *Vermont History* **70**: 19-39.
- Engel, B.J., Schaberg, P.G., Hawley, G.J., Rayback, S.A., Pontius, J., Kosiba, A.M., and Miller, E.K. 2016. Assessing relationships between red spruce radial growth and pollution critical load exceedence values. *Forest Ecol Manag* **359**: 83-91.

- Foster, J.R., and D'Amato, A.W. 2015. Montane forest ecotones moved downslope in northeastern U.S. in spite of warming between 1984 and 2011. *Global Change Biol* **21**(12): 4497-4507. doi: 10.1111/gcb.13046.
- Gavin, D.G., and Peart, D.R. 1993. Effects of beech bark disease on the growth of American beech (*Fagus grandifolia*). *Can J Forest Res* **23**(8): 1566-1575. doi: 10.1139/x93-197.
- Girardin, M.P., Bouriaud, O., Hogg, E.H., Kurz, W., Zimmermann, N.E., Metsaranta, J.M., de Jong, R., Frank, D.C., Esper, J., Büntgen, U., Guo, X.J., and Bhatti, J. 2016. No growth stimulation of Canada's boreal forest under half-century of combined warming and CO₂ fertilization. *P Natl Acad Sci* **113**(52): E8406-E8414. doi: 10.1073/pnas.1610156113.
- Greaver, T.L., Sullivan, T.J., Herrick, J.D., Barber, M.C., Baron, J.S., Cosby, B.J., Deerhake, M.E., Dennis, R.L., Dubois, J.-J.B., Goodale, C.L., Herlihy, A.T., Lawrence, G.B., Liu, L., Lynch, J.A., and Novak, K.J. 2012. Ecological effects of nitrogen and sulfur air pollution in the US: what do we know? *Front Ecol Environ* **10**(7): 365-372. doi: 10.1890/110049.
- Hansen, C.F. 2015. LiDAR remote sensing of forest canopy structure and integrity: Trophic interactions among forest health, productivity, insects and birds. M.Sc. thesis, Rubenstein School of Environment and Natural Resources, University of Vermont, VT, USA.
- Holmes, R.L. 1983. Computer-assisted quality control in tree-ring dating and measurement. *Tree-ring Bull* **43**: 69-78.

- Hornbeck, J.W., Smith, R.B., and Federer, C.A. 1987. Growth decline in red spruce and balsam fir relative to natural processes. *Water Air Soil Poll* **31**(1-2): 425-430. doi: 10.1007/BF00630860.
- Hornbeck, J.W., and Smith, R.L. 1985. Documentation of red spruce growth decline. *Canadian Journal of Forest Research* **15**: 1199-1201.
- Hufkens, K., Friedl, M.A., Keenan, T.F., Sonnentag, O., Bailey, A., O'Keefe, J., and Richardson, A.D. 2012. Ecological impacts of a widespread frost event following early spring leaf-out. *Global Change Biol* **18**(7): 2365-2377. doi: 10.1111/j.1365-2486.2012.02712.x.
- Iverson, L.R., Prasad, A.M., Matthews, S.N., and Peters, M. 2008. Estimating potential habitat for 134 eastern U.S. tree species under six climate scenarios. *Forest Ecol Manag* **254**(3): 390-406.
- Johnson, A., Cook, E., and Siccama, T. 1988. Climate and red spruce growth and decline in the northern Appalachians. *P Natl A Sci* **85**(15): 5369-5373.
- Johnson, A.H., Friedland, A.J., and Dushoff, J.G. 1986. Recent and historic red spruce mortality: Evidence of climatic influence. *Water Air Soil Poll* **30**(1): 319-330. doi: 10.1007/bf00305203.
- Johnson, A.H., and Siccama, T.G. 1983. Acid deposition and forest decline. *Envir Sci Tech* **17**(7): 294A-305A. doi: 10.1021/es00113a001.
- Körner, C. 2003. Carbon limitation in trees. *J Ecol* **91**(1): 4-17. doi: 10.1046/j.1365-2745.2003.00742.x.

- Körner, C., Asshoff, R., Bignucolo, O., Hättenschwiler, S., Keel, S.G., Peláez-Riedl, S., Pepin, S., Siegwolf, R.T., and Zotz, G. 2005. Carbon flux and growth in mature deciduous forest trees exposed to elevated CO₂. *Science* **309**(5739): 1360-1362.
- Kosiba, A.M., Schaberg, P.G., Hawley, G.J., and Hansen, C.F. 2013. Quantifying the legacy of foliar winter injury on woody aboveground carbon sequestration of red spruce trees. *Forest Ecol Manag* **302**: 363-371.
- Kunkel, K.E., Stevens, L., Stevens, S., Sun, L., Janssen, E., Wuebbles, D., Rennells, J., DeGaetano, A., and Dobson, J. 2013. NOAA Technical Report NESDIS. Regional climate trends and scenarios for the U.S. National Climate Assessment. Part 1: Climate of the Northeast U.S.
- Lazarus, B.E., Schaberg, P.G., DeHayes, D.H., and Hawley, G.J. 2004. Severe red spruce winter injury in 2003 creates unusual ecological event in the northeastern United States. *Can J Forest Res* **34**(8): 1784-1788.
- LeBlanc, D.C. 1992. Spatial and temporal variation in the prevalence of growth decline in red spruce populations of the northeastern United States. *Can. J. For. Res.* **22**: 1351–1363.
- Likens, G. 2010. Chemistry of Steamwater at HBEF WS-1. Available from <http://www.hubbardbrook.org/data/dataset.php?id=3> [accessed 14 October 2014].
- Long, R.P., Horsley, S.B., Hallett, R.A., and Bailey, S.W. 2009. Sugar maple growth in relation to nutrition and stress in the northeastern United States. *Ecol Appl* **19**(6): 1454-1466. doi: 10.1890/08-1535.1.
- McLaughlin, S.B., Downing, D.J., Blasing, T.J., Cook, E.R., and Adams, H.S. 1987. An analysis of climate and competition as contributors to decline of red spruce in

- high elevation Appalachian forests of the eastern United States. *Oecologia* **72**(4): 487-501. doi: 10.1007/BF00378973.
- Minocha, R., Long, S., Thangavel, P., Minocha, S.C., Eagar, C., and Driscoll, C.T. 2010. Elevation dependent sensitivity of northern hardwoods to Ca addition at Hubbard Brook Experimental Forest, N.H., U.S.A. *Forest Ecol and Manag* **260**(12): 2115-2124. doi: <http://dx.doi.org/10.1016/j.foreco.2010.09.002>.
- National Atmospheric Deposition Program. 2014. NADP/NTN Monitoring Location VT99. U.S. Geological Survey. Available from <http://nadp.sws.uiuc.edu/sites/siteinfo.asp?id=VT99&net=NTN> [accessed 10 May 2014].
- National Drought Mitigation Center. 2014. Drought Risk Atlas. Available from <http://droughtatlas.unl.edu/Data.aspx> [accessed 10 October 2013].
- NOAA Earth System Research Laboratory Global Monitoring Division. 2013. Monthly Average Mauna Loa CO₂. Available from <http://www.esrl.noaa.gov/gmd/ccgg/trends/> [accessed 6 October 2013].
- NOAA National Weather Service. 2014. Monthly total growing degree days for Burlington, VT. Available from http://www.weather.gov/btv/climo_gdd [accessed 23 April 2014].
- Ollinger, S.V., Aber, J.D., Lovett, G.M., Millham, S.E., Lathrop, R.G., and Ellis, J.M. 1993. A spatial model of atmospheric deposition for the northeastern U.S. *Ecol Appl* **3**(3): 459-472.
- Oren, R., Ellsworth, D.S., Johnsen, K.H., Phillips, N., Ewers, B.E., Maier, C., Schäfer, K.V., McCarthy, H., Hendrey, G., and McNulty, S.G. 2001. Soil fertility limits

- carbon sequestration by forest ecosystems in a CO₂-enriched atmosphere. *Nature* **411**(6836): 469-472.
- Pederson, N., Bell, A.R., Cook, E.R., Lall, U., Devineni, N., Seager, R., Eggleston, K., and Vranes, K.J. 2013. Is an epic pluvial masking the water insecurity of the greater New York City region? *J Climate* **24**(6): 1339-1354. doi: 10.1175/jcli-d-11-00723.1.
- PRISM Climate Group. 2004. Available from <http://prism.oregonstate.edu/> [accessed 10 May 2014].
- R Development Core Team. 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rennenberg, H., Loreto, F., Polle, A., Brilli, F., Fares, S., Beniwal, R., and Gessler, A. 2006. Physiological responses of forest trees to heat and drought. *Plant Biol* **8**(05): 556-571.
- Rhoads, A.G., Hamburg, S.P., Fahey, T.J., Siccama, T.G., Hane, E.N., Battles, J., Cogbill, C., Randall, J., and Wilson, G. 2002. Effects of an intense ice storm on the structure of a northern hardwood forest. *Can J Forest Res* **32**(10): 1763-1775. doi: 10.1139/x02-089.
- Salzer, M.W., Hughes, M.K., Bunn, A.G., and Kipfmüller, K.F. 2009. Recent unprecedented tree-ring growth in bristlecone pine at the highest elevations and possible causes. *Proceedings of the National Academy of Sciences* **106**(48): 20348-20353. doi: 10.1073/pnas.0903029106.
- Schaberg, P.G., and DeHayes, D.H. 2000. Physiological and environmental causes of freezing injury to red spruce. *In* Responses of Northern U.S. Forests to

- Environmental Change. *Edited by* R.A. Mickler and R.A. Birdsey and J.L. Hom. Springer-Verlag, New York. pp. 181-227.
- Schaberg, P.G., Tilley, J.W., Hawley, G.J., DeHayes, D.H., and Bailey, S.W. 2006. Associations of calcium and aluminum with the growth and health of sugar maple trees in Vermont. *Forest Ecol Manag* **223**(1–3): 159-169. doi: <http://dx.doi.org/10.1016/j.foreco.2005.10.067>.
- Siccama, T.G. 1971. Presettlement and present forest vegetation in northern Vermont with special reference to Chittenden County. *Am Midl Nat* **85**(1): 153-172.
- Silva, L.C.R., and Anand, M. 2013. Probing for the influence of atmospheric CO₂ and climate change on forest ecosystems across biomes. *Global Ecology and Biogeography* **22**(1): 83-92. doi: 10.1111/j.1466-8238.2012.00783.x.
- Silva, L.C.R., Anand, M., and Leithead, M.D. 2010. Recent widespread tree growth decline despite increasing atmospheric CO₂. *PLoS ONE* **5**(7): e11543. doi: 10.1371/journal.pone.0011543.
- Soil Survey Staff. 2015. Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey. Available from <http://websoilsurvey.nrcs.usda.gov/> [accessed 15 August 2015].
- Soule, P.T., and Knapp, P.A. 2006. Radial growth rate increases in naturally occurring ponderosa pine trees: a late-20th century CO₂ fertilization effect? *New Phytol* **171**(2): 379-390. doi: 10.1111/j.1469-8137.2006.01746.x.
- Speer, J.H. 2010. Fundamentals of tree-ring research. University of Arizona Press.
- Stokes, M., and Smiley, T. 1968. An Introduction to Tree-Ring Dating. University of Chicago Press.

- The National Drought Mitigation Center. 2012. SPEI. University of Nebraska, Lincoln.
Available from <http://droughtatlas.unl.edu/About.aspx> [accessed 15 March 2015].
- van Doorn, N.S., Battles, J.J., Fahey, T.J., Siccama, T.G., and Schwarz, P.A. 2011. Links between biomass and tree demography in a northern hardwood forest: a decade of stability and change in Hubbard Brook Valley, New Hampshire. *Can J Forest Res* **41**(7): 1369-1379. doi: 10.1139/x11-063.
- Vermont Agency of Natural Resources Department of Forest Parks & Recreation. 2010. Forest insects and disease conditions in Vermont 2010. Waterbury, VT.
- Vermont Agency of Natural Resources Department of Forest Parks & Recreation. 2013. Combined Master Dataset of Aerial Defoliation Surveys, 1985 to 2013. Hosted by Vermont Monitoring Cooperative. Available from <http://www.uvm.edu/vmc/research/data.php?id=66> [accessed 25 July 2014].
- Waterhouse, J.S., Switsur, V.R., Barker, A.C., Carter, A.H.C., Hemming, D.L., Loader, N.J., and Robertson, I. 2004. Northern European trees show a progressively diminishing response to increasing atmospheric carbon dioxide concentrations. *Quaternary Sci Rev* **23**(7–8): 803-810. doi.org/10.1016/j.quascirev.2003.06.011.
- West, P.W. 1980. Use of diameter increment and basal area increment in tree growth studies. *Can J Forest Res* **10**(1): 71-77. doi:10.1139/x80-012.
- Weverka, A.S. 2012. Remote sensing of productivity in northeastern forests. M.Sc. thesis, Rubenstein School of Environment and Natural Resources, University of Vermont, V.T.
- Whitney, G. 1994. From coastal wilderness to fruited plain: An environmental history of the eastern U.S. 1500 to present. Cambridge University, Cambridge, MA.

World Data Center for Paleoclimatology and NOAA Paleoclimatology Program. Law

Dome ice core data. Available from:

<ftp://ftp.ncdc.noaa.gov/pub/data/paleo/icecore/antarctica/law/law2006.txt>

[accessed 10 October 2016].

Yamaguchi, D.K. 1991. A simple method for cross-dating increment cores from living trees. *Can J Forest Res* **21**(3): 414-416. doi: 10.1139/x91-053.

Zang, C., and Biondi, F. 2015. treeclim: an R package for the numerical calibration of proxy-climate relationships. *Ecography* **38**(4): 431-436. doi: 10.1111/ecog.01335.

Zhang, Y., Bergeron, Y., Zhao, X.-H., and Drobyshev, I. 2015. Stand history is more important than climate in controlling red maple (*Acer rubrum* L.) growth at its northern distribution limit in western Quebec, Canada. *J Plant Ecol* **8**(4): 368-379. doi: 10.1093/jpe/rtu029.

Table 1: Significant correlations from stationary climate-growth assessments for all species combined per elevation for trees from Mt. Mansfield, VT, as evaluated using Principal Component analysis (1925-2012). Significance based on 99% CI.

Elevation ^a	Positive		Negative	
	Variable ^{b,c}	Cor. Coef.	Variable ^b	Cor. Coef.
High	Water year T _{mean}	0.34	HDD	-0.38
	pWinter T _{max}	0.25		
	pFall T _{min}	0.40		
	Winter T _{mean}	0.26		
	Spring T _{mean}	0.27		
	CDD	0.37		
	GDD	0.39		
	PDSI	0.36		
Mid	Summer T _{mean}	0.33	Mar SPEI	-0.27
	May-June T _{mean}	0.32		
	Jun-Sep T _{mean}	0.36		
	CDD	0.34		
	pFeb SPEI	0.43		
	pDec SPEI	0.38		
Low	pSpring T _{max}	0.25	Mar SPEI	-0.27
	Summer T _{mean}	0.38		
	May-Aug T _{mean}	0.28		
	pMar SPEI	0.28		
	NH ₄ ⁺	0.35		

^a Low = 450-650 m a.s.l., mid = 750-850 m, and high = 900-1000 m

^b p = previous year, T = temperature, Min = minimum, GDD = growing degree days (cumulative degrees > 5°C [41°F]), HDD = heating degree days (cumulative days < 18.3°C [65°F]), CDD = cooling degree days (cumulative degrees > 18.3°C [65°F]), CDD = cooling degree days (cumulative degrees > 18.3°C [65°F]), SPEI = Standardized Precipitation-Evapotranspiration Index, Winter: previous December–February, Spring: March–May, Summer: June–August, Fall: September–November. For more information on variables see methods.

^c Not all variables span the entire chronology (1925-2012, 1945-2012, or 1965-2012). See methods.

Table 2: Correlations from stationary climate-growth assessments from Mt. Mansfield, VT, shown by species and elevation for the entire chronology (1925-2012) and by chronology quartile. Coefficients in bold text are significant at 99% CI.

Species	Elevation ^a	Variable ^b	Overall	Quartiles			
			1925-2012 ^c	1925-1946	1947-1968	1969-1990	1991-2012
Balsam fir	High	pFall T _{min}	0.38	0.29	0.48	0.44	0.30
		Spring T _{min}	0.24	0.17	0.59	0.19	0.08
		pCDD	-0.37		-0.47	-0.17	-0.88
		pGDD	-0.31		-0.48	0.03	-0.88
Red maple	Low	CDD	0.39		0.24	0.34	0.65
		Summer T _{mean}	0.30	0.37	0.22	0.31	0.55
		Jun-Sep T _{mean}	0.26	0.43	0.28	0.26	0.43
		pCDD	-0.34		-0.59	-0.27	-0.17
Red spruce	Low	Water year T _{mean}	0.43	0.44	0.52	0.24	0.48
		pFall T _{min}	0.32	0.19	0.41	0.23	0.53
		Winter T _{mean}	0.26	0.14	0.39	-0.04	0.52
		Spring T _{mean}	0.36	0.34	0.37	0.26	0.46
		May-Aug T _{mean}	0.28	0.27	0.29	0.16	0.37
		Jun-Sep T _{mean}	0.43	0.31	0.48	0.01	0.13
		pCDD	-0.35		-0.40	-0.31	-0.39
		HDD	-0.37		-0.46	-0.05	-0.50
		GDD	0.38		0.46	0.21	0.37
		PDSI	0.36		0.44	-0.56	0.10
		pNO ₃ ⁻	-0.37		-0.164	-0.17	-0.46
	Mid	Water year T _{mean}	0.27	0.35	0.53	0.17	0.42
		pWinter T _{max}	0.27	0.38	0.55	-0.19	0.37
		Jun-Sep T _{mean}	0.26	0.29	0.49	0.12	0.17
		pCDD	-0.41		-0.38	-0.51	-0.40
		GDD	0.34		0.44	0.13	0.39
	High	May SPEI	-0.30		-0.15	-0.29	-0.44
		Water year T _{mean}	0.36	0.28	0.47	0.27	0.41
		pWinter T _{max}	0.23	0.36	0.46	-0.07	0.24
		pFall T _{min}	0.32	0.05	0.53	-0.28	0.52
		Winter T _{mean}	0.26	0.16	0.22	0.13	0.41
		Summer T _{max}	0.27	-0.03	0.31	0.38	0.45
		Fall T _{mean}	0.23	0.39	0.28	0.25	0.11
		Jun-Sep T _{mean}	0.31	0.06	0.55	0.48	0.41
		May-Aug T _{mean}	0.33	0.04	0.39	0.46	0.20
		pCDD	-0.32		-0.44	-0.29	-0.37
		CDD	0.42		0.36	0.43	0.50
		HDD	-0.34		-0.45	0.07	-0.44
		GDD	0.45		0.50	0.30	-0.49
Sugar maple	Low		N/S				
	Mid	pSummer T _{mean}	-0.25	-0.12	-0.40	-0.23	-0.22
		pSummer T _{max}	-0.27	-0.25	-0.38	-0.28	-0.09
		Summer T _{mean}	0.26	0.19	0.45	0.08	0.33
Yellow birch	Mid	pCDD	-0.38		-0.55	-0.29	-0.23
Yellow birch	Mid	Summer T _{mean}	0.29	0.37	0.29	0.29	0.29

^a Low = 450-650 m a.s.l., mid = 750-850 m, and high = 900-1000 m

^b p = previous, T = temperature, Max = maximum, Min = minimum, PDSI = Vermont Palmer Drought Severity Index, GDD = growing degree days (cumulative degrees > 5°C [41°F]), HDD = heating degree days (cumulative days < 18.3°C [65°F]), CDD = cooling degree days (cumulative degrees > 18.3°C [65°F]), SPEI = Vermont Standardized Precipitation-Evapotranspiration Index, Winter: previous December–February, Spring: March–May, Summer: June–August, Fall: September–November. For more information on variables see methods.

^c Not all variables span the entire chronology (1925-2012, 1945-2012, or 1965-2012). See methods.

Figure captions

Fig. 1: Map of plot locations in Mt. Mansfield State Forest, VT, USA, showing (A) three watershed transects, each with three plots located within low, mid, and high elevational zones, (B) species sampled per elevational band, and (C) study site location within the region.

Fig. 2: Biweight robust mean basal area increment (BAI, $\text{cm}^2 \pm \text{SD}$ shown with grey bars) by (A) species and (B) elevation from 1925-2012. A non-parametric Kruskal-Wallis test, followed by Wilcoxon Each Pair test, using species, elevation, and year as factors was significant for the three factors ($P < 0.001$). Different letters following species and elevations categories denote significant differences overall ($P < 0.001$). The model was also significant for each of the four 21-year quartiles (1925-1946, 1947-1986, 1969-1990, 1991-2012; denoted with grey dashed line, $P < 0.001$), for species and elevation, but year was only significant for the period 1925-1946 (BF: balsam fir, RM: red maple, RS: red spruce, SM: sugar maple, YB: yellow birch, L: low elevation [450-650 m a.s.l.], M: mid [750-800 m], and H: high [900-1000 m]). Bonferonni adjusted P -values were used.

Fig. 3: Mean basal area increment (BAI, cm^2 , solid black line) ($\pm \text{SD}$) chronologies per species and elevational zone: (A) low 450-650, (B) mid 750-800, and (C) high 900-1000 m a.s.l. elevation from 1925-2012 in Mt. Mansfield State Forest, VT. Number of trees contributing the mean are shown with grey bars. Red circles designate the five highest growth years per chronology.

Fig. 4: Comparisons of mean growth (BAI $\pm \text{SD}$) for red spruce, sugar maple, and yellow birch from this study on Mt. Mansfield to (A) sugar maple and (B) yellow birch at Hubbard Brook Experimental Forest (HBEF), NH, and (C) red spruce in the region (34 plots in VT, NH, MA). Spearman correlation coefficients (R) and associated P -values are shown in each figure. Chronologies at HBEF were limited to 1950-2012.

Figures

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Supplemental Materials

Fig. S1: (A) Mean seasonal (Winter [p Dec-Feb], Spring [Mar-May], Summer [Jun-Aug], and Fall [Sep-Nov]) temperatures (°C; 1925-2012) for the study location (see Methods for more details). (B) Seasonal Standardized Precipitation-Evapotranspiration Index (SPEI; 1945-2012) for Vermont (VT). (C) Annual total precipitation (cm) for the study location (1925-2012); cooling degree days (CDD, measured as degrees > 18.3°C, 1945-2012), heating degree days (HDD, < 18.3°C, 1945-2012) and growing degree days (GDD, > 5°C, 1945-2012) for VT. (D) Atmospheric CO₂ concentration (ppm, 1945-2012) from Mauna Loa, HI; deposition (1965-2012, mg/L, by water year) of ammonium (NH₄⁺), nitrate (NO₃⁻), and sulfate (SO₄⁻) from Hubbard Brook Experimental Forest (Thorton, NH) and Mt. Mansfield (VT).

Table S1: Raw tree ring data and statistics listed by species per elevational zone across the three watersheds.

Species	Elev. Zone ^a	N trees	Mean DBH ± SD ^b	Age at BH ± SD ^c	Mean BAI ± SD ^d	Inter-series corr. ^e	Mean Sens. ^e	Max. BAI ± SD ^f
Balsam fir	High	34	23.4 ± 3.4	77.9 ± 20.4	7.9 ± 3.4	0.583	0.209	10 ± 2.2 (2001)
Red maple	Low	28	39.6 ± 8.4	90.0 ± 25.2	19.6 ± 11.2	0.514	0.225	29.8 ± 5.8 (2012)
Red spruce	Low	31	34 ± 9.3	142.7 ± 55.8	11.3 ± 9.3	0.519	0.257	18.1 ± 3.4 (2012)
	Mid	34	41.5 ± 10.2	165.5 ± 51	16.1 ± 10.2	0.559	0.254	27.9 ± 4.0 (2012)
Sugar maple	High	41	23.5 ± 8.2	86.7 ± 32.8	8.1 ± 4.1	0.579	0.236	13.2 ± 3.4 (2012)
	Low	29	32.5 ± 5.8	95.9 ± 20.3	9.9 ± 5.8	0.474	0.304	14.2 ± 1.1 (1994)
Yellow birch	Mid	34	28.2 ± 6.8	93.1 ± 23.8	7 ± 3.4	0.466	0.325	10.5 ± 2.3 (1983)
	Mid	34	43.2 ± 10.2	101.1 ± 27.2	18 ± 10.2	0.467	0.286	22.7 ± 0.6 (1982)

^a Low = 450-650 m a.s.l., mid = 750-850 m, and high = 900-1000 m
^b Mean diameter (± SD)(cm) measured at breast height (DBH 1.37 m).
^c Age in years is approximate, based on number of rings at breast height and distance to pith.
^d Mean basal area increment (cm²) calculated using the following formula $BAI = \pi(R_n^2 - R_{n-1}^2)$ (West 1980).
^e Interseries correlation and mean sensitivity, from COFECHA output.
^f Maximum basal area increment (cm²) and year of occurrence.

Table S2: Chronology statistics from Mt. Mansfield, VT, detrended with a 67% cubic smoothing spline and prewhitened, computed for 30-year time periods with 15-year overlap from 1923-2012.

Species	Elevation ^a	Stat ^b	Time period				
			1923-1952	1938-1967	1953-1982	1968-1997	1983-2012
Balsam fir	High	n cores	40	53	59	59	59
		n trees	24	30	34	34	34
		Rbar	0.22	0.44	0.38	0.30	0.31
		EPS	0.82	0.95	0.95	0.94	0.94
		SNR	4.67	18.83	18.68	14.52	15.26
Red maple	Low	n cores	63	77	86	86	86
		n trees	39	46	51	51	51
		Rbar	0.15	0.21	0.16	0.13	0.08
		EPS	0.82	0.91	0.91	0.88	0.82
		SNR	4.57	10.06	8.55	7.11	4.69
Red spruce	Low	n cores	54	56	56	56	56
		n trees	31	31	31	31	31
		Rbar	0.28	0.40	0.34	0.26	0.28
		EPS	0.91	0.95	0.94	0.92	0.93
		SNR	10.13	19.89	16.13	10.70	12.27
	Mid	n cores	74	78	78	78	78
		n trees	38	39	39	39	39
		Rbar	0.45	0.47	0.41	0.35	0.43
		EPS	0.97	0.97	0.96	0.96	0.97
		SNR	29.05	33.09	26.51	21.00	29.01
	High	n cores	60	72	78	78	78
		n trees	33	40	41	41	41
		Rbar	0.46	0.43	0.41	0.31	0.41
		EPS	0.95	0.96	0.96	0.95	0.97
		SNR	17.36	23.69	26.39	18.65	28.03
Sugar maple	Low	n cores	45	47	47	48	48
		n trees	30	31	31	31	31
		Rbar	0.16	0.12	0.13	0.22	0.34
		EPS	0.80	0.80	0.82	0.90	0.94
		SNR	3.90	3.97	4.39	8.55	15.89
	Mid	n cores	46	56	57	57	57
		n trees	29	34	34	34	34
		Rbar	0.30	0.25	0.26	0.24	0.19
		EPS	0.89	0.91	0.92	0.92	0.89
		SNR	8.44	9.51	11.64	10.82	8.20
Yellow birch	Mid	n cores	46	56	59	59	59
		n trees	29	32	34	34	34
		Rbar	0.26	0.25	0.16	0.22	0.20
		EPS	0.90	0.91	0.86	0.90	0.90
		SNR	8.74	9.59	6.14	9.45	8.65

^a Low = 450-650 m a.s.l., mid = 750-850 m, and high = 900-1000 m^b R bar = Effective R bar, a weighted average of RBT (R bar between trees) and RWT (R bar within trees) and thus a measure of correlation among and between cores, EPS = expressed population signal, SNR = signal to noise ratio.