Canadian Journal of Microbiology
Revue canadienne de microbiologie

**A review of the current state of antimicrobial susceptibility
test methods for Brachyspira**

Journal:	<i>Canadian Journal of Microbiology</i>
Manuscript ID	cjm-2016-0756.R1
Manuscript Type:	Review
Date Submitted by the Author:	09-Feb-2017
Complete List of Authors:	Kulathunga, Dharmasiri; Western College of Veterinary Medicine, Veterinary Microbiology Rubin, Joseph E.; University of Saskatchewan, Veterinary Microbiology
Keyword:	Brachyspira, Susceptibility test, Method standardization, Veterinary diagnostics, Swine dysentery

A review of the current state of antimicrobial susceptibility test methods for *Brachyspira***D.G.R. S. Kulathunga¹, J.E. Rubin^{1*}**¹Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, CanadaD.G.R.S. Kulathunga - rsd351@mail.usask.caJ.E. Rubin - joe.rubin@usask.ca

*Corresponding Author

52 Campus Drive

Department of Veterinary Microbiology

Phone: (306) 966-7246

Fax: (306) 966-7244

Email: joe.rubin@usask.ca

Draft

Abstract

The re-emergence of swine dysentery (*Brachyspira*-associated muco-haemorrhagic colitis) since the late 2000's has illuminated diagnostic challenges associated with this genus. The methods used to detect, identify and characterize *Brachyspira* from clinical samples have not been standardized and laboratories frequently rely heavily on in-house techniques. Particularly concerning is the lack of standardized methods for determining and interpreting the antimicrobial susceptibility of *Brachyspira* spp. The integration of laboratory data into a treatment plan is a critical component of prudent antimicrobial usage, the lack of standardized methods is therefore an important limitation to the evidence based use of antimicrobials. This review will focus on describing the methodological limitations and inconsistencies between current susceptibility testing schemes employed for *Brachyspira*, provide an overview of what we do know about the susceptibility of these organisms and suggest future directions to improve and standardize diagnostic strategies.

Keywords: *Brachyspira*, susceptibility test, method standardization, veterinary diagnostics, swine dysentery

1.0 Introduction

Brachyspira are a genus of Gram-negative, oxygen-tolerant anaerobic spirochetes which colonize the intestinal tracts of wild and domestic animals and humans (Hampson and Ahmed 2009). Seven species with standing in nomenclature are currently recognized within the genus *Brachyspira*, including the causative agents of swine dysentery and porcine, avian and human intestinal spirochetosis. Swine dysentery, characterized by profuse muco-hemorrhagic colitis, was first described in 1921 in the United States, although the causative agent (*B. hyodysenteriae*) was not isolated until the early 1970s (Harris et al. 1972; Whiting et al. 1921). Since the late 2000s, several additional distinct taxa ('*Brachyspira hampsonii*' and '*Brachyspira suanatina*') have been found to causally associated with this syndrome (Burrough 2017). Swine dysentery, which primarily affects grower-finisher pigs, is the most economically damaging disease associated with *Brachyspira* attributable to both mortality and production limiting sub-optimal feed conversion (Alvarez-Ordóñez et al. 2013). *Brachyspira pilosicoli* has a broad host spectrum including pigs, other domestic animals, wildlife and humans (Trivett-Moore et al. 1998). In pigs, intestinal spirochetosis caused by *Brachyspira pilosicoli* typically affects pigs within 7 to 14 days post-weaning (Trott et al. 1996). Porcine intestinal spirochetosis is characterized by diarrhea often described as having a "wet cement" consistency (Hampson and Duhamel 2006). In chickens, *Brachyspira intermedia* and *pilosicoli* are associated with diarrhea, decreased production and fecal staining of eggs (Mapple et al. 2014). Finally, human intestinal spirochetosis caused by *B. pilosicoli* or *B. aalborgi*, is an infrequent cause of diarrhea most commonly affecting children and immunocompromised individuals (Helbling et al. 2012; Tateishi et al. 2015).

Since the late 2000s the re-emergence of *Brachyspira* associated disease in pigs in North America and the emergence of '*Brachyspira hampsonii*' in Europe have been described (Mahu et al. 2014; Martinez-Lobo et al. 2013; Rubin et al. 2013a). This re-emergence has included the identification of novel taxa causing a dysentery like disease ('*Brachyspira hampsonii*' and '*Brachyspira suanatina*') which have been effectively but not validly described (Chander et al. 2012; Mirajkar et al. 2016b; Mushtaq et al. 2015; Perez et al. 2016). The specific etiological diagnosis of *Brachyspira* associated disease is challenging and relies heavily on PCR and culture based assays; it was therefore not surprising that the emergence of '*B. hampsonii*' has been associated with substantial diagnostic challenges. The economic and animal welfare implications of *Brachyspira* associated disease have led to renewed interest in improving the diagnostic methods used for *Brachyspira*, including antimicrobial susceptibility test methods to facilitate the evidence based use of antimicrobials. The purpose of this review is therefore to summarize the current state of knowledge regarding the susceptibility of *Brachyspira* to antimicrobials, the intrinsic methodological limitations associated with *Brachyspira* and to highlight future research avenues to improve the reproducibility and reliability of *Brachyspira* antimicrobial susceptibility testing.

2.0 General information about *Brachyspira*

2.1 Taxonomy

The genus *Brachyspira* is within the order Brachyspirales, and was recognized as a distinct genus from *Treponema* in 1983 (Gupta et al. 2013; Hovind-Hougen et al. 1982; Oren and Garrity 2014). There are currently seven species with standing in the nomenclature: *Brachyspira hyodysenteriae*, *B. pilosicoli*, *B. murdochii*, *B. innocens*, *B. intermedia*, *B. aalborgi* and *B.*

alvinipulli (Euzeby 1997; Parte 2017). *Brachyspira* require specialized 'non-standard' culture conditions, don't typically form colonies on agar, grow unpredictably in broth and are poorly characterized phenotypically compared to other organisms such as *E. coli*; accurate etiological identification is therefore challenging. In the 1970s pathogenic *Brachyspira* (then *Treponema hyodysenteriae*) were identified as strongly hemolytic on blood containing media while weakly hemolytic spirochetes were considered to be non-pathogens (Kinyon et al. 1977). The identification of multiple, phylogenetically distinct weakly hemolytic *Brachyspira* highlights the limitations of this approach. Molecular tools have proven to be extremely useful diagnostically, although care must be taken to interpret test results considering the limitations of the test such as primer sensitivity and specificity. Recently, MALDI-TOF has been explored for the identification of *Brachyspira* species and shows promise (Calderaro et al. 2013; Prohaska et al. 2014; Warneke et al. 2014). Identification by phylogenetic analysis, particularly of the NADH oxidase (*nox*) gene has proven useful and provides comparable, objective data which is portable between labs. Sequence analysis at present is considered the gold standard for species level identification (Perez et al. 2016; Rohde et al. 2014). DNA sequencing has revealed phylogenetic diversity within the genus *Brachyspira*, including the novel strain 'B. hampsonii', and a large number of other strains which have not yet been characterized but appear to be distinct from recognized species (Patterson et al. 2013; Rubin et al. 2013b).

2.2 Treatment

In pigs *Brachyspira* associated disease has been effectively treated with the pleuromutilins, macrolides/lincosamides and carbadox while metronidazole has been effectively used in people with intestinal spirochetosis (Hampson 2012; Helbling et al. 2012). However, the ability to use

some of these agents in food producing animals is restricted in many jurisdictions; metronidazole is banned for use in food animals in North America and the European Union, and carbadox is banned in Canada, the EU and the UK and recently had its approval rescinded in the United States (FDA 2016). The pleuromutilins (tiamulin and valnemulin) and the macrolide/lincosamides (tylosin, lincomycin and tylvalosin) are the mainstay of anti-*Brachyspira* therapy in swine medicine, although a number of other antimicrobials including bacitracin, virginiamycin and gentamicin are labeled for the treatment or prevention of *Brachyspira* associated diseases in pigs (Table 1) (Hampson 2012). The heavy reliance on mechanistically similar drugs, to which resistance in other organisms has been shown to develop by a common mechanism highlights the potential impact of resistance emergence and the clinical value of laboratory test guided therapeutic selection. Although there is no consensus method for determining the antimicrobial susceptibility of *Brachyspira*, trends of decreasing susceptibility among *B. hyodysenteriae* for the macrolides and pleuromutilins have been described in Europe and the United States (Aarestrup et al. 2008; Lobova et al. 2004; Mirajkar et al. 2016a; Prasek et al. 2014; Rugna et al. 2015).

3.0 Antimicrobial susceptibility testing

3.1 Importance of susceptibility testing to the prudent use of antimicrobials

3.1.1 Antimicrobial resistance

The term "antimicrobial resistance" is often used imprecisely. The categorization of an organism as susceptible or resistant based on an *in vitro* test is designed to be clinically predictive, viewing bacterial drug susceptibility from the patient's perspective. Susceptibility indicates a high likelihood of clinical success, while resistance suggests a low likelihood of treatment success.

Clinical breakpoints are therefore necessarily predicated on assumptions including the species being treated, target pathogen, site of infection and treatment regimen (dose, dosing frequency and route of administration) (Dalhoff et al. 2009; Mouton et al. 2012).

Bacterial drug resistance can be intrinsic or acquired, and mechanistically falls into one of four broad categories: (i) enzymatic inactivation of the antibiotic (ii) decreased permeability of the cell wall (iii) active efflux of the antimicrobial from within the cell and (iv) alteration or absence of drug binding sites (Boerlin and White 2013). Intrinsic resistance is constitutive to the physiology of a particular organism, examples include penicillin resistance among Enterobacteriaceae and cephalosporin resistance in *Enterococcus* spp. Conversely, organisms with a susceptible wild-type can become resistant as a result of chromosomal mutation or the acquisition of resistance genes through transformation, transduction or conjugation (Boerlin and White 2013).

3.1.2 Importance of susceptibility testing for resistance surveillance

Beyond the clinical applicability of antimicrobial susceptibility test results, there is value in collecting this data for resistance surveillance. The purpose of resistance surveillance is to identify temporal changes in antimicrobial susceptibility and to detect the emergence of novel resistance phenotypes or genotypes. Globally, harmonization of test methodology, drug panels tested and interpretive criteria, is recognized as a critical step to ensure that data generated in different laboratories is comparable (OIE 2003). Substantial obstacles facing the detection of emerging resistance in *Brachyspira* therefore include the lack of standardized test methods and interpretive criteria, and consensus regarding which drugs should be included in test panels and

what concentrations should be tested. The description of the emergence of resistance to anti-*Brachyspira* therapies has at best relied upon small scale, geographically limited, single laboratory studies, and at worse the spurious comparison of data generated using different methods.

Epidemiological cut-offs based on distributions of MICs of bacterial populations, may prove to be useful for resistance surveillance. Epidemiological cut-offs, based on MIC distributions of bacterial populations, are useful for detecting organisms with acquired resistance that differentiates them from the wild type population but does not address questions of clinical efficacy because they do not consider host factors (Dalhoff et al. 2009).

3.2 General information on susceptibility testing methods

Early in the antibiotic era, the development of antimicrobial susceptibility test methods grew from the need to select the most appropriate therapy in the face of emerging resistance and the availability of new, mechanistically distinct compounds. The requirement for high quality information led to the standardization of test methods which improved comparability of results between labs, and allowed the development of uniform interpretive criteria (when to classify an isolate as susceptible or resistant) (Ericsson and Sherris 1971). Regularly updated internationally recognized standardized methods are now published by the Clinical and Laboratory Standards Institute (CLSI) in the United States, and the European Committee on Antimicrobial Susceptibility Testing (EUCAST) in Europe. Because test results vary with divergences from standard methodologies, test guidelines are necessarily prescriptive in terms of the density and growth phase of the test culture, test media composition including pH and cation concentration,

incubation temperature, time and atmosphere and endpoint definition (CLSI 2012). The effects of test chemistry on MIC are well recognized but vary by drug and organism. For example test media with too low a pH may yield decreased penicillin MICs or increased quinolone MICs (CLSI 2017). Other test factors are intuitively more universal; for instance low MICs may be an artifact of too low a test inoculum, while high MICs result from testing too many organisms (CLSI 2017). Although test methods specific to a wide variety of anaerobic and fastidious organisms (ex. HACEK group organisms, *Helicobacter*, *Listeria*, *Moraxella*, *Campylobacter*, *Actinobacillus pleuropneumoniae*) are available, no standardized protocols (human or veterinary) addressing the unique requirements of *Brachyspira* have been published (CLSI 2015, 2016, 2017). The methods described by the CLSI and EUCAST all rely on phenotypic indicators of susceptibility, either growth or inhibition of growth in the presence of antimicrobial. Unambiguous and standardized endpoint definition is therefore essential (Jenkins and Schuetz 2012).

Molecular techniques (PCR) are increasingly used to identify the presence of resistance genes and therefore infer resistance. PCR has the advantage of detecting resistance without requiring classical phenotypic susceptibility test methods, a potential benefit when characterizing *Brachyspira*. However, the use of molecular methods requires knowledge of resistance genes or resistance conferring mutations. It must also be recognized that while these methods can identify resistance, they are insufficient to confirm susceptibility due to the potential presence of novel resistance genes or resistance conferring mutations not included in the assay.

4.0 Specific challenges to susceptibility testing of *Brachyspira*

The ability to assess the antimicrobial susceptibility of *Brachyspira* is hindered by two proximate causes: 1. the growth conditions required for *Brachyspira* are different than those standardized for susceptibility testing and 2. the inability to use typical susceptibility test endpoints. Although *Brachyspira* are reported to grow within the range of conditions prescribed by the CLSI for antimicrobial susceptibility testing, no single set of conditions have been standardized for testing these organisms. The unusual growth characteristics of *Brachyspira* sp. are at the root of many of these challenges. On solid media *Brachyspira* do not form distinct colonies, growth is typically inferred by the presence of hemolytic zones on blood containing agar (Figure 1). This property complicates/precludes the application of the concept of the colony forming unit, a foundational bacteriological principle which allows viable cells to be enumerated, and genetic clones to be isolated. The inability to differentiate whether apparently distinct hemolytic zones represent unique founding organisms or multiple individuals is a critical limitation. Pure cultures are a prerequisite for antimicrobial susceptibility testing, tests of mixed cultures yield cumulative results which may not reflect the susceptibility of either organism individually. Furthermore, this characteristic potentially precludes reliable test endpoint definition; whereas an antimicrobial inhibiting the formation of a colony is taken to indicate growth inhibition, failure to observe haemolysis in the presence of an antimicrobial may simply reflect inhibition of haemolysin production with or without affects on microbial growth. This is particularly germane for *Brachyspira* where protein synthesis inhibiting drugs are the primary agents of interest. Creative approaches to address endpoint validity including microscopic evaluation or sub-culture of media may prove useful in identifying viable organisms in media containing supra-MIC drug concentrations.

Broth culture of *Brachyspira* spp. also presents some unique challenges. Anecdotal observations suggest that broth cultures require a high starting inoculum and so failure to grow an organism in broth may simply reflect an insufficient initial inoculum density. This complicates two steps of the susceptibility testing process: 1. growing an initial broth culture used to inoculate test media, and 2. interpreting broth based susceptibility tests - did the culture not grow because it was inhibited by the drug, or because the culture was insufficiently dense?

5.0 Summary of susceptibility test methods which have been used for *Brachyspira*

The English language literature through 2017 was searched and studies describing antimicrobial susceptibility testing of *Brachyspira* were reviewed. Both agar and broth dilution techniques were used although there was substantial variability in the methods reported (Table 2).

Incubation temperatures ranged from 37 – 42 °C with incubation times ranging from 2 to 5 days for agar dilution, and 37 – 38 °C and 3-5 days for broth dilution respectively. Where reported, highly variable inoculum densities were described: 1×10^5 - 5×10^6 CFU/ml for broth dilution and 10^4 - 10^6 CFU/spot for agar dilution tests. Finally, where reported atmospheres with varying composition were used. According to the manufacturers all anaerobic systems yield a final O₂ concentration of $\leq 1\%$, but yield highly variable CO₂ concentrations (7- >15%). A 2016 study has also described the use of doxycycline gradient strips although this method is not widely used (Mirajkar and Gebhart 2016). Further complicating the interpretation of susceptibility test results, is the lack of agreed upon interpretive criteria. With the lack of CLSI and EUCAST resistance breakpoints, investigators have relied upon a number of divergent sets of researcher proposed breakpoints (Burch 2005; Duhamel et al. 1998; Pringle et al. 2012; Ronne and Szancer 1990; SVARM 2014).

The introduction of the VetMIC *Brachyspira* broth micro-dilution system, first described in 2001, has led to substantial progress towards test standardization (Karlsson et al. 2004a; Karlsson et al. 2001). This product includes serial two-fold dilutions of dried antimicrobials in tissue culture trays. Testing is done by inoculating wells with 500µl of broth culture and incubating anaerobically at 37°C for 4 days with agitation (SVA 2011). Antimicrobial MICs as determined using this method have been shown to be reproducible, although interestingly typically yield a doubling dilution lower MIC than agar dilution tests (Rohde et al. 2004).

6.0 Challenges in comparing susceptibility data generated using non-standard tests

Because, there are no established standard test methods, interpretive criteria comparison of results generated by different labs must only be done with extreme caution. This inability to compare data was exemplified by a 2005 study where eight European laboratories participated in a ring test of *Brachyspira* diagnostics (Rasback et al. 2005). Each laboratory received samples of pig feces seeded with a previously identified *Brachyspira* isolate at specific concentrations and two pure cultures of known *Brachyspira* for susceptibility testing. Antibiotic susceptibility test methods differed by media composition and type, incubation temperature and time. Only a very limited description of the methods used by each laboratory for susceptibility testing and interpretation was presented. Not surprisingly results were highly variable; errors including failure to detect or identify isolates or discordant susceptibility test results occurred 32% of the time (Rasback et al. 2005). The varying concentration of antimicrobials tested made comparison of MICs between labs impossible even when the results were not inconsistent (ex. different labs reported lincomycin MIC of ≤ 4 , ≤ 2 and ≤ 0.5 for the same isolate). The lack of validated

Brachyspira specific susceptibility test methods has resulted in the development of a variety of "in-house" techniques, consequently susceptibility data are not reliably comparable between labs. Further confounding susceptibility data portability is the lack of a consensus scheme for determining species level identification (Rasback et al. 2005). Researchers have used combinations of conventional methods (ex. culture and biochemical tests) and a variety of molecular methods (ex. *nox* and 16S-rDNA sequencing and *nox*-RFLP) which are recognized to have varying levels of sensitivity and specificity.

7.0 Objective measures of resistance

Genetic associations with decreased susceptibility to a number of drugs have been elucidated in several *Brachyspira* spp. (Table 3). Most published studies have focused on the association between single nucleotide polymorphisms (SNPs) in genes encoding ribosomal RNA and ribosomal proteins and increased MIC to drugs which act by inhibiting protein synthesis inhibitors including macrolides, lincosamides, pleuromutilins, doxycycline, chloramphenicol and florfenicol (Hidalgo et al. 2011; Hillen et al. 2014; Pringle et al. 2007; Pringle et al. 2004; Verlinden et al. 2011). Narrow spectrum β -lactamases including 13 enzymes related to OXA-63 have also been identified in *B. pilosicoli* (La et al. 2015; Meziane-Cherif et al. 2008; Mortimer-Jones et al. 2008). Differences in intrinsic resistance, even between closely related bacteria can inform treatment decisions based on accurate organism identification. For example, differences in the intrinsic resistance to vancomycin of *Enterococcus gallinarum* but not *Enterococcus faecalis* or *faecium* (CLSI 2017). Among *Brachyspira* species differences in susceptibility have also been observed, one study reported significant differences between species (Clothier et al.

2011). Although this study was small, it highlights the possibility of identifying taxa-specific differences in intrinsic resistance which may prove useful to clinicians for therapeutic selection.

8.0 Summary and future directions

The lack of standardized antimicrobial susceptibility test methods and interpretive criteria for *Brachyspira* is an important limitation to the evidence based use of antimicrobials for treating clinical disease. These diagnostic limitations have been made acutely problematic by the re-emergence of *Brachyspira* associated disease in North America, and the emergence of novel taxa in Europe. Furthermore, the increasing public scrutiny of antimicrobial usage in agriculture provides constant pressure to ensure that prescribing practices are evidence based and prudent. Unfortunately, for the reasons described here, we do not recommend that diagnostic laboratories routinely report categorical susceptibility test results for *Brachyspira* spp. Instead, we encourage our colleagues to continue their research describing temporal changes in MIC and archiving *Brachyspira* culture collections to facilitate the eventual development of agreed upon test methodology and interpretive criteria.

The lack of consensus on how to identify an isolate to the species level is the most immediate obstacle to improving laboratory diagnostics for *Brachyspira*. A common ontology is critical for researchers and diagnosticians to make simple comparisons of laboratory prevalence, describe the distribution of particular taxa or to compare the drug susceptibility of two organisms. The next most urgent need is an enriched understanding the genetic determinants of resistance. These genes or mutations have the potential to serve as an objective data point against which phenotypic susceptibility can be benchmarked, potentially serving as a pillar of quality control.

Finally, we call for additional collaboration in the global *Brachyspira* diagnostic/research communities. Collaboration and consensus building is going to be a critical step in identifying and recommending best practices for *Brachyspira*. This leadership was exemplified by Rasback et al., 2005 where the results of a multi-centre ring test were described; such investigations will be essential not only for ensuring method portability and objectivity, but also as a feedback mechanism to identify areas for improvement in individual laboratories.

Acknowledgments

The authors would like to thank Elanco (formerly Novartis Animal Health) and Agriculture and Agri-Food Canada through Swine Innovation Porc for funding this investigation and providing stipend support for DGRSK. Stipend support from the Department of Veterinary Microbiology devolved fund is also appreciated. We would like to acknowledge Mo Patterson for the photograph depicted in Figure 1. Finally the authors would like to thank the veterinarians and producers of the Canadian swine industry for their continued collaboration and support of our research.

References Cited

- Aarestrup, F.M., Oliver Duran, C., and Burch, D.G. 2008. Antimicrobial resistance in swine production. *Anim Health Res Rev* **9**(2): 135-148. doi: 10.1017/S1466252308001503.
- Alvarez-Ordóñez, A., Martínez-Lobo, F.J., Arguello, H., Carvajal, A., and Rubio, P. 2013. Swine dysentery: aetiology, pathogenicity, determinants of transmission and the fight against the disease. *Int J Environ Res Public Health* **10**(5): 1927-1947. doi: 10.3390/ijerph10051927.
- Boerlin, P., and White, D.G. 2013. Antimicrobial resistance and its epidemiology. *In* Antimicrobial Therapy in Veterinary Medicine. Edited by S. Giguere and J.F. Prescott and P.M. Dowling. John Wiley & Sons, Inc, Ames, Iowa. pp. 21-40.
- Burch, D.G.S. 2005. Pharmacokinetic, pharmacodynamic and clinical correlations relating to the therapy of colonic infections in the pig and breakpoint determinations. *The Pig Journal* **56**: 8-24.
- Burrough, E.R. 2017. Swine Dysentery. *Vet Pathol* **54**(1): 22-31. doi: 10.1177/0300985816653795.
- Calderaro, A., Piccolo, G., Montecchini, S., Buttrini, M., Gorrini, C., Rossi, S., Arcangeletti, M.C., De Conto, F., Medici, M.C., and Chezzi, C. 2013. MALDI-TOF MS analysis of human and animal *Brachyspira* species and benefits of database extension. *J Proteomics* **78**: 273-280. doi: 10.1016/j.jprot.2012.09.027.
- Chander, Y., Primus, A., Oliveira, S., and Gebhart, C.J. 2012. Phenotypic and molecular characterization of a novel strongly hemolytic *Brachyspira* species, provisionally designated "*Brachyspira hampsonii*". *J Vet Diagn Invest* **24**(5): 903-910. doi: 1040638712456975 [pii] 10.1177/1040638712456975.
- Clothier, K.A., Kinyon, J.M., Frana, T.S., Naberhaus, N., Bower, L., Strait, E.L., and Schwartz, K. 2011. Species characterization and minimum inhibitory concentration patterns of *Brachyspira*

species isolates from swine with clinical disease. J Vet Diagn Invest **23**(6): 1140-1145. doi: 23/6/1140 [pii]

10.1177/1040638711425580.

CLSI. 2012. M07-A9 Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard - ninth edition. Clinical and Laboratory Standards Institute.

CLSI. 2015. VET01S Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals. Clinical and Laboratory Standards Institute.

CLSI. 2016. M45-Ed3 Methods for antimicrobial dilution and disk susceptibility testing of infrequently isolated or fastidious bacteria. Clinical and Laboratory Standards Institute.

CLSI. 2017. M100-S27 Performance standards for antimicrobial susceptibility testing. Clinical and Laboratory Standards Institute.

Dalhoff, A., Ambrose, P.G., and Mouton, J.W. 2009. A long journey from minimum inhibitory concentration testing to clinically predictive breakpoints: deterministic and probabilistic approaches in deriving breakpoints. Infection **37**(4): 296-305. doi: 10.1007/s15010-009-7108-9.

Duhamel, G.E., Kinyon, J.M., Mathiesen, M.R., Murphy, D.P., and Walter, D. 1998. In vitro activity of four antimicrobial agents against North American isolates of porcine *Serpulina pilosicoli*. J Vet Diagn Invest **10**(4): 350-356.

Ericsson, H.M., and Sherris, J.C. 1971. Antibiotic sensitivity testing. Report of an international collaborative study. Acta Pathol Microbiol Scand B Microbiol Immunol **217**: Suppl 217:211+.

Euzeby, J.P. 1997. List of Bacterial Names with Standing in Nomenclature: a folder available on the Internet. Int J Syst Bacteriol **47**(2): 590-592. doi: 10.1099/00207713-47-2-590.

FDA, U.S. 2016. FDA takes steps to withdraw approval of the swine drug carbadox due to safety concerns. United States Food and Drug Administration, Silver Spring, MD.

- Gupta, R.S., Mahmood, S., and Adeolu, M. 2013. A phylogenomic and molecular signature based approach for characterization of the phylum Spirochaetes and its major clades: proposal for a taxonomic revision of the phylum. *Front Microbiol* **4**: 217. doi: 10.3389/fmicb.2013.00217.
- Hampson, D.J. 2012. Brachyspiral colitis. *In Diseases of Swine. Edited by J.J. Zimmerman and L.A. Karriker and A. Ramirez and K.J. Schwartz and G.W. Stevenson.* John Wiley & Sons. pp. 680-696.
- Hampson, D.J., and Ahmed, N. 2009. Spirochaetes as intestinal pathogens: lessons from a *Brachyspira* genome. *Gut Pathog* **1**(1): 10. doi: 10.1186/1757-4749-1-10.
- Hampson, D.J., and Duhamel, G.E. 2006. Porcine colonic spirochetosis / intestinal spirochetosis. *In Diseases of Swine. Edited by B.E. Straw and J.J. Zimmerman and S. D'Allaire and D.J. Taylor.* Blackwell Publishing, Ames, IO. pp. 755-767.
- Harris, D.L., Kinyon, J.M., Mullin, M.T., and Glock, R.D. 1972. Isolation and propagation of spirochetes from the colon of swine dysentery affected pigs. *Can J Comp Med* **36**(1): 74-76.
- Helbling, R., Osterheld, M.C., Vaudaux, B., Jatton, K., and Nydegger, A. 2012. Intestinal spirochetosis mimicking inflammatory bowel disease in children. *BMC Pediatr* **12**: 163. doi: 10.1186/1471-2431-12-163.
- Herbst, W., Schlez, K., Heuser, J., and Baljer, G. 2014. Antimicrobial susceptibility of *Brachyspira hyodysenteriae* determined by a broth microdilution method. *Vet Rec* **174**(15): 382. doi: 10.1136/vr.102169.
- Hidalgo, A., Carvajal, A., Garcia-Feliz, C., Osorio, J., and Rubio, P. 2009. Antimicrobial susceptibility testing of Spanish field isolates of *Brachyspira hyodysenteriae*. *Res Vet Sci* **87**(1): 7-12. doi: 10.1016/j.rvsc.2008.10.017.

- Hidalgo, A., Carvajal, A., Vester, B., Pringle, M., Naharro, G., and Rubio, P. 2011. Trends towards lower antimicrobial susceptibility and characterization of acquired resistance among clinical isolates of *Brachyspira hyodysenteriae* in Spain. *Antimicrob Agents Chemother* **55**(7): 3330-3337. doi: 10.1128/AAC.01749-10.
- Hillen, S., Willems, H., Herbst, W., Rohde, J., and Reiner, G. 2014. Mutations in the 50S ribosomal subunit of *Brachyspira hyodysenteriae* associated with altered minimum inhibitory concentrations of pleuromutilins. *Vet Microbiol* **172**(1-2): 223-229. doi: 10.1016/j.vetmic.2014.04.021.
- Hovind-Hougen, K., Birch-Andersen, A., Henrik-Nielsen, R., Orholm, M., Pedersen, J.O., Teglbjaerg, P.S., and Thaysen, E.H. 1982. Intestinal spirochetosis: morphological characterization and cultivation of the spirochete *Brachyspira aalborgi* gen. nov., sp. nov. *J Clin Microbiol* **16**(6): 1127-1136.
- Jenkins, S.G., and Schuetz, A.N. 2012. Current concepts in laboratory testing to guide antimicrobial therapy. *Mayo Clin Proc* **87**(3): 290-308. doi: 10.1016/j.mayocp.2012.01.007.
- Kajiwara, K., Kozawa, M., Kanazawa, T., Uetsuka, K., Nakajima, H., and Adachi, Y. 2015. Drug-susceptibility of isolates of *Brachyspira hyodysenteriae* isolated from colonic mucosal specimens of pigs collected from slaughter houses in Japan in 2009. *J Vet Med Sci*. doi: 10.1292/jvms.15-0608.
- Karlsson, M., Aspan, A., Landen, A., and Franklin, A. 2004a. Further characterization of porcine *Brachyspira hyodysenteriae* isolates with decreased susceptibility to tiamulin. *J Med Microbiol* **53**(Pt 4): 281-285. doi: 10.1099/jmm.0.05395-0.

- Karlsson, M., Fellstrom, C., Gunnarsson, A., Landen, A., and Franklin, A. 2003. Antimicrobial susceptibility testing of porcine *Brachyspira (Serpulina)* species isolates. *J Clin Microbiol* **41**(6): 2596-2604.
- Karlsson, M., Fellstrom, C., Heldtander, M.U., Johansson, K.E., and Franklin, A. 1999. Genetic basis of macrolide and lincosamide resistance in *Brachyspira (Serpulina) hyodysenteriae*. *FEMS Microbiol Lett* **172**(2): 255-260.
- Karlsson, M., Fellstrom, C., Johansson, K.E., and Franklin, A. 2004b. Antimicrobial resistance in *Brachyspira pilosicoli* with special reference to point mutations in the 23S rRNA gene associated with macrolide and lincosamide resistance. *Microb Drug Resist* **10**(3): 204-208. doi: 10.1089/mdr.2004.10.204.
- Karlsson, M., Gunnarsson, A., and Franklin, A. 2001. Susceptibility to pleuromutilins in *Brachyspira (Serpulina) hyodysenteriae*. *Anim Health Res Rev* **2**(1): 59-65.
- Karlsson, M., Oxberry, S.L., and Hampson, D.J. 2002. Antimicrobial susceptibility testing of Australian isolates of *Brachyspira hyodysenteriae* using a new broth dilution method. *Vet Microbiol* **84**(1-2): 123-133.
- Kinyon, J.M., Harris, D.L., and Glock, R.D. 1977. Enteropathogenicity of various isolates of *Treponema hyodysenteriae*. *Infect Immun* **15**(2): 638-646.
- La, T., Neo, E., Phillips, N.D., and Hampson, D.J. 2015. Genes encoding ten newly designated OXA-63 group class D beta-lactamases identified in strains of the pathogenic intestinal spirochaete *Brachyspira pilosicoli*. *J Med Microbiol* **64**(11): 1425-1435. doi: 10.1099/jmm.0.000162.

- Lim, S., Lee, H., Nam, H., Cho, Y.S., Jung, S., and Joo, Y. 2012. Prevalence and antimicrobial susceptibility of *Brachyspira* species in pigs in Korea. Korean Journal of Veterinary Research **52**(4): 253-257.
- Lobova, D., Smola, J., and Cizek, A. 2004. Decreased susceptibility to tiamulin and valnemulin among Czech isolates of *Brachyspira hyodysenteriae*. J Med Microbiol **53**(Pt 4): 287-291. doi: 10.1099/jmm.0.05407-0.
- Mahu, M., de Jong, E., De Pauw, N., Vande Maele, L., Vandenbroucke, V., Vandersmissen, T., Miry, C., Pasmans, F., Haesebrouck, F., Martel, A., and Boyen, F. 2014. First isolation of "*Brachyspira hampsonii*" from pigs in Europe. Vet Rec **174**(2): 47. doi: 10.1136/vr.101868.
- Mappley, L.J., La Ragione, R.M., and Woodward, M.J. 2014. *Brachyspira* and its role in avian intestinal spirochaetosis. Vet Microbiol **168**(2-4): 245-260. doi: 10.1016/j.vetmic.2013.11.019.
- Martinez-Lobo, F.J., Hidalgo, A., Garcia, M., Arguello, H., Naharro, G., Carvajal, A., and Rubio, P. 2013. First identification of "*Brachyspira hampsonii*" in wild European waterfowl. PLoS One **8**(12): e82626. doi: 10.1371/journal.pone.0082626.
- Meziane-Cherif, D., Lambert, T., Dupechez, M., Courvalin, P., and Galimand, M. 2008. Genetic and biochemical characterization of OXA-63, a new class D beta-lactamase from *Brachyspira pilosicoli* BM4442. Antimicrob Agents Chemother **52**(4): 1264-1268. doi: 10.1128/AAC.00684-07.
- Mirajkar, N.S., Davies, P.R., and Gebhart, C.J. 2016a. Antimicrobial susceptibility patterns of *Brachyspira* species isolated from swine herds in the United States. J Clin Microbiol. doi: 10.1128/JCM.00834-16.

- Mirajkar, N.S., and Gebhart, C.J. 2016. Comparison of agar dilution and antibiotic gradient strip test with broth microdilution for susceptibility testing of swine *Brachyspira* species. J Vet Diagn Invest **28**(2): 133-143. doi: 10.1177/1040638716629154.
- Mirajkar, N.S., Phillips, N.D., La, T., Hampson, D.J., and Gebhart, C.J. 2016b. Characterization and Recognition of *Brachyspira hampsonii* sp. nov., a Novel Intestinal Spirochete That Is Pathogenic to Pigs. J Clin Microbiol **54**(12): 2942-2949. doi: 10.1128/JCM.01717-16.
- Mortimer-Jones, S.M., Phillips, N.D., La, T., Naresh, R., and Hampson, D.J. 2008. Penicillin resistance in the intestinal spirochaete *Brachyspira pilosicoli* associated with OXA-136 and OXA-137, two new variants of the class D beta-lactamase OXA-63. J Med Microbiol **57**(Pt 9): 1122-1128. doi: 10.1099/jmm.0.2008/001552-0.
- Mouton, J.W., Brown, D.F., Apfalter, P., Canton, R., Giske, C.G., Ivanova, M., MacGowan, A.P., Rodloff, A., Soussy, C.J., Steinbakk, M., and Kahlmeter, G. 2012. The role of pharmacokinetics/pharmacodynamics in setting clinical MIC breakpoints: the EUCAST approach. Clin Microbiol Infect **18**(3): E37-45. doi: 10.1111/j.1469-0691.2011.03752.x.
- Mushtaq, M., Zubair, S., Rasback, T., Bongcam-Rudloff, E., and Jansson, D.S. 2015. *Brachyspira suanatina* sp. nov., an enteropathogenic intestinal spirochaete isolated from pigs and mallards: genomic and phenotypic characteristics. BMC Microbiol **15**: 208. doi: 10.1186/s12866-015-0537-y.
- Ohya, T., and Sueyoshi, M. 2010. In vitro antimicrobial susceptibility of *Brachyspira hyodysenteriae* strains isolated in Japan from 1985 to 2009. J Vet Med Sci **72**(12): 1651-1653.
- OIE. 2003. Terrestrial Animal Health Code Guidelines for the harmonisation of antimicrobial resistance surveillance and monitoring programs. In OIE International Standards on Antimicrobial Resistance. OIE (World Organization for Animal Health), Paris, France. pp. 5-27.

- Oren, A., and Garrity, G.M. 2014. List of new names and new combinations previously effectively, but not validly, published. Validation List no. 156. *Int J Syst Evol Microbiol* **64**: 693-696.
- Parte, A.C. 2017. Genus *Brachyspira*. Available from <http://www.bacterio.net/brachyspira.html> [accessed February 7, 2017 2017].
- Patterson, A.H., Rubin, J.E., Fernando, C., Costa, M.O., Harding, J.C., and Hill, J.E. 2013. Fecal shedding of *Brachyspira* spp. on a farrow-to-finish swine farm with a clinical history of "*Brachyspira hampsonii*"-associated colitis. *BMC Vet Res* **9**: 137. doi: 1746-6148-9-137 [pii] 10.1186/1746-6148-9-137.
- Perez, J.B.D.S., Rubin, J.E., Fernando, C., Harding, J.C.S., and Hill, J.E. 2016. Characterization of "*Brachyspira hampsonii*" clades I and II isolated from commercial swine in Western Canada. *FACETS* **1**: 163-172. doi: 10.1139/facets-2016-0018.
- Prasek, J., Sperling, D., Lobova, D., Smola, J., and Cizek, A. 2014. Antibiotic susceptibility of *Brachyspira hyodysenteriae* isolates from Czech swine farms: a 10-year study. *Acta Veterinaria Brno* **83**: 3-7. doi: doi:10.2754/avb201483010003.
- Pringle, M., Fellstrom, C., and Johansson, K.E. 2007. Decreased susceptibility to doxycycline associated with a 16S rRNA gene mutation in *Brachyspira hyodysenteriae*. *Vet Microbiol* **123**(1-3): 245-248. doi: 10.1016/j.vetmic.2007.02.019.
- Pringle, M., Landen, A., and Franklin, A. 2006. Tiamulin resistance in porcine *Brachyspira pilosicoli* isolates. *Res Vet Sci* **80**(1): 1-4. doi: 10.1016/j.rvsc.2005.02.004.
- Pringle, M., Landen, A., Unnerstad, H.E., Molander, B., and Bengtsson, B. 2012. Antimicrobial susceptibility of porcine *Brachyspira hyodysenteriae* and *Brachyspira pilosicoli* isolated in Sweden between 1990 and 2010. *Acta Vet Scand* **54**: 54. doi: 10.1186/1751-0147-54-54.

- Pringle, M., Poehlsgaard, J., Vester, B., and Long, K.S. 2004. Mutations in ribosomal protein L3 and 23S ribosomal RNA at the peptidyl transferase centre are associated with reduced susceptibility to tiamulin in *Brachyspira* spp. isolates. *Mol Microbiol* **54**(5): 1295-1306. doi: 10.1111/j.1365-2958.2004.04373.x.
- Prohaska, S., Pfluger, V., Ziegler, D., Scherrer, S., Frei, D., Lehmann, A., Wittenbrink, M.M., and Huber, H. 2014. MALDI-TOF MS for identification of porcine *Brachyspira* species. *Lett Appl Microbiol* **58**(3): 292-298. doi: 10.1111/lam.12189.
- Rasback, T., Fellstrom, C., Bergsjö, B., Cizek, A., Collin, K., Gunnarsson, A., Jensen, S.M., Mars, A., Thomson, J., Vyt, P., and Pringle, M. 2005. Assessment of diagnostics and antimicrobial susceptibility testing of *Brachyspira* species using a ring test. *Vet Microbiol* **109**(3-4): 229-243. doi: S0378-1135(05)00185-9 [pii]
10.1016/j.vetmic.2005.05.009.
- Rohde, J., Habighorst-Blome, K., and Seehusen, F. 2014. "*Brachyspira hampsonii*" clade I isolated from Belgian pigs imported to Germany. *Vet Microbiol* **168**(2-4): 432-435. doi: 10.1016/j.vetmic.2013.11.016.
- Rohde, J., Kessler, M., Baums, C.G., and Amtsberg, G. 2004. Comparison of methods for antimicrobial susceptibility testing and MIC values for pleuromutilin drugs for *Brachyspira hyodysenteriae* isolated in Germany. *Vet Microbiol* **102**(1-2): 25-32. doi: 10.1016/j.vetmic.2004.03.023.
- Ronne, H., and Szancer, J. 1990. *In vitro* susceptibility of Danish field isolates of *Treponema hyodysenteriae* to chemotherapeutics in swine dysentery (SD) therapy. Interpretation of MIC results based on the pharmacokinetic properties of the antimicrobial agents. In 11th International Pig Veterinary Society, Lausanne, Switzerland.

Rubin, J.E., Costa, M.O., Hill, J.E., Kittrell, H.E., Fernando, C., Huang, Y., O'Connor, B., and Harding, J.C. 2013a. Reproduction of mucohaemorrhagic diarrhea and colitis indistinguishable from swine dysentery following experimental inoculation with "*Brachyspira hampsonii*" strain 30446. PLoS One **8**(2): e57146. doi: 10.1371/journal.pone.0057146

PONE-D-12-28322 [pii].

Rubin, J.E., Harms, N.J., Fernando, C., Soos, C., Detmer, S.E., Harding, J.C., and Hill, J.E. 2013b. Isolation and characterization of *Brachyspira* spp. including "*Brachyspira hampsonii*" from lesser snow geese (*Chen caerulescens caerulescens*) in the Canadian Arctic. Microb Ecol **66**(4): 813-822. doi: 10.1007/s00248-013-0273-5.

Rugna, G., Bonilauri, P., Carra, E., Bergamini, F., Luppi, A., Gherpelli, Y., Magistrali, C.F., Nigrelli, A., Alborali, G.L., Martelli, P., La, T., Hampson, D.J., and Merialdi, G. 2015. Sequence types and pleuromutilin susceptibility of *Brachyspira hyodysenteriae* isolates from Italian pigs with swine dysentery: 2003-2012. Vet J **203**(1): 115-119. doi: 10.1016/j.tvjl.2014.10.033.

Sperling, D., Smola, J., and Cizek, A. 2011. Characterisation of multiresistant *Brachyspira hyodysenteriae* isolates from Czech pig farms. Vet Rec **168**(8): 215. doi: 10.1136/vr.c4247.

SVA. 2011. VetMIC Brachy for antimicrobial susceptibility testing of *Brachyspira* spp. [Product insert]. Available from

http://www.sva.se/globalassets/redesign2011/pdf/analyser_produkter/vetmic/vetmic_brachy.pdf [accessed May 5, 2016 2016].

SVARM. 2014. 2013 SWEDRES SVARM. Use of antimicrobials and occurrence of antimicrobial resistance in Sweden.

Tateishi, Y., Takahashi, M., Horiguchi, S., Funata, N., Koizumi, K., Okudela, K., Hishima, T., and Ohashi, K. 2015. Clinicopathologic study of intestinal spirochetosis in Japan with special

reference to human immunodeficiency virus infection status and species types: analysis of 5265 consecutive colorectal biopsies. BMC Infect Dis **15**: 13. doi: 10.1186/s12879-014-0736-4.

Trivett-Moore, N.L., Gilbert, G.L., Law, C.L., Trott, D.J., and Hampson, D.J. 1998. Isolation of *Serpulina pilosicoli* from rectal biopsy specimens showing evidence of intestinal spirochetosis. J Clin Microbiol **36**(1): 261-265.

Trott, D.J., Stanton, T.B., Jensen, N.S., Duhamel, G.E., Johnson, J.L., and Hampson, D.J. 1996. *Serpulina pilosicoli* sp. nov., the agent of porcine intestinal spirochetosis. Int J Syst Bacteriol **46**(1): 206-215.

Uezato, K., Kinjo, E., and Adachi, Y. 2004. In vitro susceptibility of 21 antimicrobial agents to 37 isolates of *Brachyspira hyodysenteriae* isolated from pigs in Okinawa Prefecture. J Vet Med Sci **66**(3): 307-309.

Verlinden, M., Boyen, F., Pasmans, F., Garmyn, A., Haesebrouck, F., and Martel, A. 2011. Antimicrobial susceptibility pattern of *Brachyspira intermedia* isolates from European layers. Microb Drug Resist **17**(3): 485-488. doi: 10.1089/mdr.2011.0082.

Warneke, H.L., Kinyon, J.M., Bower, L.P., Burrough, E.R., and Frana, T.S. 2014. Matrix-assisted laser desorption ionization time-of-flight mass spectrometry for rapid identification of *Brachyspira* species isolated from swine, including the newly described "*Brachyspira hampsonii*". J Vet Diagn Invest **26**(5): 635-639. doi: 10.1177/1040638714541114.

Whiting, R.A., Doyle, L.P., and Spray, R.S. 1921. Swine Dysentery. Bulletin(257): 3-15.

Zmudzki, J., Szczotka, A., Nowak, A., Strzelecka, H., Grzesiak, A., and Pejsak, Z. 2012. Antimicrobial susceptibility of *Brachyspira hyodysenteriae* isolated from 21 Polish farms. Pol J Vet Sci **15**(2): 259-265.

Figure 1.

Hemolytic zones associated with the growth of *Brachyspira hyodysenteriae* ATCC 27164 on CVS agar. Note the lack of colony formation on the surface of the media.

Draft

Table 1. Antimicrobials with label claims for the treatment/prevention/control of *Brachyspira* associated disease

Drug Name	United States	Dosage*	Canada	Dosage‡	European Union	Dosage‡
Tylosin	YES	Water - 250mg/gallon Feed - 40-100g/kg feed	YES	Feed - 44-110 mg/kg	NO	
Lincomycin	YES	Feed - 40-100g/kg	YES	Feed - 44-110 mg/kg	NO	
Tylvalosin	NO		NO		YES	Water - 5mg/kg Feed - 4.25mg/kg
Tiamulin	YES	Feed - 200g/ton	YES	Feed - 31.2-220.1 mg/kg	NO	
Valnemulin	NO		NO		YES	Feed - 1-4mg/kg
Virginiamycin	YES	Feed - 25-100g/ton	YES	Feed - 55-110 mg/kg	NO	
Carbadox	NO		NO		NO	
Bacitracin	YES	Feed - 250g/ton	NO		NO	
Gentamicin	YES	Water - 50mg/gallon	NO		NO	

*administered dosage is per mass or volume of feed or warer, ‡ administered dosage is per mass of pig receiving medication

Table 2. Chronologicakl summary of antimicrobial susceptibility test methodologies utilized in published investigations

Method	Country	Year	Organism	Incubation Temp	Incubation Time	Final Inoculum	Atmosphere	Media	References
Agar Dilution	Sweden	2003	Various	37°C	96 hours	1 - 5 x 10 ⁵ / spot	GENbox	TSA5OX WC5SB	(Karlsson et al. 2003)
	Japan	2004	BH	Unclear, 37-38°C	72 hours	10 ⁴ - 10 ⁵ / spot	BBD GasPak	TSA5SB	(Uezato et al. 2004)
	Czech Republic	2004	BH	37°C	72 - 96 hours	10 ⁴ / spot	Anaerobic jar using gas generating kit (BR38, Oxoid)	WC5SB	(Lobova et al. 2004)
	Germany	2004	BH	42°C	48 hours	3X 10 ⁵ / spot	No details reported	TSA10BB	(Rohde et al. 2004)
	Japan	2010	BH	37°C	72 hours	10 ⁵ / spot	No details reported	TSA5SB	(Ohya and Sueyoshi 2010)
	USA	2011	Various	42°C	48-96 hours	1.6 - 4 x 10 ⁴ / spot	GasPak	TSA5SB	(Clothier et al. 2011)
	Korea	2012	Various	37°C	72 - 120 hours	1.5 x 10 ⁶ / spot	No details reported	MHA5SB	(Lim et al. 2012)
	Czech Republic	2011	BH	37°C	72-96 hours	Undefined	AnaeroGen	WC5SB	(Sperling et al. 2011)
	Czech Republic	2014	BH	37°C	72-96 hours	10 ⁵ / spot	AnaeroGen	WC5SB	(Prasek et al. 2014)
	Japan	2015	BH	37°C	72 hours	10 ⁴ - 10 ⁵ / spot	No details reported	TSA5SB	(Kajiwara et al. 2015)
	USA	2016	Various	37°C	96 hours	1 x 10 ⁵ / spot	No details reported	TSA5SB0	(Mirajkar et al. 2016a)
Broth Dilution	Sweden	1999	BH	37°C	up to 96 hours	5 x 10 ⁵ - 10 ⁶ CFU/ml	No details reported	BHIS10	(Karlsson et al. 1999)
	Sweden	\$2001	BH	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	No details reported	BHIS10	(Karlsson et al. 2001)
	Australia	2002	BH	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Karlsson et al. 2002)
	Sweden	\$2003	Various	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Karlsson et al. 2003)
	Sweden	\$2004	BP	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Karlsson et al. 2004b)
	Germany	2004	BH	37°C	96 hours	3 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Rohde et al. 2004)
	European	\$2004	BH	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Karlsson et al. 2004a)
	Sweden	\$2006	BP	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	BBD GasPak	BHIS10	(Pringle et al. 2006)
	Spain	\$2009	BH	38°C	72-120 hours	1 - 5 x 10 ⁶ CFU/ml	GENbox	BHIS10	(Hidalgo et al. 2009)
	Spain	\$2011	BH	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	No details reported	BHIS10	(Hidalgo et al. 2011)
	Sweden	\$2012	BH, BP	37°C	96 hours	1 - 5 x 10 ⁶ CFU/ml	No details reported	BHIS10	(Pringle et al. 2012)
	Poland	\$2012	BH	37°C	96 hours	10 ⁶ CFU/ml	GENbox	BHIS10	(Zmudzki et al. 2012)
	Germany	2014	BH	37°C	120 hours	10 ⁵ GFU ₅₀ /ml*	GENbox	BHIS20	(Herbst et al. 2014)
	Italy	\$2015	BH			No details reported			(Rugna et al. 2015)
	USA	\$2016	Various	37°C	96 hours	10 ⁶ CFU/ml	No details reported	BHIS10	(Mirajkar et al. 2016a)

BH - *B. hyodysenteriae*; BP - *B. pilosicoli*
Media: Tryptic soy agar + 5% sheep blood: TSA 5 SB; Tryptic soy agar + 10% sheep blood: TSA 10 SB; Tryptic soy agar + 5% ox blood: TSA 5 OX; Tryptic soy agar + 10% bovine blood: TSA10BB; Wilkins-Charlgren + 5% sheep blood: WC 5SB; Mueller-Hinton agar + 5% sheep blood: MHA 5 SB; Brain heart infusion + 10% fetal calf serum: BHIS 10; Brain heart infusion + 20% fetal calf serum: BHIS 20;
Atmosphere: AnaeroGen Sachets Oxoid: AnaeroGen; Anaerobic gas pack GasPak BD: GasPak; anaerobic atmosphere generator biomerieux GENbox: GENbox; Anaerobic gas generator envelopes BBL gas pak: BBD GasPak
*GFU - authors report "growth forming units" as opposed to CFU\$Studies which reported the use of the VetMIC Brachy panel

Table 3. Genetic associations with decreased antimicrobial susceptibility in *Brachyspira*

Organisms Investigated	Phenotypic Resistance	Genetic Mechanism	References
<i>B. hyodysenteriae</i> <i>B. pilosicoli</i>	macrolides, lincosamides, streptogramins, tiamulin	Single nucleotide polymorphisms at positions 2032 and 2058 of the 23S ribosomal subunit	(Hidalgo et al. 2011; Karlsson et al. 1999; Karlsson et al. 2004b)
<i>B. hyodysenteriae</i>	tiamulin	Mutation in ribosomal protein L2, L3, L4, L22 and 23S rRNA in the peptidyl transferase region	(Hillen et al. 2014; Pringle et al. 2004)
<i>B. hyodysenteriae</i> <i>B. intermedia</i>	doxycycline	Single nucleotide polymorphism at position 1058 of the 16S ribosomal subunit	(Pringle et al. 2007; Verlinden et al. 2011)
<i>B. pilosicoli</i>	penicillin, ampicillin and oxacillin	β -lactamases (OXA-63, OXA-136, OXA-137 and OXA-470 - OXA-479)	(La et al. 2015; Meziane-Cherif et al. 2008)

Hemolytic zones associated with the growth of *Brachyspira hyodysenteriae* ATCC 27164 on CVS agar. Note the lack of colony formation on the surface of the media.

180x179mm (300 x 300 DPI)