Recorded captures of American lobster *Homarus americanus* in Swedish waters and an observation of predation on the European lobster, *Homarus gammarus*

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0532.R2
Manuscript Type:	Rapid Communication
Date Submitted by the Author:	29-May-2017
Complete List of Authors:	Øresland, Vidar; Swedish University of Agricultural Sciences, Institute of Marine Research, Department of Aquatic Resources, Ulmestrand, Mats; Swedish University of Agricultural Sciences, Department of Aquatic Resources Agnalt, Ann-Lisbeth; Institute of Marine Research Oxby, Gert; private company, Gert Oxby Lys Air & Dive AB
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	COASTAL WATERS < Environment/Habitat, LOBSTERS < Organisms, PREDATION < General, HYBRIDISATION < General

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

RAPID COMMUNICATIONS

Recorded captures of American lobster *Homarus americanus* in Swedish waters and an observation of predation on the European lobster, *Homarus gammarus*

Vidar Øresland and Mats Ulmestrand

Institute of Marine Research, Department of Aquatic Resources, , Swedish University of Agricultural Sciences, Turistgatan 5, SE-453 30 Lysekil, Sweden

Ann-Lisbeth Agnalt

Institute of Marine Research, Post Box 1870, Nordnes, Bergen 5817, Norway

Gert Oxby

Lys Air & Dive AB, Läkarvillan, Valbogatan 52, SE-453 34 Lysekil, Sweden.

Key words: *Homarus americanus*, *Homarus gammarus*, lobster, distribution, non-native species, Skagerrak, hybrid, predation

The intentional or unintentional release of American lobsters (*Homarus americanus*), imported through the European seafood trade, has resulted in findings of *H. americanus* in European waters. A

total of 36 specimens was recorded within a smaller area, half way along the Swedish west coast, between 2008 and 2016. The lobsters were caught by both recreational and professional fishermen and delivered to the Institute of Marine Research in Lysekil where species identification, measurements and genetic sampling were performed. Twenty-five of these lobsters were caught inside the sill of the Gullmar Fjord, most of them close to Lysekil harbor, where escapes from an illegal holding is suspected. Six females had eggs which in two cases were hybrids. While there is no evidence to suggest that a population has been established in the Gullmar Fjord, or elsewhere in Swedish waters, serious concerns must be raised regarding the possible consequences for the native species. An American lobster has for the first time been filmed eating a killed European lobster in Swedish waters.

Corresponding author: Vidar Øresland; vidar.oresland@slu.se

In recent years, several specimens of the American lobster *Homarus americanus* have been caught along the Swedish west coast. Since both the recreational and commercial lobster fisheries are very important in Sweden, the consequences the new species might have for the native species *H. gammarus* are of great concern. An information campaign along the Swedish west coast, initiated in 2008, on how to identify and report on findings of the American lobster resulted in an increased awareness among the public and lobster fishermen and hopefully also among fishmongers that keep live lobsters. This paper presents all reported captures in Swedish waters up to 2016 and in particular the unique concentration of findings of *H. americanus* inside the sill of the 120 m deep Gullmar Fjord. The presence of and concerns regarding American lobsters in European waters have earlier been discussed by e.g. Stebbing et al. (2012) and Jørstad et al. (2011).

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70

Methods

The lobsters were caught by both recreational and professional fishermen using lobster pots or Norway lobster pots and one lobster was caught by a trawler and one by a diver. The fishermen were given a reward for each reported and delivered lobster. The delivery was normally on the day of the catch. Specimens were frozen after first being photographed, measured, sexed, weighed and visually identified (all our data and calculations can be found in supplementary materials). All the lobsters received had typical morphological characteristics of the American lobster that often or always differ from the European lobster (e.g. sub-rostral spines, red spines on chelipeds, carapace with some yellowish spots, brownish uropods and telson). Primary visual identification was later confirmed genetically. Four specimens were not sent for genetic identification because they were kept by the fishermen. Muscle and egg samples for genetic analyses were preserved in 96% ethanol and sent to the Institute of Marine Research in Bergen, Norway. Details of the genetic method used are given in Jørstad et al. (2007). An underwater video (see supplementary materials) showing an American lobster eating a European lobster was taken during scuba diving (8 Oct. 2016 at 13.00 h) using a GoPro camera Hero 4 Black Edition. In our information campaign for fishermen we set up posters in, e.g., harbours and in shops selling lobster equipment and lobster bait from the Norwegian border to the city of Halmstad in the south and we distributed around 70 000 cards (with photograph of an American lobster, phone numbers and a reward for reporting catches) in shops and supermarkets within 15 km from the coastline along the Swedish west coast. We estimate that we covered least 95 % (280 Km) of the coastline where lobsters are fished. We also appeared numerous times in media to inform the public of our information campaign.

Results

Fig. 1 shows the position of each finding (2008-2016). The first lobster (1) was caught by a trawler at 160 m depth in the Skagerrak in May 2008. Later the same year, during the lobster fishing season, 3 more lobsters were reported (2-4) outside a fish auction in the village of Smögen. Another one was found in 2010 off the Bornö Islands far inside the Gullmar Fjord and 3 more specimens were found in the same area in 2014, 2015 and 2016 (the 2016 position mark hides the findings in 2010 and 2015 in Fig.1). After 2010 no lobsters were reported until 2014 when 23 lobsters were caught (18 just close to Lysekil harbour (Fig. 2)). The southernmost finding (24) was caught 47 km from Lysekil (Fig. 1). In 2015 and 2016 only 5 and 3 lobsters, respectively, were reported in total. Four of these lobsters (26-30) were caught offshore >10 km from Lysekil harbour. No. 31 was caught 2 km south of the Kåvra lobster reserve (see Fig.1). Pot depths were between 9 and 38 m. The carapace lengths were between 80 and 121 mm (mean 101 mm) and there were 58 % males and 42 % females (6 females with eggs of which 2 had hybrid eggs). The weights were between 520 and 1480 gram (mean 877 gram).

The male *H. americanus* (No 36, Fig. 2 and 3) eating a *H. gammarus* was video-filmed by Gert Oxby during scuba diving at a depth of 19 m off Släggö Island 100 m from Lysekil harbour (8 Oct. 2016 at 13.00 h, T = 15 °C). To our knowledge this feeding behaviour has never before been observed. The video shows that this was not an empty exoskeleton but a dead individual (confirmed by the diver). The video shows clearly that the carapace itself was cut in two at its lower end (not between the carapace and the abdomen, which can be seen on the video), the muscles were still present in the carapace and appeared fresh (also observed by the diver) indicating that the death was recent. The muscles in the front part of the abdomen were absent. During a dive the following day neither the remains of the dead lobster nor the predator could be found. Catching lobsters during scuba diving is

95 not allowed without special permission. However, the lobster had preliminarily been identified as an
96 American lobster from the video and it was therefore caught with permission by a diver approximately
97 40 m away from the video location 18 days later and brought to the laboratory. Its carapace was 115
98 mm and its weight 1400 g. From dive photographs and photographs taken in the laboratory, especially
99 of the crusher claw (the tip was broken and the “teeth” worn down), we could conclude that we had
100 caught the individual appearing in the video.

101

102

103 **Discussion**

104 The catches of American lobster in Scandinavian waters were reported in e.g. Jørstad et al. (2011).
105 However, since that time more lobsters have been caught, especially in Swedish waters. The lobster
106 catches in Norwegian waters were more spread out along the coastline compared to the Swedish
107 catches. Stebbing et al. (2012) reported catches from Scottish waters down to the English Channel. In
108 contrast, all the American lobsters in Swedish waters were caught within a smaller area, half way
109 along the Swedish west coast, and especially within the Gullmar Fjord. Although we regard our
110 information campaign as rather successful, some catches might not have been reported due to some
111 fishermen being unaware of the campaign, especially during its first years. However, both recreational
112 and professional fishermen have indeed been very positive to the campaign. For comparison with catch
113 locations of American lobster, the European lobster is distributed, and fished, along the entire Swedish
114 west coast, between the Norwegian border and the Sound (between Sweden and Denmark) in the
115 south. Lobsters are not found in the Baltic Sea where the salinity is low ($< 10 \text{ ‰}$).

116

The lobsters reported just outside the auction hall in Smögen in 2008 may have come from an escape event from an illegal holding in the sea in the same area (two specimens had rubber bands around the claws). The large numbers found in 2014 close to Lysekil harbour most probably came from illegal holdings inside the harbour area (three specimens had rubber bands). Fishing for lobster is not allowed in the harbour area. We will not speculate as to who is responsible for these events. We have no idea how many lobsters in total have escaped into the sea and thus do not know what proportion of them has been caught. There is a high fishing pressure on lobster in the Gullmar Fjord, where in 2014 V. Øresland recorded 1977 pots (unpublished report in 2017 to the Swedish Agency for Marine and Water Management). The number of reported American lobsters decreased during 2015 and 2016. This may be due partly to the previous year's fishing and to the fact that the lobsters are spreading offshore and along the coast, as indicated by the findings, where they get harder to catch. The offshore spread is not surprising since the American lobster is known to migrate long distances (see e.g. Campbell, A. 1986 and Maine DMR 2015). In contrast, the European lobster appears to be much more stationary (Øresland and Ulmestrand 2013) and is, at least in Swedish waters, seldom caught below 40 m. The lobster fishing in Swedish waters normally occurs at depths between 10-35 m. Therefore, American lobsters that migrate to deeper waters are less likely to be fished. In addition, our ongoing information campaign has hopefully made fishmongers that keep live lobsters more aware of the problems associated with illegal holdings in the sea and this be the reason why no new large escape events have been detected since 2014.

The weight data (520 - 1480 g) reported here (see supplementary materials) show that the specimens are above normal import size (generally between 500 and 530 g). That indicates that the larger reported *H. americanus* had been in the sea before the escape event in 2014 and that minor escape events have occurred before 2014. We did get one lobster in 2010 in the Gullmar Fjord. However, we cannot rule out the possibility that some of the lobsters that escaped in 2014 were above normal import size.

Lobsters that are above the normal import size are usually delivered as special orders and in very small quantities and are therefore less likely to be kept in illegal holding tanks in the sea (pers.com. Christer Wrisemo, ABC Food). Fishermen are not allowed to keep *H. gammarus* females carrying eggs (berried). Consequently, European females with hybrid eggs will not be reported by the fishermen. However, if a fisherman reports an *H. americanus* female with eggs while still at sea, permission to land and deliver the lobster to the laboratory is given directly. Of particular note is the finding of American lobster close to the Kåvra lobster reserve, which may be the oldest in Europe (established in 1989). Lobsters have been tagged for many years in this area (Øresland and Ulmestrand 2013) and now one can also study very large lobsters that occur in high concentrations and any interaction between the two species would therefore be of great concern.

Whether or not American lobsters will establish permanent sub populations in Swedish waters is still open to speculation. Larvae of *H. gammarus* in the Kåvra lobster reserve were found at a mean depth of around 16 m, below the sharp halocline. The larvae were retained locally due to a combination of larval depth and a persistent retention of water masses below the halocline (Øresland and Ulmestrand 2013). If such larval depth distribution below the halocline is true also for *H. americanus* in the Gullmar Fjord, the establishment of subpopulations would be facilitated. The number of findings within a short period, typical lobster habitats, a sill and deep water circulation below a strong halocline make the Gullmar Fjord a possible candidate for a permanent subpopulation. However, the considerable lobster fishing in the fjord and migration offshore are two important factors that would act against the establishment of a permanent population in the fjord. The findings of hybrid eggs are of great concern but we have up to now no data on bottom-living hybrid lobsters and we have no data on hybrid eggs among *H. gammarus* females (it is forbidden to catch berried females).

We are convinced that the *H. americanus* found eating a *H. gammarus* also killed its prey and that the *H. gammarus* did not die while moulting. This is based on the diver observations of remaining muscles inside the abdomen and that the carapace appeared fresh, that the lobster was cut in two through the carapace and that its exoskeleton looked normal and healthy. We believe the actual killing process is quick and therefore extremely difficult to capture on video during diving. *H. americanus* can grow much larger than *H. gammarus*, which would naturally be an advantage in a predatory situation as in this case. This observation should lead us to ask also how common cannibalism is (within both species) and especially within lobster reserves where large lobsters are numerous.

Acknowledgements

We thank Eva Farestveit for genetic analyses, Patrik Jonsson for making Fig 1-2, Lars Hernroth for comments on the manuscript and M. Eddison-Bergquist for linguistic assistance. The maps were obtained from ICES spatial facility <http://geo.ices.dk/>. We thank the four anonymous reviewers for much appreciated suggestions. This project was funded by the Swedish Agency for Marine and Water Management (HaV) under contract no. 1084-16 and the Brita och Sven Rahmns Stiftelse that supported our scuba diving for lobsters (Divers & Scientists West Coast Sweden).

References

- 188 Campbell, A. 1986. Migratory movements of Ovigerous lobsters, *Homarus americanus*,
189 tagged off Grand Manan, Eastern Canada. Can. J. Fish. Aquat. Sci. 43 2197-2205
- 190 ICES. Spatial facility <http://geo.ices.dk/>.
- 191 Jørstad. K.E., P.A. Prodohl, A.-L. Agnalt, M. Hughes, E. Farestveit and A.F. Ferguson, 2007.
192 Comparison of genetic and morphological methods to detect the presence of American
193 lobster, *Homarus americanus* H. Milne Edwards, 1837 (Astacidea: Nephropidae) in
194 Norwegian waters. Hydrobiologia 590:103-114.
- 195 Jørstad K.E., A.-L. Agnalt, E. Farestveit, 2011. The introduced American lobster, *Homarus*
196 *americanus* in Scandinavian waters. In: Galil BS, Clark PF, Carlton JT (eds), In the wrong
197 place: alien marine crustaceans – distribution, biology and impacts. Springer series in
198 Invasion Ecology 6: pp. 625-638. Springer-Verlag, Dordrecht.
- 199 Maine DMR 2015: <http://www.maine.gov/dmr/commercial-fishing/index.html> (assessed
200 1/23/17).
- 201 Øresland, V., and M. Ulmestrand. 2013. European lobster subpopulations from limited adult
202 movements and larval retention. ICES Journal of Marine Science, 70: 532–539.
- 203 Stebbing, P., P. Johnson, A. Delahunty, P. F. Clark, T. McCollin. C. Hale, S. Clark, 2012.
204 Reports of American lobsters, *Homarus americanus* (H. Milne Edwards, 1837), in British
205 waters. BioInvasions Records, Vol. 1, Issue 1: 17–23.

Figure 1. Locations where 36 American lobsters (*Homarus americanus*) were caught between 2008 and 2016 inside and outside the Gullmar Fjord on the Swedish West coast (ICES). K = Kåvra lobster reserve.

Figure 2. Locations of American lobsters (*Homarus americanus*) close to Lysekil harbour in the Gullmar Fjord (ICES).

219 **Figure 3.** An American lobster (*Homarus americanus*) eating a European lobster (*Homarus*
220 *gammarus*) in the Gullmar Fjord. Photo: Gert Oxby, Divers & Scientist West Coast Sweden.

221

222

223

Draft