Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

Evolution of phenology in a salmonid population: a potential adaptive response to climate change

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0028.R1
Manuscript Type:	Article
Date Submitted by the Author:	24-May-2017
Complete List of Authors:	Joyce, John; Auke Bay Fisheries Laboratory, National Marine Fisheries Service - NOAA Manhard, Christopher; California Cooperative Fish and Wildlife Research Unit, ; U.S. Fish and Wildlife Service, Arcata Field Office, Gharrett, Anthony; University of Alaska Fairbanks, School of Fisheries & Ocean Sciences
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	contemporary evolution, phenology, local adaptation, SALMON < Organisms, CLIMATE CHANGE < General

SCHOLARONE™
Manuscripts

Evolution of phenology in a salmonid population: a potential adaptive response to climate change

Christopher V. Manhard¹, John E. Joyce², and Anthony J. Gharrett¹

¹College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Juneau Center, 17101 Point Lena Loop Road, Juneau, Alaska 99801, USA

²National Marine Fisheries Service, Auke Bay Laboratories, Ted Stevens Marine Research Institute, 17109 Point Lena Loop Road, Juneau, Alaska 99801, USA

Email addresses: CVM – cvmanhard@alaska.edu, JEJ – john.joyce@noaa.gov, AJG – a.gharrett@alaska.edu

Keywords: contemporary evolution, climate change, phenology, local adaptation, sea-surface temperature, salmonids

Correspondence: Christopher V. Manhard, California Cooperative Fish and Wildlife Research Unit, Humboldt State University, 1 Harpst Street, Arcata, CA 95521, USA
Telephone: 361 563 1149 e-mail: cvmanhard@alaska.edu

20 **Abstract**

21 Accumulating evidence has indicated that many fish populations are responding to
22 climate change through shifts in migration time, but genetic data identifying the role of evolution
23 in these shifts are rare. One of the first demonstrations of evolution of migration time was
24 produced by monitoring allozyme alleles that were experimentally manipulated to genetically
25 mark late-migrating pink salmon. Here, we extend that research by using observations of the
26 marker alleles in fry to demonstrate that these changes in migration time were caused by
27 directional selection against the late-migrating phenotype during the oceanic phase of the
28 salmonid life cycle. The selective event, which appeared to be driven by early vernal warming of
29 the nearshore marine environment and consequent decreased survival of late-migrating fry
30 relative to early-migrating fry, decreased the late-migrating phenotype from more than 50% to
31 approximately 10% of the total fry abundance in only one generation. These demographic
32 changes have persisted over the subsequent thirteen generations and suggest that a larger trend
33 toward earlier migration time in this population may reflect adaptation to warming sea-surface
34 temperatures.

35 Introduction

36 A primary mechanism by which animal populations can respond to environmental change
37 is by shifting the timing of life history events, known as phenology, to track optimal
38 environmental conditions in time (Bradshaw and Holzapfel 2006; Bradshaw and Holzapfel 2008;
39 Bellard et al. 2012). Phenology is of critical importance for migratory fish, such as Pacific
40 salmon (*Oncorhynchus* spp.), which must initiate an array of physiological, morphological, and
41 behavioral changes at precise times during their life cycle (reviewed by Groot and Margolis
42 1991). The precision with which salmonids time migration can constrain gene flow between
43 seasonally distinct spawning segments, thereby enabling local adaptation of phenology (Quinn et
44 al. 2000; Fillatre et al. 2003; Gharrett et al. 2013). Evidence of local adaptation of phenology in
45 salmonids has been provided by comparisons of seasonally distinct groups of Chinook salmon
46 (*O. tshawytscha*) that spawn in the same river, which demonstrated significant genetic
47 divergence at three circadian clock genes, but not at neutral markers (O'Malley et al. 2013). The
48 tight coupling of salmonid ecology and phenology, along with the generally high heritability of
49 phenological traits in salmonids (median $h^2 = 0.51$; Carlson and Seamons 2008), suggests that
50 salmonid populations may respond to periodic environmental fluctuations or persistent climatic
51 trends through contemporary evolution of phenology.

52 An increasing body of literature has indicated that many salmonid populations are
53 responding to climate change through changes in phenology. Earlier migration times have now
54 been documented in populations of pink (*O. gorbuscha*), coho (*O. kisutch*), and chum (*O. keta*)
55 salmon in Southeast Alaska (Kovach et al. 2015), sockeye salmon (*O. nerka*) in the Fraser River
56 (Cooke et al. 2004), steelhead trout (*O. mykiss*) in the Columbia River (Robards and Quinn
57 2002), and Atlantic salmon (*Salmo salar*) in Northern Ireland (Kennedy and Crozier 2010).

There is reason to expect that observed changes in migration time may be adaptive in many salmonid populations. Vernal warming is a primary determinant of the time at which the nearshore marine habitat can support growth of juvenile salmon (Friedland et al. 2000; Mortensen et al. 2000), and the ability of juveniles to time their emigrations with the onset of favorable growth conditions is an important component of fitness (Beamish and Mahnken 2001). Therefore, under a scenario in which earlier vernal warming is occurring in the nearshore marine habitat, earlier juvenile emigration times would presumably be adaptive. Because adult migration time is highly heritable in salmonids (Carlson and Seamons 2008), selection that favors earlier juvenile emigration times would be expected to result in earlier adult migration times, and the presence of earlier migration times in both life stages may therefore be a joint response that is being driven by the juvenile stage. However, in populations where adults enter freshwater after stream temperatures peak in late summer, a shift toward earlier adult migration times may be maladaptive, particularly if stream temperatures are warming beyond thermal optima. Such life history tradeoffs are probably integral in determining the scope by which salmonid populations can adaptively respond to climate change.

It remains unclear whether the widespread changes in salmonid phenology time generally reflect phenotypic plasticity, which is defined by environmentally-induced phenotypic variation of a given genotype, or genetic adaptation, which is defined by differential fitness of distinct genotypes in a given environment (Gienapp et al. 2008). Genetic monitoring, a technique in which putatively neutral molecular markers that are associated with phenotypic traits are tracked through time to identify adaptive genetic changes in relation to a changing environment, provides a means of distinguishing between plastic and adaptive responses. Hansen et al. (2012) argued that convincingly demonstrating adaptive genetic change based on molecular markers requires

that (1) the monitored traits exhibit sufficiently high heritability; (2) the traits are relevant to specific selective agents; (3) the traits are monitored over time; (4) selection is tested; (5) shifts in traits are consistent with the expected response to environmental changes; and (6) replacement by a genetically different population has been ruled out.

Migration time of pink salmon that spawn in Auke Creek, Alaska is a suitable trait for evaluating adaptive change through genetic monitoring because of its high heritability and because it is likely to respond to changes in water temperature. Genetically segregated even- and odd-year populations of pink salmon (Heard 1991) exist at Auke Creek, and adults within each population return to Auke Creek in two distinct segments; ‘early-migrating’ adults return between mid- and late August, and ‘late-migrating’ adults return between early and mid-September (Taylor 1980). These segments can be thought of as two subpopulations that are partially genetically isolated, with stream temperature modulating the amount of gene flow between subpopulations each year through its influence on stream entry time. The structuring of the spawning migration is maintained in the presence of gene flow by the high heritability of migration time within each of these subpopulations ($h^2 = 0.4$) and by potential dispersive selection against intermediate migration time (Smoker et al. 1998). Migration time is involved in synchronizing the timing of life history events with optimal environmental conditions and may therefore be subjected to selective pressures as water temperatures warm. The importance of water temperature as an adaptive driver is supported by the genetically distinct embryonic development patterns exhibited in the early and late subpopulations (Hebert et al. 1998).

During the past 40 years, trends toward earlier migrations of adults and fry have accompanied trends toward warmer water temperatures in the Auke Creek system. Because water temperature is an important component of phenology in this population, these changes in

104 migration timing may be an adaptive response to climate warming (Taylor 2008). The basis of
105 these changes was evaluated by monitoring an isolocus pair that was experimentally manipulated
106 through artificial selection to genetically mark the late-migrating subpopulation (Lane et al.
107 1991). This locus pair presumably has little genetic influence on migration time. However,
108 because of its high level of correlation with the late-migrating phenotype, changes in its
109 frequencies can be used to infer genetic changes in migration time. Genetic monitoring of this
110 late-migrating marker locus (LMML) demonstrated that a selective event, first evident in
111 broodyear 1989, had reduced the abundance of the late-migrating subpopulation and that these
112 demographic changes have persisted over the subsequent thirteen generations. These results
113 provided compelling evidence that the shifting phenology of Auke Creek pink salmon has a
114 genetic basis (Kovach et al. 2012).

115 Here we investigate the genetic basis of change in phenology in Auke Creek pink salmon
116 and its potential link to climate change by synthesizing previously detailed observations of
117 genetically marked adults with newly detailed observations of emigrating fry. The primary
118 questions that these analyses addressed were (1) Have fry exhibited changes at the LMML that
119 parallel those observed in adults? (2) Is there evidence that natural selection underlies the genetic
120 changes? (3) Is low fitness of late-migrating fish from broodyear 1989 the result of poor
121 reproductive success or low marine survival? and (4) Is there evidence that the selection event is
122 linked to environmental change?

127 **Methods**

128 *Field and laboratory methods*

129 *Study site* – Auke Creek, a 323-m outlet of Auke Lake that drains into Auke Bay, Alaska,
130 is a migratory corridor and spawning habitat of pink salmon which has produced returns of
131 adults that ranged from 1,500 to 28,000 fish over the past 40 years. The U.S. National Marine
132 Fisheries Service operates Auke Creek Research Station, a permanent fish-counting weir and
133 experimental salmon hatchery located at the head of tidewater, which enables complete censuses
134 of spawning adults and emigrating fry.

135 *Genetic marking of late-migration time* – A selective mating experiment was conducted
136 in 1979 with the goal of incorporating a putatively neutral genetic marker into the late-migrating
137 subpopulation of the odd-year population in Auke Creek (Lane et al. 1990). During the spawning
138 migration of 1979, skeletal muscle samples were collected from 3,906 pink salmon that entered
139 Auke Creek after 15 September and, therefore, represented the latest migrants that year. Each
140 sample was subsequently genotyped by electrophoretically screening the malate dehydrogenase
141 (*MDH-B1,2**) is locus pair (Allendorf and Thorgaard 1984). Adults ($n = 407$) which had an
142 *MDH-B1,2*70* allele but not an *MDH-B1,2*130* allele were artificially spawned, and the
143 embryos were incubated in the Auke Creek Hatchery. Genetically marked fry ($n = 178,219$)
144 were released into Auke Creek between 7 April and 7 May in 1980 to coincide with the peak
145 emigration of wild fry. Interbreeding between experimentally-marked adults and wild adults in
146 the 1981 spawning season produced changes in the allele frequencies of *MDH-B1,2** in the late-
147 migrating subpopulation. Specifically, the frequency of the *70** allele increased from 0.027 in
148 1979 to 0.134 in 1983, and the frequency of the *130** allele decreased from 0.022 in 1979 to
149 0.010 in 1983. Monitoring of subsequent spawning migrations from 1983 to 1989 revealed that

the allele frequencies at *MDH-B1,2** were stable and differed between samples of early- and late-migrating fish, thereby demonstrating that the experimental manipulations effectively marked genetic material that is correlated with migration time (Gharrett et al. 2013).

Census and genetic data – During each spring from 1984-1992, daily counts were made of all pink salmon fry that passed the Auke Creek weir during their downstream migration. Daily samples of fry (typically 10-20 fish) were collected for genotype determination. Genotyping was accomplished by using starch gel protein electrophoresis to resolve allozyme banding patterns at the LMML (*MDH-B1,2**).

Auxiliary environmental data – The Auke Bay Laboratories Climatological Series is a data set consisting of environmental measurements recorded at the Auke Bay Marine Station since February 1963. These data include measurements of sea-surface temperatures of Auke Bay. Interannual variation in sea-surface temperature is a primary determinant of marine survival of fry (Manhard et al. 2017). Hence, these environmental data may provide evidence of an environmental basis changes in migration time.

Statistical methods

Graphical and statistical comparisons of temporal patterns - Because the *MDH-B1,2*130* allele underwent relatively small manipulative changes and exhibited allele frequencies close to zero, it provided little power for detecting genetic changes. Consequently, the statistical methods that we used to characterize the temporal patterns of the LMML included only the *MDH-B1,2*70* allele, which we refer to as the late-migrating marker allele (LMMA). Temporal patterns of the LMMA frequencies were monitored during five odd-year fry emigration periods (1984-1992). Graphical comparisons of five-day running averages of frequencies of the LMMA were made among brood years to describe interannual changes in

genetic differences between subpopulations. A two-sample test of equality of proportions was used to determine whether these subpopulations differed at the LMMA. Only data from the calendar dates of the first and last 100 fish sampled during each migration period were included in the test.

Overall late-marker allele frequency – Interannual patterns in the overall (population-wide) frequency of the LMMA in the odd-year population can provide insight into whether allele frequency changes are most likely caused by increased gene flow between subpopulations or reduced fitness of late-migrating fish. Gene flow between subpopulations within the stream should have caused the frequency of the LMMA to decline within the late-migrating subpopulation, but remain stable throughout the population. Conversely, reduced fitness of late-migrating fish should have caused a reduction in the population-wide frequency of the LMMA. Estimating the overall frequency of the LMMA and its statistical certainty is complicated by the variation of daily return numbers and unequal sample representation throughout the migratory periods. These issues were addressed with a parametric bootstrap algorithm (Kovach et al. 2012) that resampled alleles at the LMML (1,000 iterations) in running five-day pools of genetic samples collected during each fry migration. The algorithm divided the migration distribution each year into n five day periods, beginning on the first date in which genetic samples were collected. Within each i^{th} period, the maximum likelihood frequency of the LMMA (f_i) was estimated from the genetic samples that were collected during that period. Each iteration of the algorithm began with a random draw of alleles (x_i) from a binomial distribution,

$$x_i \sim \text{Bin}(f_i, a_i), \quad (1)$$

where a_i was the number of alleles sampled within a period (2 * the number of genetically sampled fish). In each year, the overall frequency of the LMMA (f) was estimated as,

$$f = \frac{\sum_{i=1}^n \frac{x_i}{a_i} 2N_i}{\sum_{i=1}^n 2N_i}, \quad (2)$$

where N_i was the census number of fry observed during a period. The bootstrap simulation provided a point estimate and 95% confidence interval for the overall LMMA frequency each year.

Stock separation algorithm – A stock separation algorithm (Pella and Milner 1987), which used allele frequencies of the LMML in collections of early- and late-migrating adults from 1983 as a baseline, estimated the number of migrant adults and fry that belonged to each subpopulation. The baseline consisted of 564 adults collected on 10 and 18 August and 161 adults collected on 21 September, which represented the unmarked early-migrating subpopulation and the marked late-migrating subpopulation, respectively. Statistical replication of the stock separation algorithm was accomplished with a non-parametric bootstrap simulation that resampled alleles (20,000 iterations) in both the baseline and in running five-day pools of fish. On each iteration of the bootstrap simulation an expectation-maximizing algorithm estimated the proportion of fish that belonged to each subpopulation by comparing running five-day averages of allele frequencies of genetic samples to allele frequencies of the baseline. The estimated proportion of fish from each subpopulation was then multiplied by the number of fish observed on that date to produce daily estimates of the number of early- and late-migrating fish. This process was repeated on each day of the migration period. Daily estimates were used to plot the annual migration distributions of each subpopulation. The bootstrap simulation provided point estimates and 95% confidence intervals for the annual abundance of adults and fry from

each subpopulation, as well as two metrics of fitness: the number of fry produced per adult (freshwater productivity) and the number of returned adults per emigrant fry (marine survival). Comparisons of freshwater productivity and marine survival between subpopulations and among brood years provided insight into the role of natural selection in promoting genetic changes in Auke Creek.

Environmental conditions – Graphical depictions of sea-surface temperatures experienced by fry shortly after entering the ocean were used to qualitatively explore potential environmental drivers of natural selection. The incipient marine residence period is critical to the survival of juvenile salmon (Beamish and Mahnken 2001; Farley et al. 2007, Manhard et al. 2017). Observations of tagged fry in Auke Bay have demonstrated that the duration of this period varies seasonally and interannually (Mortensen et al. 2000). Given our uncertainty of the duration of this critical period, we focused on environmental conditions experienced by fry during the first few days in Auke Bay, when they are most susceptible to mortality from predation and starvation. Demographic plots of the number of early- and late-migrating fry residing in Auke Bay on each date were made with estimates from the stock separation algorithm and included only fish that had emigrated within the past seven days. Thermal-gradient plots of sea-surface temperatures, which were superimposed over the demographic plots, depicted conditions encountered by fry during their first week in the marine environment.

Results

Monitoring of the late-marker allele – The temporal characteristics of the LMMA in adults were described previously (Kovach et al. 2012), and we briefly present those results here so that comparisons can be made between adults and fry. In each generation from 1985-1989, the

LMMA frequency increased substantially after the mid-point of the migration (Figure 1), and there was a significant difference ($P < 0.001$) in allele frequencies between samples of early- and late-migrating fish. Beginning in the adult migration of 1991, there was a marked decline in the magnitude of the LMMA frequency increase over the migration period and the LMMA frequencies of samples of early- and late-migrating adults were nearly identical ($P = 0.820$). Fry exhibited patterns in the LMMA that were similar to those observed in adults. During each generation from 1984-1990, there was a rapid increase in the LMMA frequency near the mid-point of the migration (Figure 2). In each of these generations, the LMMA frequency was approximately 0.08 - 0.10 higher in fry sampled from the late-migrating segment than in the early-migrating segment, which was a significant difference ($P < 0.001$). A substantial decline in the LMMA frequency was first evident in migrating fry in spring of 1992. The LMMA frequency was only 0.02 higher in fry sampled from the late-migrating segment that year, which was not a significant difference ($P = 0.456$).

The bootstrap analysis demonstrated that the overall frequency of the LMMA declined substantially in adults that returned to Auke Creek in 1991 after being relatively stable during the previous three generations (Figure 3). Similarly, the overall frequency of the LMMA declined substantially in fry that emigrated in 1992 relative to the previous four generations, in which the overall frequency was relatively stable. These patterns indicated that the LMMA frequency was reduced by a selective event following the fry emigration of 1990.

Patterns in population demography – The stock separation algorithm produced estimates of early- and late-migrating population components in fry. During the first four generations (1984-1990), late-migrating fry accounted for an estimated 51-68% of the total fry abundance. However, beginning in the migration of 1992, the proportion of late-migrating fry declined to

only 12%. Bootstrap replication estimated marine survival and freshwater productivity in the early- and late-migrating subpopulations from 1983-1989 (Figure 4). In each brood year, marine survival of late-migrating fry was significantly lower than in early-migrating fry. This discrepancy was particularly pronounced in fry that emigrated in 1990, in which the survival rate of late migrants was only 1.4 percent. This survival rate was the lowest observed over the monitoring period and was less than one tenth of the survival rate of early migrants from the same brood year. By contrast, freshwater productivity of late-migrating adults was significantly higher than that of early-migrating adults in each generation from 1985-1989 (Figure 4). The migration of 1991 was distinct in that the mean estimate of freshwater productivity was similar between early- and late-migrating adults. However, aspects of the sampling design produced a high standard error for the estimated freshwater productivity of late migrants that brood year.

Environmental conditions – The estimated fry migration distributions indicated that early-migrating fry entered Auke Bay between 7 and 15 days earlier than late-migrating fry and, consequently, encountered cooler sea-surface temperatures each year, although the extent of this thermal discrepancy varied from year to year (Figure 5). The migrations of 1986 and 1990 are particularly noteworthy because of the contrasting temperatures experienced by early- and late-migrating fry in those years. In 1986, sea-surface temperatures remained relatively cool throughout April, and early-migrating fry experienced an average temperature of 5.4 °C during their first week in Auke Bay, whereas late-migrating fry began to enter Auke Bay in peak numbers during a warming event in the beginning of May and experienced an average temperature of 8.8 °C. In 1990, an early warming period that began on 15 April and coincided with the peak emigration of early-migrating fry provided an average temperature of 7.0 °C, which was only slightly cooler than that experienced by late-migrating fry (7.7 °C) during their

first week of marine residency. The migrations of 1986 and 1990 respectively marked the largest and smallest thermal discrepancies encountered between runs over the monitoring period.

Discussion

During the first four years of genetic monitoring, the patterns of the LMMA in fry paralleled those observed in adults in a previous study (Kovach et al. 2012). Both life history stages exhibited a pronounced increase in the LMMA frequency beginning near the midpoint of the migration, and fish sampled from the early- and late-migrating segments displayed significantly different allele frequencies, thereby demonstrating considerable genetic structure (Figures 1, 2). The presence of parallel seasonal patterns in adults and fry demonstrates that the phenology of these two life history stages is tightly coupled and indicates that natural selection upon one stage would likely produce an evolutionary response in the other. While the fingerprint of the LMMA was strong in fry that emigrated in spring of 1990 (broodyear 1989), the adult migration of that generation in 1991 was the first instance in which a substantial mid-season increase in the LMMA frequency and significant allelic differences between samples of early- and late-migrating fish were not observed. Further, similar changes in the characteristics of the LMMA were evident in the fry emigration of the following spring (1992), wherein samples of early- and late-migrating fry did not exhibit significant differences in the LMMA frequency. Hence, it appears that this population experienced genetic changes in phenology at two different life history stages, which were first evident following the oceanic phase of broodyear 1989. Subsequent observations of this population in 1993, 2001, and 2011, which also failed to demonstrate a fingerprint of the LMMA in late-migrating fish, suggest that these changes have persisted over multiple generations (Kovach et al. 2012).

Analyses of genetic data at the LMMA were used to investigate causative agents of genetic change, which may include gene flow between subpopulations, gene flow from nearby populations, and genetic drift. We evaluated the plausibility gene flow between subpopulations as a causal agent by iteratively simulating the overall frequency of the LMMA and observed that the overall frequency was stable in fry during the first four years of genetic monitoring, before undergoing a substantial, population-wide decline in 1992. This pattern, when considered alongside the similar pattern observed in adults (Figure 3), demonstrates that gene flow between subpopulations was unlikely to have caused the changes. Gene flow among populations was quantified by analyzing LMMA frequencies from nearby pink salmon populations in an island-continent model. The model estimated that the migration rate (m) necessary to produce the observed genetic changes ranged from 0.69-0.85 (Kovach et al. 2012), which was substantially higher than estimates of straying rates of Auke Creek pink salmon ($m = 0.02$ -0.04; Mortensen et al. 2002) and gene flow between proximate populations and Auke Creek ($m = 0.0015$; Gharrett et al. 2001), thereby demonstrating that interpopulation gene flow was inadequate to produce the observed changes. Genetic drift was addressed by quantifying expected changes from genetic drift with estimates of the effective population size (N_e). Analyses of genotype data over 23 microsatellite loci estimated values of N_e in the odd-year population that ranged between 788 and 6005 adults (Kovach et al. 2013a), which indicated that we could expect a change of 0.003-0.007 in the LMMA frequency from genetic drift. The observed change between the adult migrations of 1989 and 1991 was substantially higher than this range ($\Delta\text{LMMA} = 0.020$). Observations of another putatively neutral control locus, which exhibited minimal change during the monitoring period, provided additional evidence that genetic drift was not strong enough to cause the observed genetic changes (Kovach et al. 2012). Collectively, these results demonstrate that

genetic change from gene flow and genetic drift was likely minimal, leaving natural selection as the most likely causative agent.

Iterative simulations of the early- and late-migrating subpopulations with a stock separation algorithm provided evidence that the primary cause of genetic change was a natural selection event in broodyear 1989, which resulted from relatively low marine survival of late-migrating fish that emigrated in 1990. The relative fitness of these two subpopulations appears to be balanced by life history tradeoffs; late-migrating adults generally produced more fry per adult, but late-migrating fry consistently exhibited lower marine survival rates during the monitoring period (Figure 4). Higher freshwater productivity of late-migrating adults can be attributed to mortality of early-spawned embryos from redd superimposition by late-migrating adults (Fukushima et al. 1998), whereas lower marine survival of late-migrating fry can be attributed to their comparatively smaller body size and, by extension, higher predator vulnerability during the incipient marine residence period (Mortensen et al. 2000). Although these tradeoffs appear to balance fitness in most years, broodyear 1989 was unique in that the marine survival rate of late-migrating fry was less than one-tenth that of early-migrating fry; in comparison, the marine survival rate of late-migrating fry ranged between one-fourth and one-half that of early-migrating fry in the other years. Since there was little evidence of a compensating increase in freshwater productivity of late-migrating adults from broodyear 1989, this marine survival disparity was likely sufficient to convey a lifetime fitness advantage to the early-migrating subpopulation. Further demographic declines were evident in the late-migrating subpopulation in the following brood year (1991), in which late-migrating adults apparently did not have their characteristic freshwater productivity advantage. This decline in relative freshwater fitness may be attributed to

increased overlap of spawning activity between early- and late-migrating adults that year and resulting competition for limited spawning habitat.

The incipient marine residence period is the primary determinant of lifetime fitness of Auke Creek pink salmon (Manhard et al. 2017) and, consequently, this stage has the potential to drive natural selection. Because their later ocean entry time confers a shorter growth period prior to seasonal increases in nearshore predator abundance, late-migrating fry are smaller, and therefore more vulnerable to size-selective predation (Parker 1971; Hargreaves and LeBrasseur 1986), than early-migrating fry during this stage (Mortensen et al. 2000). Additionally, late migrants may be at a disadvantage in competing with larger early migrants for food. The extent of this size disparity is governed by physical conditions that modulate the potential for growth in the nearshore environment. In Auke Bay, temperature is the primary factor controlling growth of fry because of its effects on their physiology and because it drives the abundance and size composition of their planktonic prey (Bienfang and Ziemman 1995). Although their consistently higher survival indicates that early-migrating fry benefit from being on the leading edge of the migration, low temperatures and sparse prey abundance in Auke Bay limit the growth potential of early migrants in many years (Mortensen et al. 2000). Hence, the balancing of the benefit of leading the migration with the cost of poor growth conditions largely determines the relative marine fitness of these subpopulations and is likely an outcome of seasonal temperature patterns. Comparisons of vernal sea-surface temperatures during the fry emigrations of 1986 and 1990 supported this (Figure 5). In 1986, sea-surface temperatures remained relatively cool as early-migrating fry were entering the ocean, and substantial vernal warming did not occur until late-migrating fry began emigrating in large numbers. In contrast, an early vernal warming event in 1990 likely provided favorable growth conditions for early-migrating fry. Consistent with these

temperature patterns, early-migrating fry exhibited their smallest marine survival advantage in 1986 and their largest advantage in 1990.

It is generally recognized that there is an optimal window for ocean entry in many salmonid populations (Cury and Roy 1989), and that the ability of fry to synchronize their emigration with this window is a critical component of fitness. Our results provide evidence of a mechanism by which adaptation of earlier migration time can occur under an advancing vernal warming period and associated optimal window. Furthermore, although we do not have information on trends in zooplankton phenology in Auke Bay, trends toward earlier onsets of spring zooplankton blooms have been observed in other systems (Edwards and Richardson 2004; Thackeray et al. 2010), and an earlier plankton bloom would reinforce a shift in the optimal window. Because migration times of fry and adults are connected through the heritability of this trait, selection against late-migrating fry would be expected to produce earlier average migration times in adults. Therefore, our hypothesis provides a unifying explanation for the coinciding trends toward earlier migrations of adults and fry (Kovach et al. 2013b) and warmer vernal sea-surface temperatures of Auke Bay (Figure 6) that have been observed in this system over the past four decades. An advance in the optimal growth window, without a corresponding plastic response, would be expected to cause directional selection against late-migration time, thereby reducing phenotypic variation in this trait. Hence, this hypothesis is also consistent with the declines in phenotypic variation of migration time that have been observed in adults and fry in the even- and odd-year populations at Auke Creek (Kovach et al. 2013b). Importantly, our study is limited to an observation of natural selection in a single generation, whereas the long-term trend in median migration time spans 17 generations at Auke Creek. The rapid genetic change observed in this population suggests that, in some cases, phenotypic change from climate-

induced selection may be extreme and episodic, rather than gradual. However, because the long-term census data reveal a nearly continuous shift in median migration time (rather than rapid truncation in one generation), plasticity must also be a substantial component of the pattern.

Long-term abundance data collected from populations of pink, coho, and chum salmon in rapidly warming Southeast Alaska demonstrated that 16 out of 19 observed populations are migrating into freshwater earlier than they historically did (Kovach et al. 2015). Stream temperature is an important factor governing freshwater entry in salmonids, which require sufficiently cool water temperatures for developing embryos (generally $< 14^{\circ}\text{C}$; Velsen 1987). The median migration times of these three species generally occur after stream temperatures have peaked, suggesting that selection or plasticity should favor later stream entry times that enable adults to avoid increasingly warm temperatures. However, the regional temporal pattern of earlier phenology in these species contrasts such a response of adults to warming stream temperatures, suggesting that these phenological changes may instead reflect an adaptive response of fry to an earlier optimal window for oceanic entry. Moreover, that study observed that the largest changes in median migration time were correlated with decreases in the duration of the migration, indicative of a loss of phenotypic variation resulting from natural selection against late-migrating fish. Collectively, these regional patterns of phenological and environmental change are consistent with an adaptive response of multiple salmonid species to warming oceanic conditions, as we have hypothesized for Auke Creek pink salmon.

We have presented an example of contemporary evolution of phenology in a salmonid population that meets all the criteria necessary to demonstrate adaptive genetic change (Hansen et al. 2012). Despite apparently strong natural selection against late-migrating fish and resulting demographic changes, the odd-year population has exhibited sustained levels of adult

420 recruitment over the past seventeen generations. Similarly, the even-year population, which has
421 exhibited parallel changes in phenology, has also sustained its recruitment levels over the same
422 period (Kovach et al. 2013b). This indicates that genetically based variation in phenology has
423 supported sustained productivity in Auke Creek during rapid climate change, an observation that
424 underscores the importance of life history variation to the resilience of fish populations (Hilborn
425 et al. 2003; Greene et al. 2010). In populations where this adaptive variation does not exist, the
426 limited scope of phenotypic plasticity may provide the only means of shifting phenology in
427 response to climate change. Collectively, our results emphasize the importance of tailoring
428 fisheries management programs to recognize and conserve seasonal genetic variation in the face
429 of inexorable, yet unpredictable climate change.

Acknowledgements

This manuscript represents a portion of Christopher Manhard’s doctoral work at the University of Alaska Fairbanks. The research was sponsored by Alaska Sea Grant with funds from the National Oceanic and Atmospheric Administration Office of Sea Grant, Department of Commerce, under grant no. NA060AR4170013 project no. R/31-13, and from the University of Alaska with funds appropriated by the state. We thank Ryan Kovach, USGS, for providing a review of this manuscript.

Draft

Cited literature

- Allendorf, F.W., and Thoraard, G.H. 1984. Tetraploidy and the evolution of salmonid fishes. *In* Evolutionary genetics of fishes. *Edited by* B.J. Turner. Plenum Publishing, New York, USA. pp. 1–53.
- Beamish, R.J., and Mahnken, C. 2001. A critical size and period hypothesis to explain natural regulation of salmon abundance and the linkage to climate and climate change. *Prog. Oceanogr.* **49**: 423–437.
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., and Courchamp, F. 2012. Impacts of climate change on the future of biodiversity. *Ecol. Lett.* **15**: 365–377.
- Bienfang, P.K., and Ziemann, D.A. 1995. APPRISE: A multi-year investigation of environmental variation and its effects on larval recruitment. *In* Climate Change and Northern Fish Populations. *Edited by* R.J. Beamish. Canadian Special Publications of Fisheries and Aquatic Sciences, Ottawa, Canada. pp. 483–487.
- Bradshaw, W.E., and Holzapfel, C.M. 2006. Evolutionary Response to Rapid Climate Change. *Science* **312**: 1477.
- Bradshaw, W.E., and Holzapfel, C.M. 2008. Genetic response to rapid climate change: it's seasonal timing that matters. *Molecular Ecology* **17**: 157-166.
- Carlson, S.M., and Seamons, T.R. 2008. A review of quantitative genetic components of fitness in salmonids: implications for adaptation to future change. *Evol. Appl.* **1**: 222–238.
- Cooke, S.J., Hinch, S.G., Farrell, A.P., Lapointe, M.F., Jones, S.R.M., Macdonald, J.S., Patterson, D.A., Healey, M.C., and G. Van Der Kraak, 2004. Abnormal migration timing and high en route mortality of sockeye salmon in the Fraser River, British Columbia. *Fisheries* **29**: 22–33.

- 462 Cury, P., and Roy, C. 1989. Optimal environmental window and pelagic fish recruitment success
463 in upwelling areas. *Can. J. Fish. Aquat. Sci.* **46**: 670–680.
- 464 Edwards, M., and Richardson, A.J. 2004. Impact of climate change on marine pelagic
465 phenology. *Nature* **430**: 881–884.
- 466 Farley, E.V., Murphy, J.M., Adkison, M.D., Eisner, L.B., Helle, J.H., Moss, J.H., and Nielsen, J.
467 2007. Early marine growth in relation to marine-stage survival rates for Alaskan sockeye salmon
468 (*Oncorhynchus nerka*). *Fish. Bull.* **105**(1): 121–130.
- 469 Fillatre, E.K., Etherton, P., and Heath, D.D. 2003. Bimodal migrating distribution in a northern
470 population of sockeye salmon (*Oncorhynchus nerka*): life history and genetic analysis on a
471 temporal scale. *Mol. Ecol.* **12**: 1793–1805.
- 472 Fukushima, M., Quinn, T.J., and Smoker, W.W. 1998. Estimation of egg loss from the spawning
473 ground of pink salmon. *Can. J. Fish. Aquat. Sci.* **55**: 618–625.
- 474 Friedland, K.D., Hansen, L.P., Dunkley, D.A., and MacLean, J.C. 2000. Linkage between ocean
475 climate, post-smolt growth, and survival of Atlantic salmon (*Salmo salar* L.) in the North Sea
476 area. *ICES J. Mar. Sci.* **57**(2): 419–429.
- 477 Gharrett, A.J., Lane, S., McGregor, A.J., and Taylor, S.G. 2001. Use of a genetic marker to
478 examine genetic interaction among subpopulations of pink salmon (*Oncorhynchus gorbuscha*).
479 *Genetica* **111**: 259–267.
- 480 Gharrett, A.J., Joyce, J.E., and Smoker, W.W. 2013. Fine-scale temporal adaptation within a
481 salmonid population: mechanism and consequences. *Mol. Ecol.* **22**: 4457–4469.
- 482 Gienapp, P., Teplitsky, C., Alho, J.S., Mills, J.A., and Merila, J. 2008. Climate change and
483 evolution: disentangling environmental and genetic responses. *Mol. Ecol.* **17**: 167–178.

- 484 Greene, C.M., Hall, J.E., Guibault, K.R., and Quinn, T.P. 2010. Improved viability of
485 populations with diverse life-history portfolios. *Biol. Letters* **6**: 382–386.
- 486 Groot, C., and Margolis, L. 1991. Pacific salmon life histories. University of British Columbia
487 Press, Vancouver, Canada.
- 488 Hansen, M.M., Olivier, I., Waller, D.M., Nielsen, E.E., and The GeM Working Group. 2012.
489 Monitoring adaptive genetic responses to environmental change. *Mol. Ecol.* **21**: 1311–1329.
- 490 Hargreaves, N.B., and LeBrasseur, R.J. 1986. Size selectivity of coho (*Oncorhynchus kisutch*)
491 preying on fry chum salmon (*O. keta*). *Can. J. Fish. Aquat. Sci.* **43**: 581–586.
- 492 Heard, W.R. 1991. Life history of pink salmon (*Oncorhynchus gorbuscha*). *In* Pacific salmon
493 life histories. *Edited by* C. Groot, and L. Margolis. University of British Columbia Press,
494 Vancouver, Canada. pp. 199–230.
- 495 Hebert, K.P., Goddard, P.L., Smoker, W.W., and Gharrett, A.J. 1998. Quantitative genetic
496 variation and genotype by environment interaction of embryo development rate in pink salmon
497 (*Oncorhynchus gorbuscha*). *Can. J. Fish. Aquat. Sci.* **55**: 2048–2057.
- 498 Hilborn, R., Quinn, T.P., Schindler, D.E., and Rogers, D.E. 2003. Biocomplexity and fisheries
499 sustainability. *P. Natl. Acad. Sci.* **100**: 6564–6568.
- 500 Kennedy, R.J., and Crozier, W.W. 2010. Evidence of changing migratory patterns of wild
501 Atlantic salmon *Salmo salar* smolts in the River Bush, Northern Ireland, and possible
502 associations with climate change. *J. Fish. Biol.* **76**: 1786–1805.
- 503 Kovach, R.P., Gharrett, A.J., and Tallmon, D.A. 2012. Genetic change for earlier migration
504 timing in a pink salmon population. *P. R. Soc. B* **279**: 3870–3878.

- 505 Kovach, R.P., Joyce, J.E., Echave, J.D., Lindberg, M.S., and Tallmon, D.A. 2013a. Temporal
506 patterns of genetic variation in a salmon population undergoing rapid change in migration
507 timing. *Evol. Appl.* **6**(5): 795–807.
- 508 Kovach, R.P., Gharrett, A.J., and Tallmon, D.A. 2013b. Earlier migration timing, decreasing
509 phenotypic variation, and biocomplexity in multiple salmonid species. *PLoS ONE* **8**(1): e53807.
- 510 Kovach, R.P., Ellison, S.C., Pyare, S., and Tallmon, D.A. 2015. Temporal patterns in adult
511 salmon migration timing across Southeast Alaska. *Glob. Chang. Biol.* **21**(5): 1821–1833.
- 512 Lane, S., McGregor, A.J. Taylor, S.G., and Gharrett, A.J. 1990. Genetic marking of an Alaskan
513 pink salmon population, with an evaluation of the mark and the marking process. *Am. Fish. S. S.*
514 **7**: 395–406.
- 515 Manhard, C.V., Joyce, J.E., Smoker, W.W., and Gharrett, A.J. 2017. Ecological factors
516 influencing lifetime productivity of pink salmon (*Oncorhynchus gorbuscha*) in an Alaskan
517 stream. *Can. J. Fish. Aquat. Sci.* Advance online publication. doi: 10.1139/cjfas-2016-0335.
- 518 Mortensen, D., Wertheimer, A., Taylor, S., and Landingham, J. 2000. The relation between early
519 marine growth of pink salmon, *Oncorhynchus gorbuscha*, and marine water temperature,
520 secondary production, and survival to adulthood. *Fish. Bull.* **98**: 319–335.
- 521 Mortensen, D.G., Wertheimer, A.C., Maselko, J.M., and Taylor, S.G. 2002. Survival and
522 straying of Auke Creek, Alaska, pink salmon marked with coded wire tags and thermally
523 induced marks. *T. Am. Fish. Soc.* **131**: 13–26.
- 524 O'Malley, K.G., Ford, M.J., and Hard, J.J. 2010. 2013. Adaptive genetic markers discriminate
525 migratory runs of Chinook salmon (*Oncorhynchus tshawytscha*) amid continued gene flow.
526 *Evol. Appl.* **6**(8): 1184–1194.

- 527 Parker, R.R. 1971. Size selective predation among fry salmonid fishes in a British Columbia
528 inlet. J. Fish. Res. Board. Can. **28**: 1503–1510.
- 529 Pella, J.J., and Milner, G.B. 1987. Use of genetic marks in stock composition analysis. *In*
530 Population genetics and fisheries management. *Edited by* N. Ryman, and F. Utter. University of
531 Washington Press, Seattle, USA. pp. 247–276.
- 532 Quinn, T.P., Unwin, M.J., and Kinnison, M.T. 2000. Evolution of temporal isolation in the wild:
533 genetic divergence in timing of migration and breeding by introduced Chinook salmon
534 populations. *Evolution* **54**: 1372–1385.
- 535 Robards, M.D. and Quinn, T.P. 2002. The migratory timing of adult summer-migrating steelhead
536 trout (*Oncorhynchus mykiss*) in the Columbia River: six decades of environmental change. T.
537 Am. Fish. Soc. **131**: 523–536.
- 538 Smoker, W.W., Gharrett, A.J., and Stekoll, M.S. 1998. Genetic variation of return date in a
539 population of pink salmon: a consequence of fluctuating environment and dispersive selection?
540 Al. Fish. Res. Bull. **5**: 46–54.
- 541 Taylor, S.G. 1980. Marine survival of pink salmon from early and late spawners. T. Am. Fish.
542 Soc. **109**: 79–82.
- 543 Taylor, S.G. 2008. Climate warming causes phenological shift in pink salmon, *Oncorhynchus*
544 *gorbuscha*, behavior at Auke Creek, Alaska. *Glob. Change Biol.* **14**: 229–235.
- 545 Thackeray, S.J., Sparks, T.H., Frederiksen, M., Burthe, S., Bacon, P.J., et al. 2010. Trophic level
546 asynchrony in rates of phenological change for marine, freshwater, and terrestrial environments.
547 *Glob. Change Biol.* **16**: 3304–3313.
- 548

549 Velsen, F.P.J. 1987. Temperature and incubation in Pacific salmon and rainbow trout:
550 compilation of data on median hatching time, mortality, and embryonic staging. Canadian Data
551 Report of Fisheries and Aquatic Science, Number 626. Minister of Supply and Services, Ottawa,
552 Canada. pp. 58.

Draft

Figures

Figure 1 – Five-day running averages of the frequency of the late-migrating marker allele (*MDH-B1,2*70*) in adult pink salmon sampled in each odd-year migration from 1985-1991. Median migration dates are indicated by a dashed line, and the 1983 late-migrating marker allele frequencies are depicted for early- (circle) and late- (triangle) migrating fish.

561
562
563 **Figure 2** – Five-day running averages of the frequency of the late-migrating marker allele (*MDH-B1,2*70*) in pink salmon fry
564 sampled in each odd-broodyear migration from 1984-1992. Median migration dates are indicated by a dashed line.

Figure 3 – Annual bootstrap estimates (20,000 iterations) of the population-wide frequency of the late-migrating marker allele (*MDH-B1,2*70*) in migrating adults (solid circle) and fry (open circle). Error bars are the 95% confidence intervals of each estimate.

571
572
573 **Figure 4** – Bootstrap estimates (20,000 iterations) of marine survival and freshwater productivity (fry produced per adult) over four
574 brood years in the early- (open circle) and late- (closed circle) migrating subpopulations. Error bars are the 95% confidence intervals
575 of each estimate. Shading emphasizes the comparison of subpopulations within each brood year.
576
577
578

Figure 5 – Sea-surface temperatures experienced by pink salmon fry from the early- (solid line) and late-migrating (dashed line) subpopulations during their first week of marine residency. Assignment to subpopulations was made with a stock separation algorithm that used allele frequencies at the late-migrating marker locus. Three-day running averages of Auke Bay sea-temperatures (°C) are depicted by a gradation scheme.

Figure 6 – Temporal trend in spring sea-surface temperature of Auke Bay (1975-2011). Sea-surface temperatures were averaged from 1 April to 31 May. The shaded area depicts the 95% confidence interval.