Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

**Hot eats and cool creeks: juvenile Pacific salmonids use
mainstem prey while in thermal refuges**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0395.R1
Manuscript Type:	Article
Date Submitted by the Author:	29-Nov-2016
Complete List of Authors:	Brewitt, Kimberly S.; University of South Carolina Upstate, Division of Natural Sciences and Engineering; Southwest Fisheries Science Center Fisheries Ecology Division Danner, Eric M.; Southwest Fisheries Science Center Fisheries Ecology Division Moore, Jonathan; Simon Fraser University, Earth to Ocean Research Group
Keyword:	Chinook salmon, Klamath River, radio telemetry, steelhead, thermal refuges

SCHOLARONE™
Manuscripts

TITLE

Hot eats and cool creeks: juvenile Pacific salmonids use mainstem prey while in thermal refuges

AUTHORS

Kimberly S. Brewitt¹, Eric M. Danner, Jonathan W. Moore

Kimberly S. Brewitt. Fisheries Ecology Division, NOAA National Marine Fisheries Service, Southwest Fisheries Science Center, 110 Shaffer Road, Santa Cruz, California 95060 USA. (e-mail: kbrewitt@uscupstate.edu)

Eric M. Danner. Fisheries Ecology Division, NOAA National Marine Fisheries Service, Southwest Fisheries Science Center, 110 Shaffer Road, Santa Cruz, California 95060 USA. (e-mail: eric.danner@noaa.gov)

Jonathan W. Moore. Earth to Ocean Research Group, Simon Fraser University, Burnaby, BC, Canada. (e-mail: jwmoore@sfu.ca)

Corresponding author: Kim Brewitt. (e-mail: kbrewitt@uscupstate.edu)

Telephone: 978-660-1476. Fax: 864-503-5366

¹ Current affiliation: Division of Natural Sciences and Engineering, University of South Carolina Upstate, Spartanburg, South Carolina 29302 USA.

1 ABSTRACT

2 Thermal refuges form important habitat for coldwater fishes in the face of rising temperatures.

3 As fish become concentrated in refuges, food resources may become depleted. In this study, we

4 used invertebrate drift sampling and fish density surveys to quantify potential in-refuge food

5 limitation, temperature-sensitive radio-tagging studies to quantify thermal habitat use, and

6 isotopic analyses to determine diet sources for juvenile Pacific salmonids using thermal refuges

7 on California's Klamath River. Juvenile salmonids using refuges formed by tributary junctions

8 with the mainstem river obtained the majority (range = 47-97%) of their diet from mainstem prey

9 sources. Mean steelhead body temperatures were significantly cooler ($\sim 3.5^{\circ}\text{C}$) than diet-inferred

10 foraging temperatures. Thus, while fish seek cooler habitat for physiological benefits, they rely

11 primarily on mainstem prey. Moreover, consistently high densities of fish in refuges (mean = 3.5

12 fish m^{-2}) could lead to density-dependent food limitation. Thus, mobile consumers like fish can

13 exploit existing heterogeneity associated with coldwater refuges by gaining thermal benefits

14 from a food-limited coldwater habitat while deriving the majority of their prey from the warm

15 mainstem river.

16
17 KEYWORDS: Chinook salmon; Klamath River; radio telemetry; steelhead; thermal refuges

18 INTRODUCTION

19 Habitat heterogeneity provides organisms with a choice of disparate habitats, and mobile
20 consumers can integrate across a heterogeneous landscape, thereby balancing trade-offs in
21 resources (Bridges 2002, Hohnsbova et al. 2003, Scheuerell and Schindler 2003, Sims et al.
22 2006, Loose and Dawidowicz 2014). For coldwater fish, thermal refuges provide physiological
23 relief from stressful in-stream temperatures, but they are also heterogeneous environments where
24 the distribution of other resources within and surrounding a refuge (e.g., prey availability,
25 predation risk) could influence the relative costs and benefits of refuge use. Thermal refuges,
26 coldwater habitat that allows coldwater fish to escape the sub-lethal and lethal effects of hot
27 temperatures (Torgersen et al. 2012), are often formed by tributaries that are colder than the
28 mainstem river and create thermally heterogeneous mixing zones at their confluences. Most
29 studies on thermal refuges focus on their thermal benefits (Matthews et al. 1994, Nielsen et al.
30 1994, Torgersen et al. 1999, Ebersole et al. 2001); fish exploit in-stream thermal heterogeneity
31 by moving into cooler water (i.e., thermal refuges) to relieve heat stress and reduce metabolic
32 demand (Berman and Quinn 1991, Baird and Krueger 2003). Thermal refuges form increasingly
33 important habitat as river temperatures continue to rise (Isaak et al. 2011, Ruesch et al. 2012,
34 Davis et al. 2013), and these habitats have the potential to enable population persistence in what
35 may eventually become thermally uninhabitable ecosystems (Sutton et al. 2007, Loarie et al.
36 2008). However, there may be trade-offs to refuge use; for example, adult steelhead (*O. mykiss*)
37 using coolwater refuges in the Columbia River basin experienced increased harvest mortality
38 (Keefer et al. 2009). A better understanding of how individuals use thermal refuges may help us
39 understand the potential trade-offs and limits of these habitats, and the consequences for the
40 dynamics and persistence of populations in a warming world.

While fish gain thermal benefits from coolwater refuges, these areas may be sub-optimal in terms of prey availability (Sutton et al. 2007). Studies on fish using thermal refuges in warm rivers have recorded cyclic movements out of refuges into thermally stressful mainstem habitat, presumably to gain some non-thermal benefit (Kaeding 1996, Ebersole et al. 2001, Belchik 2003, Brewitt and Danner 2014). In-stream prey availability in river ecosystems is highly variable (Allan and Russek 1985, Shearer et. al 2002, Neale et. al 2008). If prey are more abundant in areas where temperatures are sub-optimal, individuals may forage in warmer water but move to cooler water to digest, thereby decreasing metabolic costs (Bevelhimer and Adams 1993, Sims et. al 2006).

Competition for prey resources between individuals using refuges likely varies as a result of both naturally heterogeneous in-stream prey availability and fish density (Shearer et al. 2002, Rosenfeld et al. 2005, Hayes et al. 2007). Artificial stream channel experiments simulating refuges have found that density-dependent effects can lead to decreased growth rates, and higher mortality and emigration rates (Keeley 2001, Armstrong and Griffiths 2001, Imre et al. 2004). Within-refuge fish densities increase with rising mainstem temperatures, often reaching high levels during hot summer months (Sutton et al. 2007, Mather et al. 2008, Brewitt and Danner 2014). We hypothesize that salmonids in the Klamath Basin gain thermal benefits from a food-limited coldwater habitat while deriving the majority of their prey from the much warmer food-rich but thermally stressful mainstem river. In this study, we examine how steelhead and Chinook salmon respond to the heterogeneous thermal and prey landscape surrounding thermal refuges at tributary confluences of a seasonally warm river system.

In the Klamath River in northern California, steelhead trout (*O. mykiss*) and Chinook salmon (*O. tshawytscha*) populations are reduced to approximately 5% of historical levels, and

juvenile salmonids increasingly rely on refuges for suitable thermal habitat during hot summer months (Nehlsen et al. 1991, Brewitt and Danner 2014). Given possible trade-offs to refuge use, including density-dependent food limitation, we sought to identify mechanisms of food limitation in refuges and to assess the diet and thermal habitat use of juvenile salmonids using thermal refuges. We therefore asked, 1) Are there differences in invertebrate prey availability between the tributaries and the larger mainstem system? 2) Are fish densities within refuges higher than in the adjacent mainstem river? 3) What proportion of juvenile steelhead and Chinook salmon diet is derived from the mainstem river versus tributary, and are there any seasonal, size-dependent, or species differences in diet? And 4) Are juvenile steelhead foraging primarily in the same areas they are holding (i.e., thermal habitat use)?

METHODS

We used a multi-faceted approach to quantify the thermal and prey landscapes surrounding thermal refuges (see Table 1). We sampled invertebrate drift (2010 and 2011) to quantify fish prey sources and assess whether invertebrate energy delivery rates ($\text{Joules m}^{-2} \text{s}^{-1}$) could be a mechanism of food limitation within thermal refuges. To assess how fish use the landscape surrounding thermal refuges at the individual and population level, we used snorkel surveys (2012) to estimate fish density in and out of refuges, temperature-sensitive radio tagging studies (2010-2012) to quantify thermal habitat use, and isotopic analyses (2012) to estimate juvenile salmonid prey sources.

Study system

The Klamath River in northern California is 423 km long, and drains approximately 41,440 km² of southern Oregon and northern California (National Research Council 2008) (Fig. 1). It has six mainstem dams, the lowest of which, Iron Gate Dam (rkm 306), acts as a migration barrier to anadromous fish, cutting off hundreds of kilometers of native anadromous salmonid habitat (Nehlsen et al. 1991). The dam, along with large-scale watershed alteration (e.g., irrigation), contributes to elevated water temperatures for the downstream habitat (Bartholow 2005, National Research Council 2008). During the summer and fall, the Iron Gate Reservoir undergoes seasonal thermal stratification, and the dam releases warm surface water rather than releasing colder water from below the thermocline (National Research Council 2004).

Below this barrier, the Klamath River supports migrations of steelhead trout (*O. mykiss*), Chinook salmon (*O. tshawytscha*), coho salmon (*O. kisutch*), and chum salmon (*O. keta*). Chinook salmon are the most abundant anadromous fish in the basin, with both fall- and spring-run life-history types, but populations have declined and are far below historical levels. Both summer and winter steelhead runs have declined from historical levels, and coho salmon were listed as threatened under the U.S. Endangered Species Act in 1997 (National Research Council 2008). Juvenile steelhead rear in the river for 1-3 years, and juvenile Chinook salmon rear for 5-12 months, before out-migrating to the ocean. During summer months the Klamath mainstem river reaches temperatures that can be thermally inhospitable to salmonids—mean daily mainstem temperatures at the study sites ranged from 15-25°C between July-September 2010-2012—and juveniles behaviorally thermoregulate by seeking out thermal refuges, usually at tributary confluences (Sutton et al. 2007, Brewitt and Danner 2014).

We conducted this research on the lower Klamath River in July-September 2010-2011, and July-August 2012. We chose four study sites on the river below Iron Gate Dam, each located

at a major coldwater tributary confluence: Beaver Creek (rkm 259), Grider Creek (rkm 210), Fort Goff Creek (rkm 204), and Thompson Creek (rkm 198). We chose these tributaries based on the presence of coolwater refuges during summer and early fall months, and the presence of juvenile steelhead and Chinook salmon. Hereafter, we refer to ‘refuges’ as the coolwater area that includes both the tributary and thermal mixing zone (i.e., the area where mainstem and tributary water mix, creating an area of heterogeneous temperature).

Water temperatures

To determine the difference in mean water temperature between the tributaries and mainstem river, we recorded water temperatures at 15-minute intervals throughout July-August 2012 at each study site. We used 2-4 Hobo pendant data loggers (Onset Computer Corporation, Pocasset, MA; $\pm 0.5^{\circ}\text{C}$) in both the coldwater tributary and mainstem river at point locations approximately 10 m upstream of each confluence. Recorded temperatures therefore represent the minimum and maximum temperatures available to fish. All data loggers were placed near the riverbed to minimize thermal input from direct sunlight, at depths ranging from 0.5–2 m.

Mainstem versus tributary invertebrate energy delivery rates

To assess whether fish may be experiencing food limitation in thermal refuges, we collected invertebrate drift samples and calculated energy delivery rates ($\text{Joules m}^{-2} \text{second}^{-1}$) as an estimate of food availability in both the mainstem river and coldwater tributaries. We collected monthly samples (June-August) at all four study sites in 2010, and weekly samples (July-August) at Beaver Creek (farthest upstream) in 2011; samples were collected from the mainstem and tributary approximately 15 m upstream of the confluence, and upstream of where

high densities of fish were observed in the tributaries (Fig. 2a). We suspended mesh drift nets (23x31 cm, 500 μ m) approximately 5 cm above the substrate for 5-10 minutes, and measured water velocity directly in front of the net at 60% of water column depth at both the beginning and end of the sample period to estimate mean flow. We took samples at dawn and dusk to capture temporal variation in drift, and collected approximately two samples per thermal environment during each sampling event to capture spatial variation in drift (Appendix: Table A1&A2). Invertebrate samples were identified, counted, and measured to the nearest millimeter. We estimated invertebrate biomass (mg dry mass) from taxon-specific length–mass regressions (Rogers et al. 1977, Hodar 1996, Burgherr and Meyer 1997, Kawabata and Urabe 1998, Benke et al. 1999, Miserendino 2001, Sabo et al. 2002, Baumgartner and Rothhaupt 2003), and calculated invertebrate energy content (Joules / mg) for each sample using taxon-specific caloric values in the literature (Cummins and Wuycheck 1971, Driver et al. 1974, Driver 1981, Ciancio and Pacual 2006).

We calculated energy delivery rates (EDR ; Joules m^{-2} second $^{-1}$) as an estimate of salmonid prey availability, defined as the product of energy density and water velocity:

$$(1) \quad EDR = (E / V) * v$$

where E is the total energy in a sample, V is the water volume of the sample, and v is the mean water velocity in front of the drift net. Energy delivery rate explicitly takes into account water velocity, making it a better metric than energy density for assessing how much food per unit time is reaching a fish holding in a specific location. Since there was a high degree of variability in the number of samples collected per month at each study site, we used a meta-analytical approach to assess whether there was any difference between mainstem and tributary drift delivery rates across sites and years. We treated each group of monthly samples taken at a site as

a separate analysis, and determined the effect size between mainstem and tributary energy delivery rates; an effect size of zero indicates no difference. While invertebrate drift data were taken in 2010 and 2011, and all isotopic samples and fish density data were collected in 2012, we are using these drift data simply to draw inferences about general patterns of relative food supply in thermal refuges.

Fish density in and out of refuges

We quantified fish densities in the mainstem river and refuge (mixing zone and tributary) to assess whether fish density could be a possible source of in-refuge food limitation. We conducted weekly snorkel surveys at three times of day (9:00, 12:00, 16:00) at the Beaver Creek study site throughout August 2012. We established nine 1x2 m² plots, three in the tributary (T), four in the mixing zone (MZ), and two in the mainstem (MS) (Fig. 2a); all plots were marked with orange flagging on corner rocks. To estimate fish densities, the snorkeler approached the plot from downstream to avoid spooking the fish, and held a position in the water near the plot for approximately ten minutes. We calculated mean steelhead and Chinook salmon densities for all plots across the month. While the limited spatial extent of these surveys may have underestimated mainstem river salmonid densities, where fish may be more patchily distributed, this survey design was adequate for establishing relative densities of fish in thermal refuges and the adjacent mainstem river.

Quantifying juvenile salmonid diet sources

We used stable isotope analyses to quantify the proportion of mainstem versus tributary prey contributing to the diet of juvenile steelhead and Chinook salmon using thermal refuges at

the two upstream study sites (Beaver and Grider Creek confluences) in 2012. We chose two sampling periods (early July and late August) in order to assess whether there were any seasonal (i.e., temperature-dependent) changes in fish diet. We chose these sampling periods because mainstem temperatures rise with decreased flows and increased air temperatures, and are therefore usually cooler in early July than in late August.

To determine stable isotope values of aquatic prey, we collected benthic invertebrate samples from the tributary and mainstem at both sites. We chose taxa from the benthic samples that were representative of the most common invertebrate families found in fish diets and in the drift for both thermal environments (tributary and mainstem), and that represented a range of functional groups (see Appendix). We performed invertebrate sampling twice during the two-week period prior to fish sampling in early July and late August 2012, to capture the integration window for fish fin tissue (Heady and Moore 2012). During each sampling event, we collected samples of benthic invertebrates by scrubbing the substrate at 4-7 point locations between 5-15 m upstream of the confluence, and collecting aquatic invertebrates with a 500 μ m net. We identified invertebrates to family under a stereomicroscope, and analyzed 2-3 samples of each taxon.

We took caudal fin tissue samples to measure stable isotope ratios for juvenile steelhead and Chinook salmon. Fish were caught within 50 m of the tributary confluence (in either the tributary, thermal mixing zone, or mainstem) using a combination of angling, electro-fishing, and seining at Beaver Creek in July ($n = 30$ steelhead; $n = 18$ Chinook salmon) and August ($n = 23$ steelhead; $n = 7$ Chinook salmon), and Grider Creek in July ($n = 14$ steelhead) and August ($n = 18$ steelhead) 2012. A small subset of the fish at Beaver Creek ($n = 6$) were caught in early August for radio tagging, but these fish showed no statistical differences in inferred diet from the

fish caught in July, and were therefore grouped with the early July fish. We weighed and measured (fork length; FL) all fish prior to taking fin clips.

We used stable isotopes of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) to investigate the contribution of mainstem versus tributary prey to the diet of juvenile salmonids. Prior to isotopic analyses, invertebrate samples were frozen, and all samples were subsequently dried at 15°C in a drying oven. For small invertebrates (e.g., chironomids), multiple individuals were pooled into one sample and homogenized. All samples were analyzed using a PDZ Europa ANCA-GSL elemental analyzer interfaced with a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK) at the University of California, Davis Stable Isotope Facility. Isotopic composition is expressed in δ values (parts per thousand (‰) different from a standard), calculated as:

$$(2) \quad \delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1] * 1000$$

where X is the element, R is the ratio of the heavy to light element, and the standards are Vienna Pee Dee Belemnite limestone (V-PDB) and air for carbon and nitrogen, respectively.

We used MixSIAR, a Bayesian stable isotope mixing model (Stock and Semmens 2013), to determine the probability distributions for the relative contribution of different prey sources to salmonid diet. Within MixSIAR, Gibbs sampling was performed for each of three parallel chains. We used a burn-in phase of 10,000 iterations, and ran the model for 100,000 iterations, retaining every 15th posterior sample, resulting in 6,000 independent draws for the posterior distribution. We used Gelman-Rubin diagnostics to confirm model convergence (Gelman and Rubin 1992). We ran the model separately for each study site, and included sampling month as a categorical variable in the model, and individuals as a random effect. We incorporated a trophic

discrimination factor of 3.2 ± 0.2 for $\delta^{15}\text{N}$ and 1.9 ± 0.5 for $\delta^{13}\text{C}$ ($\pm 1\text{SD}$) into the mixing model (McCutchan et al. 2003).

We collected juvenile steelhead diet samples to qualitatively assess whether fish diet composition was similar to the species composition of the drift. Fish were sampled by hook and line from Beaver and Fort Goff sites in 2010 ($n = 22$) and from Fort Goff in 2011 ($n = 12$). (Appendix: Table A3 and Fig. A3&A4).

Fish thermal habitat versus foraging habitat

We used a two-part approach to assess whether juvenile steelhead were foraging primarily in the same areas they were holding (i.e., thermal habitat use). First, we used an individual-level analysis for the small number of steelhead ($n = 9$) for which we had both body temperature data and isotopic diet analyses; we compared mean thermal habitat use derived from radio-tag data to the mean water temperature where individuals were foraging, inferred from isotopic diet analyses (hereafter, diet-inferred foraging temperatures). Since this first analysis only allowed us to analyze a small number of individuals, we then used a population-level analysis to address the same question, comparing diet-inferred foraging temperatures for all steelhead for which we had isotopic diet data (July-August 2012; $n = 53$ at Beaver Creek; $n = 32$ at Grider Creek) to thermal habitat use across three years of temperature sensitive radio-tagging data (June-August 2010-2012; $n = 76$ at Beaver Creek; $n = 40$ at Grider Creek).

We used temperature-sensitive radio tags to track the body temperature of juvenile steelhead caught at Beaver Creek and Grider Creek study sites. Size limitations prevented tagging juvenile Chinook salmon. We surgically implanted the tags (Lotek's MST-720T temperature-sensor transmitter tags; 1.3g dry weight; $\pm 0.8^\circ\text{C}$) following the tagging protocol

described in Brewitt and Danner (2014). The life expectancy of the radio tags was 42 days. We used data-logging receivers (Lotek SRX_400A and SRX600) to log fish body temperatures at 5-second intervals for any tagged fish within approximately 100 m of the tributary confluence.

For the individual-level analysis of steelhead radio-tagged at Beaver Creek in 2012, we compared the proportion of time spent in mainstem water (derived from radio-tag data) to the mean contribution of mainstem prey sources to fish diet (derived from isotopic diet analyses). For this analysis, we used only fish for which we had both temperature data and isotopic diet data, and that remained at the refuge for a minimum of seven days post-tagging ($n = 9$). We used a mass-balance equation to translate time-series of fish body temperatures into the mean proportion of time an individual held in mainstem water (P_m). The equation we used was:

$$(3) \quad P_m = \text{mean} ((T_f - T_t) / (T_m - T_t))$$

where T_f represents time-series of fish temperatures, and T_t and T_m represent simultaneous time-series of tributary and mainstem water temperatures, respectively. In making this comparison, we are assuming that these individuals' behavior was similar before and after tagging, since the fish temperature time-series represent data collected in the two weeks after fin samples were taken for isotopic diet analysis; this constraint was due to our inability to recapture radio tagged fish post-tagging. We believe this is a reasonable assumption, since time-series of fish temperature were fairly consistent for each individual tagged, until the fish left the study area.

We then used a population-level analysis to determine whether the diet-inferred foraging temperatures of all fish for which we had isotopic diet data (2012; $n = 53$ at Beaver Creek; $n = 32$ at Grider Creek) matched the holding temperatures of all fish for which we had radio-tagging data (2010-2012; $n = 76$ at Beaver Creek; $n = 40$ at Grider Creek). We used a mass-balance equation to calculate the water temperature where fish were foraging (diet-inferred foraging

temperature; T_d) based on their isotope-inferred diet. We used time-series of mainstem and tributary temperatures from the two weeks prior to fish sampling to characterize the water temperature profile experienced by individuals during the fin tissue integration window (Heady and Moore 2012), and multiplied the time-series through by the isotope-inferred proportion mainstem diet for each fish. The diet-inferred foraging temperature for an individual fish can therefore be expressed as:

$$(4) \quad T_d = \text{mean} ((D_m * T_m) + ((1 - D_m) * T_t))$$

where D_m is the proportion of mainstem diet for an individual, and T_t and T_m represent time-series of mainstem and tributary temperatures. Thus, T_d represents a metric of the temperature experienced by the fish when they were obtaining their prey. For the purposes of this calculation, we assumed that mainstem and tributary prey in the mixing zone are being mixed just like water, since steelhead primarily feed on invertebrates in the water column (i.e., drift) that use passive dispersal mechanisms. In addition, we are assuming that local prey production in the mixing zone does not contribute significantly to fish diet, given that the volume of the mixing zone is minimal compared to the larger mainstem and tributary water bodies. While T_d is a useful index of potential foraging temperatures, there may be sources of uncertainty (e.g., non-random daily foraging patterns) that we have not taken into consideration in these estimates.

To quantify thermal habitat use for the population of fish using the thermal refuge at Beaver Creek and Grider Creek study sites, we used data from a larger temperature-sensitive radio-tagging study (Brewitt and Danner 2014). We calculated the fish body temperature distribution for the sample of steelhead tagged at each site during the study period 2010-2012 at Beaver Creek ($n = 76$) and 2010-2011 at Grider Creek ($n = 40$), using all fish detections subsampled at 5-minute intervals, and weighting all fish equally. We used a Kolmogorov-Smirnov

two-sample test to test for differences between the thermal and foraging temperature distributions at each site.

RESULTS

Water temperatures

Mainstem temperatures were consistently warmer than tributary temperatures at Beaver Creek (mean difference = $6.8 \pm 0.5^{\circ}\text{C}$) and Grider Creek (mean difference = $7.1 \pm 0.9^{\circ}\text{C}$) throughout July and August 2012 (Appendix: Fig. A1). Mean mainstem temperatures for the two weeks prior to fish isotope sampling in early July and late August 2012 were 22.5°C and 22.0°C at Beaver Creek, and 21.8°C and 21.6°C at Grider Creek, respectively. In contrast, mean temperatures in the coldwater refuges during these same periods ranged from $15.5 - 19.5^{\circ}\text{C}$ at Beaver Creek and $13.1 - 18.8^{\circ}\text{C}$ at Grider Creek.

Mainstem versus tributary invertebrate energy delivery rates

Invertebrate energy delivery rates were highly variable across sites, and were not consistently higher in the larger mainstem river than the tributaries. In 2010, energy delivery rates ranged from $0.9\text{-}43.3 \text{ Joules m}^{-2} \text{ s}^{-1}$ in the mainstem and $0.4\text{-}37.5 \text{ Joules m}^{-2} \text{ s}^{-1}$ in the tributary across the four study sites (Appendix: Fig. A2). In 2011, energy delivery rates at Beaver Creek site ranged from $13.2\text{-}86.7 \text{ Joules m}^{-2} \text{ s}^{-1}$ in the mainstem and $11.6\text{-}828.9 \text{ Joules m}^{-2} \text{ s}^{-1}$ in the tributary. The mean ($\pm$ SD) effect size between the mainstem and tributaries across all sites and years was $0.102 \pm 1.18 \text{ Joules m}^{-2} \text{ s}^{-1}$; an effect size of zero indicates no difference. Thus, there was no significant difference between mainstem and tributary prey supply per unit area.

Fish density in and out of refuges

Steelhead and Chinook salmon densities were approximately eight times higher in the refuge than in the mainstem river at Beaver Creek throughout August 2012 (Fig. 2b,c). Mean ($\pm$ SD) juvenile steelhead densities (fish m^{-2}), aggregated across hourly and weekly observations, were higher in the tributary (mean = 2.0 ± 1.3) and mixing zone (mean = 1.9 ± 1.6) than in the mainstem plots (mean = 0.4 ± 0.4). The plot with the consistently highest density of steelhead (4.0 fish m^{-2}) was located near the top of the mixing zone, just below the tributary confluence with the mainstem, and had a mean temperature of 19.8°C . Mean ($\pm$ SD) juvenile Chinook salmon densities were also higher in the tributary (mean = 1.7 ± 2.3) and mixing zone (mean = 1.5 ± 1.9) than in the mainstem plots (mean = 0 ± 0.1). The plot with the consistently highest density of Chinook (3 fish m^{-2}) was located at the downstream portion of the mixing zone, in an area of lower velocity flow, and had a mean temperature of 20.5°C . Fish density within a thermal environment did not vary by time of day for either steelhead ($F = 1.97$; $p = 0.16$) or Chinook salmon ($F = 0.047$; $p = 0.83$).

Quantifying juvenile salmonid diet sources

Mainstem invertebrate prey sources were $\delta^{15}\text{N}$ -enriched and $\delta^{13}\text{C}$ -enriched compared to tributary invertebrate prey sources. Juvenile steelhead and Chinook salmon isotope signatures ranged between tributary and mainstem prey source signatures, indicating diet variation between individuals (Fig. 3a,b). There were no seasonal differences in juvenile steelhead or Chinook salmon diet between the two sampling periods (early July and late August).

Juvenile salmonids using thermal refuges obtained over 50% of their diet from mainstem prey sources (Fig. 4). The mean proportion mainstem diet for juvenile steelhead caught in the

mainstem (Beaver = 0.82 ± 0.13 standard deviation (SD); Grider = 0.97 ± 0.02) and mixing zone (Beaver = 0.75 ± 0.13 ; Grider = 0.94 ± 0.11) was much higher than for tributary-caught steelhead (Beaver = 0.57 ± 0.20 ; Grider = 0.60 ± 0.38). However, tributary-caught steelhead and Chinook salmon (0.47 ± 0.15) still obtained approximately half their diet from mainstem prey sources (Fig. 4a,b).

Tributary-caught steelhead had a smaller mean fork length (Beaver = 148 ± 40 mm; Grider = 134 ± 34 mm) than steelhead caught in the mixing zone (Beaver = 168 ± 26 mm; Grider = 218 ± 31 mm) or mainstem (Beaver = 212 ± 80 mm; Grider = 203 ± 71 mm). The proportion of diet obtained from mainstem sources increased with increasing steelhead body size (fork length) ($R^2 = 0.30$ and 0.23 for Beaver and Grider Creeks, respectively; linear regression).

Fish thermal habitat versus foraging habitat

Steelhead body temperatures describing thermal habitat use were significantly cooler ($\sim 3.5^\circ\text{C}$) than diet-inferred steelhead foraging temperatures, indicating a mismatch in thermal and foraging habitat for fish using thermal refuges. Fish temperatures for the nine radio-tagged steelhead at Beaver Creek indicate that fish used a range of thermal habitat (range in mean body temperature across sample period: $15.7^\circ\text{C} - 22.5^\circ\text{C}$). Based on these body temperatures and river temperatures over the same time period, the mean ($\pm$ SD) proportion of time individuals spent in mainstem water (0.22 ± 0.23) was relatively low, whereas the mean proportion of prey these same individuals obtained from mainstem sources (0.72 ± 0.15) was higher than would be expected if fish were feeding in the area in which they were holding based on body temperatures (Fig. 5).

In the population-level analysis comparing thermal habitat use and diet-inferred foraging temperatures, fish body temperatures at both Beaver Creek (mean = $17.5 \pm 2.3^{\circ}\text{C}$) and Grider Creek (mean = $17.8 \pm 2.3^{\circ}\text{C}$) were significantly cooler than diet-inferred foraging temperatures at Beaver Creek (mean = $20.7 \pm 1.2^{\circ}\text{C}$) and Grider Creek (mean = $21.0 \pm 2.6^{\circ}\text{C}$) (Kolmogorov-Smirnov two-sample test, $p < 0.001$ at both sites; Fig. 6).

DISCUSSION

We found that while juvenile steelhead and Chinook salmon consistently used thermal refuges to behaviorally thermoregulate, they obtained the majority of their prey from mainstem sources, even during thermally stressful summer months when the mainstem neared upper incipient lethal temperatures. Steelhead body temperatures centered around $18\text{--}19^{\circ}\text{C}$, the optimal growth temperature given sufficient food availability (Myrick and Cech 2000); this suggests that steelhead use refuges to thermoregulate and possibly optimize growth, but can still obtain much of their prey from warmer mainstem water (mean diet-inferred foraging temperature $\cong 21\text{--}22^{\circ}\text{C}$). Thus, juvenile salmonids can exploit the heterogeneity at tributary junctions forming thermal refuges by using refuges to thermoregulate while still accessing the mainstem river prey base.

Our results suggest that juvenile salmonids using coolwater refuges shift laterally between the refuges and the mainstem river in order to access mainstem prey. While we did not measure fish movement directly, steelhead body temperatures used as a proxy for location indicate that fish move in and out of thermal refuges (Brewitt and Danner 2014). The high proportion of mainstem prey in isotope-inferred fish diets therefore implies that juvenile steelhead and Chinook salmon move out of refuges to obtain mainstem prey. There is evidence of fish in other systems using diel vertical or horizontal migration as a strategy to maximize

access to food availability and metabolic efficiency across heterogeneous thermal landscapes (Wurtsbaugh and Neverman 1988, Bevelhimer and Adams 1993, Armstrong et al. 2013). For example, Armstrong and Schindler (2013) found that juvenile coho salmon forage in colder water but shift to warmer water to digest, thereby increasing digestive capacity; conversely, Sims et. al (2006) found that a benthic shark (*Scyliorhinus canicula*) lowered daily metabolic costs by hunting in warmer water and shifting to cooler water to digest.

Juvenile salmonid movement out of thermal refuges into sub-optimal mainstem temperatures suggests that refuges may be food-limited, likely due to high fish densities causing competition for prey. Invertebrate energy delivery rates were highly variable, and there was no consistent difference between energy delivery rates in the tributaries and mainstem river; invertebrate drift densities alone are therefore unlikely to drive fish to consistently forage outside of thermal refuges. However, fish densities were approximately eight times higher in the refuge (mean = 3.5 fish m⁻²) than the adjacent mainstem river throughout August 2012, and competition for prey would therefore be much higher in the refuge than the mainstem. These densities were much higher than stream-dwelling steelhead densities in other systems (Everest and Chapman 1972, Harvey and Nakamoto 1996); for example, Keeley and McPhail (1998) measured juvenile steelhead densities of approximately 1.9 fish m⁻² (mean FL $\cong$ 70 mm) in two Canadian rivers. As mainstem temperatures rise, movement of individuals into thermal refuges will increase in-refuge fish densities, likely reducing per capita prey availability. Density-dependent processes may therefore be driving individuals to venture into warmer water to access more prey; however, access to the mainstem prey base could allow thermal refuges to support higher densities of fish.

Steelhead size appears to mediate how individuals negotiate the variability in temperature and prey availability in refuges, as indicated by size-based variation in steelhead isotope-inferred

diet. Larger juvenile steelhead (FL ~ 200 mm) caught in the mainstem and mixing zone were more likely to rely on mainstem prey (~75-95% mean proportion mainstem prey) than smaller steelhead (FL ~ 150 mm) caught in the tributary, who had a lower proportion (~60% mean proportion mainstem prey) of mainstem prey in their isotope-inferred diet composition (see Fig. 4). Given sufficient food availability, the mixing zone and boundary habitat (i.e., boundary between the refuge and mainstem river) is likely optimal habitat for fish growth; mixing zone temperatures were often at the bioenergetic optima for steelhead (~18-19°C) (Brett et al. 1982; Heady 2012; Myrick and Cech 2000). Larger juveniles may be holding dominant positions near the edge of thermal refuges where temperature and prey availability can be optimized, while smaller fish hold in cooler tributary waters. Both observational and radio-tagging studies on juvenile steelhead in Klamath River refuges have observed fish (especially larger juveniles) concentrated along the edge of refuges (Sutton et al. 2007, Brewitt and Danner 2014). An alternate hypothesis is that larger individuals are able to make longer feeding forays into the mainstem river before incurring thermal stress, since it takes longer for the body temperature of larger fish to equilibrate to ambient temperatures (Pepino et al. 2015). Balancing resource needs may therefore be easier for larger juvenile steelhead that can establish dominant positions on the mixing zone boundary habitat, and likely forage in the mainstem river at longer time scales, while smaller tributary-based juveniles that still need access to mainstem prey would need to move further to gain sufficient resources.

Juvenile steelhead and Chinook salmon did not always hold in the coldest areas of the refuge, and both species tended to cluster in areas that met their species-specific habitat requirements (Everest and Chapman 1972, Quinn 2005). Chinook salmon were caught in either the tributary or the lower mixing zone (~50 m below the confluence) in slower velocity areas,

rather than the higher velocity riffle forming the top of the mixing zone where larger steelhead were found. Juvenile Chinook salmon also exhibited size-based variation in holding position within the refuge; Chinook salmon caught in the lower mixing zone ($\sim 20.5^{\circ}\text{C}$) were larger (mean FL = 88.3 mm) than those caught in the tributary ($\sim 16.8^{\circ}\text{C}$) (mean FL = 83.3 mm). The preference of larger Chinook salmon for warmer water suggests that, as with steelhead, larger individuals may be holding dominant positions nearer the boundary of thermal refuges where temperature and prey availability can be optimized; given sufficient prey availability, Chinook salmon growth potential is optimal at approximately 19°C (Brett et al. 1982, McCullough 1999). Maintaining habitat heterogeneity in thermal refuges could provide the necessary temperature and prey resources for different species and size-classes of juveniles present, and facilitate co-existence due to differential habitat selection and foraging strategies.

Our results emphasize that not only are thermal refuges at tributary junctions important for fish seeking thermal respite, but the efficacy of refuges may depend on food web dynamics in both tributary and mainstem habitats. Even if the mainstem river is apparently too hot for fish to reside in, it still provides key food resources for fish in thermal refuges. While the scope of this study was limited to the Klamath River, the isotopic values for fish at both study sites make a compelling case that fish using thermal refuges are foraging in the mainstem; moreover, the population-level analysis of steelhead shows a clear mismatch between diet-inferred foraging temperatures and thermal habitat use (Fig. 6), indicating that fish are disproportionately foraging in the mainstem river. Rivers along the west coast are experiencing warming trends similar to the Klamath River (Knowles and Cayan 2002, Payne et al. 2004, Bartholow 2005, Isaak et al. 2012), making thermal refuges essential habitat for the over-summer survival of coldwater fish such as stream-rearing salmonids. As mainstem temperatures rise, any density-dependent food limitation

452 in refuges will likely be exacerbated by higher in-refuge fish densities (Ebersole et. al 2001,
453 Belchik 2003, Sutton et al. 2007, Brewitt and Danner 2014), as well as by low tributary flows
454 that could cut off connectivity from the mainstem river (Van Kirk and Naman 2009). While
455 coldwater tributaries are also susceptible to climate-induced warming, proper riparian shading
456 can maintain cooler temperatures and protect thermal refuges formed by tributary junctions with
457 the mainstem river (Boughton et. al 2012). Management that targets maintaining robust thermal
458 refuges at tributary junctions with the mainstem river will provide essential habitat for juvenile
459 salmonids seeking to exploit a heterogeneous thermal and prey landscape.

Draft

ACKNOWLEDGEMENTS

We thank Pete Raimondi and Dave Rundio for discussions regarding data analysis, and Mark Carr for helpful comments on this manuscript. We also thank the many individuals who helped with fieldwork, including Kacey Munson, Jeremy Notch, Kyle Swann, Jordan Sheffield, and the Karuk Tribe field crews. Funding for this research was provided by the U.S. Environmental Protection Agency Science to Achieve Results (STAR) program, NOAA Grant NA07NMF4540337, and NOAA Fisheries, Southwest Fisheries Science Center. Additional support was provided by Friends of Long Marine Lab Student Research Award, the Davis Scholarship Fund, the UC Santa Cruz Ecology and Evolutionary Biology Department, and the Myers Oceanographic and Marine Biology Trust.

Draft

LITERATURE CITED

- Achord, S., P. S. Levin, and R. W. Zabel. 2003. Density-dependent mortality in Pacific salmon: the ghost of impacts past? *Ecology Letters* 6:335–342.
- Armstrong, J., and S. Griffiths. 2001. Density-dependent refuge use among over-wintering wild Atlantic salmon juveniles. *Journal of Fish Biology* 58:1524–1530.
- Armstrong, J. B. and D. E. Schindler. 2013. Going with the flow: spatial distributions of juvenile coho salmon track an annually shifting mosaic of water temperature. *Ecosystems* 16(8):1429–41.
- Armstrong, J., D. Schindler, C. Ruff, G. T. Brooks, K. E. Bentley, and C. E. Torgersen. 2013. Diel horizontal migration in streams: juvenile fish exploit spatial heterogeneity in thermal and trophic resources. *Ecology* 94:2066–2075.
- Baird, O. E., and C. C. Krueger. 2003. Behavioral thermoregulation of brook and rainbow trout: comparison of summer habitat use in an Adirondack River, New York. *Transactions of the American Fisheries Society* 132:1194–1206.
- Bartholow, J. M. 2005. Recent water temperature trends in the Lower Klamath River, California. *North American Journal of Fisheries Management* 25:152–162.
- Baumgartner, D. and K. Rothhaupt. 2003. Predictive length–dry mass regressions for freshwater invertebrates in a pre-alpine lake littoral. *Hydrobiology* 88(5): 453–463.

- Belchik, M. 2003. Use of thermal refugial areas on the Klamath River by juvenile salmonids; Summer 1998. Yurok Tribal Fisheries Program Technical Report 8-FG-20-17510. Klamath, California, USA.
- Benke, A.C., A.D. Huryn, L.A. Smock, and J.B. Wallace. 1999. Length-mass relationships for freshwater macroinvertebrates in North America with particular reference to the southeastern United States. *Journal of the North American Benthological Society* 18:308–343.
- Bentley, K. T., D. E. Schindler, T. J. Cline, J. B. Armstrong, D. Macias, L. R. Ciepiela, and R. Hilborn. 2014. Predator avoidance during reproduction: diel movements by spawning sockeye salmon between stream and lake habitats. *Journal of Animal Ecology*. In press.
- Berman, C., and T. P. Quinn. 1991. Behavioural thermoregulation and homing by spring chinook salmon, *Oncorhynchus tshawytscha* (Walbaum), in the Yakima River. *Journal of Fish Biology* 39:301–312.
- Bevelhimer, M. S., and S. M. Adams. 1993. A bioenergetics analysis of diel vertical migration by kokanee salmon, *Oncorhynchus nerka*. *Canadian Journal of Fisheries and Aquatic Sciences* 50:2336–2349.
- Blumstein, D., and D. Pelletier. 2005. Yellow-bellied marmot hiding time is sensitive to variation in costs. *Canadian Journal of Zoology* 367:363–367.
- Boughton, D. A., C. Hatch, and E. Mora, Identifying distinct thermal components of a creek. *Water Resources Research*. 48(9):W09506.

- Breau, C., R. A. Cunjak, and G. Bremset. 2007. Age-specific aggregation of wild juvenile Atlantic salmon *Salmo salar* at cool water sources during high temperature events. *Journal of Fish Biology* 71:1179–1191.
- Brett, J., W. Clarke, J. Shelbourn, and P. Station. 1982. Experiments on thermal requirements for growth and food conversion efficiency of juvenile chinook salmon *Oncorhynchus tshawytscha*. Canadian Technical Report of Fisheries and Aquatic Sciences No. 1127. Nanaimo, British Columbia, Canada.
- Brewitt, K. S. and E. M. Danner. 2014. Spatio-temporal temperature variation influences juvenile steelhead (*Oncorhynchus mykiss*) use of thermal refuges. *Ecosphere*. 5(7):1-14.
- Burgherr, P., and E.I. Meyer. 1997. Regression analysis of linear body dimensions vs. dry mass in stream macroinvertebrates. *Archiv für Hydrobiologie* 139:101–112.
- Bridges, C. M. 2002. Tadpoles balance foraging and predator avoidance: Effects of predation, pond drying, and hunger. *Journal of Herpetology* 36:627–634.
- Ciancio, J. and M. Pacual. 2006. Energy density of freshwater Patagonian organisms. *Ecología Austral* 16:xxx-xxx.
- Christy, T. 2003. Calif. Dept. of Fish and Wildlife and Pacific States Marine Fisheries. Biogeographic Information and Observation System (BIOS). Map layer Cdfg_100k_2003_6.shp. Retrieved Nov 11, 2011 from <http://bios.dfg.ca.gov>

- Cummins, K. W., and J. C. Wuycheck. 1971. Caloric equivalents for investigations in ecological energetics. Internationale Vereinigung für Theoretische und Angewandte Limnologie, Mitteilung 1818:1–158.
- Davis, J., A. Pavlova, R. Thompson, and P. Sunnucks. 2013. Evolutionary refugia and ecological refuges: key concepts for conserving Australian arid zone freshwater biodiversity under climate change. *Global Change Biology* 19:1970–1984.
- Deas, M., S. K. Tanaka, and J. C. Vaughn. 2006. Klamath River thermal refugia study: Flow and temperature characterization. Watercourse Engineering, Inc., prepared for U.S. Bureau of Reclamation in cooperation with the Karuk Tribe and Yurok Tribe. Klamath Falls, Oregon.
- Dhondt, A. A., B. Kempenaers, and F. Adriaensen. 1992. Density-dependent clutch size caused by habitat heterogeneity. *Journal of Animal Ecology* 61:643–648.
- Driver, E.A., L.G. Sugden, and R.J. Kovach. 1974. Calorific, chemical and physical values of potential duck foods. *Freshwater Biology* 4(3):281–292.
- Driver, E.A. 1981. Calorific values of pond invertebrates eaten by ducks. *Freshwater Biology* 11(6):579–581.
- Ebersole, J., W. Liss, and C. Frissell. 2001. Relationship between stream temperature, thermal refugia and rainbow trout *Oncorhynchus mykiss* abundance in arid-land streams in the northwestern United States. *Ecology of Freshwater Fish* 10:1–10.

- Ebersole, J., W. J. Liss, and C. A. Frissell. 2003. Thermal heterogeneity, stream channel morphology, and salmonid abundance in northeastern Oregon streams. *Canadian Journal of Fisheries and Aquatic Sciences* 60:1266–1280.
- Everest, F., and D. Chapman. 1972. Habitat selection and spatial interaction by juvenile chinook salmon and steelhead trout in two Idaho streams. *Journal of the Fisheries Research Board of Canada* 29:91–100.
- Gelman, A., and D. B. Rubin. 1992. Inference from iterative simulation using multiple sequences. *Statistical Science* 7:457–472.
- Grant, J. W. A., S. Ó. Steingrímsson, E. R. Keeley, and R. A. Cunjak. 1998. Implications of territory size for the measurement and prediction of salmonid abundance in streams. *Canadian Journal of Fisheries and Aquatic Sciences* 55:181–190.
- Greene, C., and T. Beechie. 2004. Consequences of potential density-dependent mechanisms on recovery of ocean-type Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* 61:590–602.
- Gregory, J., and J. Griffith. 1996. Aggressive behaviour of underyearling rainbow trout in simulated winter concealment habitat. *Journal of Fish Biology* 49:237–245.
- Hanson, P., T. Johnson, J. Kitchell, and D.E. Schindler. 1997. Fish bioenergetics 3.0. University of Wisconsin Sea Grant Institute, Madison, Wisconsin, USA.
- Harvey, B. C. and R. J. Nakamoto. 1996. Effects of Steelhead Density on Growth of Coho

- Salmon in a Small Coastal California Stream. Transactions of the American Fisheries Society. 125:237-243.
- Hayes, J. W., N. F. Hughes, and L. H. Kelly. 2007. Process-based modelling of invertebrate drift transport, net energy intake and reach carrying capacity for drift-feeding salmonids. Ecological Modelling 207:171–188.
- Heady, W., and J. Moore. 2012. Tissue turnover and stable isotope clocks to quantify resource shifts in anadromous rainbow trout. Oecologia 172:21–34.
- Heady, W. N. 2012. Relationships among environment, movement, growth and survival of coastal rainbow trout (*Oncorhynchus mykiss*). Dissertation. University of California Santa Cruz, Santa Cruz, California, USA.
- Hodar, J. 1996. The use of regression equations for estimation of arthropod biomass in ecological studies. Acta Oecologia 17:421–433.
- Hohausova, E., G. Copp, and P. Jankovsky. 2003. Movement of fish between a river and its backwater: diel activity and relation to environmental gradients. Ecology of Freshwater Fish 12:107–117.
- Huntsman, B. M., and J. T. Petty. 2014. Density-dependent regulation of brook trout population dynamics along a core-periphery distribution gradient in a complex Appalachian watershed. PLoS ONE 9:e91673.

- Imre, I., J. W. A. Grant, and E. R. Keeley. 2004. The effect of food abundance on territory size and population density of juvenile steelhead trout (*Oncorhynchus mykiss*). *Oecologia* 138:371–8.
- Isaak, D. J., S. Wollrab, D. Horan, and G. Chandler. 2012. Climate change effects on stream and river temperatures across the northwest U.S. from 1980–2009 and implications for salmonid fishes. *Climatic Change* 113:499–524.
- Jobling, M. 1994. *Fish Bioenergetics*. Chapman & Hill, London.
- Kaeding, L. 1996. Summer use of coolwater tributaries of a geothermally heated stream by rainbow and brown trout, *Oncorhynchus mykiss* and *Salmo trutta*. *American Midland Naturalist* 135:283–292.
- Kawabata, K. and J. Urabe 1998. Length–weight relationships of eight freshwater planktonic crustacean species in Japan. *Freshwater Biology* 39(2): 199–205.
- Keefer, M. L., C. A. Peery, and B. High. 2009. Behavioral thermoregulation and associated mortality trade-offs in migrating adult steelhead (*Oncorhynchus mykiss*): variability among sympatric populations. *Canadian Journal of Fisheries and Aquatic Sciences* 66:1734–1747.
- Keeley, E., and J. McPhail. 1998. Food abundance, intruder pressure, and body size as determinants of territory size in juvenile steelhead trout (*Oncorhynchus mykiss*). *Behaviour* 135:65–82.
- Keeley, E. R. 2001. Demographic responses to food and space competition by juvenile steelhead trout. *Ecology* 82:1247.

- Kiffney, P., C. Greene, J. Hall, and J. Davies. 2006. Tributary streams create spatial discontinuities in habitat, biological productivity, and diversity in mainstem rivers. *Canadian Journal of Fisheries and Aquatic Sciences* 63:2518–2530.
- Knowles, N. and D. R. Cayan. 2002. Potential effects of global warming on the Sacramento/San Joaquin watershed and the San Francisco estuary. *Geophysical Research Letters*. 29(18):38-1 - 38-4.
- Lawler, J. J., D. White, R. P. Neilson, and A. R. Blaustein. 2006. Predicting climate-induced range shifts: model differences and model reliability. *Global Change Biology* 12:1568–1584.
- Loarie, S. R., B. E. Carter, K. Hayhoe, S. McMahon, R. Moe, C. A. Knight, and D. D. Ackerly. 2008. Climate change and the future of California's endemic flora. *PLoS ONE* 3:e2502.
- Loose, C. J., and P. Dawidowicz. 2014. Trade-offs in diel vertical migration by zooplankton: the costs of predator avoidance. *Ecology* 75:2255–2263.
- Mather, M. E., D. L. Parrish, C. A. Campbell, J. R. McMenemy, and J. M. Smith. 2008. Summer temperature variation and implications for juvenile Atlantic salmon. *Hydrobiologia* 603:183–196.
- Matthews, K. R., N. H. Berg, D. L. Azuma, and T. R. Lambert. 1994. Cool water formation and trout habitat use in a deep pool in the Sierra Nevada, California. *Transactions of the American Fisheries Society*. 123:549–564.

- McCullough, D. A. 1999. A review and synthesis of effects of alterations to the water temperature regime on freshwater life stages of salmonids, with special reference to Chinook Salmon. U.S. Environmental Protection Agency. Report 910-R-99-010, Seattle.
- McCutchan, J., W. Lewis, C. Kendall, and C. McGrath. 2003. Variation in trophic shift for stable isotope ratios of carbon, nitrogen, and sulfur. *Oikos* 2:378–390.
- Miserendino, M.L. 2001. Length-mass relationships for macroinvertebrates in freshwater environments of Patagonia (Argentina). *Ecología Austral* 11:3-8.
- Myrick, C. A., and J. J. Cech. 2000. Temperature influences on California rainbow trout physiological performance. *Fish Physiology and Biochemistry* 22:245–254.
- Myrick, C., and J. Cech. 2005. Effects of temperature on the growth, food consumption, and thermal tolerance of age-0 nimbus-strain steelhead. *North American Journal of Aquaculture* 67:324–330.
- Mac Nally, R., A. F. Bennett, J. R. Thomson, J. Q. Radford, G. Unmack, G. Horrocks, and P. A. Vesk. 2009. Collapse of an avifauna: climate change appears to exacerbate habitat loss and degradation. *Diversity and Distributions* 15:720–730.
- National Research Council. 2004. Endangered and threatened fishes in the Klamath River basin. National Academies Press, Washington DC, USA.
- National Research Council. 2008. Hydrology, ecology, and fishes of the Klamath River basin. National Academies Press, Washington DC, USA.

- Nehlsen, W., J. E. Williams, and J. A. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. *Fisheries* 16:4–21.
- Pärn, H., T. H. Ringsby, H. Jensen, and B.-E. Sæther. 2011. Spatial heterogeneity in the effects of climate and density-dependence on dispersal in a house sparrow metapopulation. *Proceedings of the Royal Society B* 279:144–152.
- Payne, J. T., A. W. Wood, A. F. Hamlet, R. N. Palmer, and D. P. Lettenmaier. 2004. Mitigating the effects of climate change on the water resources of the Columbia River Basin. *Climatic Change*. 62(1):233–56.
- Pearson, R. G., and T. P. Dawson. 2003. Predicting the impacts of climate change on the distribution of species: Are bioclimate envelope models useful? *Global Ecology and Biogeography* 12:361–371.
- Pepino, M., K. Goyer, and P. Magnan. 2015. Heat transfer in fish: are short excursions between habitats a thermoregulatory behaviour to exploit resources in an unfavourable thermal environment? *Journal of Experimental Biology* 218: 3461-3467.
- Petty, J. T., J. L. Hansbarger, B. M. Huntsman, and P. M. Mazik. 2012. Brook trout movement in response to temperature, flow, and thermal refugia within a complex Appalachian riverscape. *Transactions of the American Fisheries Society* 141:1060–1073.
- Quinn T. P. 2005. The behavior and ecology of Pacific salmon and trout. American Fisheries Society. Bethesda, Maryland, in association with University of Washington Press. Seattle, Washington.

- Rangeley, R., and D. Kramer. 1998. Density-dependent antipredator tactics and habitat selection in juvenile pollock. *Ecology* 79:943–952.
- Richter, A., and S. A. Kolmes. 2005. Maximum temperature limits for chinook, Coho, and Chum salmon, and steelhead trout in the Pacific Northwest. *Reviews in Fisheries Science* 13:23–49.
- Rogers, L. E., R. L. Buschbom, and C. R. Watson. 1977. Length-weight relationships of shrub-steppe invertebrates. *Annals of the Entomological Society of America* 70:51–53.
- Rosenfeld, J. S., T. Leiter, G. Lindner, and L. Rothman. 2005. Food abundance and fish density alters habitat selection, growth, and habitat suitability curves for juvenile coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences* 62:1691–1701.
- Ruesch, A. S., C. E. Torgersen, J. J. Lawler, J. D. Olden, E. E. Peterson, C. J. Volk, and D. J. Lawrence. 2012. Projected climate-induced habitat loss for salmonids in the John Day River network, Oregon, U.S.A. *Conservation Biology* 26:873–82.
- Sabo, J. L., J. L. Bastow, and M. E. Power. 2002. Length-mass relationships for adult aquatic and terrestrial invertebrates in a California watershed. *Journal of the North American Benthological Society* 21:336–343.
- Scheuerell, M. D., and D. E. Schindler. 2003. Diel vertical migration by juvenile sockeye salmon: empirical evidence for the antipredation window. *Ecology* 84:1713–1720.

- Shearer, K., J. Hayes, and J. Stark. 2002. Temporal and spatial quantification of aquatic invertebrate drift in the Maruia River, South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36:529–536.
- Sims, D. W., V. J. Wearmouth, E. J. Southall, J. M. Hill, P. Moore, K. Rawlinson, N. Hutchinson, G. C. Budd, D. Righton, J. D. Metcalfe, J. P. Nash, and D. Morritt. 2006. Hunt warm, rest cool: bioenergetic strategy underlying diel vertical migration of a benthic shark. *Journal of Animal Ecology* 75:176–190.
- Stock, B. C., and B. X. Semmens. 2013. MixSIAR GUI user manual, version 1.0. Accessible online at: <http://conserver.iugo-cafe.org/user/brice.semmens/MixSIAR>
- Sutton, R., M. Deas, S. Tanaka, T. Soto, and A. Corum. 2007. Salmonid observations at a Klamath River thermal refuge under various hydrological and meteorological conditions. *River Research and Applications* 23:775–785.
- Thomas, C. D., A. M. A. Franco, and J. K. Hill. 2006. Range retractions and extinction in the face of climate warming. *Trends in Ecology and Evolution* 21:415–6.
- Torgersen, C., J. Ebersole, and D. Keenan. 2012. Primer for identifying cold-water refuges to protect and restore thermal diversity in riverine landscapes. EPA scientific guidance handbook 910-C-12-001. U.S. Environmental Protection Agency, Seattle, Washington, USA.
- Travis, J. M. J. 2003. Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proceedings of the Royal Society B* 270:467–73.

Van Kirk, R. W. and S. W. Naman. 2009. Relative Effects of Climate and Water Use on Base-Flow Trends in the Lower Klamath Basin. *Journal of the American Water Resources Association*. 44(4): 1035-1052.

Vitousek, P. M. 1997. Human domination of Earth's ecosystems. *Science* 277:494–499.

Willis, K. J., and S. A. Bhagwat. 2009. Biodiversity and climate change. *Science* 326:806–807.

Wurtsbaugh, W., and D. Neverman. 1988. Post-feeding thermotaxis and daily vertical migration in a larval fish. *Nature* 333:846–848.

Draft

Table I. Summary of field methods and modeled variables.

Variable	Method	Year and Study Site	Purpose of data
Water temperature (°C)	Collected using in-stream temperature loggers in the tributary and mainstem river.	2012 (S1&2) ^a	To quantify the difference in water temperature between the coldwater tributary and mainstem river; recorded temperatures represent the minimum and maximum temperatures available to fish.
Energy delivery rate (Joules m ⁻² s ⁻¹) of invertebrate prey	Modeled using data from invertebrate drift sampling in the tributary and mainstem river (see eq. 1).	2010 (S1-4) 2011 (S1)	To quantify energy delivery rates in the different thermal habitats at refuges (tributary and adjacent mainstem river), in order to assess potential sources of in-refuge food limitation.
Fish density (fish m ⁻²)	Collected using snorkel surveys in the tributary, mixing zone, and mainstem river.	2012 (S1)	To quantify fish density in the different thermal habitats at refuges, in order to assess potential sources of in-refuge food limitation.
Isotope-inferred fish diet	Mixing model analysis using data from fish tissue samples and invertebrate prey sources (collected via benthic sampling in tributary and mainstem at thermal refuges).	2012 (S1&2)	To quantify the proportion of mainstem vs. tributary invertebrate prey contributing to the diet of fish using thermal refuges.
Fish body temperatures (°C)	Collected using continuous recordings of steelhead body temperatures from temperature-sensitive radio tags.	2010-2012 (S1&2)	To quantify thermal habitat use for juvenile steelhead using thermal refuges, so as to assess whether fish were foraging primarily in the same areas that they were holding.
Diet-inferred foraging temperatures (°C)	Modeled using isotope-inferred diet and water temperature data (see eq. 4).	2010-2012 (S1&2)	To model the water temperatures where fish were foraging, so as to assess whether fish using thermal refuges were foraging primarily in the same areas that they are holding.

^a Sites indicated as follows: S1 (Beaver Creek), S2 (Grider Creek), S3 (Fort Goff Creek), and S4 (Thompson Creek)

FIGURE CAPTIONS

Figure 1. Study sites located at tributary confluences on the Klamath River. (1 = Beaver Creek; 2 = Grider Creek; 3 = Fort Goff Creek; 4 = Thompson Creek). Inset shows location of study area within the watershed. Map layer from the California Department of Fish and Game, Pacific States Marine Fisheries Commission (Christy 2003).

Figure 2. (A) Map of snorkel survey plots (open squares) and invertebrate drift sampling locations (filled triangles) at Beaver Creek confluence. (B) Steelhead and (C) Chinook salmon densities (m^{-2}) in plots within each thermal environment at Beaver Creek refuge in August 2012. Boxplots are coded by thermal environment (tributary (blue) = A, B, C; mixing zone (purple) = D, E, F, G; mainstem (red) = H, I). Labels above boxplots indicate mean water temperatures in each plot across the sampling period.

Figure 3. Isotope signatures of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ for juvenile salmonids and mainstem (red) and tributary (blue) invertebrate prey sources (mean $\pm$ SD). (A) Beaver Creek steelhead ($n = 53$) indicated by solid gray points and Chinook ($n = 25$) indicated by open squares. (B) Grider Creek steelhead ($n = 32$) indicated by solid gray points. Isotope signatures shown are unadjusted, but it worth noting that trophic enrichment is approximately 3.2 for $\delta^{15}\text{N}$ and 1.9 for $\delta^{13}\text{C}$.

Figure 4. Posterior estimates from MixSIAR of the contribution of mainstem prey sources to fish diet plotted against fish body size (fork length). Mean diet estimates and 90% credible interval (CI) for individual steelhead at (A) Beaver Creek, (B) Grider Creek, and (C) Chinook at Beaver

Creek. Symbols are coded by the thermal environment in which individuals were caught (tributary = blue squares; mixing zone = purple triangles; mainstem = red circles).

Figure 5. Posterior estimates of the mean contribution of mainstem prey sources to fish diet versus the proportion of time spent in mainstem water for individual juvenile steelhead radio tagged at Beaver Creek, for which there were >7 days of temperature data ($n = 9$).

Figure 6. Distribution of fish body temperatures (blue) and diet-inferred foraging temperatures (green) for steelhead sampled at (A) Beaver Creek and (B) Grider Creek in 2012. Thermal distributions are derived from temperature-sensitive radio tag data, and diet-inferred foraging temperature distributions are derived from the mass balance equation (Eq. 4).

FIGURES

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

APPENDIX FIGURE CAPTIONS

Figure A1. Mainstem (red) and tributary (blue) water temperatures at (A) Beaver Creek and (B) Grider Creek in 2012. Black dashed vertical lines indicate fish sampling events, and gray shaded rectangles indicate the two week period prior to fish sampling when benthic invertebrates were collected.

Figure A2. Invertebrate energy delivery rates (mean $\pm$ SD) at all sampled sites in 2010 and 2011 (mainstem = red; tributary = blue).

Figure A3. Relative abundance of invertebrate species (identified to family) in ten-minute drift samples (mainstem = red; tributary = blue). Taxa included represent most abundant represented in drift (mean abundance across samples >40 per family).

Figure A4. Relative abundance of invertebrate species (identified to family) in juvenile steelhead diets in 2010 and 2011.

APPENDIX FIGURES

Fig. A1

Figure A2.

Figure A3.

Figure A4.

APPENDIX TABLES

Table A1. Sample sizes for invertebrate drift samples in 2010. Samples were collected monthly at each site; thermal environment indicates samples taken from either the mainstem river (MS) or tributary (T).

Site	Sample Month	Thermal Environment	N
Beaver	August	MS	4
Beaver	August	T	3
Beaver	July	MS	6
Beaver	July	T	6
Beaver	June	MS	6
Beaver	June	T	6
Fort Goff	August	MS	4
Fort Goff	August	T	3
Fort Goff	July	MS	4
Fort Goff	July	T	4
Grider	August	MS	4
Grider	August	T	4
Grider	July	MS	4
Grider	July	T	4
Thompson	August	MS	4
Thompson	August	T	4
Thompson	July	MS	2
Thompson	July	T	4

Table A2. Sample sizes for invertebrate drift samples in 2011. Samples were collected weekly (Week indicates sampling date), at both dawn and dusk, but only the dawn samples were analyzed due to processing time constraints.

Site	Week	Thermal Environment	N
Beaver	7-05	MS	2
Beaver	7-05	T	2
Beaver	7-12	T	2
Beaver	7-19	MS	1
Beaver	7-19	T	2
Beaver	7-25	MS	2
Beaver	7-25	T	2
Beaver	8-01	MS	2
Beaver	8-01	T	2
Beaver	8-05	MS	2
Beaver	8-05	T	2
Beaver	8-16	MS	2
Beaver	8-16	T	2
Beaver	8-24	MS	2
Beaver	8-24	T	2
Beaver	9-01	MS	2
Beaver	9-01	T	2

Table A3. Sample sizes for diet samples of juvenile steelhead taken in 2010 and 2011.

Year and Site	Date	N
2010		
Beaver	29-Jun	2
Beaver	27-Jul	10
Fort Goff	4-Aug	10
2011		
Fort Goff	14-Oct	12

Draft

Table A4. Invertebrate families selected from benthic samples and used to determine isotopic signatures of juvenile salmonid prey sources.

Mainstem samples	Tributary samples
Chironomidae	Chironomidae
Baetidae	Baetidae
Hydrophychidae	Hydrophychidae
Simuliidae	Simuliidae
Brachycentridae	Glossosomatidae
	Perlidae

Draft