**Fen restoration: defining a reference ecosystem using
paleoecological stratigraphy and present-day inventories**

Journal:	<i>Botany</i>
Manuscript ID	cjb-2016-0281.R1
Manuscript Type:	Article
Date Submitted by the Author:	06-Feb-2017
Complete List of Authors:	Bérubé, Vicky; Université Laval, Plant Sciences Rochefort, Line; Université Laval, Plant Sciences Lavoie, Claude; Université Laval, Centre de recherche en aménagement et développement
Please Select from this Special Issues list if applicable:	N/A
Keyword:	rehabilitation, minerotrophic peatland, plant assemblages, plant dynamic, indicator value index(IndVal)

SCHOLARONE™
Manuscripts

Fen restoration: defining a reference ecosystem using paleoecological stratigraphy and present-day inventories

Vicky Bérubé¹, Line Rochefort¹ and Claude Lavoie²

¹Département de phytologie and Centre d'études nordiques, Université Laval, 2425, rue de l'Agriculture, Québec, QC G1V 0A6, Canada

²École supérieure d'aménagement du territoire et de développement régional and Centre d'études nordiques, Université Laval, 2325, rue des Bibliothèques, Québec, QC G1V 0A6, Canada;

e-mail addresses: vicky.berube.1@ulaval.ca, line.rochefort@fsaa.ulaval.ca, claudio.lavoie@esad.ulaval.ca

Corresponding author :

Vicky Bérubé

2425, rue de l'Agriculture,

Université Laval,

Québec (Québec)

G1V 0A6, Canada

Phone: 1-418-656-2131 ext. 3595

Fax: 1-418-656-7856

vicky.berube.1@ulaval.ca

Abstract

Choosing past and present-day indicators could strengthen the reference ecosystem used for ecological restoration projects. Based on the paleoecological analysis of four peat cores and the characterization of 13 contemporary natural sites, the reference ecosystem for minerotrophic peatlands in southeastern Canada is composed of two types of plant assemblages described as tall-sedge and *Sphagnum-Thuja*/brown moss broad categories. In paleoecological peat profiles, tall-sedge communities were found at the transition between aquatic and terrestrial and were associated with high graminoid production, riverine peatlands and elevated water table in the present-day analyses. *Sphagnum-Thuja* communities resemble present-day vegetation found in natural basin type peatlands. Except for *Sphagnum warnstorffii*, these communities, with high taxonomical diversity, contain more generalist species like *Rhododendron groenlandicum*, *Thuja occidentalis*, *Linnaea borealis* and *Maianthemum trifolium* from boreal peatland vegetation. They grow in dryer or shady habitats like hummocks or forest understory. The importance of brown mosses was revealed by paleoecological analysis. Overall, findings from both approaches are complementary: paleoecological stratigraphy informs us about past ecosystem dynamics, while present-day inventories allow us to define current plant communities and their major environmental characteristics. The range of variability of vegetation and environmental variables found in these studies are essential tools for fen restoration projects.

Keywords: rehabilitation, minerotrophic peatland, plant assemblages, plant dynamic, indicator value index(IndVal).

Résumé

Lors d'un projet de restauration écologique, la description de l'écosystème de référence est renforcée par l'utilisation d'indicateurs combinant les études temporelles et spatiales. Se basant sur une étude paléoécologique et la caractérisation de treize sites naturels, la référence pour une tourbière minérotrophe modérément riche située dans le sud-est du Canada se compose de deux types d'assemblages d'espèces végétales : l'assemblage des Grandes-cypéracées et l'assemblage général de *Sphagum-Thuja*/mousse brune. Dans l'étude paléoécologique, la communauté des Grande-cypéracées est observée à la transition entre les écosystèmes de marais et de tourbières. Dans l'étude des sites naturels, cette communauté est associée à une production de biomasse élevée, un niveau d'eau élevé et aux fens de type riparien. L'assemblage *Sphagum-Thuja* est semblable à ceux inventoriés dans les fens de type basin. En faisant l'exception du *Sphagnum warnstofii*, cet assemblage, très diversifié, inclut surtout des espèces généralistes provenant des tourbières boréales telles le *Rhododendron groenlandicum*, le *Thuja occidentalis*, le *Linnaea borealis* et le *Maianthemum trifolium*. L'importance des mousses brunes est révélée dans l'analyse paléoécologique. En somme, les deux approches sont complémentaires : la paléoécologie nous informe sur la dynamique de l'écosystème tandis que les inventaires des sites naturels nous a permis de définir les assemblages et leurs associations aux conditions environnementales. Les étendues de variabilité définies dans ce travail sont un outil à la décision essentiel dans un projet de restauration écologique de fen.

Mots-clés : réhabilitation, tourbière minérotrophe, assemblage de végétaux, dynamique des communautés, index valeur indicatrice (IndVal)

Introduction

Reference ecosystems act as a standard of comparison and evaluation for any restoration project (Society for Ecological Restoration International Science and Policy Working Group 2004; Clewell and Aronson 2013; Shackelford et al. 2013). Selecting references among other similar ecosystems located in the same region is a usual starting point for establishing standards (White and Walker 1997; SERI 2004; Hallet et al. 2013). However, the absence or limited knowledge of pre-disturbance states, ecosystem dynamics and cultural services may be responsible for an inappropriate reference selection (Aronson 1995; Birks 1996; Higgs 1997; Jackson and Hobbs 2009; Balaguer et al. 2014). An in-depth knowledge of ecosystem variability with temporal and spatial perspectives is thus recommended for making better choices. Moreover, multiple sources of information improve the quality and accuracy of the data that can be used to build a reliable picture of a reference ecosystem (White and Walker 1997; SERI 2004), and may help to better define realistic restoration objectives (Ehrenfeld 2000; SERI 2004; Hallet et al. 2013; Shackelford et al. 2013).

Data useful for defining a reference ecosystem can be collected using historical records, by studying similar undisturbed ecosystems, or both (Birks 1996; White and Walker 1997; Balaguer et al. 2014). Historical tools include paleoecological studies, past plant or faunal surveys, and aerial and *in situ* photographs. One of the advantages of using historical data is that they provide insights into ecosystem dynamics over time (Jackson and Hobbs 2009; Balaguer et al. 2014). They also help to choose the developmental stage at which the ecosystem should be restored. Paleoecological studies are widely used in peatlands to assess past vegetation patterns and succession dynamics (e.g. Glaser and Janssen 1986; Kurhy et al. 1993; Bauer et al. 2003) or climate change (e.g. Halsey et al. 1998; Yu 2006). They proved to be useful to provide a better understanding of the ecosystem functioning over the long term, which can help to evaluate the time frame over which restoration objectives could reasonably be achieved (Lavoie et al. 2001; Blundell and

Holden 2015). The response of ecosystems to past disturbance events can also inspire restoration engineers by providing clues on natural recovery processes (Benscoter and Vitt 2008). On the other hand, present-day studies on the existing composition and distribution of species are essential to complete the reference picture. They represent a timescale snapshot but can nevertheless give a precise overview of the spatial distribution of species along a wide range of biotic and abiotic conditions.

In North America, most peatlands that have actively been restored are cutover bogs (ombrotrophic peatlands; Poulin et al. 2013; González et al. 2014). The peat was extracted for horticultural purposes. The most widely used extraction method (vacuum) consists of draining the peatland, removing the living vegetation, and harrowing and vacuuming a thin layer of dried peat. Peat extraction activities cease when peat quality does not correspond to industrial standards (too decomposed, too woody, etc.) and, until recently, at least 50 cm of *Sphagnum* peat is left on site to ensure enough material is left for restoration. However, in some cases, the peat industry produces specific growing substrates with the deeper, older peat layers of peatlands. In such cases, ancient minerotrophic peat layers composed of the remains of sedges and brown moss become exposed to the surface (Graf et al. 2008). Since water and peat chemical properties are then similar to fen (Wind-Mulher and Vitt 2000), the rehabilitation of the peatland to a minerotrophic developmental stage could be considered if an ecological restoration is sought. The challenge in those conditions is to picture a reference ecosystem from a series of developmental stages that the ecosystem has experienced in the past and could experience in the future.

The goal of this study was to build a reference ecosystem in the context of fen restoration in southeastern Canada using paleoecological studies and current regional plant surveys. Specifically, we wanted (i) to describe the historical trajectory of a minerotrophic peatland via its developmental states and its associated plant assemblages; (ii) to assess if modern plant assemblages are representative of the past; (iii) to evaluate the importance of bryophytes for fen development; (iv) to study natural disturbance regimes and

the resilience process, and finally (v) to find which state the disturbed peatland has regressed to after the cessation of peat harvesting activities. If the present plant assemblage is representative of the past, the modern plant surveys should complete the reference by determining the variability of species composition and environmental condition of moderately rich fens. Our overall intent with this study was to demonstrate the usefulness of both approaches to establish precise restoration objectives.

Materials and methods

Study sites

All the peatlands investigated in this study were located near the south shore of the St. Lawrence River, in the Bas-Saint-Laurent region, Québec, Canada (Fig. 1). The area was covered by a glacier until 14 000 years BP (before present), and then by the Goldthwait Sea until 10 000 years BP (Locat 1977), leaving clay, sand and gravel as surficial marine deposits over the St. Lawrence River Lowlands (Fulton 1995). The paleoecological study was conducted in the Bic-Saint-Fabien (BSF) peatland (48°19'22" N, 68°49'52" W). It is part of a peatland complex that developed in a narrow valley (Fig. 2). BSF, which is now included in the Bic National Park, is bordered by steep calcareous hills (340 m a.s.l.) to the north and by agricultural fields to the south. A large pond (about 40 ha) was located in the central part of the peatland but was drained in 1954. The section of the peatland east of the pond was formerly a *Sphagnum* bog (Scoggan 1942) with a well-defined lagg on its margins. This eastern section had largely been used for peat extraction. The section west of the pond is now a mosaic of peat extraction and few natural (no major disturbance) sites. Peat extraction began in 1946 and ended in the 1990s. The 15 ha area that was targeted for a restoration project has a peat substrate with the following minerotrophic characteristics, as assessed in summer 2011 (L. Rochefort, unpublished data; n = 111): the water pH was 6.0 ± 0.4 (mean $\pm$ SD) and the electroconductivity was $229 \pm 141 \mu\text{S cm}^{-1}$. The central section was spontaneously revegetated, mostly

with *Typha latifolia*, *Scirpus cyperinus*, *S. atrocinctus* and *Salix* species, essentially along former drainage ditches. Bare peat covered the last four peat fields of this eastern section.

The northern and eastern sections of BSF bordering the extraction zone still support a moderately rich fen plant community (*sensu* Chee and Vitt 1989 and Locky et al. 2005) dominated by *Thuja occidentalis* (Gauthier and Grandtner 1975). In these sections, the water pH was 6.5 ± 0.8 (mean $\pm$ SD) and the electroconductivity $237 \pm 136 \mu\text{S cm}^{-1}$ (L. Rochefort, unpublished data; $n = 41$).

Paleoecological analyses

For this study, only the eastern section of the BSF peatland complex has been considered, excluding the former lakebed. Efforts for an ecological restoration project were concentrated in this section (Fig. 2). The condition of the disturbed sector was heterogeneous if we consider the time of abandonment, elevation and spontaneous vegetation cover. To get a good representation of the site, we divided the area into three more homogeneous subsectors for sampling. One peat core was extracted from each of the three subsectors (R for restored section: R1, R2 and R3). The peat cores were respectively about 700, 950 and 1200 m from the location of the drained pond (Fig. 2). A fourth core (N for natural section) was sampled at a 1300 m distance but in the part of the peatland with no peat extraction.

A side-wall peat corer (Jowsey 1966) was used to extract the peat cores, from the soil surface to the top of the surficial marine deposit. Each core was cut every 5 cm into subsamples, and each subsample was washed through a series of sieves (2, 1 and 0.5 mm meshes) and dried. Fossil pieces were sorted and identified to the finest taxonomical level (family, genus or species when possible). Nomenclature followed the Integrated Taxonomic Information System (2015) for vascular plants and animals, Faubert (2007) for bryophytes, and LoBuglio (1999) for fungi. Pieces were counted under a low power (50 $\times$) microscope. When the fossil pieces were too numerous, a subsample of 0.5 g of peat was extracted, the pieces counted,

and the total number of pieces was estimated using the total dry weight of the sample. All the results were adjusted to a volume 100 cm³ of peat. Macrofossil zones were defined visually according to changes in the relative abundance of plant remains. Samples at the limits of the zones and at the surface of the peat (for R1, R2 and R3) were dated using accelerator mass spectrometry (AMS) radiocarbon dating. Samples were prepared by the Radiochronology lab at Université Laval, QC, and analyzed by Keck Carbon Cycle AMS Facility (University of California, Irvine) Radiocarbon dates were calibrated with CALIB REV 7.0.1 software (Stuiver et al. 2005).

Natural site surveys

A total of 13 fens were found in the vicinity of the BSF peatland (from a 5–30 km distance), with the help of a local bryologist and by looking at satellite maps (Fig. 1). The final selection was based on the presence of indicator plant species of moderately rich fens and water pH above 5.5 (Chee and Vitt 1989). A survey of modern vegetation communities and associated environmental conditions was made during summer 2012. Six of the 13 peatlands were subdivided into two homogeneous units on the basis of their tree cover (with or without) or because they represent different sectors within a peatland complex. The number of quadrats depended on the area of the unit. Between 10 and 12 circular quadrats of 70 cm diameter were positioned equidistant along a linear transect passing throughout the longest axis of the fen (unit), for a total of 200 quadrats. The cover occupied by each species present within the quadrats was visually estimated with the accuracy of $\pm 5\%$ (for more details, see Rochefort et al. 2013). Environmental variables were measured for each peatland unit. The sampling was located in the central part of the peatland unit. The species richness is defined as the number of species in each quadrat and the diversity was also calculated for each quadrat with Shannon diversity index (Whittaker 1972). All variables are described in Table 1 along with analytical methods.

Data processing and statistical analyses

The cluster method, specifically K-means partitioning, and the indicator value index (IndVal) method (Borcard et al. 2011) were used to define modern plant communities and to identify their associated indicator species. K-means are based on species similarity found in each quadrat. To achieve this goal, a truncated species dataset was used for all statistical analyses to avoid biases potentially caused by rare species (Legendre and Gallagher 2001; Borcard et al. 2011); only species with a frequency (occurrence among all the quadrats from all peatlands) and/or a mean cover (among the quadrats with the species) > 5% were kept. Hellinger's transformation was conducted to take into account the importance of double zeros and common species (Legendre and Gallagher 2001). Rank transformations on all numerical environmental variables, except Shannon diversity index, were performed to achieve normality.

The Calinski-Harabasz criterion (Milligan 1996) was used to determine the adequate number of clusters (plant assemblages). To facilitate the categorization of assemblages, indicator species in each group were identified with the indicator value index (IndVal) developed by Dufrêne and Legendre (1997). This index computes the product of specificity (abundance of a species in all the quadrats of a cluster) and fidelity (presence of a species in a majority of quadrats in a cluster) for every species in each cluster previously delineated with k-means partitioning. IndVal values range from 0 to 1; a value close to 1 for a particular species indicates that the species is present exclusively in this cluster and has a high occurrence. Only species with IndVal values > 0.25, and an associated $P < 0.001$, were used to identify plant communities.

A Redundancy Analysis (RDA) was performed to understand the influence of environmental variables on plant communities. A forward selection was applied to reduce type I error and select the significant environmental variables. Cumulative r^2 adjusted from this selection must be lower than the r^2 adjusted from the RDA model previously ran with all the variables to respect the two stopping criteria (Blanchet et al. 2008). The significant environmental variables issued from the forward selection were used to perform a

second RDA and improve model variation. To view the position of the k-means clusters, equiprobability ellipses were created, including 95% of the sites from the same plant community. Finally, the RDA model was submitted to a permutation test with 9999 randomized runs to assess its significance (Legendre and Legendre 2012). All statistical analyses were conducted using R software version 2.14.1 (R Development Core Team 2011), with the function RDA, Ordistep and CascadeKM from the vegan package (Oksanen et al. 2011), and IndVal from the labdsv (Roberts 2013) and ellipse package (Murdoch and Chow 2007).

Results

Paleoecological analyses

Four major vegetation zones were identified in N and R3 while only two were observed in R1 and R2. Limnic deposit known as gyttja or coprogenous earth (Groupe de travail sur la classification des sols 2002; Meier-Uhlherr et al. 2015) started to accumulate over a sand and clay marine deposit between 8037 years BP at R1 and 7090 years BP at N (Fig. 3; Table 2; Supplementary file). The closer a peat core was to the pond of BSF peatland, the later peat accumulation was initiated. Rates of peat accumulation for R3 and N varied between 0.31 to 1.12 mm per year depending on the core sampling location and vegetation assemblage. Peat composed of brown mosses (zones II and IV) accumulated faster than peat composed only of wood remains (zone III).

At least three major fire events occurred. The peatland burned about 4870 years BP and 2700 years BP at the R3 location. The first fire coincided with the initiation of peat accumulation at this location, and the second with the end of zone II and the beginning of zone III. Around 1930, a fire burned part of the peatland (Scoggan 1942) and left charcoal pieces near the surface of N core. High abundance of *Campylium stellatum* and *Scorpidium cossonii* remains was noticed after this fire. *Thuja occidentalis* macrofossils also appeared after the 1930 fire.

The four peat cores followed a similar trajectory concerning the evolution of vegetation assemblages (Fig. 3 and supplementary file). Consequently, descriptions of each homogeneous zone identified through time are grouped together to lighten the text and avoid repetition.

Zone I (R1, R2, R3, N)

Gyttja sediments attested to the terrestrialization (in-filling) origin of the peatland (Łachacz et al. 2009). The jelly-like sediments included plant species usually found in a shallow water habitat, like *Najas flexilis*, *Nymphaea* spp., *Potamogeton* spp., *Sparganium* spp. and the bryozoan *Cristatella mucedo*. An assemblage composed of emergent species characterized the end of this zone. Marsh species (*Bidens cernua*, *Dulichium arundinaceum* and *Sagittaria latifolia*) were sporadically found. The abundance of *Carex* cf. *diandra*, *Carex rostrata* and *Schoenoplectus* s.l. *acutus* suggested a transition from a shallow lake ecosystem to a fen.

Zone II (R1, R2, R3, N)

Bryophytes were the main components of zone II. High abundance of *Paludella squarrosa*, *Calliergon* spp. and other brown moss fragments from Amblystegiaceae and Calliergonaceae families were recovered. The peat cores closer to the ancient pond (R1 and R2) also included *Sphagnum* stems from Acutifolia section species. At the beginning of zone II, the abundance of remains of mat-forming species (*Menyanthes trifoliata*, *Comarum palustre*, *Bidens cernua*) decreased. Cyperaceae species like *Carex diandra*, *C. rostrata* and *Rhynchospora alba*, along with other unidentified *Carex* species with planoconvex and trigonous shape seeds, were well represented for the entire zone.

An abrupt change in plant assemblages was detected in this moss-rich zone of R3, and was characterized by (1) a decrease in the number of *Calliergon* sp., *Ptychostomum* s.l. *pseudotriquetrum* and *Drepanocladus* sp. s.l. remains, followed by (2) the disappearance of *Paludella squarrosa*, and then by

(3) the establishment of other brown moss species (Amblystegiaceae spp.) dominated by *Campylium stellatum* and *Pseudocalliergon trifarium*.

Zone III (R3, N)

A major change in the vegetation composition of the peatland occurred about 2665 years BP at R3 and 4182 years BP at N, when bryophytes suddenly disappeared. The highly decomposed peat was mainly composed of small woody debris and no seeds or discernable plant parts were noticed, except sclerotia of *Cenococcum geophilum*.

Zone IV (R3, N)

A rapid increase in the abundance of bryophytes characterizes the beginning of zone IV. *Sphagnum* spp. were especially abundant at R3, which was also the case for *Paludella squarrosa* at the upper end of the profile. N was dominated by brown mosses (*Campylium stellatum*, *Calliergon giganteum* and *Straminergon cf. stramineum*). The vascular plant assemblages included *Rhynchospora alba*, *Trichophorum alpinum*, *Eleocharis cf. tenuis*, *Triglochin maritima* and *Andromeda polifolia* var. *latifolia*.

Present-day fen surveys

The 19 sampled fens (and units) had a peat thickness of 345 ± 169 cm (mean $\pm$ SD), water table depth 10 ± 13 cm below the soil surface, pH 6.7 ± 0.6 , corrected electroconductivity 159 ± 53 $\mu\text{S cm}^{-1}$ and Ca concentration 18.1 ± 11.3 mg L⁻¹. Based on The Canadian Wetland System Classification (NWWG 1997), hydrological regime (terrigenous or littogenous) and topography, there were two main types of fens: basin (n = 8 peatlands or units) and riverine (n = 11). The basin fens were richer in species (17 versus 11), had a deeper water table depth (18 versus 5 cm), a lower pH (6.4 versus 7.0) and a lower electroconductivity (143 versus 171 $\mu\text{S m}^{-1}$) than riverine fens. Water element concentrations were also 35% lower (on average) in

the basin than in riverine fens, except for P and Na that had a higher concentration in basin fens. Basin fens also occupied a smaller area ($3 \text{ ha} \pm 3 \text{ [SD]}$ versus $5 \pm 4 \text{ [SD]}$) than riverine fens.

Plant species found in the quadrats ($n = 200$) surveyed in 2012 were grouped into two vegetation communities (Fig. 4; the complete list of recorded species is available in Appendix A1), according to the k-means analysis with the highest Calinsky-Harabasz criterion. RDA also divided plant quadrats into two distinct groups (Fig. 5). The first group, named *tall-sedge community*, was the most widely represented with 126 quadrats out of 200. *Campylium stellatum*, *Carex lasiocarpa* and *C. aquatilis* were the indicator species of this group, as revealed by the IndVal analysis (Fig. 4). Other associated species were *Myrica gale*, *Andromeda polifolia* var. *latifolia*, *Schoenoplectus acutus* var. *acutus* and *Scorpidium cossonii*. These plants were mainly found in sites with a water table close to the surface and open habitats. The high production of graminoid biomass ($> 120 \text{ g m}^{-2}$), which characterized this group, was positively correlated with electroconductivity, Mg, Ca and distance to the St. Lawrence River but negatively with SO_4 concentration, bulk density and the Shannon diversity index.

The second group, named *Sphagnum-Thuja* community, pooled the other 74 quadrats. *Sphagnum warnstorffii* had the highest species indicator value. Other indicator species were mainly woody species (*Linnaea borealis*, *Rhododendron groenlandicum*, *Thuja occidentalis* and *Vaccinium oxycoccus*) or forbs (*Maianthemum trifolium* and *Rubus pubescens*). These species usually grew in drier microhabitats like hummocks. *Carex leptalea* and *C. disperma*, also indicator species, were usually associated with mossy and shady forests.

Discussion

We used paleoecological tools and modern (present-day) plant surveys to gain knowledge on plant assemblages occurring in moderately rich fens of St. Lawrence Lowlands, Canada. The presence in the

gyttja sediments of species thriving in ponds and completing most of their life cycle submerged, such as *Cristatella mucedo* and *Najas flexilis* (Ricciardi and Reiswigh 1994; Wingfield et al. 2005) suggests a limnogenous origin for the Bic-Saint-Fabien peatland. Peatland development followed a successional pathway from a shallow water body to a bryophyte-dominated habitat. This hydrosere succession is a common form of autogenic development (Kuhry et al. 1993; Charman 2002). In short, paleoecological analyses revealed the existence of three distinct plant assemblages: (1) an emergent plant assemblage in the first phase of the peatland history (dominated by *Schoenoplectus acutus* or *S. tabernaemontani* and *Menyanthes trifoliata*), (2) a tall-sedge assemblage (dominated by *Carex rostrata* and *Carex diandra*), and (3) a brown moss assemblage with small herbaceous plants, including *Paludella squarrosa*, various species of Amblystegiaceae and Calliergonaceae, *Rhynchospora alba* and *Trichophorum alpinum*.

Modern plant surveys showed the existence of two distinct plant communities dominated by (1) tall sedges or (2) *Sphagnum* species with *Thuja occidentalis*. The first community was associated with *Carex aquatilis*, *Carex lasiocarpa* and *Campylium stellatum*, all species performing better in high water table and high pH conditions (Gignac 1992; Gignac et al. 2004). The analogs to the tall-sedge community were found in paleoecological assemblages, especially in zone II. The second community was strongly associated with *Sphagnum warnstorffii* along with generalist wetland boreal species found in dryer peatland microhabitats (Kenkel 1987; Locky et al. 2005).

Relevance of paleoecological analyses

Paleoecological studies are usually used to assess plant succession through the development of the peatland and to identify processes driving this succession. Here, the results were used in a restoration perspective. First, radiocarbon dates of surface peat may provide indications on the amount of time it was necessary to form the peat deposit that was extracted. The 50 years of peat harvesting activities resulted, at

R1 and R2, in a loss of about 2500 to 3200 years of peat accumulation. At R3, 10 years of peat harvesting resulted in a loss of about 800 years of peat. At the time peat harvesting stopped, the BSF site at the location of the R1 and R2 cores 'regressed', from a peat stratigraphy point of view, from a *Sphagnum* bog to a poor fen. The plant assemblage found at the top part of these cores included species representative of moderately rich to rich fen (ex. mosses from Amblystegiaceae family, *Ptychostomum pseudotriquetrum*) but also from poor fen (ex. Ericaceous species, *Comarum palustre*). The richness status could be interpreted as the transitional between the two types of fens (rich and poor). The R3 core most probably regressed to a rich fen. According to successional patterns observed in the paleoecological analyses, target plant assemblage for reference should be inspired from moderately rich fen in the eastern part (R3 location). The same assemblage could be introduced in the central part (R1 and R2) as long as the water chemistry and hydrology fall into the range seen in natural fens. As the paleoecological status was not clear-cut, if the environmental conditions at BSF are closer to poor fen characteristics, we advised targeting this specific assemblage.

High occurrence of bryophytes in zones II and IV generated a higher peat accumulation rate, averaging respectively 1.1 and 0.6 mm per year compared to 0.05 mm for zone III where bryophytes were absent. Even though brown mosses decompose at a faster pace than *Sphagnum* species (Vitt 1990), they decompose at a slower rate than vascular plants (Bartsch and Moore 1985), and thus generate significant peat accumulation. If the most important objective of a peatland restoration plan is to restore the carbon sequestration function, then particular attention should be given to the return of the bryophytes.

Natural disturbance events are of interest for restoration ecologists. By looking at the evolution of an ecosystem after disturbance, one may detect which species help system recovery and the potential trajectories that a restored ecosystem can potentially take (Robert et al. 1999). Fires are a common natural disturbance in peatlands (Turetsky and St. Louis 2006; Lavoie and Pellerin 2007) and past fire events

(charcoal layers) are easily detectable in paleoecological analyses. For instance, *Polytrichum strictum* grew in abundance after fire in ombrotrophic peatlands (as indicated by paleoecological studies: Kuhry et al. 1993, Lavoie et al. 2001, Benscoter 2006), which suggests it helps to reestablish the plant cover. The same species was also found to be an excellent nurse plant for *Sphagnum* reestablishment during the restoration of harvested peatlands (Rocheffort et al. 2003; Groeneveld et al. 2007; González et al. 2014). Fire events were not frequent at BSF, probably because the water level in the peatland was most of the time probably very close to the surface (as inferred by the species assemblages), or because the tree cover was too sparse to generate detectable amounts of charcoal. However, at N, *Campylium stellatum* appeared just after the last fire (1930). This species often colonizes bare peat surface quickly after disturbance (Tyler 1984; Arnesen 1999; Mälson et al. 2010) and is able to grow on mineral ground (Gauthier 2014; Borkenhagen and Cooper 2016). It was observed in 2012 abundantly growing in disturbed places like all-terrain vehicles trails where the surface vegetation cover disappeared. Even if there was no evidence of its role in the facilitation of plant reestablishment, *C. stellatum* could be a candidate to test in this respect.

Usefulness of modern plant surveys

We surveyed a wide range of minerotrophic peatland types in the BSF region, but only two distinct plant communities were identified, i.e., tall-sedge and *Sphagnum-Thuja* communities. These communities are similar in species composition to those found in open, moderately-rich to extremely-rich fens of western boreal Canada (Chee and Vitt 1989; Locky et al. 2005), and in the forested cedar swamp of northern Ontario (Kenkel 1987; Jeglum 1991). Although similarities have been found, reference system determination should nevertheless essentially be based on local surveys, since species habitat requirements may change from region to region (Clewett and Aronson 2013). Moreover, modern plant surveys may also highlight discrepancies between past vegetation assemblages – which are always only a partial record of real communities, since not all plants produce well-preserved macrofossils – and modern

vegetation. For instance, *Thuja occidentalis* was very abundant in the peatlands in 2012 but absent in the macrofossil analyses.

Knowledge of environmental variables of undisturbed, natural sites can be used as guidelines for evaluating the success of a restoration project (Ruiz-Jaen and Aide 2005; Shackelford et al. 2013). In Table 1 and summary Fig. 6, we showed the range of variability for several environmental variables for fens of the BSF area. The characteristics of the restored site should perhaps fall within this range. The nutrients and mineral water concentrations fall within the range seen in the literature for fen (compilation from Andersen et al. 2011), especially moderately rich fen (Chee and Vitt 1989). Among interesting observations that should be considered during the restoration, there was the higher nitrate concentrations that favored shrub growth. Although it is recognized that high nitrate concentrations can negatively impact plant diversity in fens (Drexler and Bedford 2002; Bobbink et al. 2010); that was not the case in our study. However, in the long term, shrub expansion might be detrimental to plant diversity, by diminishing moss abundance for example (Pedrotti et al. 2014). As moss cover is a difficult characteristic to re-establish in disturbed fens, nitrogen concentration should be considered carefully before implementing a restoration project. A second observation concerns the possible link between the high sulfate concentrations in water as seen in riverine fens with poor species richness. There are numerous factors that affect species richness, and we do not pretend an explanation with sulfate. However, there is a paucity of research on the direct impact of sulfate on vegetation communities in peatlands as most studies on the influence of the sulfur cycle looked at microorganisms (Andersen et al. 2013) and the impact on the C cycle (Vile and Novák 2006). Since sulfate concentrations increased with time after the restoration (Andersen et al. 2010), the interaction between sulfate and vegetation should be examined closer.

In summary, our present-day study succeeded in establishing relationships between plant assemblages and a set of environmental conditions (Fig. 6). These relationships are the foundation of reference system for a

restoration. Therefore, to expect success, the reference system for a restoration should match up with the environmental conditions in the disturbed site to be restored. In the BSF case study, for example, if a high water table is forecast, tall-sedge assemblage will be more successful than a *Sphagnum warnstorffii* dominated assemblage. Thus, it is only after evaluation of the disturbed site conditions that we can determine the best plant assemblage to introduce (Clewett and Aronson 2013).

Application of the results: BSF case-study

As seen previously in both paleoecological and present-day studies, a total of four fen plant communities were described as possible references for the BSF restoration project. We should then target one or several of these communities for this project. At BSF, the topography of the disturbed area is one of the main constraints influencing the choice of an appropriate reference. The central section of BSF, where R1 and R2 were cored, is in a lower position compared to the eastern section (R3, N). The monitoring of the water table depth and water chemical analyses indicated that the blockage of the drainage ditches in 2009 consistently maintained a high water table in the central section (Malloy and Price 2013), which was facilitated by the fact that mineral-rich water is continuously flowing toward this section (L. Rochefort, unpublished data). Spontaneous revegetation, mostly with *Typha latifolia*, *Scirpus cyperinus*, *S. atrocinctus* and *Salix* species, already occurred along former drainage ditches in the central section. Reintroducing small herbaceous plants and bryophytes could be risky since they will likely be out-competed. Paleoecological analyses would suggest the restoration of a poor fen assemblage in the central section (R1 and R2 cores), but modern data would instead point toward the restoration of a tall-sedge community. Species from this community are probably competitive enough to cope with the other species already established.

Peat harvesting did not last long in the eastern section of BSF where R3 was cored. However, negative impacts are important: peat compaction and subsidence on the first meter of the peat profile, low water

table depth, frost heaving, no spontaneous vegetation, etc. These impacts are commonly seen in post-harvested cutover peatlands (Price 1996; Poulin et al. 2005). At R3, a *Sphagnum*-brown moss community with more hummock species and with a better survival in driest conditions would consequently be an objective that can realistically be achieved with the moss layer transfer technique currently used in North American bogs (Rocheport et al. 2003).

Complementarity of both methods

The selection of species to introduce during the restoration process is the key to reach broader functional and ecosystem stability objectives (Clewett and Aronson 2013; Shackelford et al. 2013). We focused on the knowledge acquisition of minerotrophic peatland plant communities as it is becoming a more prominent goal and a tool for peatland restoration planning. Present-day studies produce more applied outcomes but only illustrated a snapshot in the sequence of changing species. Paleoecological tools informed us about the succession of communities and link them to autogenic or allogenic processes. In our study, long persistence in some communities indicated stability. Emergent species were replaced by mat species then bryophytes and small herbaceous plants. The present-day vegetation at BSF still supports this last community. Disadvantages of paleoecological analyses center on time required to process cores, and the need to train people. Therefore, paleoecological analyses are less applicable or accessible for smaller restoration projects. But once it is done in a region, one can assume similar dynamics for similar hydrogeomorphological settings of peatlands (e.g. Swinehart and Parker 2000; Arlen-Pouliot 2009). In this sense, field vegetation surveys have the benefits of being faster and have more direct applied outcomes than paleoecological studies mostly to describe environmental variables associated with vegetation. Paleoecology extrapolates peatland development from one precise point and interpreting cores are based on actual knowledge of the peatland ecology. Still, understanding ecosystem dynamics is an important component of restoration, (SERI 2004; Shackelford et al. 2013), and this understanding is best achieved

with paleoecology. For example, in our study, the structure of a plant community was shown to be stable in time where tall-sedge community did initiate peat accumulation and brown moss was an essential component. From our present-day study, we were able to compare distance between reference and disturbed sites and make clearer restoration objectives.

Acknowledgments

Financial support was provided by the Natural Sciences and Engineering Research Council of Canada, the Canadian Sphagnum Peat Moss Association and its members. Vicky Bérubé's personal grants were provided by the Fonds québécois de la recherche sur la nature et les technologies. We thank Anne-Marie Wagner, Marie-Eve Gauthier, Phan Cat Tuong and Kathy Pouliot for assistance in the field and laboratory. We are grateful to Jean Faubert who shared its knowledge of the Rimouski area as well as the landowners who allowed us on their land. We thank members of Peatland Ecology Research Group who provided comments on earlier versions of the manuscript as well as reviewers who provided constructive comments.

References

- Andersen, R., Rochefort, L., and Poulin, M. 2010. Peat, water and plant tissue chemistry monitoring: A seven-year case-study in a restored peatland. *Wetlands*, **30**: 159–170. doi:10.1007/s13157-009-0015-0.
- Andersen, R., Rochefort, L., and Landry, J. 2011. La chimie des tourbières du Québec : une synthèse de 30 années de données. *Nat. Can.* **135**(1): 5–14.
- Andersen, R., Wells, C., Macrae, M.L., and Price, J.S. 2013. Nutrient mineralisation and microbial functional diversity in a restored bog approach natural conditions 10 years post restoration. *Soil Biol. Biochem.* **64**: 37–47. doi:org/10.1016/j.soilbio.2013.04.004.
- Arlen-Pouliot, Y. 2009. Développement holocène et dynamique récente des tourbières minérotrophes structurées du haut-boréal québécois. Ph.D. thesis, Université Laval, Québec.
- Arnesen, T.A. 1999, Vegetation dynamics following trampling in rich fen at Sølendet, Central Norway; a 15 year study of recovery. *Nord. J. Bot.* **19**(3): 313–327. doi:10.1111/j.1756-1051.1999.tb01902.x.
- Aronson, J., Dhillon, S., and Le Floch, E. 1995. On the need to select an ecosystem of reference, however imperfect: A reply to Pickett and Parker. *Restor. Ecol.* **3**(1): 1–3. doi:10.1111/j.1526-100X.1995.tb00069.x.
- Balaguer, L., Escudero, A., Martín-Duque, J.F., Mola, I., and Aronson, J. 2014. The historical reference in restoration ecology: Re-defining a cornerstone concept. *Biol. Conserv.* **176**: 12–20. doi:10.1016/j.biocon.2014.05.007.
- Bartsch, L., and Moore, T.R. 1985. A preliminary investigation of primary production and decomposition in four peatlands near Schefferville, Québec. *Can. J. Bot.* **63**(7): 1241–1248. doi:10.1139/b85-171.
- Bauer, I.E., Gignac, L.D., and Vitt, D.H. 2003. Development of a peatland complex in boreal western Canada: lateral site expansion and local variability in vegetation succession and long-term peat accumulation. *Can. J. Bot.* **81**(8): 833–847. doi:10.1139/B03-076.
- Benscoter, B.W. 2006. Post-fire bryophyte establishment in a continental bog. *J. Veg. Sci.* **17**(5): 647–652.

doi:10.1111/j.1654-1103.2006.tb02488.x.

Benscoter, B.W., and Vitt, D.H. 2008. Spatial patterns and temporal trajectories of the bog ground layer along a post-fire chronosequence. *Ecosystems*, **11**(7): 1054–1064. doi:10.1007/s10021-008-9178-4.

Blanchet, F.G., Legendre, P., and Borcard, D. 2008. Forward selection of explanatory variables. *Ecology*, **89**: 2623–2632. doi:10.1890/07-0986.1.

Blundell, A., and Holden, J. 2015. Using palaeoecology to support blanket peatland management. *Ecol. Indic.* **49**: 110–120. doi:10.1016/j.palaeo.2015.11.048.

Birks, H.J.B. 1996. Contributions of Quaternary palaeoecology to nature conservation. *J. Veg. Sci.* **7**: 89–98. doi:10.2307/3236420.

Bobbink R, Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L., and De Vries, W. 2010. Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Appl.* **20**: 30–59. doi:10.1890/08-1140.1.

Borcard, D., Gillet, F., and Legendre, P. 2011. *Numerical Ecology with R*. Springer, New York.

Borkenhagen, A., and Cooper, D.J. 2016. Creating fen initiation conditions: a new approach for peatland reclamation in the oil sands region of Alberta. *J. Appl. Ecol.* **53**: 550–558. doi:10.1111/1365-2664.12555.

Charman, D.J. 2002. *Peatland and environmental change*. J. Wiley, New York.

Chee, W.-L., and Vitt, D.H. 1989. The vegetation, surface water chemistry and peat chemistry of moderate-rich fens in central Alberta, Canada. *Wetlands*, **9**(2): 227–261. doi:10.1007/BF03160747.

Clewell, A.F., and Aronson, J. 2013. *Ecological restoration principles, values, and structures of an emerging profession*. Second edition. Island Press, Washington.

- Dufrène, M., and Legendre, P. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol. Monogr.* **67**: 345–366. doi:10.1890/0012-9615(1997)067[0345:SAAIST]2.0.CO;2.
- Drexler, J.Z., and Bedford, B.L. 2002. Pathways of nutrient loading and impacts on plant diversity in a New York peatland. *Wetlands* **2**(22): 263–281. doi:10.1672/0277-5212(2002)022[0263:PONLAI]2.0.CO;2.
- Ehrenfeld, J.G. 2000. Defining the limits of restoration: the need for realistic goals. *Restor. Ecol.* **8**: 2–9. doi:10.1046/j.1526-100x.2000.80002.x.
- Faubert, J. 2007. Catalogue des bryophytes du Québec et du Labrador. *Provancheria : Mémoire de l'Herbier Louis-Marie*, no 30, Université Laval, Québec.
- Fulton, R.J. 1995. Surficial materials of Canada. Map 1880A. Geological Survey of Canada, Ottawa.
- Gauthier, M.-E. 2014. Restoring peatland plant communities on mineral well pads. M.Sc. thesis. Université Laval, Québec.
- Gauthier, R., and Grandtner, M.M. 1975. Étude phytosociologique des tourbières du Bas Saint-Laurent, Québec. *Nat. Can.* **102**(1): 109–153.
- Gignac, L.D. 1992. Niche structure, resource partitioning, and species interactions of mire bryophytes relative to climatic and ecological gradients in Western Canada. *Bryologist*, **95**(4): 406–418. doi:10.2307/3243564.
- Gignac, L.D., Gauthier, R., Rochefort, L., and Bubier, J. 2004. Distribution and habitat niches of 37 peatland Cyperaceae species across a broad geographic range in Canada. *Can. J. Bot.* **82**(9): 1292–1313. doi:10.1139/B04-081.
- Glaser, P.H., and Janssens, J.A. 1986. Raised bogs in eastern North America: transitions in landforms and gross stratigraphy. *Can. J. Bot.* **64**(2): 395–415. doi:10.1139/b86-056.
- González, E., Rochefort, L., Boudreau, S., and Poulin, M. 2014. Combining indicator species and key

environmental and management factors to predict restoration success. *Ecol. Indic.* **46**: 156–166.
<http://dx.doi.org/10.1016/j.ecolind.2014.06.016>.

Gouvernement du Québec, Ministère de l'Énergie et des Ressources naturelles. 2001. Orthophoto of the Province of Québec 1/40 000 01211-085. Black and white, TIFF format, 61 Mo, Géoboutique Québec, Que.

Gouvernement du Québec, Ministère des Ressources naturelles et de la Faune. 2006. Carte du Québec méridional, carte muette 1/5 000 000. PDF format, 1.2 Mo, Géoboutique Québec, Que.

Graf, M.D., Rochefort, L., and Poulin, M. 2008. Spontaneous revegetation of cutaway peatlands of North America. *Wetlands*, **28**(1): 28–39. doi:10.1672/06-136.1.

Groeneveld, E.V.G., Massé, A., and Rochefort, L. 2007. *Polytrichum strictum* as a nurse-plant in peatland restoration. *Restor. Ecol.* **15**(4): 709–719. doi:10.1111/j.1526-100X.2007.00283.x.

Groupe de travail sur la classification des sols. 2002. Le système canadien de classification des sols. Troisième édition. Publication 1646. Direction générale de la recherche, Agriculture et Agro-Alimentaire Canada, Ottawa.

Hallett, M.L., Diver, S., Eitzel, M.V., Olson, J.J., Ramage, B.S., Sardinas, H., Statman-Weil, Z., and Suding, K.N. 2013. Do we practice what we preach? Goal setting for ecological restoration. *Restor. Ecol.* **21**: 312–319. doi:10.1111/rec.12007.

Halsey, L.A., Vitt, D.H., and Bauer, I.E. 1998. Peatland initiation during the Holocene in continental Western Canada. *Clim. Chang.* **40**(2): 315–342. doi:10.1023/A:1005425124749.

Higgs, E.S. 1997. What is good ecological restoration? *Conserv. Biol.* **11**: 338–348.

Integrated Taxonomic Information System (ITIS) on-line database [online]. Available from <http://www.itis.gov> [accessed on 4 April 2015].

Jackson, S.T., and Hobbs, R.J. 2009. Ecological restoration in the light of ecological history. *Science*, **325**: 567–569. doi:10.1126/science.1172977.

Journal of Association of Official Analytical Chemistry (JAOAC). 1967. Ash of Peat. Official Method 967.04. AOAC **50**: 394.

Jeglum, J.K. 1991. Definition of trophic classes in wooded peatlands by means of vegetation types and plant indicators. Ann. Bot. Fenn. **28**: 175–192.

Jowsey, P.C. 1966. An improved peat sampler. New Phytol. **65**: 245–248. doi:10.1111/j.1469–8137.1966.tb06356.x.

Kenkel, N.C. 1987. Trends and interrelationships in boreal wetland vegetation. Can. J. Bot. **65**(1): 12–22. doi:10.1139/b87-002.

Kuhry, P., Nicholson, B.J., Gignac, L.D., Vitt, D.H., and Bayley, S.E. 1993. Development of *Sphagnum*-dominated peatlands in boreal continental Canada. Can. J. Bot. **71**(1): 10–22. doi:10.1139/b93-002.

Łachacz, A., Nitkiewicz, M., and Pisarek, W. 2009. Soil conditions and vegetation on gyttia lands in the Masurian Lakeland. In Contemporary problems of management and environmental protection. Edited by A. Łachacz. Uniwersytecie Warmińsko-Mazurskim w Olsztynie, Olsztyn, Poland. pp. 61–94.

Lavoie, C., and Pellerin, S. 2007. Fires in temperate peatlands (southern Quebec): past and recent trends. Can. J. Bot. **85**(3): 263–272. doi:10.1139/B07-012.

Lavoie, C., Zimmermann, C., and Pellerin, S. 2001. Peatland restoration in southern Québec (Canada): a paleoecological perspective. Écoscience, **8**: 247–258.

Legendre, P., and Gallagher, E.D. 2001. Ecologically meaningful transformations for ordination of species data. Oecologia, **129**: 271–280. doi:10.1007/s004420100716.

Legendre, P., and Legendre, L. 2012. Numerical ecology, third English edition. Elsevier Science, Amsterdam.

LoBuglio, K.F. 1999. Cenococcum. In Ectomycorrhizal fungi: key genera in profile. Edited by J.W.G. Cairney and S.M. Chambers. Springer-Verlag, Berlin. pp. 287–309.

- Locat, J. 1977. L'émersion des terres dans la région de Baie-des-Sables/Trois-Pistoles, Québec. *Géogr. Phys. Quatern.* **31**: 297–306. doi:10.7202/1000279ar.
- Locky, D.A., and Bayley, S.E. 2005. Plant diversity, composition, and rarity in the southern boreal peatlands of Manitoba, Canada. *Can. J. Bot.* **84**: 940–955. doi:10.1139/b06-049.
- Malloy, S., and Price, J.S. 2014. Fen restoration on a bog harvested down to sedge peat: A hydrological assessment. *Ecol. Eng.* **64**: 151–160. <http://dx.doi.org/10.1016/j.ecoleng.2013.12.015>.
- Meier-Uhlherr, R., Schulz, C., and Luthardt, V. 2015. Steckbriefe Moorsubstrate. 2., unveränd. Aufl., HNE Eberswalde (Hrsg.), Berlin.
- Milligan, G.W. 1996. Clustering validation: results and implications for applied analyses. *In* Clustering and Classification. *Edited by* P. Arabie, L.J. Hubert and G. De Soete. World Scientific Publication, River Edge NJ. pp. 341–375.
- Murdoch, D., and Chow, E.D. (porting to R by Jesus M. Frias Celayeta). 2007. Ellipse: Functions for drawing ellipses and ellipse-like confidence regions. R package version 0.3-5. <http://CRAN.R-project.org/package=ellipse>.
- Mälson, K, Sundberg, S., and Rydin, H. 2010. Peat disturbance, mowing and ditch blocking as tools in rich fen restoration. *Restor. Ecol.* **18**(S2): 469–478. doi:10.1111/j.1526-100X.2009.00563.x.
- National Wetlands Working Group. 1997. The Canadian Wetland Classification System. 2nd Edition. *Edited by* B.G. Warner and C.D.A. Rubec. University of Waterloo, Ontario.
- Oksanen, J.F., Blanchet, G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P.M., Stevens, H.H., and Wagner, H. 2011. Vegan: Community Ecology Package. R package version 2.0-2. <http://CRAN.R-project.org/package=vegan>.

- Pedrotti, E., Rydin, H., Ingmar, T., Hytteborn, H., Turunen, P., and Granath, G. 2014. Fine-scale dynamics and community stability in boreal peatlands: revisiting a fen and a bog in Sweden after 50 years. *Ecosphere*, **5**(10): 133. doi:10.1890/ES14-00202.1.
- Poulin, M., Rochefort, L., Quinty, F., and Lavoie, C. 2005. Spontaneous revegetation of mined peatlands in eastern Canada. *Can. J. Bot.* **83**(5): 539–557. doi:10.1139/b05-025.
- Poulin, M., Andersen, R., and Rochefort, L. 2013. A new approach for tracking vegetation change after restoration: A case study with peatlands. *Restor. Ecol.* **21**(3): 1–9. doi:10.1111/j.1526-100X.2012.00889.x.
- Price, J.S. 1996. Hydrology and microclimate of partly restored cutover bog, Québec. *Hydrol. Process.* **10**: 1263–1272. doi:10.1002/(SICI)1099-1085(199610)10:10<1263::AID-HYP458>3.0.CO;2-1.
- R Development Core Team. 2008. R: A language and environment for statistical computing. Vienna, Austria. [online]. Available from <http://www.R-project.org/>
- Ricciardi, A., and Reiswich, H.M. 1994. Taxonomy, distribution, and ecology of the freshwater bryozoans (Ectoprocta) of eastern Canada. *Can. J. Zool.* **72**: 339–359. doi:10.1139/z94-048.
- Robert, E.C., Rochefort, L., and Garneau, M. 1999. Natural revegetation of two block-cutting post-mined peatlands in eastern Canada. *Can. J. Bot.* **77**(3): 447–459. doi:10.1139/b99-019.
- Roberts, D.W. 2013. Ordination and Multivariate Analysis for Ecology. R package version 1.6-1. [online]. Available from <http://CRAN.R-project.org/package=labdsy>.
- Rochefort, L., Quinty, F., Campeau, S., Jonhson, K., and Malterer, T. 2003. North American approach to the restoration of *Sphagnum* dominated peatlands. *Wetl. Ecol. Manag.* **11**(1-2): 3–20. doi:10.1023/A:1022011027946.
- Rochefort, L., Isselin-Nondedeu, F., Boudreau, S., and Poulin, M. 2013. Comparing survey methods for monitoring vegetation change through time in a restored peatland. *Wetl. Ecol. Manag.* **21**: 71–85. doi:10.1007/s11273-012-9280-4.

- Ruiz-Jaen, M.C., and Aide, T.M. 2005. Restoration success: how is it being measured? *Restor. Ecol.* **13**(3): 569–577. doi:10.1111/j.1526-100X.2005.00072.x.
- Scoggan, H.J. 1942. Ecological studies of the arctic-alpine flora of the Gaspé Penninsular and of Bic. Ph.D. thesis, McGill University, Montréal.
- Shackelford, N., Hobbs, R.J., Burgar, J.M., Erickson, T.E., Fontaine, J.B., Laliberté, E., Ramalho, C.E., Perring, M.P., and Standish, R.J. 2013. Primed for change: Developing ecological restoration for the 21st century. *Restor. Ecol.* **21**: 297–304. doi:10.1111/rec.12012.
- Sjörs, H. 1950. On the relation between vegetation and electrolytes in North Swedish mire waters. *Oikos* **2**: 241–259. doi:10.2307/3564795.
- Society for Ecological Restoration International Science and Policy Working Group. 2004. The SER International Primer on Ecological Restoration. Society for Ecological Restoration International, Tuscon, AZ. [online]. Available from www.ser.org.
- Stuiver, M., Reimer, P.J., and Reimer, R.W. 2005. CALIB 5.0 [WWW program and documentation]. [online]. Available from <http://radiocarbon.pa.qub.ac.uk/calib/manual/reference.html>.
- Swinehart, A.L., and Parker, G.R. 2000. Palaeoecology and development of peatlands in Indiana. *Am. Midl. Nat.* **143**(2): 267–297.
- Turetsky, M.R., and St. Louis, V.L. 2006. Disturbance in boreal peatlands. *In* Boreal peatland ecosystems. Edited by R.K. Wieder and D.H. Vitt. Ecological Studies, vol. 188. Springer-Verlag, Berlin. pp. 359–379.
- Tyler, C. 1984. Calcareous fens in South Sweden. Previous use, effects of management and management recommendations. *Biol. Conserv.* **30**: 69–89. doi:10.1016/0006-3207(84)90020-X.
- Vile, M.A., and Novák, M. 2006. Sulfur cycling in boreal peatlands: from acid rain to global climate change. *In* Boreal peatland ecosystems. Edited by R.K. Wieder and D.H. Vitt. Ecological Studies, vol. 188. Springer-Verlag, Berlin. pp. 259–288.

- Vitt, D.H. 1990. Growth and production dynamics of boreal bryophytes over climatic, chemical and topographical gradients. *Bot. J. Linn. Soc.* **104**: 35–59. doi:10.1111/j.1095-8339.1990.tb02210.x.
- Wind-Mulder, H.L., and Vitt, D.H. 2000. Comparison of water and peat chemistries of a post-harvested and undisturbed peatland with relevance to restoration. *Wetlands*, **20**: 616–628. doi:10.1672/0277-5212(2000)020[0616:COWAPC]2.0.CO;2.
- Wingfield, R., Murphy, K., and Gaywood, M. 2005. Lake habitat suitability for the rare European macrophyte *Najas flexilis* (Willd.) Rostk. & Schmidt. *Aquat. Conserv.* **15**(3): 227–241. doi:10.1002/aqc.673.
- Whittaker, R.H. 1972. Evolution and measurement of species diversity. *Taxon*, **21**: 213–251.
- White, P.S., and Walker, J.L. 1997. Approximating nature's variation: selecting and using reference information in restoration ecology. *Restor. Ecol.* **5**(4): 338–349. doi:10.1046/j.1526-100X.1997.00547.x.
- Yu, Z. 2006. Holocene carbon accumulation of fen peatlands in boreal Western Canada: A complex ecosystem response to climate variation and disturbance. *Ecosystems*, **9**(8): 1278–1288. doi:10.1007/s10021-006-0174-2.

Tables

Table 1 Description of 23 environmental variables, code names and units measured in each of the fens (units) surveyed in the Bas-Saint-Laurent region in 2012. All measurements were conducted at the end of July 2012, except for water sampling in Plourde and Aqueduc peatlands in mid-July 2009. Water chemistry samples were analyzed at Laval University. Asterisk indicates variables kept for the RDA model after forward selection.

Group	Variables	Code	Unit	Range	Notes
Descriptive	Peatland classification*	type	factor	riverine, basin	Estimated in the field (NWWC 1997).
	Distance to the St. Lawrence River*	dist_river	km	1 to 21	Estimation based on satellite photographs (Google earth V7.1.5.1557).
	Disturbance*	disturb	factor	none, limited	Estimated in the field.
Physical	Water level	WT	cm	-35 to 17	Used a ruler within a dug hole
	Peat depth*	depth	cm	42 to 612	Metal rod inserted in the central part of the peatland (unit).
	Bulk density*	BD	g ml ⁻¹	0.13 to 0.30	Peat samples dried and weighed in the laboratory.
	Organic mater	OM	%	78.6 to 91.9	Dry-ash method (JAOAC 1967)
Water chemistry	pH	pH	-	5.8 to 8.1	Measured in the laboratory (Fisher Scientific Accumet AB150).
	Electroconductivity*	EC	µS cm ⁻¹	94 to 275	Measured in the laboratory (Orion Model 122) and corrected according to Sjörs (1950).
	P	P	mg L ⁻¹	0 to 1.6	ICP†
	Ca*	Ca	mg L ⁻¹	7.16 to 45.75	ICP
	Mg*	Mg	mg L ⁻¹	0.75 to 7.01	ICP
	Fe	Fe	mg L ⁻¹	0 to 0.26	ICP
	K*	K	mg L ⁻¹	0.21 to 7.90	ICP
	Na*	Na	mg L ⁻¹	1.82 to 12.22	ICP
	NH ₄	NH ₄	mg L ⁻¹	0.05 to 5.58	FIA‡
	NO ₃ *	NO ₃	mg L ⁻¹	0.01 to 0.14	FIA
	SO ₄ *	SO ₄	mg L ⁻¹	1.13 to 21.64	FIA
Vegetation	Shannon diversity index*	shannon	—	0 to 2.53	H' = $\sum [(p_i) \cdot \log_2(p_i)]$ calculated with all species per quadrat.
	Richness*	rich	—	4 to 22	Number of species per quadrat
	Biomass - graminoid*	b_gram	g m ² ⁻¹	2.92 to 296.50	All living vegetation cut from two 50×50 cm quadrats randomly placed in a representative
	Biomass - forbs*	b_forb	g m ² ⁻¹	2.92 to 63.71	

Group	Variables	Code	Unit	Range	Notes
	Biomass - shrubs*	b_shrub	g m ² ⁻¹	0 to 84.50	homogenous section, dried and weighed.
	General tree cover*	cover	factor	variable, open	Estimated in the field for each quadrat.

† ICP = Inductively Coupled Plasma spectroscopy (Optima 4300 DV from Perkin Elmer).
‡ FIA = Flow Injection Analysis (Quikchem 4000, Lâchât Instrument Division, Milwaukee, WI)

Draft

Table 2 Radiocarbon dates from the four peat cores extracted at the Bic-Saint-Fabien (BSF) peatland.

Core site and depth (cm)	Laboratory no.	Age (¹⁴ C year BP)	1-sigma range (cal year BP)	Calibrated age (cal year BP)	1-sigma range (cal year)	Calibrated age	Material dated
R1, 2-4	ULA-1627/UCIAMS-76677	3015 ± 15	3175-3221	3198	1272-1226 BC	1249 BC	Moss fragments
R1, 18-20	ULA-1617/UCIAMS-76675	3300 ± 15	3492-3533	3513	1584-1543 BC	1564 BC	Moss fragments
R1, 35-40	ULA-1616/UCIAMS-76674	3360 ± 15	3575-3614	3595	1665-1627 BC	1646 BC	Moss fragments
R1, 155-160	ULA-1626/UCIAMS-76676	7235 ± 20	8001-8052	8027	6103-6052 BC	6078 BC	Wood fragments
R2, 2-4	ULA-1615/UCIAMS-76673	2425 ± 15	2376-2465	2421	516-427 BC	472 BC	Moss fragments
R2, 16-18	ULA-1614/UCIAMS-76672	2660 ± 15	2754-2769	2762	820-805 BC	813 BC	Moss fragments
R2, 165-170	ULA-1582/UCIAMS-76658	7260 ± 20	8019-8055	8037	6106-6070 BC	6088 BC	Wood fragments
R3, 2-4	ULA-1613/UCIAMS-76671	820 ± 15	702-735	719	1215-1248 AD	1232 AD	Moss fragments
R3, 75-80	ULA-1581/UCIAMS-76657	2460 ± 15	2635-2694	2665	754-686 BC	720 BC	Wood fragments
R3, 185-190	ULA-1584/UCIAMS-76660	3955 ± 20	4412-4431	4422	2489-2461 BC	2475 BC	Moss fragments
R3, 235-240	ULA-1564/UCIAMS-75742	4340 ± 20	4854-4886	4870	2937-2905 BC	2921 BC	Moss fragments
R3, 320-325	ULA-1583/UCIAMS-76659	5915 ± 20	6716-6752	6734	4803-4767 BC	4785 BC	Gyttja sediments
N, 105-110	ULA-1567/UCIAMS-75745	1300 ± 15	1256-1281	1269	669-674 AD	672 AD	Moss fragments
N, 195-200	ULA-1566/UCIAMS-75744	3815 ± 15	4158-4205	4182	2256-2209 BC	2233 BC	Moss fragments
N, 275-280	ULA-1565/UCIAMS-75743	4370 ± 15	4927-4960	4944	3011-2978 BC	2995 BC	Moss fragments
N, 335-340	ULA-1563/UCIAMS-75741	4815 ± 20	5488-5505	5497	3556-3539 BC	3548 BC	Wood fragments
N, 445-450	ULA-1562/UCIAMS-75740	6205 ± 20	7067-7112	7090	5163-5118 BC	5141 BC	Wood fragments

Figures captions

Figure 1 Locations of the 13 sites used for present-day surveys of fen vegetation in the Bas-Saint-Laurent region, Québec (Canada). Map was re-used with the permission of ©Gouvernement du Québec (2006).

Figure 2 Aerial photograph of the Bic-Saint-Fabien (BSF) peatland complex and of the surrounding landscape. The peatland developed in a valley surrounded by a mountain in the north (dark green), part of Bic National Park, and agricultural lands in the south (pale yellow). An ancient lake located in the central part of the peatland, as seen on an aerial photograph taken in 1925, is overlaid in blue. The lake was drained in 1954 and is now covered, in the sections that have not been used for peat extraction, by dense shrubs and wet meadows. The dotted white line delimits the 15 ha area claimed for a restoration project. The peat core locations (R1, R2, R3, N) are also indicated. Source : Orthophoto was used with the permission of the ©Gouvernement du Québec (2001).

Figure 3 Macrofossil diagrams (selected taxa) of the Bic-Saint-Fabien BSF peatland. (a) R1, (b) R2, (c) R3 and (d) N. The dotted lines mark the different vegetation zones.

Figure 4 The 40 most frequent species found in fens of the Bas-Saint-Laurent region, classified by life form and grouped into two communities obtained by K-means partitioning: (a) tall-sedge and (b) *Sphagnum-Thuja* communities. Frequency (bar chart) represents, in percentage, the occurrence of each species in all quadrats. Abundance (dots), also in percentage, is the average cover ($\pm$ SD) per quadrat when species is present. Significant indicator species obtained with the IndVal method are in bold. We retained species with an IndVal>0.25 and $P < 0.001$ of each

community; ($***P < 0.001$, $**P < 0.01$, $*P < 0.05$). The full name of species, coded here by the first three letters for the genus and species, are available in Appendix A1.

Figure 5 RDA triplot performed with 17 significant environmental variables. RDA explained 33.1% of the total variation ($r^2 = 0.36$, $F = 6.348$, $P < 0.001$). Open circles represent centroids of the ellipses and include 95% of the quadrats from the two plant communities obtained by k-means partitioning. Only the 16 species with the highest scores (most significant) are shown for clarity. Codes name for species are available in Appendix A1 and for environmental variables in Table 1.

Figure 6 Schematic representations of the plant assemblages and their associated habitats within a moderately rich fen ecosystem as described by the paleoecological stratigraphies and present-day studies. The range of variability of the main water chemical elements and biomass are also summarized for two main fen types.

MOSSES

TREES and SHRUBS Botany

ERICACEOUS SHRUBS

MYCORRHIZAE BRYOZOA CHARCOAL

CYPERACEAE

FORBS

b

MOSSES

TREES and SHRUBS

ERICACEOUS SHRUBS

MYCORRHIZAE BRYOZOA

CHARCOAL

CYPERACEAE

FORBS

Number of macrofossil pieces / 100 cm³ of peat

Legend

Moderately decomposed peat with moss and sedge, Moderately decomposed peat with moss and wood remains, Gytja, Sand and clay

MOSSES

TREES and SHRUBS

ERICACEOUS SHRUBS

MYCORRHIZAE

BRYOZOA CHARCOAL

CYPERACEAE

FORBS

Number of macrofossil pieces / 100 cm³ of peat

b Sphagnum-Thuja community

Appendix A1

Title: List of species identified during present-day inventories in 19 moderately rich to rich minerotrophic peatlands (and units) located around Rimouski, QC. Nomenclature followed the Integrated Taxonomic Information System (2015) for vascular plants and Faubert (2007) for bryophytes.

Form	Code	Species names	Authority names
Bryophyte	aul_pal	<i>Aulacomnium palustre</i>	(Hedw.) Schwaegr.
Bryophyte	pty_pse	<i>Ptychostomum s.l. pseudotriquetrum</i>	(Hedw.) J.R.Spence & H.P.Ramsay
Bryophyte	bryo_sp	<i>Bryophyte</i> sp.	
Bryophyte	cal_cus	<i>Calliergonella cuspidata</i>	(Hedw.) Loeske
Bryophyte	cal_gig	<i>Calliergon giganteum</i>	(Schimp.) Kindb.
Bryophyte	cal_lin	<i>Calliergonella lindbergii</i>	(Mitt.) Hedenäs
Bryophyte	calli_sp	<i>Calliergon s.l. sp.</i>	(Sull.) Kindb.
Bryophyte	cam_ste	<i>Campylium stellatum</i>	(Hedw.) Lange & C.E.O. Jensen
Bryophyte	cli_den	<i>Climacium dendroides</i>	(Hedw.) Web. & Mohr
Bryophyte	dic_pol	<i>Dicranum polysetum</i>	Sw. ex anon.
Bryophyte	dre_adu	<i>Drepanocladus aduncus</i>	(Hedw.) Warnst.
Bryophyte	dre_unc	<i>Sanionia uncinata</i>	(Hedw.) Loeske
Bryophyte	drep_sp	<i>Depanocladus s.l. sp.</i>	(C. Müll.) G. Roth
Bryophyte	fis_sp	<i>Fissidens</i> sp.	Hedw.
Bryophyte	ham_ver	<i>Hamatocaulis vernicosus</i>	(Mitt.) Hedenäs
Bryophyte	hel_bla	<i>Helodium blandowii</i>	(Web. & Mohr) Warnst.
Bryophyte	hep_sp	<i>Hepatic</i> sp.	
Bryophyte	hyl_spl	<i>Hylocomium splendens</i>	(Hedw.) Schimp.
Bryophyte	hypn_sp	<i>Hypnaceae</i> sp.	
Bryophyte	mni_pun	<i>Rhizomnium punctatum</i>	(Hedw.) T.J.Kop.
Bryophyte	mni_sp	<i>Mnium</i> sp.	Hedw.
Bryophyte	pal_squ	<i>Paludella squarrosa</i>	(Hedw.) Brid.
Bryophyte	ple_sch	<i>Pleurozium schreberi</i>	(Brid.) Mitt.
Bryophyte	pol_str	<i>Polytrichum strictum</i>	Brid.
Bryophyte	pti_cri	<i>Ptilium crista-castrensis</i>	(Hedw.) De Not.
Bryophyte	rhy_tri	<i>Rhytidiadelphus triquetrus</i>	(Hedw.) Warnst.
Bryophyte	sco_cos	<i>Scorpidium cossonii</i>	(Schimp.) Hedenäs
Bryophyte	sco_sco	<i>Scopidium scorpioides</i>	(Hedw.) Limpr.
Bryophyte	sph_ang	<i>Sphagnum angustifolium</i>	(C. Jens. ex Russ.) C. Jens.
Bryophyte	sph_cen	<i>Sphagnum centrale</i>	C. Jens
Bryophyte	sph_fim	<i>Sphagnum fimbriatum</i>	Wils.

Form	Code	Species names	Authority names
Bryophyte	sph_fus	<i>Sphagnum fuscum</i>	(Schimp.) Klinggr.
Bryophyte	sph_mag	<i>Sphagnum magellanicum</i>	Brid.
Bryophyte	sph_pal	<i>Sphagnum palustre</i>	L.
Bryophyte	sph_squ	<i>Sphagnum squarrosum</i>	Crome.
Bryophyte	sph_sub	<i>Sphagnum subsecundum</i>	Nees.
Bryophyte	sph_ter	<i>Sphagnum teres</i>	(Schimp.) Angstr.
Bryophyte	sph_war	<i>Sphagnum warnstorffii</i>	Russ.
Bryophyte	thu_del	<i>Thuidium delicatulum</i>	(Hedw.) Schimp.
Bryophyte	tom_nit	<i>Tomenthypnum nitens</i>	(Hedw.) Loeske.
Bryophyte	tor_tor	<i>Tortella tortuosa</i>	(Hedw.) Limpr.
Graminoid	cal_can	<i>Calamagrostis canadensis</i>	(Michx.) Nutt.
Graminoid	car_aqu	<i>Carex aquatilis</i>	Wahl.
Graminoid	car_aur	<i>Carex aurea</i>	Nutt.
Graminoid	car_dia	<i>Carex c.f. diandra</i>	Schrank
Graminoid	car_dis	<i>Carex disperma</i>	Dewey
Graminoid	car_ech	<i>Carex echinata</i>	Murray
Graminoid	car_exi	<i>Carex exilis</i>	Dewey.
Graminoid	car fla	<i>Carex flava</i>	L.
Graminoid	car_gyn	<i>Carex gynocrates</i>	Wormsk. ex Drejer
Graminoid	car_int	<i>Carex interior</i>	L.H. Bailey
Graminoid	car_lac	<i>Carex lacustris</i>	Willd.
Graminoid	car_las	<i>Carex lasiocarpa</i>	Ehrh.
Graminoid	car_lep	<i>Carex leptalea</i>	Wahl.
Graminoid	car_lim	<i>Carex limosa</i>	L.
Graminoid	car_mag	<i>Carex magellanica</i>	Lam.
Graminoid	car_pra	<i>Carex prairea</i>	Dewey ex Alph. Wood
Graminoid	car_pse	<i>Carex pseudocyperus</i>	L.
Graminoid	car_sp	<i>Carex sp.</i>	L.
Graminoid	car_vag	<i>Carex vaginata</i>	Tausch
Graminoid	eri_viri	<i>Eriophorum viridicarinarum</i>	(Engelm.) Fernald.
Graminoid	gly_can	<i>Glyceria canadensis</i>	(Michx.) Trin.
Graminoid	mul_glo	<i>Muhlenbergia glomerata</i>	(Willd.) Trin.
Graminoid	poa_sp	<i>Poaceae sp.</i>	
Graminoid	rhy_alb	<i>Rhynchospora alba</i>	(L.) Vahl.
Graminoid	rhy_fus	<i>Rhynchospora fusca</i>	(L.) W.T. Aiton
Graminoid	sci_acu	<i>Schoenoplectus acutus</i> var. <i>acutus</i>	(Muhl. ex Bigelow) Á. Löve & D. Löve
Graminoid	tri_alp	<i>Trichophorum alpinum</i>	L.
Graminoid	tri_ces	<i>Trichophorum cespitosum</i>	L.
Forb	gal_rot	<i>Galearis rotundifolia</i>	(Banks ex Pursh) R.M. Bateman
Forb	ara_nud	<i>Aralia nudicaulis</i>	L.
Forb	are_bul	<i>Arethusa bulbosa</i>	L.

Form	Code	Species names	Authority names
Forb	ast_novb	<i>Symphyotrichum novi-belgii</i> var. <i>novi-belgii</i>	(L.) G.L. Nesom
Forb	ast_pun	<i>Symphyotrichum puniceum</i>	(L.) Á. Löve & D. Löve
Forb	ast_sp	<i>Aster s.l.</i> sp.	
Forb	doe_umb	<i>Doellingeria umbellata</i>	(Mill.) Nees
Forb	ath_fil	<i>Athyrium filix-femina</i>	(L.) Roth
Forb	cal_pal	<i>Calla palustris</i>	L.
Forb	cic_bul	<i>Cicuta bulbifera</i>	L.
Forb	cli_bor	<i>Clintonia borealis</i>	(Aiton) Raf.
Forb	con_chi	<i>Conioselinum chinense</i>	(L.) Britton, Sterns & Poggenb.
Forb	cop_tri	<i>Coptis trifolia</i>	L.
Forb	cyp_par	<i>Cypripedium parviflorum</i>	Salisb.
Forb	cyp_reg	<i>Cypripedium reginae</i>	Walter
Forb	dro_int	<i>Drosera intermedia</i>	Hayne.
Forb	dro_rot	<i>Drosera rotundifolia</i>	L.
Forb	epi_pal	<i>Epilobium palustre</i>	L.
Forb	equ_pal	<i>Equisetum palustre</i>	L.
Forb	eup_mac	<i>Eutrochium maculatum</i>	(L.) E.E. Lamont
Forb	fra_vir	<i>Fragaria virginiana</i>	Duchesne.
Forb	gal_lab	<i>Galium labradoricum</i>	(Wiegand) Wiegand
Forb	gal_sp	<i>Galium</i> sp.	L.
Forb	gal_tri	<i>Galium triflorum</i>	Michx.
Forb	geu_riv	<i>Geum rivale</i>	L.
Forb	tri_vir	<i>Triadenum virginicum</i>	(L.) Raf.
Forb	iri_ver	<i>Iris versicolor</i>	L.
Forb	neo_con	<i>Neottia convallariodes</i>	(Sw.) Rich.
Forb	lob_kal	<i>Lobelia kalmii</i>	L.
Forb	lyc_uni	<i>Lycopus uniflorus</i>	Michx.
Forb	mai_can	<i>Maianthemum canadense</i>	Desf.
Forb	mal_uni	<i>Malaxia unifolia</i>	Michx.
Forb	men_tri	<i>Menyanthes trifoliata</i>	L.
Forb	mit_nud	<i>Mitella nuda</i>	L.
Forb	nup_var	<i>Nuphar variegata</i>	Engelm.
Forb	orch_sp	<i>Orchidaceae</i>	
Forb	osm_cin	<i>Osmundastrum cinnamomea</i>	(L.) C. Presl
Forb	osm_reg	<i>Osmunda regalis</i>	L.
Forb	par_gla	<i>Parnassia glauca</i>	Raf.
Forb	phr_aut	<i>Phragmites australis</i> ssp. <i>americanus</i>	(Cav.) Trin. ex Steud.
Forb	gym_cla	<i>Gymnadeniopsis clavellata</i>	(Michx.) Rydb.
Forb	pla_dil	<i>Platanthera dilatata</i>	(Pursh) Lindl. ex L.C. Beck
Forb	pog_oph	<i>Pogonia ophioglossoides</i>	(L.) Ker-Gawl.
Forb	per_amp	<i>Persicaria amphibia</i>	(L.) DelTree

Form	Code	Species names	Authority names
Forb	com_pal	<i>Comarum palustre</i>	L.
Forb	pota_sp	<i>Potamogeton</i> sp.	L.
Forb	pyr_asa	<i>Pyrola asarifolia</i>	Michx.
Forb	pyr_ell	<i>Pyrola elliptica</i>	Nutt.
Forb	ran_acr	<i>Ranunculus acris</i>	A. Gray
Forb	rub_pub	<i>Rubus pubescens</i>	Raf.
Forb	sag_lat	<i>Sagittaria latifolia</i>	Willd.
Forb	san_can	<i>Sanguisorba canadensis</i>	L.
Forb	sar_pur	<i>Sarracenia purpurea</i>	L.
Forb	scu_gal	<i>Scutellaria galericulata</i>	L.
Forb	sel_sel	<i>Selaginella selaginoides</i>	(L.) Link.
Forb	mai_tri	<i>Maianthemum trifolium</i>	L.
Forb	sol_uli	<i>Solidago uliginosa</i>	Nutt.
Forb	tha_pub	<i>Thalictrum pubescens</i>	Pursh
Forb	the_pal	<i>Thelypteris palustris</i>	Schott
Forb	thy_lat	<i>Typha latifolia</i>	L.
Forb	tof_glu	<i>Tofieldia glutinosa</i>	(Michx.) Pers.
Forb	tri_bor	<i>Trientalis borealis</i>	Raf.
Forb	tri_mar	<i>Triglochin maritima</i>	L.
Forb	utr_int	<i>Utricularia intermedia</i>	Hayne.
Forb	val_uli	<i>Valeriana sitchensis</i> ssp. <i>uliginosa</i>	(Torr. & A. Gray) F.G. Mey.
Forb	vic_cra	<i>Vicia cracca</i>	L.
Forb	vio_sp	<i>Viola</i> sp.	L.
Shrub	and_pol	<i>Andromeda polifolia</i> var. <i>latifolia</i>	Aiton
Shrub	cha_cal	<i>Chamaedaphne calyculata</i>	(L.) Moench.
Shrub	cor_can	<i>Cornus canadensis</i>	L.
Shrub	cor_ser	<i>Cornus sericea</i>	L.
Shrub	gau_his	<i>Gaultheria hispidula</i>	(L.) Muhl. ex Bigelow
Shrub	gay_bac	<i>Gaylussacia baccata</i>	(Wang.) K. Koch.
Shrub	kal_ang	<i>Kalmia angustifolia</i>	L.
Shrub	kal_pol	<i>Kalmia polifolia</i>	Wang.
Shrub	rho_gro	<i>Rhododendron groenlandicum</i>	(Oeder) Kron & Judd
Shrub	lin_bor	<i>Linnaea borealis</i>	L.
Shrub	lon_vil	<i>Lonicera villosa</i>	(Michx.) R. & S.
Shrub	myr_gal	<i>Myrica gale</i>	L.
Shrub	nep_muc	<i>Nemopanthus mucronatus</i>	(L.) Trel.
Shrub	dai_fru	<i>Dasiphora fruticosa</i>	(L.) Rydb.
Shrub	rha_aln	<i>Rhamnus alnifolia</i>	L'Hér.
Shrub	rib_lac	<i>Ribes lacustre</i>	(Pers.) Poir.
Shrub	ros_bla	<i>Rosa palustris</i>	Marshall
Shrub	sal_beb	<i>Salix bebianna</i>	Sarg.
Shrub	sal_dis	<i>Salix discolor</i>	Mühl.

Form	Code	Species names	Authority names
Shrub	sal_sp	<i>Salix</i> sp.	L.
Shrub	spi_lat	<i>Spiraea latifolia</i>	(Ait.) Borkh.
Shrub	vac_ang	<i>Vaccinium angustifolium</i>	Ait.
Shrub	vac_oxy	<i>Vaccinium oxycoccos</i>	L.
Shrub	vib_cas	<i>Viburnum cassinoides</i>	L.
Tree	abi_bal	<i>Abies balsamea</i>	(L.) Mill.
Tree	ace_pen	<i>Acer pensylvanicum</i>	L.
Tree	ace_rub	<i>Acer rubrum</i>	L.
Tree	aln_rug	<i>Alnus incana</i>	(L.) Moench.
Tree	bet_pap	<i>Betula papyrifera</i>	Marsh.
Tree	fra_nig	<i>Fraxinus nigra</i>	Marsh.
Tree	lar_lar	<i>Larix laricina</i>	(DuRoi) Koch.
Tree	pic_mar	<i>Picea mariana</i>	(Mill.) BSP.
Tree	sor_dec	<i>Sorbus decora</i>	(Sarg.) C.K. Schneid.
Tree	thu_occ	<i>Thuja occidentalis</i>	L.

References

Faubert, J. 2007. Catalogue des bryophytes du Québec et du Labrador. Provancheria : Mémoire de l'Herbier Louis-Marie, no 30, Université Laval, Québec.

Integrated Taxonomic Information System (ITIS) on-line database [online]. Available from <http://www.itis.gov> [accessed on 4 April 2015].