Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

**Bipartite networks improve understanding of the effects of
waterbody size and angling method on angler-fish
interactions**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0176.R3
Manuscript Type:	Article
Date Submitted by the Author:	09-Mar-2017
Complete List of Authors:	Chizinski, Christopher; University of Nebraska, School of Natural Resources Martin, Dustin; University of Nebraska-Lincoln, Nebraska Cooperative Fish, and Wildlife Research Unit and School of Natural Resources Shizuka, Daizaburo ; University of Nebraska-Lincoln, School of Biological Sciences Pope, Kevin; University of Nebraska-Lincoln, U.S. Geological Survey— Nebraska Cooperative Fish and Wildlife Research Unit, and School of Natural Resources
Please Select from this Special Issues list if applicable:	
Keyword:	social-ecological systems, interaction networks, fisheries management, RECREATIONAL FISHERIES < General, BYCATCH < General

SCHOLARONE™
Manuscripts

Bipartite networks improve understanding of effects of waterbody size and angling method on angler-fish interactions

Christopher J. Chizinski*
School of Natural Resources, University of Nebraska, Lincoln, NE 68583, USA. cchizinski@unl.edu

Dustin R. Martin^
Nebraska Cooperative Fish and Wildlife Research Unit, and School of Natural Resources, University of Nebraska, Lincoln, NE 68583, USA. dustin.martin@huskers.unl.edu

Daizaburo Shizuka
School of Biological Sciences, University of Nebraska, Lincoln, NE 68588, USA. dshizuka2@unl.edu

Kevin L. Pope
U.S. Geological Survey—Nebraska Cooperative Fish and Wildlife Research Unit, and School of Natural Resources, University of Nebraska, Lincoln, NE 68583, USA. kpope2@unl.edu

*Corresponding author: email: cchizinski2@unl.edu, phone 402-405-4809, fax: 402-472-2946

^Current address: ReelSonar, Inc., Seattle, WA 98109, USA. dustin@reelsonar.com

Running head: Angler-sportfish bipartite networks

For re-submission to Canadian Journal of Fisheries and Aquatic Sciences

Abstract

Networks used to study interactions could provide insights to fisheries. We compiled data from 27297 interviews of anglers across waterbodies that ranged in size from 1 to 12113 ha. Catch rates of fish species among anglers grouped by species targeted generally differed between angling methods (bank or boat). We constructed angler-catch bipartite networks (angling-method specific) between anglers and fish, and measured several network metrics. There was considerable variation in networks among waterbodies, with multiple metrics influenced by waterbody size. Number of species-targeting angler groups and number of fish species caught increased with increasing waterbody size. Mean number of links for species-targeting angler groups and fish species caught also increased with waterbody size. Connectance (realized proportion of possible links) of angler-catch interaction networks decreased slower for boat anglers than bank anglers with increasing waterbody size. Network specialization (deviation of number of interactions from expected) was not significantly related to waterbody size or angling methods. Application of bipartite networks in fishery science requires careful interpretation of outputs, especially considering the numerous confounding factors prevalent in recreational fisheries.

Introduction

In recreational fisheries, anglers' influence extends beyond the species that they target (Cooke and Cowx 2004, 2006; Lewin et al. 2006). The influence of recreational angling on targeted species is well known and includes decreases in abundance, changes in age structure and size structure, and changes in species composition (Blaber et al. 2000). Anglers tend to prefer certain species because of their value for food and angling challenge (Lewin et al. 2006). However, the angling population is not comprised of a single, homogeneous group, but is rather a heterogeneous group made up of numerous subgroups (O'Neill 2001; Arlinghaus et al. 2008; Hutt and Jackson 2008; Beardmore et al. 2015). Each subgroup of anglers can vary in their specialization, motivations, choice of species targeted, angling method, and tackle choices, among a multiplicity of other factors. Thus, during any given period at a waterbody there is a heterogeneous suite of anglers fishing for a multitude of species (Chizinski et al. 2014a, 2014b), using various gears and approaches (e.g., angling from a boat or bank). Furthermore, no angling tackle or bait is exclusive to the species being targeted. Anglers' decisions on where and how to fish, along with decisions on tackle choice, create a potential subset of the fish community that could be caught (whether intended or not). Pope et al. (2016) provided evidence that anglers tend to catch and harvest the species being targeted, but there was also considerable recreational bycatch (species caught but not targeted). Further, Pope et al. (2016) indicated that there was a large segment of the angling community that were target generalists (i.e., targeting all species), who tended to have the second greatest catch rate of each species.

Interaction networks are increasingly used to study dynamics within populations, communities, and ecosystems (Montoya et al. 2006; Kéfi et al. 2015). A primary objective of analyzing ecological networks is to provide an understanding of the complexity in the natural world that ultimately affects ecosystem functioning (Ings et al. 2009). Ecological interactions need to be assessed at the community level to draw valid conclusions about the ecological and evolutionary mechanisms underlying these

interactions (Vázquez et al. 2009). This understanding is essential in predicting ecological responses to disturbances, habitat loss, climate change, and invasive species (Ings et al. 2009).

Ecological interaction networks are representations of associations (links) between species (nodes). A particular type of ecological network, the bipartite network, describes the interactions between two levels in the community (Vázquez et al. 2007; Eklöf et al. 2013), which includes plant-animal interactions, plant-pollinator interactions, and parasite-host interactions. Recreational fisheries, like other complex social-ecological systems (Mitchell 2009), are well suited to assess interactions between anglers and fish because the relationships between anglers and fish are complex and dynamic (Pope et al. 2014; Beardmore et al. 2015; Mee et al. 2016), made further so by multiple angler groups interacting with multiple fish species at any given time. Network analysis could provide unique understanding of the underlying structure of a social-ecological system that could inform detailed mechanistic models and provide managers better information on basic patterns of fishing activity (Martin 2013). Bipartite networks can be extended from an ecological context to investigate the socio-ecological interaction between anglers grouped by species targeted (i.e., the predator) and captured fish species (i.e., the prey). In theory, this “angler-catch interaction network” could range in structure from relatively simple networks in which anglers grouped by species targeted catch only the species targeted to random networks in which each species-targeting angler group captures a random set of fish species (Fig. 1). As we will show, real networks display complex structure in which each species-targeting angler group catches a mixture of fish species, but to different degrees (Fig. 1). The purpose of this study is threefold: 1) illustrate application of bipartite networks to recreational fisheries, 2) compare angler-catch interaction networks between angling methods (i.e., anglers fishing from the bank versus anglers fishing from a boat), and 3) describe the influence of waterbody size on the angler-catch interaction networks.

The species-area relationship predicts that there would be increasing numbers of species with increasing size of area (Connor and McCoy 1979), and this could have complex effects on the interactions between anglers and fish. As the number of fish species in a waterbody increases with size (Eadie et al. 1986), we expect a greater number of anglers and hence a greater number of targeting groups. With

increasing number of groups, we may observe patterns in the interactions associated with waterbody size (i.e., decreased connectivity, greater network complexity) among these groups. Further, with increasing waterbody size there is often a transition from littoral- to pelagic-dominated fish communities (Mehner et al. 2005) and thus, we might expect differences in the networks (e.g., number of species caught, number of angler targeting groups, network connectance) with waterbody size between anglers that are fishing from the shore and anglers fishing from a boat.

Materials and methods

Angler survey data

We compiled creel surveys from 2009 to 2013 for 42 waterbodies in Nebraska (Table 1). Data were collected through in-person interviews to document angler-participation patterns, species targeted, catch and harvest at reservoirs across Nebraska. Though interviews took place throughout the year, we limited data to interviews that occurred during May-August. One angler, the representative of the party, completed the survey; thus, all data were collected at the party (i.e., a group of individuals travelling together for fishing) level. All complete trips and incomplete trips > 30 mins were included in the analysis. Bank anglers were defined as those fishing from the shoreline and boat anglers were those using a floating vessel (e.g., kayak, float tube, or motorized boat). A stratified multi-stage probability-sampling regime (Malvestuto 1996) was used to determine days of interviews. At each waterbody, we calculated the mean catch per unit effort (1 hr of fishing) for each species for all species present in the waterbody for each species-targeting angler group. The result was a matrix of catch per unit effort with rows representing each species caught and columns representing the species sought by anglers. For waterbodies that had data for several years, we took the mean across the years.

Similarity between angling method

Before building the angler-catch networks we needed to assess the similarity between angling methods (bank or boat) within a waterbody to provide justification on whether the catches by angler groups were different enough to warrant different networks for bank and boat anglers. Strong correlation between the two matrices would indicate similar catches and at similar rates between boat and bank anglers, and suggest that a network could be created with catches combined across angling method; lack of correlation would indicate that the catches were different enough that we should construct separate networks for bank and boat anglers. To do this, we scaled and centered by columns (angler targeting groups). Scaling and centering allowed us to determine the relative influence of each fish species by each angler group rather than differences in magnitude in those values. We then used Euclidean distances and Mantel's test to assess the correlation between the two matrices using the *ade4* package (Dray et al. 2007) in R. Significance of Mantel's r was determined from 999 iterations.

Angler-catch interaction networks

We created networks in which fish taxa and angler groups are represented as "nodes" (or vertices), and the reported harvest of a fish taxon by an angler group as an "edge" (or link) that connects these two nodes. This network is considered a "bipartite network" because it only contains edges between two distinct types of nodes ("anglers" and "fish"). Edges in bipartite networks represent interactions or strengths of associations between the two different groups (Michailidis 2008). In the context of an angler-catch interaction network and for the approach we used (i.e., using catch per unit effort [CPUE]), we have the anglers grouped by species targeted as the upper nodes and the fish species caught as the lower nodes (Fig. 1). The width of the node (related to node strength) for an angler group is the relative CPUE for all species caught by that angler group, whereas the width of the node for a fish species caught is the relative CPUE for that species caught by all anglers. The width of an edge (bar connecting the upper

and lower nodes) is the proportion of total CPUE for the species-targeting angler group comprised of the individual fish species. The bipartite networks are arranged such that nodes in the middle have more connections (i.e., anglers grouped by species targeted that caught more fish species and fish species caught by more angler groups) and nodes at the periphery have fewer connections. Weighted bipartite networks were created for each lake using the bipartite package (Dormann et al. 2008) in R (R Development Core Team 2015); this bipartite package generates a visualization of the network.

For each network, we extracted several network- (Dormann et al. 2009), group-, and species-level (Dormann et al. 2008) metrics. Network-level metrics describe the entire network structure, group-level metrics describe the properties of upper (i.e., species-targeting groups or predators) and lower (i.e., species caught or prey) nodes, and species-level metrics describe individual nodes in the upper and lower levels. The specific network-level metrics that we examined were connectance, which is the realized proportion of possible links (Dunne et al. 2002), and H_2' , which is the network-level measure of specialization that is based on the deviation of a species' realized number of interactions from that expected for each species' total number of interactions (Blüthgen et al. 2006). The H_2' index is scaled such that its values range from 0 in networks with completely random interactions to 1 in networks with complete specialization. The specific group-level metrics that we examined were the number of upper (i.e., anglers grouped by species targeted) and lower (i.e., fish species caught) nodes, and the mean number of links per node for the upper and lower levels. For the species-level metrics, we identified the node with the greatest species strength, which is the sum of dependencies of each node (Bascompte et al. 2006)—i.e., either the number of fish species that are caught by species-targeting angler groups or the number of species-targeting angler groups that caught a particular fish species.

The above metrics were chosen to describe angler-catch interaction networks for several reasons. Connectance is the most commonly used and simplest metric to describe the density of links between nodes in an interaction network and can be interpreted as the degree of generalization in the system (i.e., high connectance = high generalization) (Blüthgen et al. 2006, 2008). An angler-catch interaction network with a high connectance would indicate that many of the angler groups catch many of the fish

species, and likewise that many of the fish species are being caught by many of the angler groups. Although similar to connectance in describing a degree of generalization of a network, $H2'$ is a finer description of the nature of the connectance in an interaction network. As such, $H2'$ describes specialization in terms of exclusiveness, where $H2' = 1$ represents the most nested scenario possible given the frequency distribution, and $H2' = 0$ describes the greatest deviation from that nested assemblage (i.e., completely random interactions) (Blüthgen et al. 2006, 2008). In the context of the angler-catch interaction network, $H2'$ describes the degree of specialization (i.e., catching only the species that is being sought) across all the angler groups, where $H2' = 1$ indicates a fishery with no recreational bycatch and $H2' = 0$ indicates a fishery with extensive recreational bycatch. The species strength focuses on the architecture of the network and provides a measure of the importance of a node from the perspective of the nodes at the opposite level (i.e., the sum of the weight of all interactions of a predator on its prey) (Barrat et al. 2004; Bascompte et al. 2006). In the context of the angler-catch interaction network, at the upper level a node with the greatest strength would indicate a species-targeting angler group with high CPUE of many fish species, and at the lower level a node with the greatest strength would indicate a fish species that has a high CPUE among many angler groups.

To illustrate how the metrics described above change with varying network complexity, we will refer to our conceptual networks described previously (Fig. 1). In this conceptualization, all the simulated networks had 10 upper nodes and 10 lower nodes. The simple network with no interaction (Fig. 1A) has a connectance of 0.10 (10 connections of a possible 100), an $H2'$ of 1 (complete specialization), a mean number of links of upper nodes of 1, and a mean number of links of lower nodes of 1. In this simple network, species strength was constant across the upper nodes (1 for each node) and the lower nodes (1 for each node), indicating that all nodes were equally influential. The network with intermediate complexity (Fig. 1B) has a connectance of 0.38 (38 connections of a possible 100), an $H2'$ of 0 (random interactions), a mean number of links of upper nodes of 4.47, and a mean number of links of lower nodes of 4.37. In this intermediate network, species strength varied across the upper nodes and across the lower nodes, indicating that node U6 was the most influential on the upper level and node L6 was the most

influential on the lower level. The complex network (Fig. 1C) has a connectance of 0.29 (29 connections of a possible 100), an $H2'$ of 0.85 (somewhat specialized), a mean number of links of upper nodes of 2.93, and a mean number of links of lower nodes of 2.95. In this network, species strength varied across the upper nodes and across the lower nodes, indicating that node U2 was the most influential on the upper level and node L5 was the most influential on the lower level.

Statistical analyses

We were interested in how the angler-catch interaction networks varied over waterbody size and angling method. For the network-level and group-level metrics, we assessed the relationship of the metric (dependent) with surface area (log transformed), angling method (bank or boat), and the interaction between surface area and angling method using linear regression. If the interaction term was not significant ($P > 0.05$), we dropped the interaction and ran the model with the main effects only. For each network metric, we present the final model (i.e., with or without the interaction term) following standard nomenclature, including F statistic with model and residual degrees of freedom, model significance and variance explained (i.e., R^2). In addition, we provide the coefficients for the final linear regression models with coefficient estimate, standard error, t statistic (i.e., coefficient estimate divided by the standard error), and associated p value (i.e., upper-tail probability of achieving a t value from a t distribution with degrees of freedom equal to the residual degrees of freedom from the model). We log-transformed surface area to reduce the spread in our surface area (i.e., most lakes were < 1000 ha with a few lakes > 5000 ha), which qualitatively had no influence on our results. For the influential species metric, we identified the node with the greatest strength for both anglers grouped by species targeted and fish species for both angling methods.

Results

We compiled interview data from 27297 interviews of angler parties for waterbodies that ranged in size from 1 to 12113 ha (median size = 19 ha; Table 1). The across-year mean proportion of bank anglers ranged from 0.12 to 1.00, with lower proportions of bank anglers associated with larger waterbodies.

We assessed the similarity in the angler-catch between bank and boat anglers for 36 of the 42 waterbodies. Six waterbodies had data only for bank anglers; these waterbodies were excluded from the Mantel's test. Only four (Fremont Lake 20 [Mantel test $r = 0.977$, $p = 0.003$], Harlan County Reservoir [Mantel test $r = 0.728$, $p = 0.001$], Fremont Lake 11 [Mantel test $r = 0.716$, $p = 0.014$], and Willow Creek Lake [Mantel test $r = 0.592$, $p = 0.050$]) of the 36 waterbodies had a significant correlation between bank and boat anglers grouped by species targeted and fish species caught, which suggests that the angler-catch interaction networks were different between bank and boat anglers, and warranted separate analyses. The number of angler groups ranged from 3 to 14 for bank anglers and from 2 to 12 for boat anglers. The number of fish species caught ranged from 3 to 22 for bank anglers and from 2 to 19 for boat anglers. In our assessments, 59% of bank and 71% of the boat angler-catch interaction networks had more fish species caught than were targeted. The most frequently sought fish species among bank anglers were anything (98% of waterbodies), channel catfish (93%), and largemouth bass *Micropterus salmoides* (88%), and among boat anglers were largemouth bass (97%), anything (86%), and crappie *Pomoxis* spp. (83%). There were five fish species (rainbow trout *Oncorhynchus mykiss*, bullhead *Ameiurus* spp., grass pickerel *Esox americanus vermiculatus*, buffalo *Ictiobus* spp., and gizzard shad *Dorosoma cepedianum*) sought by bank anglers that were not sought by boat anglers. Conversely, there were four fish species (longnose sucker *Catostomus catostomus*, tiger muskellunge *E. masquinongy* $\times$ *E. lucius*, carp sucker *Carpiodes carpio*, and rudd *Scardinius erythrophthalmus*) sought by boat anglers that were not sought by bank anglers. There were two species (shorthead redhorse *Moxostoma macrolepidotum* and smallmouth buffalo *I. bubalus*) caught by bank anglers that were not caught by boat anglers. Conversely, there were eight fish species (reardear sunfish *Lepomis microlophus*, saugeye *Sander vitreus* $\times$ *S. canadensis*, goldeye *Hiodon alosoides*, yellow bass *Morone mississippiensis*, blue sucker *Cycleptus elongatus*, carp sucker, and grass pickerel) caught by boat anglers that were not caught by bank anglers.

We illustrate the utility of bipartite networks for assessing angler-targeting and catch data using boat angler-catch networks from Fremont Lake 15 (simple network) and Merritt Reservoir (complicated network) (Fig. 2). Fremont Lake 15 is a 22-ha waterbody in eastern Nebraska and Merritt Reservoir is a 1093-ha waterbody in north central Nebraska. At Fremont Lake 15, there were three upper-level (i.e., angler targeting groups) and four lower-level groups (i.e., species caught), a connectance of 0.50, an $H2'$ of 0.97, a mean number of species caught per species-targeting angler group of 2.26, and a mean number of species-targeting angler groups per species of 1.35. The largemouth bass-targeting angler was the most influential upper-level group and channel catfish *Ictalurus punctatus* was the most influential lower-level group on the angler-catch network in Fremont Lake 15. At Merritt Reservoir, there were 12 upper-level and 18 lower-level groups, a connectance of 0.52, an $H2'$ of 0.50, a mean number of species caught per species-targeting angler group of 9.44, and a mean number of species-targeting angler groups per species of 7.97. The anything-targeting angler group (anglers that indicated they targeted any fish species) was the most influential upper-level group and yellow perch was the most influential lower-level group on the angler-catch network in Merritt Reservoir.

The anything-targeting angler group was most frequently (21 of the 40 waterbodies) the angler group of greatest importance to the angler-catch interaction networks, regardless of waterbody size for bank anglers (Table 1). The anything-targeting angler group was also important in the boat angler-catch interaction networks, but was less frequently the angler group of greatest importance (10 of 34 waterbodies). The crappie-targeting angler group closely followed the anything-targeting angler group in the boat angler-catch interaction networks (9 of 34 waterbodies). There was a transition in the fish species caught of greatest importance for both bank and boat angler-catch interaction networks with increasing waterbody size (Table 1). At smaller waterbodies (< 100 ha), the fish species of greatest importance were littoral-associated species (e.g., bluegill *Lepomis macrochirus*, crappie, and largemouth bass). In contrast, there was a transition to pelagic species (white bass *Morone chrysops* and walleye *Sander vitreus*) becoming the fish species of greatest importance at larger (>100 ha) waterbodies.

There was considerable variation in the complexity in the networks among the 42 waterbodies assessed, with multiple metrics influenced by waterbody size, method, and the associated interactions. The model of the number of species-targeting angler groups ($F_{3,72} = 37.08$, $P < 0.001$; $R^2 = 0.61$) included log-transformed area, angler type, and the associated interaction. The interaction in this model indicated that the number of angler groups increased with waterbody surface area slower for bank anglers than boat anglers (Table 2; Fig. 3). The model of the number of species caught ($F_{2,73} = 86.50$, $P < 0.001$; $R^2 = 0.70$) included log-transformed area and angler type, with no associated interaction. The lack of significant interaction indicated that the numbers of fish species caught increased significantly with waterbody surface area, but there was no difference between angling methods (Table 2; Fig. 3). The model of the mean number of links for species-targeting angler groups ($F_{2,73} = 34.94$, $P < 0.001$; $R^2 = 0.49$) included log-transformed area and angler type, with no associated interaction. In contrast, the model predicting the mean number of links for fish species caught ($F_{3,72} = 20.07$, $P < 0.001$; $R^2 = 0.46$) included log-transformed area, angler type, and the associated interaction. The inclusion of the interaction in the mean number of links for fish species caught but not for the mean number of links for species-targeting angler groups is opposite from what was observed in the number of angler groups and number of species caught. The interaction in the mean number of links for fish species caught indicated that number of angler groups increased with waterbody surface area slower for bank anglers than boat anglers (Table 2; Fig. 3). The model of connectance ($F_{2,73} = 10.70$, $P < 0.001$; $R^2 = 0.23$) included log-transformed area and angler type, with no associated interaction. Connectance of angler-catch interaction networks decreased slower for boat anglers than bank anglers with increasing waterbody size (Table 2; Fig. 3). The model of H2' ($F_{2,73} = 0.2882$, $P = 0.750$; $R^2 = 0.01$) included log-transformed area and angler type, with no associated interaction. Unexpectedly, network specialization (H2') was not significantly related to waterbody size or angling method (Table 2, Fig. 3).

Discussion

Research has highlighted the extent and subsequent effect on the ecosystem of commercial fishing (Cooke and Cowx 2006; Drake and Mandrak 2014), but the influence of recreational fishing is less well known (Arlinghaus et al. 2007). The angler-catch interaction networks from this study indicate that recreational fisheries are complex and extend well beyond a single angler group-fish species relationship. Anglers can potentially affect a great number of fish species beyond the target species. Many fish species are caught and released (Pope et al. 2016), and thus are subject to numerous sub-lethal and behavioral changes associated with catch and release (Cooke and Cowx 2004, 2006). Further, there is great variation in the number and type of by-catch produced across angler groups that target different fish species. Therefore, angler-fish interactions on any given waterbody can be complex (Pope et al. 2016; Arlinghaus et al. 2017). One can improve understanding of interactions between anglers and fish, whether targeted or not, by examining angler-catch interaction networks; improved understanding could lead to better fisheries management and conservation practices.

We documented a range of complexity in angler-catch interaction networks across waterbodies. For example, the boat angler-catch network in Fremont Lake 15 was relatively simple (Fig. 2A) and illustrated some of the utility of bipartite networks to assess angler groups and fish species. On the upper level of the Fremont Lake 15 angler-catch network we see that crappie-targeting anglers had the greatest relative catch rates, followed by largemouth bass-targeting anglers, and then channel catfish-targeting anglers. On the lower level of the network we see the greatest relative catch rates were for crappie (regardless of who sought them) and that relative catch rates for largemouth bass and channel catfish were similar. Angler-catch interaction networks also provide an opportunity to illustrate patterns between species-targeting angler groups and the fish species caught. The Fremont Lake 15 angler-catch interaction network illustrates that largemouth bass-targeting anglers were the only anglers to catch largemouth bass. Channel catfish-targeting anglers only caught channel catfish, but channel catfish were also caught by largemouth bass- and crappie-targeting anglers. In addition to providing an opportunity to identify which fish species are being caught by which angler groups, angler-catch interaction networks also provide the ability to identify potentially what angler groups are catching species of concern (e.g.,

invasive, nuisance species). White perch *Morone americana* is an invasive species in Nebraska (Chizinski et al. 2010) and is the focus of state management efforts to maintain populations below ecologically damaging levels (Stewart 2015). The Fremont Lake 15 angler-catch interaction network illustrates that only largemouth bass-targeting anglers are catching this invasive species; in this system, largemouth bass-targeting anglers may be the most likely to potentially remove this species from the system and thus, education material (e.g., signs, pamphlets) could be targeted to this angler group to illustrate the danger of transporting this species.

The boat angler-catch interaction network in Merritt Reservoir was much more complex, in relation to Fremont Lake 15 (Fig. 2B). This network indicates that there is considerable recreational bycatch (although relatively small in some cases) by the species-targeting angler groups in this system. The alignment of the anything-targeting anglers in the middle of the figure illustrates that this angler group had the greatest interactions with the fish species caught, whereas the alignment of the bluegill- and common carp-targeting anglers on the sides of the network illustrates that these angler groups had the least interactions with the fish species caught. Anything-targeting anglers at Merritt Reservoir primarily caught yellow perch and bluegill, but also caught 11 other species. Unfortunately, managers and scientists often exclude the anything-targeting angler group from species-specific analyses of angler effort, catch and harvest, and attitudes and opinions (Pope et al. 2016), despite having relatively moderate catch rates and interacting with numerous fish species in the community.

The position of the angler group provides insight for our conceptualization that species targeted influences the recreational bycatch caught. We expect that anglers targeting similar species groups, such as bluegill and redear sunfish, would choose approaches and gears that are similar and hence catch similar fish species; as such, bluegill- and redear sunfish-targeting angler groups would be proximate to each other in the network. In the Merritt angler-catch network, muskellunge- and northern pike-targeting angler groups were proximate to each other, as were smallmouth bass- and largemouth bass-targeting angler groups. The angler-catch interaction network could thus be used to understand and predict the appropriate subset of fish species that are vulnerable to a specific angler group.

We compared angler catch networks between bank and boat anglers and determined that in most cases there were significant differences in the catch rates of fish species among species-targeting angler groups. Our research focused on differences occurring between bank and boat anglers, and not the mechanisms behind those differences. Recent research (Pope et al. 2016) indicated that catch and harvest for bluegill, common carp *Cyprinus carpio*, and crappie were greater for bank anglers, whereas catch and harvest for channel catfish, walleye and white bass were greater for boat anglers. Perhaps this is not surprising given that the availability and catchability of a species likely differ among habitat types, especially habitats typical within littoral and limnetic zones. Further, bank anglers are often confined more than boat anglers in their choice of habitat to fish; bank anglers can only fish littoral habitat (with a possible few exceptions), whereas boat anglers can choose to fish littoral or limnetic or both zones. In addition to differences in the environmental factors influencing differences in catch composition between bank and boat anglers, there are motivational differences that could affect species caught and the rates that those species were caught. For example, Hudgins (1984) indicated that bank anglers rated the importance of harvest and privacy greater than did boat anglers, whereas beauty of the landing and trophy fish were more important to boat anglers. Further research is needed to evaluate the mechanisms that may be influencing the difference between bank and boat anglers.

A common generalization in community ecology is that, within a group of islands or other isolated habitats within a restricted latitudinal range, larger areas harbor more species (i.e., species-area relationship) (MacArthur and Wilson 1967; Connor and McCoy 1979). Similar to ecological networks within a temperate region, larger waterbodies are able to support more fish species caught and more species-targeting angler groups. Further, increasing area affects both the number of interactions and network structure in ecological networks (Sugiura 2010). In the 42 angler-catch interaction networks in Nebraska, we observed increasing numbers of species-targeting angler groups with increasing waterbody size, and this rate of increase was lesser for bank anglers than boat anglers. We also observed increasing numbers of fish species caught by anglers with greater waterbody size, but this rate was not different between bank and boat anglers. Further, there is decreasing connectance (i.e., proportion of links from all

those possible) in the angler-catch interaction network with increasing waterbody size. That is, the number of possible links between angler groups and fish species increases with waterbody size faster than the number of fish species each angler group actually captures. This property of the angler-catch interaction network follows the general rule that connectance decreases with the number of species in ecological networks (Jordano 1987; Sugiura 2010).

There was no relationship between the degree of network specialization or partitioning and waterbody size or angling method when using the H2' index, which compares the diversity of each node's connections with the null expectation given the network size and connectance. For example, in the two networks illustrated in this manuscript, we observed greater specialization in the small waterbody (i.e., Fremont Lake 15) than we did in the larger waterbody (i.e., Merritt Reservoir). Overall, we see that angler-catch interaction networks become more complicated with waterbody size, and that angler-catch interaction networks for boat anglers are more intricate (i.e., contain more species-targeting angler groups and more fish species caught) than angler-catch interaction networks for bank anglers, but we do not necessarily observe increases in network specialization. This may indicate that there is a general rule that governs the degree of network specialization (or generalization) by angler groups that is independent of the number of fish species available. In the case of angler-catch interaction networks, the combination of species-specific fishing strategies (e.g., lure type, habitat choice) is likely to constrain generalization of the fish species they catch, while ecological overlap of fish species may make some degree of bycatch inevitable. Our results suggest that this interplay between anglers and bycatch species may be indirectly related to waterbody size through the effect of waterbody size on species diversity.

Networks in this study were built on mean catch per unit effort between angler groups and fish species caught. Many other angler-fish networks could be investigated to elucidate other patterns in network structure. For example, if we looked at total catch (as opposed to CPUE) by each angler group we could identify the total influence on fish community. Unfortunately, we were unable to calculate total fishing pressure by each species-targeting angler group given the manner in which data were collected in our creel surveys and thus unable to predict the total number of fish caught by each species-targeting

angler group. Further, these networks likely would appear different if focused on species harvested instead of species caught. If we had investigated angler-harvest interaction networks, we would have observed fewer nodes and decreased connectance as anglers tend to harvest a smaller subset of the species caught (Pope et al. 2016). Two additional caveats of this study have to do with our sampling approach that could influence the interpretation of our results. First, we collected information at the party level and there is the possibility that there may have been subtle decisions at the individual level that may have been masked at the party level, though we primarily generalized across two individuals (mean $\pm$ SE party size was 1.80 ± 0.02 anglers for bank anglers and 2.24 ± 0.01 anglers for boat anglers). Second, our survey approach precluded gathering data on multiple target species. If multiple species were specified (other than “anything”) by the respondent, the creel clerk was instructed to record the first species specified; thus, bycatch as discussed in this manuscript is likely overstated.

The use of bipartite networks to assess the interactions between anglers and fish provides an important visualization of the inherent complex relationships in recreational fisheries. This application of bipartite networks requires careful interpretation of outputs, especially considering the numerous confounding factors prevalent in recreational fisheries. Effective management of social-ecological systems, like those observed in recreational fisheries, requires an understanding of the complex interactions between people and the environment (Mee et al. 2016). We can describe the interactions between the entities of the system using network theory and illustrate some of the observed emergent properties (e.g., resilience of the system [Pope et al. 2014]) of the entire system. For example, we can assess how the establishment of an invasive species affects the species-targeting angler groups and associated species caught, identifying angler groups and fish species that are resilient to the establishment of the new species. Focusing on and responding to the complexity of these systems will lead to better management of these social-ecological systems (Degnbol et al. 2006; Mee et al. 2016).

Acknowledgements

433 We thank Tyler Anderson, Zac Brashears, Phil Chvala, Chris Dietrich, Michael Dedinsky, Cameron
434 Depue, Dan Dobesh, Darrol Eichner, Brad Eifert, Holly Evans, Amber Fandrich, Ryan Foley, Andy
435 Glidden, Ron Grandi, Al Hanson, Caleb Huber, Hannah Hummel, Carla Knight, Luke Kowalewski,
436 Rhonda Lawing, Dennis Liess, Jared Lorensen, Natalie Luben, Alexis Maple, George Maynard, Jean Paul
437 Montes, Brad Newcomb, Ashley Pella, Minnie Petch, Robert Pierson, Joe Rydell, Jerry Ryschon, Jeff
438 Schuckman, Mike Smith, Phil Stolberg, and John Walrath for assistance in the field. This project was
439 funded by Federal Aid in Sport Fish Restoration project F-182-R, which was administered by the
440 Nebraska Game and Parks Commission. Any use of trade, firm, or product names is for descriptive
441 purposes only and does not imply endorsement by the U.S. Government. The Nebraska Cooperative Fish
442 and Wildlife Research Unit is jointly supported by a cooperative agreement among the U.S. Geological
443 Survey, the Nebraska Game and Parks Commission, the University of Nebraska, the U.S. Fish and
444 Wildlife Service, and the Wildlife Management Institute.

445

References

- Arlinghaus, R., Alós, J., Beardmore, B., Daedlow, K., Dorow, M., Fujitana, M., Hühn, D., Haider, W., Hunt, L. M., Johnson, B. M., Johnston, F., Kleforh, T., Matsumura, S., Monk, C., Pagel, T., Post, J. R., Rapp, T., Riepe, C., Ward, H., and Wolter, C. 2017. Understanding and managing freshwater recreational fisheries as complex adaptive social-ecological systems. *Rev. Fish. Sci. Aquac.* **35**:1-14.
- Arlinghaus, R., Bork, M., and Fladung, E. 2008. Understanding the heterogeneity of recreational anglers across an urban–rural gradient in a metropolitan area (Berlin, Germany), with implications for fisheries management. *Fish. Res.* **92**(1): 53–62.
- Arlinghaus, R., Cooke, S.J., Lyman, J., Policansky, D., Schwab, A., Suski, C., Sutton, S.G., and Thorstad, E.B. 2007. Understanding the complexity of catch-and-release in recreational fishing: an integrative synthesis of global knowledge from historical, ethical, social, and biological perspectives. *Rev. Fish. Sci.* **15**(1-2): 75–167.
- Barrat, A., Barthelemy, M., Pastor-Satorras, R., and Vespignani, A. 2004. The architecture of complex weighted networks. *Proc. Natl. Acad. Sci. U. S. A.* **101**(11): 3747–3752.
- Bascompte, J., Jordano, P., and Olesen, J.M. 2006. Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* **312**(5772): 431–433.
- Beardmore, B., Hunt, L.M., Haider, W., Dorow, M., and Arlinghaus, R. 2015. Effectively managing angler satisfaction in recreational fisheries requires understanding the fish species and the anglers. *Can. J. Fish. Aquat. Sci.* **72**(4): 500–513.
- Blaber, S., Cyrus, D., Albaret, J.-J., Ching, C.V., Day, J., Elliott, M., Fonseca, M., Hoss, D., Orensanz, J., Potter, I., and others. 2000. Effects of fishing on the structure and functioning of estuarine and nearshore ecosystems. *ICES J. Mar. Sci. J. Cons.* **57**(3): 590–602.
- Blüthgen, N., Fründ, J., Vázquez, D.P., and Menzel, F. 2008. What do interaction network metrics tell us about specialization and biological traits? *Ecology* **89**(12): 3387–3399.

- 471 Blüthgen, N., Menzel, F., and Blüthgen, N. 2006. Measuring specialization in species interaction
472 networks. *BMC Ecol.* **6**(1): 9.
- 473 Chizinski, C.J., Martin, D.R., Hurley, K.L., and Pope, K.L. 2014a. Self-imposed length limits in
474 recreational fisheries. *Fish. Res.* **155**: 83–89.
- 475 Chizinski, C.J., Martin, D.R., and Pope, K.L. 2014b. The influence of a rapid drawdown and prolonged
476 dewatering on fishing effort, catch, and harvest in a Nebraska reservoir. *Gt. Plains Res.* **24**(2):
477 145–152.
- 478 Chizinski, C., Pope, K.L., and Wilde, G. 2010. A modelling approach to evaluate potential management
479 actions designed to increase growth of white perch in a high-density population. *Fish. Manag.*
480 *Ecol.* **17**(3): 262–271.
- 481 Connor, E.F., and McCoy, E.D. 1979. The statistics and biology of the species-area relationship. *Am.*
482 *Nat.* **113**(6): 791–833.
- 483 Cooke, S.J., and Cowx, I.G. 2004. The role of recreational fishing in global fish crises. *BioScience* **54**(9):
484 857–859.
- 485 Cooke, S.J., and Cowx, I.G. 2006. Contrasting recreational and commercial fishing: searching for
486 common issues to promote unified conservation of fisheries resources and aquatic environments.
487 *Biol. Conserv.* **128**(1): 93–108.
- 488 Degnbol, P., Gislason, H., Hanna, S., Jentoft, S., Nielsen, J.R., Sverdrup-Jensen, S., and Wilson, D.C.
489 2006. Painting the floor with a hammer: technical fixes in fisheries management. *Mar. Policy*
490 **30**(5): 534–543.
- 491 Dormann, C.F., Fründ, J., Blüthgen, N., and Gruber, B. 2009. Indices, graphs and null models: analyzing
492 bipartite ecological networks. *Open J. Ecol.* **2**: 7–24.
- 493 Dormann, C.F., Gruber, B., and Fruend, J. 2008. Introducing the bipartite package: Analysing Ecological
494 Networks. *R News* **8**(2): 8–11.
- 495 Drake, D.A.R., and Mandrak, N.E. 2014. Bycatch, bait, anglers, and roads: quantifying vector activity and
496 propagule introduction risk across lake ecosystems. *Ecol. Appl.* **24**(4): 877–894.

- 497 Dray, S., Dufour, A.-B., and others. 2007. The ade4 package: implementing the duality diagram for
498 ecologists. *J. Stat. Softw.* **22**(4): 1–20.
- 499 Dunne, J.A., Williams, R.J., and Martinez, N.D. 2002. Food-web structure and network theory: the role of
500 connectance and size. *Proc. Natl. Acad. Sci.* **99**(20): 12917–12922.
- 501 Eadie, J.M., Hurly, T.A., Montgomerie, R.D., and Teather, K.L. 1986. Lakes and rivers as islands:
502 species-area relationships in the fish faunas of Ontario. *Environ. Biol. Fishes* **15**(2): 81–89.
- 503 Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N.P., Dalsgaard, B., Sassi, C., Galetti,
504 M., Guimarães, P.R., and others. 2013. The dimensionality of ecological networks. *Ecol. Lett.*
505 **16**(5): 577–583.
- 506 Hudgins, M.D. 1984. Structure of the angling experience. *Trans. Am. Fish. Soc.* **113**(6): 750–759.
- 507 Hutt, C.P., and Jackson, J.R. 2008. Implications of angler motivations and preferences for urban fisheries
508 management. *In* Urban and community fisheries programs: development, management, and
509 evaluation. *Edited by* R.T. Eades, J.W. Neal, T.J. Lang, K.M. Hunt, and P. Pajak. American
510 Fisheries Society Symposium 67, Bethesda, Maryland. pp. 63–70.
- 511 Ings, T.C., Montoya, J.M., Bascompte, J., Blüthgen, N., Brown, L., Dormann, C.F., Edwards, F.,
512 Figueroa, D., Jacob, U., Jones, J.I., and others. 2009. Review: Ecological networks—beyond food
513 webs. *J. Anim. Ecol.* **78**(1): 253–269.
- 514 Jordano, P. 1987. Patterns of mutualistic interactions in pollination and seed dispersal: connectance,
515 dependence asymmetries, and coevolution. *Am. Nat.* **129**(5): 657–677.
- 516 Kéfi, S., Berlow, E.L., Wieters, E.A., Joppa, L.N., Wood, S.A., Brose, U., and Navarrete, S.A. 2015.
517 Network structure beyond food webs: mapping non-trophic and trophic interactions on Chilean
518 rocky shores. *Ecology* **96**(1): 291–303.
- 519 Lewin, W.-C., Arlinghaus, R., and Mehner, T. 2006. Documented and potential biological impacts of
520 recreational fishing: insights for management and conservation. *Rev. Fish. Sci.* **14**(4): 305–367.
- 521 MacArthur, R.H., and Wilson, E.O. 1967. *Theory of Island Biogeography*. Princeton University Press.

- 522 Malvestuto, S. 1996. Sampling the recreational creel. *In* Fisheries techniques, 2nd edition. *Edited by* B.
523 Murphy and D. Willis. American Fisheries Society, Bethesda, Maryland. pp. 591–623.
- 524 Martin, D.R. 2013. Spatial and temporal participation in recreational fishing. University of Nebraska-
525 Lincoln.
- 526 Mee, J.A., Post, J.R., Ward, H., Wilson, K., Newton, E., and Cantin, A. 2016. Interaction of ecological
527 and angler processes: experimental stocking in an open access, spatially structured fishery. *Ecol.*
528 *Appl.* **26**(6): 1693–1707. doi:10.1890/15-0879.1.
- 529 Mehner, T., Diekmann, M., Brämick, U., and Lemcke, R. 2005. Composition of fish communities in
530 German lakes as related to lake morphology, trophic state, shore structure and human-use
531 intensity. *Freshw. Biol.* **50**(1): 70–85.
- 532 Michailidis, G. 2008. Data visualization through their graph representations. *In* Handbook of Data
533 Visualization. Edited by C. Chen, W.K. Härdle, and A. Unwin. Springer. pp. 103–120.
- 534 Mitchell, M. 2009. Complexity: A Guided Tour. Oxford University Press.
- 535 Montoya, J.M., Pimm, S.L., and Solé, R.V. 2006. Ecological networks and their fragility. *Nature*
536 **442**(7100): 259–264.
- 537 O'Neill, B.M. 2001. Market segmentation, motivations, attitudes, and preferences of Virginia resident
538 freshwater anglers. Virginia Tech, Blacksburg, Virginia.
- 539 Pope, K.L., Allen, C.R., and Angeler, D.G. 2014. Fishing for resilience. *Trans. Am. Fish. Soc.* **143**(2):
540 467–478.
- 541 Pope, K.L., Chizinski, C.J., Wiley, C.L., and Martin, D.R. 2016. Influence of anglers' specializations on
542 catch, harvest, and bycatch of targeted taxa. *Fish. Res.* **183**: 128–137.
- 543 R Development Core Team. 2015. R: A Language and Environment for Statistical Computing. R
544 Foundation for Statistical Computing, Vienna, Austria. Available from <http://www.R-project.org/>.
- 545 Stewart, N.T. 2015. Ecology and Management of Superabundant Fish Populations. University of
546 Nebraska–Lincoln, Nebraska.

547 Stone, L., and Roberts, A. 1992. Competitive exclusion, or species aggregation? *Oecologia* **91**(3): 419–
548 424.

549 Sugiura, S. 2010. Species interactions–area relationships: biological invasions and network structure in
550 relation to island area. *Proc. R. Soc. B Biol. Sci.* **277**(1689): 1807–1815.

551 Vázquez, D.P., Melián, C.J., Williams, N.M., Blüthgen, N., Krasnov, B.R., and Poulin, R. 2007. Species
552 abundance and asymmetric interaction strength in ecological networks. *Oikos* **116**(7): 1120–1127.

553

Draft

Table 1. Lake names, surface area (ha), number of angler interviews, and mean $\pm$ SE proportion of interviews for anglers fishing from the shoreline, species targeted (anglers grouped by species targeted) and species (fish species caught) of greatest importance for bank-angler and boat-angler networks. A dash indicates that the network metrics could not be calculated because of too few interviews or no fish caught by interviewed anglers. Taxa codes are in Table A1.

Name	Area	Interviews	Bank*	Bank-angler network		Boat-angler network	
				Species targeted	Species	Species targeted	Species
Fremont 5	1.1	262	0.83 $\pm$ 0.03	ANY	BLG	CRP	BLG
Fremont 13	1.9	15	0.92 $\pm$ 0.08	-	-	-	-
TaHaZouka	2.2	51	1.0	ANY	BLG	-	-
Fremont 17	2.3	252	0.93 $\pm$ 0.02	CCF	BLG	-	-
Fremont 11	2.5	221	0.94 $\pm$ 0.02	ANY	BLG	CRP	LMB
Fremont 3	2.6	378	0.99 $\pm$ 0.00	ANY	BHD	-	-
Fremont 12	2.9	74	0.77 $\pm$ 0.05	LMB	BLG	LMB	CCF
Fremont 9	4.3	62	0.95 $\pm$ 0.03	ANY	BLG	-	-
Fremont 18	4.5	139	0.95 $\pm$ 0.05	CCF	BLG	-	-
Fremont 4	4.8	50	0.95 $\pm$ 0.03	LMB	BLG	-	-
Yanney	5.0	121	1.0	ANY	LMB	-	-
Fremont 1	5.1	413	0.92 $\pm$ 0.04	CCF	BLG	CRP	LMB
Fremont 7/8	5.1	9	0.88 $\pm$ 0.13	-	-	CRP	LMB
Fremont 16	5.4	83	0.92 $\pm$ 0.02	ANY	BLG	CRP	BLG
Fremont 2	6.4	557	0.93 $\pm$ 0.02	ANY	BLG	ANY	CCF
Cottontail	10.2	36	0.47	ANY	CCF	ANY	BLG
Timber Point	11.7	43	0.72	ANY	LMB	LMB	BLG
Cottonmill	12.2	168	0.92	ANY	LMB	LMB	LMB
Fremont 10	14.8	160	0.92 $\pm$ 0.02	ANY	CCF	CCF	CRP
Skyview	16.3	167	0.98	CRP	BLG	ANY	LMB
Fremont 20	18.9	420	0.50 $\pm$ 0.07	LMB	CCF	BLG	BLG
Meadowlark	18.9	17	0.71	CRP	BLG	LMB	BLG
Fremont 15	21.7	484	0.97 $\pm$ 0.01	ANY	CCF	LMB	CCF
Holmes	39.2	247	0.14 $\pm$ 0.05	LMB	BLG	ANY	BLG
Wildwood	45.5	271	0.71 $\pm$ 0.03	CCF	BLG	CRP	BLG
Stagecoach	70.3	152	0.54 $\pm$ 0.00	ANY	LMB	ANY	BLG
Conestoga	74.7	76	0.45	LMB	CCF	ANY	BLG
Yankee Hill	79.2	123	0.62	ANY	LMB	LMB	BLG
Willow Creek	234.3	235	0.62	ANY	BLG	WAE	CCF
Pawnee	254.3	138	0.26 $\pm$ 0.02	WAE	CCP	WAE	BLG
Enders	500.5	817	0.26 $\pm$ 0.10	ANY	WHB	ANY	WHB
Box Butte	555.3	794	0.14 $\pm$ 0.06	YEP	NOP	ANY	NOP
Medicine Creek	642.5	930	0.41 $\pm$ 0.05	WAE	WHB	CRP	WHB
Branched Oak	712.6	467	0.66 $\pm$ 0.01	CCF	CCF	CCF	BLG
Johnson	907.6	1872	0.77 $\pm$ 0.01	ANY	FWD	CRP	WHB
Merritt	1093.2	3393	0.33 $\pm$ 0.05	ANY	BLG	ANY	YEP
Sherman	1173.9	2352	0.60 $\pm$ 0.04	ANY	CRP	CRP	CRP
Swanson	1657.3	1219	0.22 $\pm$ 0.06	WAE	LMB	ANY	WHB
Calamus	2054.5	2990	0.45 $\pm$ 0.05	WAE	WHB	WAE	WHB
Harlan Co.	5544.1	3690	0.21 $\pm$ 0.03	ANY	CCF	WAE	WHB

Lewis and Clark	11581.2	1621	0.36± 0.04	WAE	CCF	WAE	BLG
McConaughy	12113.0	1728	0.12± 0.04	ANY	CCF	WAE	WHB

* Mean and SE were calculated across years. Lakes without SE presented had one year of data.

Draft

Table 2. Coefficients for the linear regression models assessing network metrics to the surface area of the waterbodies. Angling method is for bank anglers relative to boat anglers.

Coefficient	Estimate	Standard error	t statistic	Residual df	P value
Number of species-targeting angler groups					
Intercept	4.711	0.468	10.063	72	< 0.001
log(Area)	0.575	0.101	5.690	72	< 0.001
Method	-2.755	0.756	-3.643	72	0.001
log(Area)*Method	0.449	0.154	2.917	72	0.005
Number of fish species caught					
Intercept	2.623	0.623	4.211	73	< 0.001
log(Area)	1.584	0.121	13.144	73	< 0.001
Method	-0.682	0.633	-1.078	73	0.285
Mean number of links for species-targeting angler groups					
Intercept	1.971	0.323	6.088	73	< 0.001
log(area)	0.513	0.063	8.187	73	< 0.001
Method	0.232	0.328	0.707	73	0.482
Mean number of links for fish species caught					
Intercept	3.398	0.371	9.168	72	< 0.001
log(Area)	0.245	0.080	3.056	72	0.003
Method	-2.138	0.589	-3.573	72	< 0.001
log(Area)*Method	0.409	0.122	3.357	72	0.001
Connectance					
Intercept	0.578	0.025	23.460	73	< 0.001
log(Area)	-0.020	0.005	-4.352	73	< 0.001
Method	0.052	0.025	-2.071	73	0.041
H2'					
Intercept	0.625	0.040	15.692	73	< 0.001
log(Area)	-0.000	0.007	-0.023	73	0.982
Method	-0.030	0.040	-0.751	73	0.455

Figure Captions

Fig. 1. Examples of weighted angler catch interactions networks displaying the range of possible network structures. A) is a simple network with complete specialization (species-targeting angler groups catch only what is sought), B) is a more complex network with specialization but with some bycatch, and C) is a complex network with low specialization (species-targeting angler groups catch multiple fish species). The orange boxes signify the anglers grouped by species-targeted nodes and blue boxes signify fish species-caught nodes. The connections between the nodes (edges) are in light gray. Node labels are denoted with a U for upper nodes and L for lower nodes. Node strength is indicated above the node label for the upper nodes and below the node label for the lower nodes. Further detail about these example networks is provided in the Methods.

Fig. 2. Examples of a simple angler-catch network (Fremont 15) and a complicated angler-catch network (Merritt Reservoir) for boat anglers. The width of an edge (bar connecting the upper and lower nodes) is the proportion of total CPUE for the species-targeting angler group comprised of the individual fish species. Taxa codes are in the Table A1.

Fig. 3. Relationship of number of species-targeting angler groups (A), number of fish species caught (B), mean number of links for species-targeting angler groups (C), mean number of links for species caught (D), connectance (E), and $H2'$ (F) to the natural log-transformed surface area. Blue circles, lines, and confidence ribbons are for bank anglers; red circles, lines, and confidence ribbons are for boat anglers. A black line and confidence ribbon indicates a significant relationship between waterbody size and the metric that was unaffected by angling method (bank or boat). No line indicates no significant relationship between waterbody size and the metric. Regression coefficients and associated statistics are in Table 2.

A

1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
U1	U2	U3	U4	U5	U6	U7	U8	U9	U10	

B

1.14	0.54	0.70	1.29	1.43	2.13	1.12	0.48	0.93	0.25
U1	U2	U3	U4	U5	U6	U7	U8	U9	U10

C

1.00	1.09	0.92	0.92	0.88	1.04	1.04	1.04	1.08	0.99
U1	U2	U3	U4	U5	U6	U7	U8	U9	U10

Draft

L1	L2	L3	L4	L5	L6	L7	L8	L9	L10
1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

L1	L2	L3	L4	L5	L6	L7	L8	L9	L10
1.78	0.75	1.26	0.34	0.70	1.88	1.13	0.89	0.93	0.34

L1	L2	L3	L4	L5	L6	L7	L8	L9	L10
1.00	0.92	1.09	1.09	1.13	0.96	0.96	0.96	0.91	1.00

A

B

Table A1. Codes, common names and scientific names for angler targeting groups and taxa caught used in the weighted angler-catch interaction networks.

Code	Common name	Scientific name
ANY	Anything	--
AWF	Alewife	<i>Alosa pseudoharengus</i>
BLG	Bluegill	<i>Lepomis macrochirus</i>
BHD	Black bullhead	<i>Ameiurus melas</i>
CCF	Channel catfish	<i>Ictalurus punctatus</i>
CCP	Common carp	<i>Cyprinus carpio</i>
CRP	Crappie	<i>Pomoxis</i> spp.
FWD	Freshwater drum	<i>Aplodinotus grunniens</i>
LMB	Largemouth bass	<i>Micropterus salmoides</i>
MSK	Muskellunge	<i>Esox masquinongy</i>
NOP	Northern pike	<i>E. lucius</i>
PKS	Pumpkinseed	<i>L. gibbosus</i>
RDR	Redear sunfish	<i>L. microlophus</i>
RKB	Rock bass	<i>Ambloplites rupestris</i>
SMB	Smallmouth bass	<i>M. dolomieu</i>
WAE	Walleye	<i>Sander vitreus</i>
WHB	White bass	<i>Morone chrysops</i>
WHP	White perch	<i>M. americana</i>
WHS	White sucker	<i>Catostomus commersonii</i>
YEP	Yellow perch	<i>Perca flavescens</i>