Canadian Journal of Earth Sciences
 Revue canadienne des sciences de la Terre

**A middle Holocene steppe bison and paleoenvironments
 from the Versleuce Meadows, Whitehorse, Yukon, Canada.**

Journal:	<i>Canadian Journal of Earth Sciences</i>
Manuscript ID	cjes-2017-0100.R1
Manuscript Type:	Article
Date Submitted by the Author:	09-Jun-2017
Complete List of Authors:	Zazula, Grant; Yukon Palaeontology Program Hall, Elizabeth; Yukon Government, Palaeontology Program, , Department of Tourism and Culture Hare, P. ; Yukon Government, Archaeology Program, Thomas, Christian ; Yukon Government, Archaeology Program, Department of Tourism and Culture Mathewes, Rolf W.; Simon Fraser University, La Farge, Catherine; University of Alberta, Department of Biological Sciences Martel, André ; Canadian Museum of Nature Heintzman, Peter; University of California Santa Cruz, Department of Ecology and Evolutionary Biology Shapiro, Beth; University of California Santa Cruz, Ecology and Evolutionary Biology
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	Quaternary, Beringia, Bison, Holocene, Yukon

SCHOLARONE™
 Manuscripts

1

1 **A middle Holocene steppe bison and paleoenvironments from the Versleuce Meadows,**
2 **Whitehorse, Yukon, Canada**

3 Grant D. Zazula^{1*}, Elizabeth Hall², P. Gregory Hare³, Christian Thomas⁴, Rolf Mathewes⁵,
4 Catherine La Farge⁶, André L. Martel⁷, Peter D. Heintzman⁸, Beth Shapiro⁹

- 5
- 6 1. Yukon Government, Palaeontology Program, Department of Tourism and Culture,
7 Whitehorse, Yukon, Canada, Y1A 2C6, ph. 1-867-667-8089; grant.zazula@gov.yk.ca
- 8 2. Yukon Government, Palaeontology Program, Department of Tourism and Culture,
9 Whitehorse, Yukon, Canada, Y1A 2C6, ph. 1-867-667-8880; elizabeth.hall@gov.yk.ca
- 10 3. Yukon Government, Archaeology Program, Department of Tourism and Culture, Whitehorse,
11 Yukon, Y1A 2C6; greg.hare@gov.yk.ca
- 12 4. Yukon Government, Archaeology Program, Department of Tourism and Culture, Whitehorse,
13 Yukon, Y1A 2C6; Christian.thomas@gov.yk.ca
- 14 5. Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia,
15 Canada, V5A 1S6; matthewes@sfu.ca
- 16 6. Department of Biological Sciences, University of Alberta, Edmonton, Alberta, T6G 2E9;
17 clafarge@ualberta.ca
- 18 7. Malacology Unit, Research and Collections, Canadian Museum of Nature, P.O. Box 3443,
19 Ottawa, Ontario, Canada, K1P 6P4; AMARTEL@mus-nature.ca

20

2

1 8. Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa
2 Cruz, CA 95064, USA; peteheintzman@gmail.com

3 9. Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa
4 Cruz, CA 95064, USA; beth.shapiro@gmail.com

5

6 **Corresponding author:**

7 Grant D. Zazula Yukon Government, Palaeontology Program, Department of Tourism and
8 Culture, Whitehorse, Yukon, Canada, Y1A 2C6, ph. 1-867-667-8089; fax: 1-867-667-5377;
9 grant.zazula@gov.yk.ca

10

11

12

13

14

15

16

17

18

1 **Abstract**

2 A partial skeleton of a bison was recovered during residential house construction in Whitehorse,
3 Yukon, Canada. The specimen represents a young (estimated 6 year old) bison individual that
4 died, was partially scavenged by carnivores, and subsequently buried by calcareous silt sediment
5 in a pond or small lake during the middle Holocene, ~ 5,400 years ago. Palaeoenvironmental
6 data, including molluscs, pollen, vascular plant, and bryophyte macrofossils demonstrate that the
7 small waterbody was surrounded by white spruce dominated boreal forest. Morphometric
8 analysis of the skeleton reveals that its taxonomic affinity is ambiguous, likely due to it
9 representing an ontogenetically young individual, though it does share some cranial and horn
10 core characteristics of named species such as *Bison occidentalis* or *Bison priscus*. Mitochondrial
11 genomic data confirm that this bison belongs to Clade 2A (northern clade), which represents
12 Pleistocene steppe bison (*B. cf. priscus*) in Beringia through the Holocene and is not represented
13 in living bison species (Shapiro et al. 2004; Heintzman et al. 2016). These data further
14 demonstrate that northern steppe bison population survived the late Pleistocene extinction event,
15 persisted locally in southern Yukon into the Holocene, and are best characterized as a species
16 with a high degree of morphological variability and ecological flexibility.

17

18

19

20

21

1 Introduction

2 Bison (*Bison*) dominate Pleistocene mammal faunas from the unglaciated regions of interior and
3 northern Alaska and Yukon, the area commonly referred to as eastern Beringia (Guthrie, 1990;
4 Harington, 2011). Fossil records suggest that once bison initially dispersed from Asia across the
5 exposed Bering Isthmus 195-135 thousand years ago (Froese et al. 2017), they subsequently
6 established themselves as the most abundant, large Pleistocene ungulate in many parts of North
7 America (Guthrie 1970, 1990, McDonald 1981). Estimates suggest that over 80% of typical late
8 Pleistocene large mammal faunas in eastern Beringia are composed of steppe bison (*B. priscus*)
9 remains (Guthrie 1990). In contrast, remains of bison dating to the Holocene in glaciated regions
10 of northern North America are relatively rare, often poorly preserved and generally lack the
11 morphologically diagnostic cranial characteristics that are often used to make taxonomic
12 identifications (Stephenson et al. 2001). As such, there is limited information on eastern
13 Beringian bison populations following deglaciation in comparison to more common subarctic
14 ungulates such as caribou (*Rangifer tarandus*) (Farnell et al. 2004). Coupled with generally poor
15 preservation conditions in subarctic soils, cultural practices such as fracturing long bones to
16 remove marrow likely resulted in significant under-representation of identified bison remains
17 from Holocene zooarchaeological assemblages.

18

19 Because of their ubiquity and abundance in the fossil, archaeological, and historical records,
20 bison are one of the most extensively studied North American Quaternary mammal species
21 (Skinner and Kaisen 1947; McDonald 1981; Guthrie 1990). However, their formal taxonomy is
22 rather confusing because of numerous named, but likely nominal species, largely defined based

on inconsistent cranial and horn core metrics and morphology. However, there has been a trend towards informal taxonomic lumping, especially in regards to Pleistocene bison from eastern Beringia which are now generally considered to all represent the Holarctic species *B. priscus* (Bojanus 1827) *sensu lato* (Guthrie 1970, 1990). Further, some have suggested based on morphological criteria that Holocene aged bison in Alaska and Yukon represent the extant wood bison (*B. bison athabasca*; Stephenson et al., 2001).

Genetic studies based on mitochondrial DNA from Quaternary bison have shown that extant bison, *B. bison*, arose during the last glacial maximum from ancestors situated south of the continental ice sheets, and that *B. bison* mitochondrial variation falls within a much more diverse bison complex (including the steppe bison, *B. priscus*) that occurred both north and south of the ice sheets during the late Pleistocene (Shapiro et al. 2004; Heintzman et al. 2016; Froese et al. 2017). As mitochondrial DNA sequences can distinguish *B. bison* from other bison taxa, they can be used to infer the systematic status and phylogeographic history of enigmatic bison specimens. The mitochondrial north-south differentiation between Pleistocene bison species is loosely supported by morphological data as well, with *B. priscus* found in the north and *B. antiquus* in the south (Shapiro et al. 2004; Wilson et al. 2008). The integration of both morphological and genetic data for ancient bison are key to resolving questions about bison dispersal history and phylogenetics especially regarding the dramatic changes following the Pleistocene-Holocene transition (Wilson et al. 2008). In Yukon and Alaska, there are relatively few well-preserved Holocene bison specimens to address questions related to the taxonomy, evolution and spatial-temporal distribution of bison in this region. Although it is beyond the

scope of the present paper to resolve, studies of North American bison are clearly hindered by the inconsistency of morphological and genetic data and future work should attempt to address the numerous species names that are presently recognized in the literature.

This paper reports a well-preserved partial skeleton of a bison including the cranium and many of the post-cranial elements from the Versleuce Meadows site in the formerly glaciated region of Whitehorse, southern Yukon, Canada (Figure 1). The specimen is described, identified, radiocarbon dated, and placed into a mitochondrial genome-based phylogenetic context. We also evaluate associated palaeoenvironmental proxies to further shed light on the palaeoecological context of this post-glacial bison in northern Canada. The primary topics addressed in this paper are:

- 1) Can morphometric data be used to confidently assign this specimen into one of the previously defined species of Quaternary bison?
- 2) What is the mitochondrial genetic affinity of the Versleuce Meadows bison?
- 3) When and how did this bison die and become preserved at Versleuce Meadows?
- 4) What was the environment surrounding Versleuce Meadows when this bison died?
- 5) What is its significance for bison taxonomy and history in the north?

1 **Materials and Methods**

2 **Study site and regional context**

3 The study site (Figure 1) is located within the city of Whitehorse, Yukon in an area known
4 locally as “Versleuce Meadows” (60° 46’ 21.9” N, 135° 08’ 26.3” W) in the neighbourhood of
5 Porter Creek. Historically, this shallow low-lying area (elevation: 715 m a.s.l.) consisted of
6 agricultural hayfields surrounding a small ephemeral pond or slough. In recent decades,
7 extensive topsoil mining in the meadow has removed the surface substrate for local gardener
8 usage. The fossil discovery site is on the periphery of a low-lying meadow, immediately
9 downslope from a ridge that forms the northern edge of a small basin. The entire area was
10 glaciated during McConnell Glaciation of the northern Cordilleran ice sheet (Figure 1), and
11 became ice free after ~11,000 years ago (Bond 2004).

12
13 The study site is situated in the Yukon Southern Lakes ecoregion (Yukon Ecoregions Working
14 Group 2004), generally characterized by rounded montane summits and broad valleys resulting
15 from the effects of Pleistocene glaciation. The vegetation in the region is dominated by open
16 coniferous and mixed forests, with lodgepole pine (*Pinus contorta*) dominating dry and recently
17 burned sites, whereas white spruce (*Picea glauca*) forests are more common on active
18 floodplains, other mesic to moist sites and serally advanced sites. Mixed aspen (*Populus*
19 *tremuloides*) and white spruce are common on sites with fine soils with variable ground shrubs
20 and lichen. Open areas and steep south and west facing slopes at low elevation commonly

1 include grasslands, while depressions have alkaline ponds and small lakes. Black spruce (*Picea*
2 *mariana*) on permafrost soils are rare in the region at low elevation.

3
4 Paleoecological data from southern Yukon demonstrate the changes in vegetation cover
5 following the retreat of the Cordilleran ice sheet after 11,000 years ago (Cwynar and Spear 1995;
6 Gajewski et al. 2014). Immediately after deglaciation, the landscape was covered by a herb
7 tundra vegetation with abundant grass (Poaceae), sage (*Artemisia*), and forbs, transitioning to a
8 more shrub-dominated tundra environment with abundant birch (*Betula*), some trembling aspen
9 (*Populus tremuloides*), and juniper (*Juniperus* spp.) in the understory, and which dominated until
10 around 10,000 or 9,000 years ago (Gajewski et al. 2014). White spruce forests were abruptly
11 established after this time and continued to be abundant for much of the Holocene. Significant
12 changes in regional vegetation composition occurred between roughly 6,000 and 5,000 years
13 ago, with a shift towards a predominance of black spruce over white spruce and with major
14 increases in alder (*Alnus crispa*). Pine trees replaced spruce as the dominant tree of open
15 woodlands around Whitehorse by around 2,000 years ago, following dispersal from the south
16 and east (Cwynar 1988; Cwynar and Spear 1995).

17
18 The composition of the present-day large mammal fauna in the region is the result of the
19 dramatic changes during the Pleistocene-Holocene transition, which included the extirpation of
20 many species of the “mammoth-steppe” such as woolly mammoth (*Mammuthus primigenius*),
21 horses (*Equus* spp.), and saiga antelope (*Saiga tatarica*). This period of faunal turnover also saw

the initial dispersal of Asian migrants such as elk (*Cervus elaphus*), moose (*Alces alces*), and people (*Homo sapiens*) into North America (Guthrie 2006). Populations of elk, bison and caribou locally survived in Yukon through this period of dramatic ecological turmoil and remained in the subarctic and arctic through much of the Holocene. Fossil and ethnohistorical evidence demonstrates that native bison populations in Yukon were regionally extirpated roughly 400 years ago (Stephenson et al. 2001; Heintzman et al. 2016). Present day populations of elk and wood bison (*Bison bison athabasca*) are now common in the boreal forest of southern Yukon and are the result of human-mediated reintroductions during the mid-20th century.

Field Excavation

On the morning of April 26, 2012, Whitehorse resident Craig Duncan encountered several large mammal bones, while excavating with heavy equipment to construct a basement on his residential property. He gathered these bones and brought them to the Government of Yukon, Heritage Resources Unit office. The bones were all readily identified as those from a bison. This small collection included a cervical vertebra, a thoracic vertebra, left metacarpal, right metatarsal and fragments of both a left and right tibia. Since it appeared that all these elements belonged to a single individual, were well-preserved, and discovered in approximate anatomical position, a small crew was immediately assembled to further investigate the site.

After initial investigation in the afternoon of April 26, 2012, it became evident the area being excavated on the periphery of Versleuce Meadow (Figure 2a) was part of an ancient lacustrine

environment. The initial trenches and excavation conducted by the equipment revealed fine grained, horizontally bedded tan colored silt (Figure 2b) with abundant mollusks at depth (Figure 2c). The excavated and disturbed areas were examined by a small group of palaeontologists and archaeologists in an attempt to recover more faunal material and determine the extent of the fossil site. A number of other elements were discovered immediately, including a pair of unfused bison innominates found together, upside down. A large-scale salvage excavation then commenced, employing the use of a backhoe that was available on site, to further reveal the site stratigraphy and look for more of the bison remains. The backhoe was used to slowly excavate a large area until a bone bed was discovered at a depth of approximately 150 cm below the surface (Figure 2d). Excavation of the remaining elements of the bison skeleton was conducted with shovel and trowel the next day with the assistance of local community volunteers. As the salvage excavation had brought construction activities to a halt it was determined that the priority was to recover as much of the faunal material as quickly as possible, with less emphasis on details of skeletal element provenience and detailed site recording. This excavation resulted in the recovery of a partial skeleton of an individual bison and an abundance of palaeoecological samples. When shovel and trowel excavations were concluded, a much larger area was subsequently excavated by backhoe to ensure that no other bones had been missed.

Due to time constraints of the field project, only one day of hasty salvage excavation was performed at the Versleuce Meadows site. Excavation of the partial skeleton was focussed on the bone bed that occurred roughly 150 cm below the surface. The skeleton was largely confined to a horizontal plane, roughly 550 cm in length. The skeleton was roughly orientated in anatomical

11

1 position with the cranium upside down. After removal of fossils, further reconnaissance was
2 performed with the backhoe below and laterally, but failed to yield any further remains. Samples
3 of the sediments associated with the bison skeleton were collected at the site for subsequent
4 analysis. All the faunal material was taken back to the Yukon Government Heritage Resource
5 Unit offices for further study.

6

7 **Metric osteological analysis**

8 Mensurational data were taken on the bison fossils using digital calipers and following the
9 methods of van den Driesch (1976), Skinner and Kaisen (1947) with others from McDonald
10 (1981) and Wilson et al. (2008). Age-of-death estimation for the bison was established following
11 the methods of Duffield (1973).

12

13 **Radiocarbon analysis**

14 A subsample of cortical bone from the diaphysis of the right tibia, specimen YG 422.21, was
15 submitted to the KECK University of California Irvine Accelerator Mass Spectrometer
16 (UCIAMS) laboratory for radiocarbon dating. Procedures used in the preparation of the
17 ultrafiltered (> 30 kD) bone collagen gelatin are detailed in Beaumont et al. (2010). Detailed
18 description of the combustion and graphitization protocols and the operation of the UCIAMS
19 instrument are described in Beverly et al. (2010).

1 **Phylogenetic analysis of mitochondrial genomes**

2 To diagnose the taxonomic identity of the Versleuce Meadows bison and place it within a
3 mitochondrial phylogenetic framework (Shapiro et al. 2004; Heintzman et al. 2016), a sample of
4 specimen YG 422.21 (right tibia shaft) was sent to the specialized ancient DNA facility at the
5 University of California, Santa Cruz. DNA extraction, library preparation, mitochondrial genome
6 target enrichment, and sequencing were all performed during a previous study using approach
7 four (Heintzman et al. 2016). We constructed a consensus sequence following Heintzman et al.
8 (2016), except that we retained the entire mitochondrial genome (GenBank accession number
9 MF134653). We compared the Versleuce Meadows bison mitogenome sequence to a previously
10 published alignment (Froese et al. 2017), which consists of mitochondrial genome sequences
11 from 46 North American or Siberian bison and four yak. We performed a phylogenetic timetree
12 analysis using BEAST (v1.8.1; Drummond et al. 2012) utilizing parameters outlined in Froese et
13 al. (2017). Briefly, we partitioned the data, used a strict molecular clock informed by radiocarbon
14 or stratigraphic data (with independent rates for each partition), and a skygrid coalescent prior
15 (Gill et al. 2013). We ran two analyses of 60 million iterations each, sampling trees every 3,000
16 iterations, and discarding the first 10% as burn-in. Sampled trees from both analyses were
17 combined and summarized as a maximum clade credibility (MCC) tree.

18

19 **Paleoecological analyses**

20 Two samples of calcareous lake sediment in association with the bison skull were analyzed for
21 pollen, spores, and algal remains to help assess the local paleoenvironment. Subsamples of 5 mL

1 were treated by standard processing techniques (Faegri et al. 1989), involving HCl, HF, KOH,
2 and acetolysis after adding exotic *Eucalyptus* marker pollen grains (1 tablet, $16,180 \pm 1460$
3 grains) to each sample in order to calculate palynomorph concentrations. Residues were mounted
4 in glycerine jelly and identified and counted at 400× magnification. Pollen sums are 540 and 571
5 grains for % calculations. As expected, the sediment analyzed was strongly calcareous, and
6 reacted violently with dilute HCl. Most of the volume, however, was siliceous and needed two
7 HF treatments and hot HCl to dissolve. White and black spruce pollen types were separated
8 using the morphological criteria of Hansen and Engstrom (1985).

9
10 For other macrofossil analyses, sediments collected in association with the bison fossils were
11 soaked in water to disaggregate the material and washed through a 425 micron mesh sieve. The
12 retained material was picked through under a dissecting microscope. Vascular plant macrofossils
13 were identified with the aid of comparative modern plant material at the Yukon Heritage
14 Resources Unit, as well as publications such as Lévesque et al. (1988). Taxonomic nomenclature
15 follows Cody (1996).

16
17 Bryophyte macrofossils were dispersed and separated from detrital material (unidentifiable
18 rootlets and degraded graminoid tissue), then examined initially in water under a dissecting
19 microscope. Bryophyte fragments were then compiled into distinct taxa, further examined with
20 the aid of a compound microscope and identified using Hedenäs (2014 a,b). Species
21 determinations were confirmed with comparison to extant material available in the Cryptogamic

Herbarium (ALTA), Department of Biological Sciences, University of Alberta. Exceptional preservation was noted for all species determined. Vouchers are deposited in the Cryptogamic Herbarium (ALTA).

Sieved gastropods and bivalves were brought to the malacology laboratory at the Canadian Museum of Nature (CMN), in Gatineau Québec, where the National mollusc collections are located. The specimens were examined using a stereomicroscope and identification was conducted using Clarke (1981) and the national mollusc collection at the CMN. Specimens are generally well-preserved, with the surface texture of the shell intact, thus facilitating the identification. Nomenclature and taxonomical authority follow the Integrated Taxonomical Information System (ITIS 2016).

Results

Site Stratigraphy

Excavation with the backhoe and shovels revealed a consistent stratigraphic profile across the site (Figure 2). Considering the area underwent extensive removal of surficial sediments and soil at various times in the past, it is difficult to assess the nature of the uppermost sediments. The observed stratigraphy is as follows:

0-20 cm (below surface): highly disturbed modern sod and organic rich soil

20-70 cm: well sorted tan colored silt, no clasts, rare gastropods and bivalves, modern roots from the surficial vegetation

15

1 70-100 cm: well sorted tan colored silt with some grey beds of clay, increasing abundance of
2 mollusc shells with depth, sediments reacted vigorously to HCl
3 100 – 170 cm: tan colored silt with grey clay beds, beds of marl consisting of dense
4 concentrations of mollusc shells, increasing shell content with depth in the sediment, rare clasts,
5 sediments reacted vigorously to HCl. This is the level which contained the bison fossils.

6
7 A partial skeleton of a bison (Figure 3; Table 1, 2) was recovered from within the silty-marl
8 sediments approximately 150-170 cm below the surface. The bison was in approximate death
9 position. All skeletal material is accessioned in the collections of the Yukon Palaeontology
10 Program (specimen acronym YG) in Whitehorse, Yukon. Description of the skeletal remains are
11 as follows.

12
13 **Systematic vertebrate palaeontology**

14 Order: Artiodactyla

15 Family: Bovidae

16 *Bison* sp.

17
18 *Cranium*. Referred specimens: YG 422.1 with both horn cores, LM1-3, LPM2-4, RM1-3, RPM3.

19 The cranium from the Versleuce Meadows bison is largely complete (Figure 4; Table 1), though
20 much of the maxilla is broken into numerous small pieces. The maxillary teeth were recovered in

the sediment surrounding the cranium. The cranium has a relatively narrow, flat frontal and slightly domed, broad nasal. The orbits are roughly parallel to the frontal plan. The horn core bases are triangular in cross section, and slightly rotated rearward. The horn cores have a dorsal groove on the proximal half and they taper gradually toward the tip which are elliptical in cross section. The anterior-posterior margin of the horn cores are weakly concave and proximally downswept with tips rising about 60 mm above the frontal plane. The horn cores are directed laterally with the distal halves backswept and with tips terminating slightly behind the occipital plan. The growth of the horn cores is relatively straight on an arched axis with negligible distal twist. The rugose burrs at the base of the horn cores have weakly developed exostoses and the frontal midline suture of the cranium is not fused, suggesting a young individual.

Mandibles. Referred specimens: YG 422.2 (R) with m1-3, i1-3, c1 and pm4 (Figure 5a); YG 422.3 (L) with m1-3 and pm2-3.

Both mandibles halves were recovered near to the cranium and are largely complete, though are not fused at the symphysis. The incisor teeth were found in the sediment next to the mandibles.

Vertebral column and ribs. Referred specimens: *atlas*: YG 422.4; *axis*: YG 422.5; *cervical vertebra*: YG 422.24 (3rd cervical: complete though centrum epiphysis is not fused to body; Figure 5b); YG 422.7 (4th cervical: transverse processes broken on distal ends); YG 422.8 (5th cervical: complete); YG 422.6 (6th cervical: complete); YG 422.9 (7th cervical: complete); *thoracic vertebra*: YG 422.25 (unfused portion of posterior centra epiphysis); YG 422.26

1 (spinous process broken on distal end, anterior epyphysis of centra missing; Figure 5c); YG
 2 422.27 (spinous process broken on distal end); YG 422.85 (fragment of unfused distal centrum
 3 epiphysis); *innominates* (Figure 5d): YG 422.32 (L: incomplete and broken in 5 pieces); YG
 4 422.31 (R: incomplete and broken in 4 pieces); *ribs*: (Figure 5e) YG 422.12; YG 422.13; YG
 5 422.14; YG 422.15; YG 422.16; YG 422.17; YG 422.17; YG 422.18; YG 422.19; YG 422.22;
 6 YG 422.13.

7 The vertebral elements are consistent in that they exhibit only partial fusion of the centra or in
 8 some, no fusion at all. The lack of complete fusion in the vertebra further suggests this was a
 9 young individual. Only eleven individual broken rib fragments were recovered. The ribs were all
 10 recovered *in situ* immediately below the left ulna and radius. The innominates are not fused and
 11 broken in several pieces, with both ilia showing evidence of carnivore gnawing. The innominates
 12 were recovered approximately 4 metres from the rest of the skeleton.

13

14 *Forelimbs*. Referred specimens: *metacarpal* (Figure 5f): YG 422.33 (R) and YG 422.40 (L);
 15 *proximal phalanx (pairs)*: YG 422.47 (R)- YG 422.48 (L) and YG 422.45 (R) – YG 422.46 (L);
 16 *medial phalanx (pairs)*: YG 422.51 (R) – YG 422.52 (L) and YG 422.49 (R) - YG 422.50 (L);
 17 *distal phalanx (pairs)*: YG 422.55 (R) - YG 422.46 (L) and YG 422.53 (R) - YG 422.54 (L);
 18 *magnum*: YG 422.34 (R) and YG 422.41 (L); *unciform*: YG 422.35 (R); *scaphoid*: YG 422.36
 19 (R) and YG 422.42 (L); *lunar*: YG 422.37 (R) and YG 422.42 (L); *cuneiform*: YG 422.38 (R)
 20 and YG 422.38 (L); *trapezoid*: YG 422.39 (R); *radius* (Figure 5g): YG 422.85 (L) and YG
 21 422.30 (R); *ulna*: YG 422.84 (L) and YG 422.29 (R); *humerus*: YG 422.10 (L; (Figure 5h)) and
 22 YG 422.11 (R); *scapula*: YG 422.28 (L; Figure 5i).

The distal forelimb elements are complete, except for missing unciform and trapezoid on the left side. The radius and ulna are not fused; both radii are complete, the left ulna is complete, while the right ulna has been broken into three pieces postmortem. Both humerii are largely complete though substantial carnivore gnawing has completely eliminated the proximal end of the right humerus, while the proximal end of the left humerus is partially removed. Several broken portions of the left scapula were recovered and the distal margins seem to have been removed by carnivores.

Hindlimbs. Referred specimens: *metatarsal* (Figure 5j): YG 422.58 (R) and YG 422.57 (L); *proximal phalanx* (pairs): YG 422.66 (R) - YG 422.67 (L) and YG 422.68 (L); *medial phalanx*: YG 422.69 (R) – YG 422.70 (L) and YG 422.71 (R) – YG 422.72 (L); *distal phalanx*: YG 422.73 (L) and YG 422.74 (R) – YG 422.75 (L); *naviculo-cuboid*: YG 422.62 (R) and YG 422.59 (L); *lateral malleolus*: YG 422.63 (R) and YG 422.60 (L); *astragalus*: YG 422.64 (R); *calcaneum*: YG 422.65 (R) and YG 422.61 (L); *tibia*: YG 422.20 (L) and YG 422.21 (R; Figure 5k).

The distal hindlimb elements are complete, except for missing right distal (3rd) phalanx, a left proximal (1st) phalanx, and a left astragalus. Both femora are missing. Both tibias are mostly complete, though there is substantial evidence of carnivore gnawing on the right tibia, and minimal gnawing marks on the left. Midshaft breakage on the left tibia seems to be from excavation by heavy equipment, though the pattern of breakage on the right tibia seems to suggest natural postmortem damage as warping of the bone prevented the broken pieces from fitting back together smoothly.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Other elements. Referred specimens: *sesamoids*: YG 422.76; YG 422.77; YG 422.78; YG 422.79; YG 422.80; YG 422.81; YG 422.82; YG 422.83.

Ontogeny

Several skeletal features enable an age estimation for this bison based on known skeletal development of extant European bison (*Bison bonasus*) (Duffield 1973). The distal epiphyses of the radius are fused (Figure 5g), and the proximal head of the humerus is fused (Figure 5h), which generally occurs early in the sixth year. Since some of the vertebral epiphyses are in early stages of fusion, especially in the cervicals, though the thoracics remain largely fused, it is likely the animal died before the start of its seventh year. This assessment is supported by the lack of frontal fusion on the cranium (Figure 4), and minimal development of exostoses on the rugose burrs of the horn core bases. We also heed Duffield’s (1973) cautions with determination of sex since this is an immature individual and there may be considerable overlap in several measurement indices between young males and females.

Radiocarbon date

Ultrafiltered collagen from the bone sample yielded a radiocarbon age of 4710 ± 15 ^{14}C yr BP (UCIAMS 114030; Heintzman et al. 2016). Calibration of this radiocarbon date using the

IntCal13 dataset (Reimer et al. 2013 Radiocarbon) places the bison between 5578 and 5327 cal. yr BP (midpoint = 5452 cal yr BP) at 95.4% confidence.

Phylogenetic analysis

Separate analyses of bison mitochondrial genomes confirm that both subspecies of extant bison (*B. bison*) form a single, well-supported cluster (Clade 1A) within the greater diversity of North American bison (Shapiro et al. 2004; Heintzman et al. 2016; Froese et al. 2017; Figure 6). To confirm the phylogenetic placement of the Versleuce Meadows bison, we added this newly generated complete mitochondrial genome sequence to a previously published data set of North American and Siberian bison mitochondrial genomes (Froese et al. 2017) and constructed a new phylogeny. The Versleuce Meadows bison clusters with a Clade 2A bison from Edmonton (PH027). This is consistent with a previous analysis that included only a ~600 base pair sequence of the mitochondrial control region (Heintzman et al. 2016) and confirms the placement of the Versleuce Meadows bison within Clade 2A. This phylogenetic position indicates that Versleuce Meadows bison did not belong to the mitochondrial lineage of bison that eventually gave rise to extant bison subspecies.

Paleoenvironmental Data

Palynomorph concentrations are very low (Table 3) but preservation was generally good. The pollen spectra are dominated by arboreal taxa such as white and black spruce, birch and alder shrubs, while sedges (Cyperaceae) dominate the herb component.

1 Plant seeds, fruits and vegetative parts from vascular plants are not abundant within the silt
2 sediments, though are generally well-preserved (Figure 7). The plants are dominated by aquatic
3 plants that presently inhabit littoral edges of ponds and lakes in the region at present, such as
4 pondweeds (*Potamogeton* sp.; Figure 7a) and common mare's tail (*Hippuris vulgaris*; Figure
5 7b), and nearshore sedges (*Carex* spp.; Figure 7 c,d) with rare *Elocharis* sp. (Figure 7e)).
6 Moderate amounts of spruce (*Picea* sp.; Figure 7f) needles were recovered, most being charred.
7 Rare terrestrial herbaceous taxa were recovered as well, including goosefoot (*Chenopodium* sp.;
8 Figure 7g), buttercup (*Ranunculus aquatilis* type), and knotweed (*Polygonum* sp.). In addition to
9 vascular plants, remains of the charophyte algae (*Chara* sp.) and statoblasts of the aquatic
10 bryozoan *Cristatella mucedo* (Figure 7h) were common.

11
12 The bryophyte macrofossils recovered from the Versleuce Meadows sediments include three
13 species (Figure 7 i,j,k). Excellent preservation included intact leaves with leaf habit and
14 attachment to stems clearly visible, in addition to detached leaves or denuded stems. *Scorpidium*
15 *scorpioides* (Hedwig) Limpricht (Figure 7i) – a submerged to emergent hydric taxon, often had
16 leaves with shredded apices, yet the ovate leaf shape; linear, smooth leaf cells, entire margins
17 and lack of a costa and differentiated alar cells are diagnostic of this common circumpolar arctic-
18 boreal wetland taxon. *Calliergon giganteum* (Schimper) Kindberg (Figure 7j) is a common
19 circumpolar arctic-boreal emergent wetland species that had broad, strongly concave leaves with
20 well preserved abruptly inflated alar cells reaching the costa and a single unbranched costa that
21 reaches the apical region of the leaf. The rounded apex and linear, smooth leaf cells help to
22 determine this taxon. *Drepanocladus longifolius* (Wilson ex Mitten) Brotherus ex Paris (Figure

7k) is one of the most common taxa preserved in sediments at northern sites (as *D. crassicostatus* Janssens in Janssens 1983). This is a distinct taxon determined by its long excurrent and stout costa. It has alar cells that form 1-2 rows along the leaf base. Its preferred habitat also includes nutrient-rich waters of pools, lake margins, oxbows, or depressions of wet meadows (Hedenäs 2014; Janssens 1983). Within the subsample *S. scorpioides* was the most abundant taxon (** 50-100%); *Drepanocladus longifolius* (** 25-50%) and *Calliergon giganteum* (* 0-25%).

An abundance of well-preserved molluscan taxa, including at least 10 species of freshwater bivalves and gastropods (Table 4; Figure 7 l,m,n), were recovered from the silty marl collected in association with the bison skeleton. All species of bivalves and gastropods found are currently found in Canada and are typical of a pond or lake environment – although some of those species also currently inhabit running water.

The largest molluscan species found is the frigid lymnaea (sometimes called Alaskan lymnaea), *Lymnaea atkaensis* and is restricted at present to British Columbia, the Northwest Territories and the Yukon Territory (Clarke 1981). It prefers cold, oligotrophic lakes and appears to have similar habitat characteristics as *Sphaerium nitidum*. Many complete and unbroken large adult specimens of *Lymnaea atkaensis* were recovered, suggesting that deposition of the shell material in the pond or lake occurred with little reworking of the sediment or mixing of the sediments. All the gastropods belong to species that are typically found in the littoral or sublittoral zone of ponds or lakes. They are known to crawl on submerged macrophytes, sometimes on the fine

sediment (sand, silt, mud), occasionally crawling on sunken debris, and less frequently on rocks or stones. Among the bivalves, the two species of *Pisidium* (Figure 7m) and the one species of *Sphaerium* collected (Table 4) belong to the Sphaeriidae family and are commonly found in small or large ponds, or lakes. They live normally just below the surface of the sediment on the bottom of the pond or lake, and are capable of moving through the sediment, using their muscular foot. These little clams use their long siphonal tubes, feeding on microscopic plants and detritus deposited on the bottom (deposit feeders). Many/most of the mollusc species found are usually associated with cold, hard waters, although some, especially *Pisidium casertanum* occurs in wide range of pH (5.50-8.34), total alkalinity (0-280 mg CaCO₃/L), total hardness (10-332 mg CaCO₃/L), and Ca hardness (Mackie and Flippance 1983).

Discussion

Bison Taxonomy

Bison taxonomy has focused traditionally on adult male cranial and horn core characteristics and measurements (Skinner and Kaisen 1947), while the post-cranial skeleton is largely considered to be unreliable for taxonomic assignments (McDonald 1981). The relative completeness of the Versluis Meadows bison cranium (Figure 4) enables comparison with other reported bison fossils, to help resolve the taxonomy of this individual. However, caution must be taken with this assessment, because cranial and post-cranial features demonstrate this is a young individual. Horn core measurements on a young bison are not directly comparable to published data from mature adults and may preclude any definitive taxonomic assignment (*sensu* McDonald 1981).

1 Even with a complete adult cranium, there are few diagnostic characters that can be used to
2 positively distinguish one named bison taxon from another, and there is demonstrated
3 mensurational overlap between most of the recognized species.

4
5 Superficially, the cranial characters and measurements (Table 1; Figure 4) from the Versluce
6 Meadows bison demonstrate a possible morphological affinity with a variety of named bison
7 species, including *Bison occidentalis* and *B. priscus* (Skinner and Kaisen 1947; McDonald
8 1981). Some of the characters considered important are the triangular horn core bases that
9 gradually taper toward the tip which are elliptical in cross section, a dorsal groove along the
10 proximal part of the horn core, proximally down-swept and distally rear-swept horn cores,
11 relatively flat frontal, and low lying orbits. Considering overall size, the Versleuce Meadows
12 bison fits within the lower size ranges reported for *Bison priscus*, and is well below the range for
13 the highly curved and robust horned *Bison alaskensis* (McDonald 1981). Based purely on cranial
14 measurements, the Versleuce Meadows specimen also fits within the range reported for large
15 sized extant wood bison (McDonald 1981). In recognition that this specimen represents a sub-
16 adult, and the inherent problems with bison taxonomy, inconsistency in morphological species
17 diagnoses, and substantial inter-species mensurational overlap, we remain conservative in our
18 approach and are hesitant to commit to any assignment apart from *Bison* sp. based purely on
19 morphological or mensurational grounds.

20 However, genomic data enable us to place this specimen within a phylogenetic model that
21 includes numerous other ancient bison from this region, assisting greatly in a taxonomic
22 assignment (Shapiro et al., 2004; Heintzman et al., 2016; Froese et al., 2017). Mitochondrial

genomic data from the Versleuce Meadows bison confirm that it belongs to Clade 2A, which includes Pleistocene *B. cf. priscus* from Beringia and northwest Canada through the Holocene and is not represented in living bison species (Shapiro et al. 2004; Heintzman et al. 2016). For this study, we refer all North American bison outside of Clade 1A as *Bison cf. priscus* regardless of morphological affinity, but acknowledge that more work needs to be done to further resolve the inconsistency between morphological and genetic data in the study of North American bison. Given the combination of genomic data placing the Versleuce Meadows specimen in the Clade 2A, while recognizing that some of the cranial characteristics and measurements of this young individual are similar to descriptions reported for steppe bison, a taxonomic assignment of *B. c.f. priscus* is warranted.

Importantly, the Versleuce Meadows bison did not belong to the lineage that gave rise to present-day wood and plains bison (Figure 6). All living bison have mitochondrial haplotypes within Clade 1A. The geographical origins of Clades 2A and 2B are of interest, as this is relevant to the timing and direction of bison dispersal between Beringia and the North American midcontinent after the recession of glacial ice (Heintzman et al. 2016). While the location of origin of the common ancestor of Clade 2B cannot be determined with confidence, phylogeographic models provide strong support for a northern or Beringian origin of Clade 2A, to which the Versleuce Meadows bison and other Holocene steppe bison from Beringia belong. We acknowledge that in the absence of nuclear genetic data, the degree of breeding between resident bison from Beringia and northward dispersing bison during post-glacial times after the opening of the ice free corridor remains unknown (Heintzman et al., 2016). While it is not possible to know how long this lineage of steppe bison persisted into the Holocene, the young radiocarbon age on the Versleuce Meadows specimen makes it one of the last surviving members.

1

2 **Versleuce Bison Skeleton Taphonomy**

3 The sediments, stratigraphy and radiocarbon age from the Versleuce Meadows site suggests that
4 the animal died in a small lake ~5,400 years ago. With much of the skeleton recovered in
5 partially articulated death position, there was minimal post-mortem movement of many of the
6 bones. Considering the presence and excellent preservation of the distal limb elements, including
7 metatarsals, metacarpals, carpals sesamoids and phalanges, it is possible that the animal's lower
8 limbs became mired in the lake bottom mud immediately prior to its death (Figure 4, 5). The
9 breakage on the distal tibia shafts perhaps occurred as the animal struggled to free itself and
10 collapsed. Much of the anterior half of the skeleton was recovered *in situ* in a confined area
11 which seemingly was the location of the animal's death (cranium, mandible, radii, ulnae, cervical
12 and thoracic vertebrae, rib fragments, metacarpals and carpals), while much of the posterior parts
13 of the skeleton (pelvis, tibia, metatarsals) were discovered a few meters away from the main
14 skeleton and accidentally discovered and recovered during the initial construction and trenching
15 at the site. Carnivores clearly were able to scavenge the more proximal parts of the post-cranial
16 skeleton as demonstrated by substantial gnawing and removal of bone on the proximal humeri
17 and tibiae. These scavengers seemingly could not access the fairly complete distal limb elements
18 (metapodials, phalanges) because they were buried in the lake mud, while other parts of the torso
19 and proximal limbs were exposed. The highly fragmented nature of the ribs and limited
20 preservation of the vertebral column suggests heavy scavenging of the internal organs and guts
21 within the abdominal cavity. Since we were not able to recover some elements such as the
22 femurs, while the pelvis was recovered some distance from the rest of the skeleton, this suggests

that much of the carcass decomposed *in situ* while the forelimbs were stuck in the lake mud, and the posterior parts were scattered by carnivores or floated away. There is no evidence of human hunting or butchering on the bison remains or at the site.

Paleoenvironment

Paleoecological data recovered from sediments in association with the bison skeleton provides details of the plant communities, molluscs and limnological conditions in and around Versleuce meadow ~5,400 years ago. The calcareous silty sediments exposed during the excavation demonstrate that a substantial body of water occupied the site during the middle Holocene. The diverse and well-preserved gastropod and bivalve assemblage supports this interpretation, as ten species recovered are presently known to inhabit permanent ponds or lakes. Recovery of the bivalve *Pisidium ventricosum* is particularly relevant as is not presently known to live in ephemeral aquatic habitats.

Vascular plant and bryophyte macrofossils and pollen also document the plant community within and immediately surrounding the pond. An aquatic environment is indicated by the abundance of pondweed (*Potamogeton*) seeds (and rare pollen), mare's tail and *Chara* macrofossils, in addition to palynomorphs of algae *Botryococcus* and *Pediastrum*. Sedges were probably very common along the marshy areas along the lake margin as revealed by the abundance of *Carex* spp. achenes and pollen. However, some sedge and sage pollen might come from open meadows, such as surrounding dry, south facing slopes around the lake basin. The hydric bryophyte species

typify an *in situ* minerotrophic, marl-rich fen, wet meadow or still oxbow pool type of aquatic environment similar to those at present in the circumpolar arctic and boreal regions.

Palynological results support previous studies and reveal a boreal forest dominated by white spruce, with lesser amounts of black spruce, birch and shrub alder, with a conspicuous absence of lodgepoll pine. Rare, highly fragmentary, spruce needles confirm that spruce trees were locally present, and their pollen was not transported to Versleuce meadows from long-distance sources. The lack of lodgepoll pine in the southern Yukon as revealed in this and previous studies provides a stark contrast to the forest community at present.

Based on paleoecological data from southern Yukon, Cwynar and Spear (1995) suggests the period between 6,000 and 5,400 years ago was characterized by relatively cooler and wetter climates than times before and after. Data recovered from Versleuce Meadows supports that interpretation and suggests that a sizable pond or small lake existed at this site during that period. A shallow lake in open spruce woodlands would have likely been an attractive focal point for a variety of fauna, including bison. This paleoecological reconstruction, however, is seemingly at odds the interpretation of data from nearby alpine non-glacial ice patches. Analysis of three representative Holocene alpine ice patches suggests there was little or no accumulation of alpine ice between 6,700 and 4,700 years ago, suggesting increased temperatures and/or reduced precipitation during this time period (Farnell et al. 2004). These differences may reflect differences associated with lowland versus upland paleoenvironmental records.

1

2 **Holocene Steppe Bison in Beringia**

3 The Versleuce Meadows bison skeleton is important for our understanding post-glacial bison in
4 the north as it may be the best preserved middle Holocene specimen that has yielded both
5 comprehensive morphometric and mtDNA data from the region. The combined morphometric
6 and genetic data are a further demonstration that there is considerable morphological variation
7 within and between previously defined species. Morphological and mensurational characteristics
8 may not be adequate alone to make taxonomic assignments for bison fossils, or useful in
9 distinguishing species from one another. As such, it is critical to recover genomic data to
10 determine whether bison from the Subarctic represent a lineage that includes steppe bison from
11 the north or the lineage that gave rise to extant bison species from the south (Clade 1A). At least
12 for bison that lived in Beringia, steppe bison *B. priscus* (*sensu lato*) may be best interpreted as
13 representing a single, morphologically variable species that dominated late Pleistocene mammal
14 communities and persisted through the Holocene in southern Yukon until their relatively recent
15 extirpation ~400 years ago (Shapiro et al. 2004; Heintzman et al. 2016). Further, the common
16 name “steppe bison” may be misleading as Holocene records of Clade 2A bison in the Subarctic
17 represent populations that inhabited boreal forests. As such, this expands our interpretation of *B.*
18 *priscus* ecology which has traditionally been thought of as grazers of the open, largely treeless,
19 “mammoth steppe” biome (Guthrie 1990). Rather, steppe bison should be considered to have
20 been a species that had a high degree of ecological flexibility, which could occupy a wide range
21 of open and forested habitats across the Arctic and Subarctic, during both glacial and interglacial
22 time periods of the Quaternary. This ecological flexibility may be the major factor which led to

the high degree of morphological variability exhibited in horn cores and crania clearly found in the paleontological record.

Conclusions

1) The Versleuce Meadows bison, a juvenile, is likely the most complete Holocene bison reported from the subarctic of northern Canada. The animal died roughly 5,400 years ago after being mired down in a pond or small lake which was surrounded by a white spruce dominated forest. Exposed parts of the carcass were scavenged by carnivores, though most of the distal limbs and crania were left unscathed. The rest of the animal was buried in silty-marl sediment.

2) Combined morphometric and mitochondrial genomic data support the assignment of this specimen to *Bison cf. priscus* and reveal it was a member of Clade 2A, which in Beringia are considered to represent steppe bison. This bison was not a member of the lineage of present-day extant plains and woods bison (Clade 1A) which are descended from populations that were restricted to the southern part of the continent during the last glaciation. Steppe bison were a morphologically variable and ecologically flexible species of bison that dominated late Pleistocene landscapes of Beringia, and survived locally through the Holocene interglaciation.

Acknowledgments

We thank Craig and Sandy Duncan for alerting the Yukon Government to their discovery and for allowing us to excavate this skeleton from their property. We recognize the Kwanlin Dün First Nation and The Ta'an Kwach'an Council for enabling us to work in their Traditional Territories and for their support of this research. Dan Chang, Joshua Kapp, and Amanda Hart provided technical assistance with the genomics research. Dr. Gerry Mackie also provided generous assistance in the identification of molluscan species. Dr. John Southon of the KECK University of California Irvine Accelerator Mass Spectrometer (UCIAMS) laboratory provided the radiocarbon date. We thank the following families and individuals for assisting with the excavation: the Deuling family, The Zazula-Castillo family, The Thomas-Coppard family, The Hall family; Ken Slonski for operation of the backhoe excavator. We appreciate all those that helped or just came out to provide support during our brief, but highly successful, field excavation of this bison skeleton.

References

- Beaumont, W., Beverly, R., Southon J., and Taylor R.E. 2010. Nuclear Instruments and Methods in Physics Research B, 268 (7-8): 906-909.
- Beverly, R.K., Beaumont, W., Tauz, D., Ormsby, K.M., von Reden, K.F., Santos, G.M., and Southon, J.R. 2010. The Keck Carbon Cycle AMS Laboratory, University of California Irvine: Status report. Radiocarbon 52(2): 301-309.
- Bond, J.D. 2004. Late Wisconsinan McConnell glaciation of the Whitehorse map area (105D), Yukon. Yukon Exploration and Geology 2003: 73-88.

- 1 Clarke, A.H. 1981. Freshwater mollusc of Canada. National Museum of Natural Sciences.
2 National Museum of Canada. Ottawa. 447 pp
- 3 Cody, W.J. 1996. Flora of the Yukon. NRC Research Press, Ottawa.
4
- 5 Cywnar, L.C. 1988. Late Quaternary vegetation history of Kettlehole Pond, southwestern Yukon.
6 Canadian Journal of Forest Research 18(10): 1270-1279.
7
- 8 Cwynar, L.C., and Spear, R.W. 1995. Paleovegetation and paleoclimatic changes in the Yukon at
9 6 ka BP. Géographie physique et Quaternaire 49(1): 29-35.
10
- 11 Dillon, R.T Jr. 2000. The Ecology of Freshwater Molluscs. Cambridge University Press.
12
- 13 Drummond, A.J., Suchard, M.A., Xie, and D., Rambaut, A. 2012. Bayesian phylogenetics with
14 BEAUti and the BEAST 1.7. Molecular Biology and Evolution 29(8): 1969-1973.
15
- 16 Duffield, L.F. 1973. Aging and sexing the post-cranial skeleton of bison. Plains anthropologist
17 18(60): 132-139.
18
- 19 Faegri, K., Iversen, J., Kaland, P.E., and Krzywinski, K., 1989. Textbook of pollen analysis, IV
20 edition. John Wiley and Sons.
21

Farnell, R., Hare, P.G., Blake, E., Bowyer, V., Schweger, C., Greer, S., and Gotthardt, R. 2004. Multidisciplinary investigations of alpine ice patches in Southwest Yukon, Canada: Paleoenvironmental and paleobiological investigations. *Arctic* 57(3): 247–259.

Froese, D.G., Zazula, G.D., Westgate, J.A., Preece, S.J., Sanborn, P.T., Reyes, A.V., and Pearce, N.J. 2009. The Klondike goldfields and Pleistocene environments of Beringia. *GSA Today* 19 (8): 4-10.

Froese, D., Stiller, M., Heintzman, P.D., Reyes, A.V., Zazula, G.D., Soares, A.E.R., Meyer, M., Hall, E., Jensen, B.J., Arnold, L.J., MacPhee, R.D.E., and Shapiro, B. 2017. Fossil and genomic evidence constrains the timing of bison arrival in North America. *Proceedings of the National Academy of Sciences*, 114(13): 3457-3462.

Gajewski, K., Bunbury, J., Vetter, M., Kroeker, N., and Khan, A.H. 2014. Paleoenvironmental studies in Southwest Yukon. *Arctic*: 2-13.

Gilbert, B.M. 1990. *Mammalian osteology*. Missouri Archaeological Society.

Gill, M.S., Lemey, P., Faria, N.R., Rambaut, A., Shapiro, B., and Suchard, M.A., 2013. Improving Bayesian population dynamics inference: a coalescent-based model for multiple loci. *Molecular Biology and Evolution*, 30(3): 713-724.

- 1 Guthrie, R.D. 2006. New carbon dates link climatic change with human colonization and
2 Pleistocene extinctions. *Nature* 441: 207-209.
- 3
- 4 Guthrie, R.D. 1990. Frozen fauna of the mammoth steppe: The story of Blue Babe. The
5 University of Chicago Press, Chicago.
- 6
- 7 Guthrie, R.D. 1970. Bison evolution and zoogeography in North America during the Pleistocene.
8 *The Quarterly Review of Biology* 45(1): 1-15.
- 9
- 10 Hansen, B.C.S., and Engstrom, D.R. 1985. A comparison of numerical and qualitative methods
11 of separating pollen of black and white spruce. *Canadian Journal of Botany* 63: 2159-2163.
- 12
- 13 Harington, C.R., 2011. Pleistocene vertebrates of the Yukon Territory. *Quaternary Science*
14 *Reviews* 30: 2341-2354.
- 15
- 16 Hedenas, L. 2014a. Amblystegiaceae. *Flora of North America Editorial Committee, eds. 1993+.*
17 *Flora of North America North of Mexico. 19+ vols. New York and Oxford. Vol. 28.*
- 18
- 19 Hedenas, L. 2014b. Calliergonaceae. *Flora of North America Editorial Committee, eds. 1993+.*
20 *Flora of North America North of Mexico. 19+ vols. New York and Oxford. Vol. 28.*
- 21
- 22 Heintzman, P.D., Froese, D., Ives, J.W., Soares, A.E., Zazula, G.D., Letts, B., Andrews, T.D.,
23 Driver, J.C., Hall, E., Hare, P.G., Jass, C.N., MacKay, G., Southon, J.R., Stiller, M., Woywitka,

- 1 R., Suchard, M.A., and Shapiro, B. 2016. Bison phylogeography constrains dispersal and
2 viability of the Ice Free Corridor in western Canada. *Proceedings of the National Academy of*
3 *Sciences*, 113(29): 8057-8063.
- 4
- 5 ITIS. 2016. Integrated Taxonomical Information System. <https://www.itis.gov/>
- 6
- 7 Lévesque, P.E.M., Dinel, H, and Larouche, A. 1988. Guide to identification of plant macrofossils
8 in Canadian peatlands. Agriculture Canada, Research Branch, Publication No. 1817.
- 9
- 10 Janssens, J.A. 1983. Past and present record of *Drepanocladus crassicosatus* sp. nov. (Musci:
11 *Amblystegiaceae*) and status of *D. trichophyllus* in North America. *The Bryologist* 86: 44-53.
- 12
- 13 Mackie, G.L., and Flippance, L.A. 1983. Intra- and interspecific variations in calcium content of
14 freshwater Mollusca in relation to calcium content of the water. *Journal of Molluscan Studies* 49:
15 204-212
- 16
- 17 McDonald, J.N. 1981. North American bison: their classification and evolution. University of
18 California Press, Berkeley, California.
- 19
- 20 Shapiro, B., Drummond, A.J., Rambaut, A., Wilson, M.C., Matheus, P.E., Sher, A.V., Pybus,
21 O.G., Gilbert, M.T.P., Barnes, I., Binladen, J., Willerslev, E., Hansen, A.J., Baryshnikov, G.F.,
22 Burns, J.A., Davydov, S., Driver, J.C., Froese, D.G., Harington, C.R., Keddie, G., Kosintsev, P.,

- 1 Kunz, M.L., Martin, L.D., Stephenson, R.O., Storer, J., Tedford, R., Zimov, S., and Cooper, A.
2 2004. Rise and Fall of the Beringian Steppe Bison. *Science* 306: 1561-1565.
- 3
- 4 Skinner, M.F. and Kaisen, O.C. 1947. The fossil bison of Alaska and preliminary revision of the
5 genus. *Bulletin of the American Museum of Natural History* 89: 123-256.
- 6
- 7 Stephenson, R.O. Gerlach, S.C., Guthrie, R.D., Harington, C.R. Mills, R.O., and Hare, G. 2001.
8 Wood bison in late Holocene Alaska and adjacent Canada: paleontological, archaeological and
9 historical records. *In People and Wildlife in Northern North America: Essays in honor of R. Dale*
10 *Guthrie. Edited by S.C. Gerlach and M.S. Murray, M.S. BAR International Series 944, Oxford,*
11 *United Kingdom. pp. 124-158.*
- 12
- 13 Yukon Ecoregions Working Group, 2004. Yukon Southern Lakes. *In Ecoregions of the Yukon*
14 *Territory: Biophysical properties of Yukon landscapes. Edited by C.A.S. Smith, J.C. Meikle, and*
15 *C.F. Roots, C.F. Agriculture and Agri-Food Canada, PARC Technical Bulletin No. 0401,*
16 *Summerland, British Columbia, pp. 207-218.*
- 17
- 18 Westgate, J.A., Preece, S.J., Kotler, E.C., and Hall, S. 2000. Dawson Tephra: a prominent
19 stratigraphic marker of late Wisconsinan age in west-central Yukon, Canada. *Canadian Journal*
20 *of Earth Sciences* 37: 621-627.
- 21

Wilson, M.C., Hills, L.V., Shapiro, B. 2008. Late Pleistocene northward-dispersing *Bison antiquus* from the Bighill Creek Formation, Gallelli Gravel Pit, Alberta, Canada, and the fate of *Bison occidentalis*. Canadian Journal of Earth Sciences 45: 827-859.

Figure Captions

Figures 1. Map of Yukon Territory including the late Pleistocene McConnell and Laurentide ice sheet limits (after Westgate et al., 2001). Inset is the Versleuce Meadows study site in relation to the City of Whitehorse and Yukon River.

Figure 2. Photographs from the Versleuce Meadows site. a) area with trenches where the initial accidental discovery of bones was made and where further excavation was focused; b) tan colored silty-marl sediments exposed by the excavation; c) mollusc rich silt sediment; d) *in situ* partial bison skeleton found roughly 150 cm below the surface.

Figure 3. An idealized skeleton of a bison showing parts that we recovered at the Versleuce Meadows site shaded in gray. Precise anatomical of some elements, such as rib fragments, were not determined but are illustrated here for interpretive purposes. Illustration after Gilbert (1990).

Figure 4. Versleuce Meadows Bison cranium (YG 422.1).

Figure 5. Versleuce Meadows bison skeletal elements. A) right mandible; B) 3rd cervical vertebra showing unfused epiphysis of the centrum; C) thoracic vertebra lacking anterior epiphysis of centra and broken spinous process; D) innominates; E) various rib fragments; F) metacarpal with complete set of phalanges; G) left radius; H) left humerus with substantial carnivore gnawing of proximal end; I) left scapula with substantial carnivore gnawing on the periphery; J) metatarsal with complete set of phalanges; K) right tibia which is heavily carnivore gnawed on proximal end and fractured on distal part of the shaft, radiocarbon sample removed from the middle shaft.

Figure 6. A Bayesian timetree of bison mitochondrial genomes, with three well-supported clades (1A, 2A, 2B; Heintzman et al. 2016) highlighted in grey. The Versleuce Meadows bison (VersMead) is highlighted in bold and belongs to clade 2A. All living bison in North America are members of Clade 1A. For this study, we refer to all bison outside of Clade 1A as *Bison* c.f. *priscus* regardless of their morphology. Four yak outgroup sequences have been removed. Branch support values are given for deep nodes and those shallower nodes with posterior probability values >0.95. Light grey node bars are 95% highest posterior density intervals for node heights and are shown for nodes with posterior probability values >0.95.

Figure 7. Macrofossils recovered from sediment associated with the bison skeleton. A) *Potamogeton* nutlet; b) *Hippuris* seed; c) *Carex* perigynia; d) *Carex* achene; e) *Eleocharis* achene; f) *Picea* needle; g) *Chenopodium* seed; h) *Cristatella mucedo* statoblast; i) *Scorpidium* scorpioides stem with leaves; j) *Calliergon giganteum* stem with leaves; k) *Drepanocladus longifolius* stem with leaves; l) *Stagnicola arctica* shell; j) *Valvata sincera* bivalve shell; k) *Pisidium casertanum* shell.

1
2
3
4
5
6
7
8
9
10
11
12
13

Tables

Table 1. Mensurational data from Versleuce Meadows bison cranium YG 422.1 with comparative metrics from other Quaternary bison species (after McDonald 1981). Measurements reported for YG 422.1 horn cores are taken from the left side.

Table 2. Mensurational data from Versleuce Meadows bison mandibles and select post-cranial elements.

Table 3. Palynomorph percentages from Versleuce Meadows site.

Table 4. Summary of mollusk taxa and preferred habitats (after Clark 1981) from Versleuce Meadows site.

133x185mm (300 x 300 DPI)

86x182mm (300 x 300 DPI)

144x114mm (300 x 300 DPI)

86x129mm (300 x 300 DPI)

179x205mm (300 x 300 DPI)

Thousands of calendar years before present (ka BP)

178x186mm (300 x 300 DPI)

Table 1.																		
Description of metric	Skinner and Kaisen (1947) metric	von den Driesch (1976) metric (Figure 8, <i>Bos taurus</i>)	YG 422.1 mm	<i>Bison antiquus</i>			<i>Bison occidentalis</i>			<i>Bison priscus</i>			<i>Bison bison athabascaae</i>			<i>Bison alaskensis</i>		
				min	mean	max	min	mean	max	min	mean	max	min	mean	max	min	mean	max
spread of horn cores, tip to tip	#1	#42	802	765	870	1067	626	779	1055	751	888	1064	542	681	848	800	1151	1540
greatest spread of cores on outside curve	#2	#43	810				735	782	892									
core length on upper curve, tip to burr	#3		282	203	279	364	186	278	392	285	362	416	165	235	323	426	553	795
core length on lower curve, tip to burr	#4	#47	332	280	336	395	275	340	405									
straight line length, tip of core to upper base at burr	#5		253	185	250	330	175	249	350	268	318	379	154	207	277	294	459	667
vertical (or least) diameter of horn core at base (at burr)	#6		93	81	102	126	70	95	114	84	98	116	81	92	106	100	117	153
greatest (transverse) diameter of horn core at base		#45	96	76	106	129	77	99	120	98	115	130	83	97	109	122	142	171
circumference of core at base (at burr)		#44	313	233	324	392	237	300	355	293	330	387	254	289	322	341	397	493
greatest width of auditory opening	#8	#25	273	251	288	318	238	262	294	248	276	310	243	274	298	268	302	340
width of condyles	#9	#26	145	132	144	161	111	135	151	127	141	165	118	130	139	133	152	175
depth from occipital crest to top of foramen magnum	#10	#41	94	94	112	134	89	104	120	91	103	119	92	100	114	97	111	134
depth from occipital crest to bottom of foramen magnum	#11	#40	135				136	158	177									
greatest width between bases of horn cores	#13		340															
least width between horn cores		#31	335															
width of cranium between horn cores and orbits	#14	#32	280	276	315	352	261	297	348	269	294	336	273	293	313	281	332	394
maximum internal width of orbit		#23	80															
greatest breadth across the orbits = greatest frontal breadth	#15	#33	331	346	353	360	311	348	394	313	350	415	326	354	384	356	396	440

least breadth between the orbits	#16	#34	252															
facial breadth	#17	#35	n/a															
rostral width at maxillary-premaxillary suture	#18		n/a				102	115	125									
least occipital breadth		#30	163															
greatest breadth across nasals		#36	94															
condylobasal length	O-P		573				505	573	630			562	579	604	602	650	676	
basal length	F-P	#3	548	520	545	560	467	516	582									
breadth across the premaxillae on the oral protruberences		#37	n/a															
greatest palatal breadth measured across outer borders of alveoli		#38	n/a															
length of cheek tooth row	#19	#20	n/a				137	147	160									
length of molar row	#20	#21	n/a	105.2	105.6	106	84	91	102									
length of premolar row		#22	n/a															
length of nasal bone	N-T	#12	219															
length, occipital crest to nasal-frontal suture	O-N	#8	244	240	295	350	229	259	273	254	272	286	240	256	276	265	298	338
length, beyond P2 to tip of premaxilla	M-P		n/a				143	153	168									
short upper cranium length = occipital crest to tip of nasals.	O-T	#10	454	482	510	527	471	493	543									
angle of posterior divergence of horncore	#A21		74°	72°	79°	86°	63°	72°	83°	63°	71°	79°	63°	71°	77°	64°	74°	85°
angle of proximal horncore depression	#A22		12°															

Table 2. Post cranial measurements	
Metacarpal (YG 422.40)	mm
overall length	225.2
proximal breadth	83.4
midshaft breadth	49.7
distal breadth	83.2
midshaft depth	31.6
proximal depth	47.2
distal depth	44.4
minimum shaft depth	29.9
mimimm shaft breadth	49.1
metatarsal (YG 422.57)	
overall length	275.2
proximal breadth	62
midshaft breadth	39
distal breadth	71.1
midshaft depth	37.1
proximal depth	61.2
distal depth	43.4
minimum shaft depth	34.3
mimimm shaft breadth	38.1
Humerus (YG 422.10)	
smallest breadth of shaft	56.3
greatest breadth of trochlea	112.4
greatest breadth of distal end	120.2
Radius (YG 422.30)	
greatest length	376.4
smallest breadth of shaft	58.2

Draft

greatest breadth of proximal end	109.4
greatest breadth of proximal articular facies	99
breath of distal end	99.7
breadth of distal articular facies	94.9
greatest proximal depth	59.6
Ulna	
(YG 422.84)	
greatest length	440.2
smallest depth of olecranon	74.8
greatest depth across the processus anconaeus	104.3
smallest depth across the processus anconaeus	66.7
Tibia	
(YG 422.20)	
greatest length	453.9
medial length	439.4
proximal depth	127.1
minimum breadth of shaft	43.2
mimum depth of shaft	38.5
greatest distal breadth	84.1
greatest depth of distal end	54.7
breadth at mid shaft	49.6
depth at mid shaft	55.2
Mandible	
(YG 422.2)	
Length from the angle: Gonion caudale - Infradentale	423
Length from the condyle: aboral border of the condyle process -Infradentale	453
Length: Gonion caudale - aboral border of the alveolus of M ₃	133
Length of the horizontal ramus: aboral border of the alveolus of M ₃ - Infradentale	291
Length: Gonion caudale - oral border of the alveolus of P ₂	287
Length: Gonion caudale - the most aboral indentation of the mental foramen	360
Length of the cheektooth row, measured along the alveoli on the buccal side	158
Length of the molar row, measured along the alveoli on the buccal side	101

Length of the premolar row, measured along the alveoli on the buccal side	58
Length (L) and breadth (B) of M ₃ , measured near the biting surface	44 (L), 16 (B)
Length of the diastema: oral border of the alveolus of P ₂ - aboral border of the alveolus of I ₄	125
Aboral height of the vertical ramus: Gonion ventrale - highest point of the condyle process	202
Middle height of the vertical ramus: Gonion ventrale - deepest point of the mandibular notch	183
Oral height of the vertical ramus: Gonion ventrale - Coronion	232
Height of the mandible behind M ₃ from the most aboral point of the alveolus on the buccal side	71
Height of the mandible in front of M ₁	50
Height of the mandible in front of P ₂	39

Draft

Table 3. Fossil palynomorphs

Taxon	Sample A % (sum=540)	Sample B % (sum = 571)
TREES total (AP)	60.7	67.1
<i>Picea glauca</i> type	38.1	42.1
<i>Picea mariana</i> type	9	13.9
<i>Pinus contorta</i> type	0.8	0
<i>Betula</i>	12.3	10.6
SHRUBS		
<i>Alnus viridis</i> type	11.5	8.5
<i>Salix</i>	2	5.6
<i>Shepherdia</i> <i>canadensis</i>	0.8	0.2
HERBS		
CYPERACEAE	22	16.7
POACEAE	0.8	0
<i>Artemisia</i>	2.3	1.8
<i>Potamogeton</i> type	0.4	0.2
SPORES		
Filicales undiff.	0.6	1.1
<i>Equisetum</i>	0.2	0
<i>Lycopodium</i>	0.4	0
ALGAE		
<i>Botryococcus</i>	2.2	1.4
<i>Pediastrum</i>	1.1	0.2
PALYNOMORPH CONC.	25,300 per mL	15,500 per mL

Table 4. Mollusks				
Class	Family	Common name	Species	Habitat
Gastropoda	Lymnaeidae	Frigid lymnaea, Alaskan lymnaea	<i>Lymnaea atkaensis</i> (Dall, 1884)	Restricted to BC, NWT, and Yukon Territory. Clarke (1980) reports it from cold, oligotrophic lakes and appears to have similar habitat characteristics as <i>Sphaerium nitidu</i> .
Gastropoda	Valvatidae	Mossy valvata (subspecies)	<i>Valvata sincera sincera</i> (Say, 1824)	Widely spread and very common when found. Most common in littoral and sublittoral zones on substrates of mud and organic debris.
Gastropoda	Valvatidae	Mossy valvata (subspecies)	<i>Valvata sincera helicoidea</i> (Dall, 1905)	A northern species (subspecies) in shallow ponds, lakes, rivers, streams, and bogs, on a variety of substrates in littoral and sublittoral zones.
Gastropoda	Planorbidae	Ash Gyro	<i>Gyraulus parvus</i> (Say, 1817)	Common in vegetation zones of permanent ponds and lakes, and in organic to mud substrates.
Gastropoda	Lymnaeidae	Arctic Pondsnaail	<i>Stagnicola arctica</i> (Lea, 1864)	A northern species in shallow ponds, lakes, rivers, streams, and bogs.
Gastropoda	Lymnaeidae	Marsh pondsnaail ?	<i>Stagnicola elodes</i> ? (Say, 1821)	
Gastropoda	Lymnaeidae	Modest Fossaria	<i>Fossaria modicella</i> (Say, 1825)	Very common everywhere, in temporary and permanent aquatic habitats. A littoral/sublittoral species common in muddy, organic substrates
Gastropoda	Lymnaeidae	Boreal Fossaria ?	<i>Fossaria decampi</i> ? (Streng, 1896) (syn. <i>Galba alba</i>)	Cold water species, most common in large lakes in southern Canada, streams and shallow lakes in northern parts of its range.
Bivalvia	Sphaeriidae	Ubiquitous peaclam	<i>Pisidium casertanum</i> (Poli, 1791)	The most ubiquitous fingernail clam, and probably freshwater mollusc in all parts of the world. Found in oligotrophic to eutrophic lakes, deep and shallow waters, sandy to muddy organic ooze, temporary and permanent aquatic habitats.
Bivalvia	Sphaeriidae		<i>Sphaerium nitidum</i>	Cold water species, most common in oligotrophic lakes, in shallow depths in North but hypolimnial depths in southern parts of Canada.
Bivalvia	Sphaeriidae		<i>Pisidium ventricosum</i>	Found principally in lakes and large rivers (never in ephemeral aquatic habitats). Large populations of the species are uncommon, but when found they are usually in fine sand or mud.