Rapid evolution meets invasive species control: the potential for pesticide resistance in sea lamprey

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0015.R1
Manuscript Type:	Perspective
Date Submitted by the Author:	24-Mar-2017
Complete List of Authors:	<p>Dunlop, Erin; Ontario Ministry of Natural Resources McLaughlin, Robert; University of Guelph, Adams, Jean; US Geological Survey Great Lakes Science Center, Jones, Michael; Michigan State University, Department of Fisheries and Wildlife Birceanu, Oana; Wilfrid Laurier University Christie, Mark; Purdue University Criger, Lori; US Fish and Wildlife Service, Marquette Hinderer, Julia; Great Lakes Fishery Commission Hollingworth, Robert; Michigan State University Johnson, Nicholas; USGS, Great Lakes Science Center, Hammond Bay Biological Station Lantz, Stephen; US Fish and Wildlife Service Li, Weiming; Michigan State University, Miller, James; Michigan State University, Department of Entomology Morrison, Bruce; Fisheries and Oceans Canada Mota-Sanchez, David; Michigan State University Muir, Andrew; Michigan State University, Fisheries and Wildlife; Geat Lakes Fishery Commission, Sepúlveda, Maria; Purdue University Steeves, Todd; Department of Fisheries and Oceans, Walter, Lisa; Great Lakes Fishery Commission Westman, Erin; McMaster University Michael G DeGroote School of Medicine Wirgin, Issac; NYU Wilkie, Michael; Wilfrid Laurier University</p>
Please Select from this Special Issues list if applicable:	N/A
Keyword:	contemporary evolution, behavioural evolution, aquatic invasive species, integrative pest management, physiological resistance

SCHOLARONE™
Manuscripts

Draft

Rapid Evolution Meets Invasive Species Control: The Potential for Pesticide

Resistance in Sea Lamprey

By Erin S. Dunlop^{1,2*}, Rob McLaughlin³, Jean V. Adams⁴, Michael Jones⁵, Oana Birceanu⁶, Mark R. Christie^{7,8}, Lori A. Criger⁹, Julia L.M. Hinderer¹⁰, Robert M. Hollingworth¹¹, Nicholas S. Johnson¹², Stephen R. Lantz⁹, Weiming Li¹³, James Miller¹⁴, Bruce J. Morrison¹⁵, David Mota-Sanchez¹⁶, Andrew Muir¹⁰, Maria S. Sepúlveda⁸, Todd Steeves¹⁵, Lisa Walter¹⁰, Erin Westman¹⁷, Isaac Wirgin¹⁸, and Michael P. Wilkie⁶.

¹Aquatic Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry, Peterborough, ON K9L 0G2, Canada.

²Environmental and Life Sciences Graduate Program, Trent University, Peterborough, ON K9L 0G2, Canada.

³Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1, Canada

⁴United States Geological Survey, Great Lakes Science Center, Ann Arbor, MI 48105, USA.

⁵Quantitative Fisheries Center, Michigan State University, East Lansing, MI 48824, USA.

⁶Department of Biology and Laurier Institute for Water Science, Wilfrid Laurier University, Waterloo, ON N2L 3C5, Canada.

⁷Department of Biological Sciences, Purdue University, West Lafayette, IN 47907, USA.

⁸Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN 47907, USA.

⁹United States Fish and Wildlife Service, Sea Lamprey Control Program, Marquette Biological Station, Marquette, MI 49855, USA.

¹⁰Great Lakes Fishery Commission, Ann Arbor, MI 48105, USA.

¹¹Center for Integrative Toxicology and Department of Entomology, Michigan State University, East Lansing, MI 48824, USA.

¹²United States Geological Survey, Great Lakes Science Center, Hammond Bay Biological Station, Millersburg, MI 49759, USA.

¹³Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI 48824, USA.

¹⁴Department of Entomology, Michigan State University, East Lansing, MI 48824, USA.

¹⁵Sea Lamprey Control Centre, Fisheries and Oceans Canada, Sault Ste Marie, ON P6A 2E5, Canada.

¹⁶Department of Entomology, Michigan State University, East Lansing, MI 48864, USA.

¹⁷M. G. DeGroote Institute for Infectious Disease Research, Department of Biochemistry and Biomedical Sciences, DeGroote School of Medicine, McMaster University, Hamilton, ON L8N 4K1, Canada.

¹⁸Department of Environmental Medicine, NYU School of Medicine, Tuxedo, NY 10987, USA.

*Corresponding author: Erin S. Dunlop (email: erin.dunlop@ontario.ca; phone: 705-755-2296).

Abstract

Rapid evolution of pest, pathogen and wildlife populations can have undesirable effects; for example, when insects evolve resistance to pesticides or fishes evolve smaller body size in response to harvest. A destructive invasive species in the Laurentian Great Lakes, the sea lamprey (*Petromyzon marinus*) has been controlled with the pesticide 3-trifluoromethyl-4-nitrophenol (TFM) since the 1950s. We evaluated the likelihood of sea lamprey evolving resistance to TFM by (1) reviewing sea lamprey life history and control; (2) identifying physiological and behavioural resistance strategies; (3) estimating the strength of selection from TFM; (4) assessing the timeline for evolution; and (5) analyzing historical toxicity data for evidence of resistance. The number of sea lamprey generations exposed to TFM was within the range observed for fish populations where rapid evolution has occurred. Mortality from TFM was estimated as 82-90%, suggesting significant selective pressure. However, 57 years of toxicity data revealed no increase in lethal concentrations of TFM. Vigilance and the development of alternative controls are required to prevent this aquatic invasive species from evolving strategies to evade control.

Keywords: contemporary evolution; behavioural evolution; toxicity; physiological resistance; aquatic invasive species; artificial selection; lampricide; niclosamide; integrative pest management; Great Lakes Fishery Commission.

59 Introduction

60 The capability of pathogens and pests to evolve resistance to control measures is a global
61 concern with implications for human health, natural resource management, and agriculture
62 (Palumbi 2001). Resistance to various chemicals has been documented in many taxa (Davies
63 and Davies 2010; Mota-Sanchez et al. 2002; Wirgin and Waldman 2004;
64 www.pesticideresistance.org). Further, the evolution of resistance can result in significant
65 economic loss to individuals, industry, and governments (Howard et al. 2003; Howard and Scott
66 2005; Alsan et al. 2015). Insecticide-resistant mosquitos carrying the malaria vector in Africa
67 provides a prominent example involving significant human suffering (Enayati and Hemingway
68 2010). Recognition of the widespread occurrence of resistance and its consequences is
69 prompting the development of specific strategies targeted at slowing or preventing its
70 occurrence (Tabashnik et al. 2014).

71 In fisheries, aquatic invasive species are non-native pests that threaten aquatic
72 ecosystems. One of the most notorious aquatic invasive species is the sea lamprey
73 (*Petromyzon marinus*), which precipitated a catastrophic collapse of economically valuable and
74 culturally significant fisheries resources when it invaded the Laurentian Great Lakes (Siefkes et
75 al. 2013). Sea lamprey were first recorded in Lake Ontario in 1888 (Eshenroder 2014) and by
76 1938 had spread to each of the Great Lakes (Christie and Goddard 2003). This invasion was
77 followed by massive declines and the extirpation of several native fish populations (Siefkes et
78 al. 2013).

79 The chemical pesticide, 3-trifluoromethyl-4-nitrophenol (TFM) has been applied to Great
80 Lakes tributaries to control sea lamprey since the 1950s (Siefkes et al. 2013). TFM is applied as

a lampricide to selectively target and kill larval sea lamprey, while leaving many other aquatic organisms unharmed (Applegate et al. 1957, 1958). The application of TFM expanded between the 1950s and 1980s (Table 1), contributing to substantial reductions in sea lamprey abundance (Lawrie 1970; Heinrich et al. 2003). A second pesticide, 2',5-dichloro-4'-nitrosalicylanilide (aka. Niclosamide, Bayluscide®), originally used as a molluscicide was discovered in 1964 as having synergistic properties with TFM (Howell et al. 1964) and was subsequently applied in small doses to treat select tributaries. Application of these two pesticides (also referred to as lampricides), and in particular TFM, remains the cornerstone of sea lamprey control today and has supported recoveries of formerly depleted fish stocks, such as Lake Superior lake trout *Salvelinus namaycush*, highly valued for their contribution to the health and economy of the Great Lakes, and their cultural importance to First Nations communities (Smith and Tibbles 1980; Christie and Goddard 2003; Goddard 2013).

In this paper, we evaluate the possibility that Great Lakes sea lamprey could evolve resistance to the pesticides used to control them. As would be expected from any source of substantial mortality acting on a population, the application of lampricides over the past 4-6 decades has potentially imposed considerable selective pressure on sea lamprey. Pesticide resistance has evolved in diverse taxa, including bacteria, insects, and vertebrates, following continued exposure to a toxin, pesticide, or contaminant. While the generation time of sea lamprey (> 4 years; see discussion below) is considerably longer than the generation times of bacteria (hours-days) or arthropod pests (days-months), numerous examples of rapid evolution contributing to human-induced trait changes in fish and wildlife populations highlight the potential threat and warrant consideration (Palkovacs et al. 2012; Sharpe and Hendry 2009).

Resistance to lampricides would lead to reduced effectiveness of sea lamprey control, resulting in increased mortality of commercial, sport and culturally significant fish populations in the Great Lakes. Furthermore, once resistance evolves it may be slow to reverse depending on associated costs or trade-offs. In fish, evolutionary recovery following the removal of anthropogenic selection has a slow reverse trajectory (Conover et al. 2009; Enberg et al. 2009), coined a “Darwinian debt” to be paid by future generations (U. Dieckmann, Financial Times, August 28 2004). However, although resistance leads to increased fitness in the presence of the chemical, it could cause reduced fitness under normal conditions; in such a case, resistance might be expected to reverse relatively quickly once the chemical is removed (Levinton et al. 2003; Mackie et al. 2010). Evaluating the likelihood of sea lamprey resistance to lampricides has broader implications for management and control of invasive species, particularly vertebrate aquatic invasive species, by providing insights on rapid evolution in animal populations and on how applied evolutionary biology can inform management of pressing global issues (Hendry et al. 2011; Carroll et al. 2014).

Our synthesis of lampricide resistance includes: (1) a review of sea lamprey life history and control in the Great Lakes; (2) identifying the opportunities for resistance to develop in sea lamprey, including physiological and behavioural strategies; (3) estimates of the strength of selection from lampricides; (4) assessment of whether sufficient time has passed for any evolution to occur; (5) analysis of lampricide toxicity data for signs of resistance; (6) identifying strategies for reducing the chances of resistance; and (7) conclusions on the implications our synthesis has for the control and management of other invasive species.

Sea Lamprey Life History and Control in the Great Lakes

Following metamorphosis and prior to maturation, sea lamprey are parasites—attaching to their fish hosts using an oral disc, puncturing the skin and muscle of the fish with their rasping tongue, and ingesting the blood and body fluids of the living animal (Renaud et al. 2009). Sea lamprey spend 12-18 months in this parasitic phase in the open waters of the Great Lakes before maturing and migrating into tributaries to spawn (Bergstedt and Swink 1995). Sea lamprey do not show natal homing behaviour (Bergstedt and Seelye 1995; Waldman et al. 2008), instead relying on odorants emitted by lamprey larvae to find suitable tributaries for spawning (Fine et al. 2004; Buchinger et al. 2013). Sea lamprey fecundity varies geographically within the Great Lakes, averaging greater than 50,000 eggs per female (Smith and Marsden 2007; Dawson et al. 2015). Fecundity is considered to be relatively high compared to other fishes, but fertilized eggs are vulnerable to predation and annual recruitment variability is high (Jones et al. 2003; Dawson et al. 2015). Following spawning, adults rapidly senesce and die (Johnson et al. 2015).

After hatching, larvae drift downstream and burrow into stream sediments where they remain as filter feeders for the majority of their life until reaching a critical length and condition factor in the fall preceding metamorphosis (Holmes and Youson 1994; Manzon et al. 2015). This ensures that they have sufficient lipid reserves to undergo the non-feeding period of metamorphosis the following summer (Manzon et al. 2015). In the Great Lakes, age at metamorphosis is reported to vary from 3 to 7 years (Potter 1980), but interpretation of these values is complicated by the challenges of accurately estimating sea lamprey age and by the

variation in age at metamorphosis among tributaries and among individuals within a cohort (Dawson and Jones 2009; Potter 1980; Potts et al. 2015).

Sea lamprey do not return to natal tributaries to spawn, which causes genetic mixing during reproduction of individuals born in different tributaries; this feature of sea lamprey life history constrains the extent of local adaptation. However, lack of genetic differentiation at a local scale does not preclude the possibility of weak spatial stock structure or subtle genetic differences between sea lamprey residing far apart in geographically different regions of the Great Lakes (e.g., Hess et al. 2013; Lanca et al. 2014; Krabbenhoft and Dowling 2015). For example, a higher probability of spawning between individuals from the tributaries of a given lake, could create some weak spatial stock structure even though some mixing also occurs.

The Great Lakes Fishery Commission (GLFC) oversees sea lamprey control in the Great Lakes (reviewed in Siefkes et al. 2013). In the 1950s, initial attempts to control sea lamprey involved using traps and weirs to target adults migrating into spawning tributaries (McLain et al. 1965). The discovery that TFM could selectively kill larval sea lamprey (Applegate et al. 1957, 1958) led to its first application in Lake Superior in 1958, with full expansion of the program to the other lakes by the 1980s (Table 1). After its discovery in 1964, niclosamide was incorporated in lesser amounts (1-2%) into some TFM applications (Howell et al. 1964). The toxicity of niclosamide is not as species-specific, but can reduce the amount of TFM required for successful treatment. A granular formulation of niclosamide is also used in some deep tributaries and lentic areas where the application of TFM is not practical (Siefkes et al. 2013). Barriers placed into tributaries have also contributed to sea lamprey control by blocking upstream spawning migration and restricting access to spawning and larval habitat (McLaughlin

et al. 2013). Other control measures have been carried out to a lesser degree including trapping and release of sterile male sea lamprey (Christie and Goddard 2003). Lampricide application, in particular the continued use of TFM, is the foundation of sea lamprey control in the Great Lakes, both historically and for the foreseeable future.

Only 500 of the approximately 5,340 Great Lakes tributaries have produced sea lamprey and 336 of these have been treated at least once with lampricide (Adair and Sullivan 2014a). Tributaries do not require treatment on an annual basis because of the time required by larval sea lamprey to grow and reach metamorphosis. Instead, each year larval assessments are undertaken by Fisheries and Oceans Canada and the U.S. Fish and Wildlife Service to monitor the presence, relative abundance, and size structure of sea lamprey larvae in Great Lakes tributaries. The larvae within a stream are monitored to determine when they have reached sizes (>100 mm) predicted to result in metamorphosis. Streams are ranked for lampricide treatment based on the abundance of large larvae relative to the costs of treating that particular stream.

Resistance Strategies

Resistance is a microevolutionary process that occurs within a population over multiple generations. We define resistance as a reduction in sea lamprey mortality from lampricides as a consequence of genetic change in the population (Sawicki 1987; Mota-Sanchez et al. 2002; Tabashnik et al. 2014). Genetic change that is heritable across generations is required for an evolved response. Our definition includes two primary types of resistance, physiological and behavioural, each of which could result in reduced effectiveness of the sea lamprey control

program (Fig. 1). Our definition does not include epigenetic forms of inheritance that do not involve changes in DNA sequences (Jablonka and Raz 2009), although we recognize that epigenetic forms of resistance would also have consequences for sea lamprey control. Our definition also does not include tolerance or acclimation arising within an individual upon prior exposure to a sub-lethal concentration of a pesticide, because such changes are not transmitted across multiple generations (Wirgin and Waldman 2004; Hua et al. 2013). These non-genetic responses could increase chances of surviving pesticide treatment, but would be quickly reversible when the selective pressure is removed.

Three prerequisites must be met for resistance to evolve and be detectable. First, selection from the control measure needs to be sufficiently strong to elicit genetic adaptation. Second, phenotypic variation in the population that has a genetic basis must provide some individuals with an increased probability of surviving control measures. Third, adequate time to detect the genetic response must pass. The selection pressure needs to be sufficiently high to kill many individuals but not so high that the few survivors cannot find and reproduce with one another (Mota-Sanchez et al. 2002). The initial frequency of individuals with resistant genotypes provides the basis on which selection acts. Resistant genotypes might be present at an initial low level within the founding population (depending on the genetic variation) or arise through mutation or immigration of new individuals. If selection is strong enough and a sufficient number of generations have passed, the frequency of resistant individuals in the population will increase over time and eventually the presence of resistance will be detectable as a reduction in the mortality from the control measure.

Most documented examples of pesticide or antibiotic resistance are physiological (Williamson et al. 1996; Wright 2007), probably related to the fact that it can be demonstrated in the laboratory by measuring lethal concentrations, such as the LC50 or LC99 (the concentrations required to kill 50 or 99% of the population, respectively). Behavioural resistance is more difficult to demonstrate because the behaviour of targeted animals is often difficult to study in the field, behavioural studies of pests typically focus on populations rather than individuals, identifying the genetic basis for behaviour can be difficult, and toxicity tests conducted in the laboratory are not usually suitable for studying complex behaviours (Sparks et al. 1989; Ranson et al. 2011; Morales et al. 2013). Both types of resistance strategies can evolve in a population simultaneously, and interactions between behaviour and physiology occur (Sparks et al. 1989).

Behavioural Resistance

As a first line of defense, organisms can evolve behaviours to avoid encounter or reduce their uptake of a pesticide (Sparks et al. 1989; Ranson et al. 2011). One of the few well-documented examples of behavioral resistance occurred in the sheep blowfly *Lucilia cuprina* when females evolved the strategy of avoiding laying eggs in the presence of an insecticide aimed at their larvae (Mariath et al. 1990).

Behavioural resistance to lampricides could arise in sea lamprey via the evolution of behaviours or life history strategies that enable sea lamprey to avoid exposure or circumvent death despite exposure to a treatment. Tributaries are typically treated with lampricide on a cyclical basis to coincide with the time it takes larvae to reach their age and size at

metamorphosis. This repeated, cyclic nature of lampricide applications could select for earlier age at metamorphosis, thereby increasing the probability of individuals leaving a tributary between treatments. Similarly, lampricide treatment could select for individuals that avoid treatment by out-migrating to lakes before metamorphosis. Normally, out-migration is timed with metamorphosis, but the movement of larvae downstream as they age (Manion and McLain 1971; Manion and Smith 1978; Dawson et al. 2015) or the presence of larvae in lentic areas at the mouths of some tributaries (Fodale et al. 2003) suggests that other variables or behaviours could be involved in out-migration timing. Whether these strategies evolve will depend on balancing any advantages gained against any potential costs of these behaviours such as increased predation pressure or reduced growth. The presence of exogenous factors such as temperature and discharge that are potentially involved with triggering downstream movement (Dawson et al. 2015) might also have a bearing on the evolution of behavioural resistance.

Repeated lampricide treatments could also select for larval lamprey that prefer local habitats where lampricide exposure is reduced. These habitats could include stream reaches above locations where treatments are initiated or secondary habitats where it is difficult to achieve lethal concentrations of lampricide, such as side channels, backwaters, and oxbows (Adair and Sullivan 2014a). Similarly, treatment could select for changes in the behavioural response to lampricide exposure. Treatment crews observe larval sea lamprey moving into side channels, still-water areas, or onto mudflats upon lampricide exposure, which might offer a selective advantage if these behaviors have a genetic basis. Lampricide-induced selection could

potentially also act on burrowing behaviour, although evidence suggests that burrows offer little protection against lampricide (Bills et al. 2003).

Body condition and the timing, amount, and level of feeding activity could also evolve in response to lampricide treatment. Larvae are more susceptible to lampricide in spring than in summer (Applegate et al. 1961; Scholefield et al. 2008), likely a result of differences in body size and glycogen and lipid stores (Wilkie et al. 2014). Large, high condition sea lamprey suffer less mortality from lampricides, possibly creating a selection gradient that could lead to evolution of life history characteristics over time. In many ways the evolution of life history traits in response to lampricide treatment is similar in concept to the rapid evolution of traits like body size or maturation schedule that have been documented in fish stocks in response to harvest (Jørgensen et al. 2007).

Physiological Resistance

Once bypassing any behavioural means of avoiding or reducing uptake of the chemical, physiological adaptation provides the next opportunity for evolving resistance. Physiological mechanisms that enable uptake rates to be reduced or slowed at the gills could provide a site for selection. Following uptake of the chemical into the body, there are then many additional routes for physiological resistance to evolve (Fig. 1), all resulting in reduced toxicity of the chemical to the organism. In sea lamprey, these routes for physiological resistance to TFM and niclosamide would likely be related to the mode of action of these pesticides, which exert their toxicity by uncoupling mitochondrial oxidative phosphorylation (Niblett and Ballantyne 1976; Birceanu et al. 2011).

Oxidative phosphorylation takes place in the mitochondria of cells, and results in the production of adenosine triphosphate (ATP), which provides the energy needed by an organism to function. The process requires oxygen and takes place on the inner mitochondrial membrane, where three of the protein complexes (I, III, IV) that make-up the electron transport chain pump protons from the matrix of the mitochondria into the inter-membrane space. This creates a gradient for protons to flow back into the matrix. Owing to the low permeability of the inner membrane, the flow of the protons back into the matrix is directed through another protein complex, the ATP synthase. As the protons move down their gradient, energy is released that is used for ATP production.

Chemicals that uncouple oxidative phosphorylation target the inner mitochondrial membrane by either increasing its permeability to protons or by acting as protonophores that shuttle protons back into the mitochondrial matrix (Terada 1990). This reduces the proton gradient for the movement of protons into the mitochondrial matrix via the ATP synthase, lowering ATP production. Eventually, this uncoupling mechanism creates a mismatch between ATP supply and demand, forcing the organism to rely on anaerobic fuels such as glycogen, glucose or phosphocreatine to generate ATP (Birceanu 2009; Clifford et al. 2012; Birceanu et al. 2014). Once these energy stores are depleted, the organism can no longer meet its energy demands and death ensues (Hollingworth 2001).

Recent evidence suggests that TFM likely uncouples oxidative phosphorylation by acting as a protonophore, reducing ATP production in a concentration-dependent fashion in sea lamprey and some other species such as rainbow trout *Oncorhynchus mykiss* (Birceanu et al. 2011). Less is known about the mode of action of niclosamide in sea lamprey, but the weight of

evidence suggests that it too uncouples oxidative phosphorylation (Ishak et al. 1970; Nettles et al. 2001; Tao et al. 2014) by acting non-specifically as a protonophore (Jurgeit et al. 2012).

Several potential routes exist for the evolution of physiological resistance to TFM (and likely also niclosamide), and we describe only a few here. One route is for resistance to develop at the level of the mitochondria. For instance, a change in mitochondrial sensitivity to TFM could evolve via a change in mitochondrial membrane lipid composition (Monteiro et al. 2011) or an increase in the number of mitochondria in a cell which changes the efficacy of the lampricide. A transporter that facilitates the shuttling of TFM across the mitochondrial membrane could be another site for resistance development, although evidence for the existence of such a transporter for TFM is lacking. Another route of physiological resistance could come about through increased detoxification capacity. Phase II metabolism via glucuronidation of TFM, catalyzed by the enzyme UDP-glucuronyl transferase (UDP-GT), appears particularly important in the detoxification process, resulting in the formation of water soluble TFM-glucuronide, which is more easily excreted and likely to be inactive as an uncoupler (Lech and Statham 1975; Kane et al. 1994). Sea lamprey have very low glucuronidation capacity compared to some other fishes, which explains their higher sensitivity to TFM (Lech and Statham 1975; Kane et al. 1994). However, genetic variation among individual sea lamprey in their glucuronidation capacity (e.g., stemming from variation in the interaction between UDP-GT and TFM) could provide a site for resistance development. Genomic analyses indicate that sea lamprey express at least two isoforms of the UDP-GT gene (Smith et al. 2013), but further research is needed to better characterize the functional properties of the isoforms in sea lamprey, if we are to predict if it is a potential target for

resistance selection. Similar arguments also apply to niclosamide, but its detoxification is more complicated, involving not only glucuronidation but also sulfation, in which another sulfate group is added to the compound to facilitate its excretion (Hubert et al. 2005).

Like other pesticides that uncouple mitochondrial oxidative phosphorylation, TFM and niclosamide lack a specific binding site, which could limit the way physiological resistance evolves. Because TFM and niclosamide act as proton shuttles across the inner mitochondrial membrane, rather than targeting a specific protein (e.g., a receptor), it is difficult to envision how naturally selected changes could be induced at this level. A mutation at the binding site of a toxic chemical, a frequent cause of physiological resistance evolution in agricultural pests and bacteria (French-Constant et al. 2004; Wright 2007), therefore would not be a likely mode of resistance development to lampricide.

Despite the lack of a binding site, however, resistance to mitochondrial uncoupling pesticides has occurred in agricultural systems (e.g., Van Leeuwen et al. 2006; Ahmad and Arif 2009; Leroux and Walker 2013,). In one example, resistance was related to increased activity of a transmembrane transporter (Leroux and Walker 2013) and in another example, resistance appeared related to enhanced activity of the esterase enzyme (Van Leeuwen et al. 2006). Organisms have used a variety of mechanisms to evolve resistance to chemicals present in their environment, including strategies that don't involve binding site mutations (Posthuma and Van Straalen 1993; Despres et al. 2007).

Examples of Resistance from Other Systems

While arthropods have a long history of evolving mechanisms to defeat toxins (Mota-Sanchez et al. 2002), fewer examples of vertebrate pests developing resistance to chemical control have been documented. The best known vertebrate example is that of rodents evolving resistance to rodenticides that inhibit blood coagulation. Application of the anticoagulating rodenticides (most notably warfarin) began in the early 1950s, with the first published evidence of resistance in 1958 (Boyle 1960) and with documented cases of resistance arising in numerous countries since (Pelz et al. 2005). Another example of resistance in vertebrate pests is that of European rabbits (*Oryctolagus cuniculus*) developing resistance to Compound 1080, a pesticide used heavily in baited traps in Australia and New Zealand since the 1950s to control non-native mammals. The first signs of reduced bait effectiveness in rabbits arose in the 1970s (Oliver et al. 1982), with follow-up toxicity tests confirming resistance that was related to the population's history of exposure in the field (Twigg et al. 2002).

Some vertebrates, particularly fishes, have developed resistance to chemical toxicants present in their environment (Wirgin and Waldman 2004). Atlantic tomcod (*Microgadus tomcod*) in the Hudson River, NY offer one of the most compelling examples of a fish population rapidly evolving resistance to chemicals. When embryonic offspring of Atlantic tomcod from more pristine environments are exposed to coplanar polychlorinated biphenyls (PCBs) or 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), they typically develop a range of developmental abnormalities that likely result from impaired heart structure and function. However, the offspring of tomcod from the Hudson River, where release of industrial pollutants began in the late 1940s do not show these responses and instead demonstrate reduced embryonic mortality to PCBs and TCDD and decreased expression of the cytochrome P4501A (CYP1A) gene. Results

from controlled laboratory crosses revealed that the resistance was heritable across at least the F_2 generation (Wirgin et al. 2011). The genetic basis of resistance was shown to arise from a single two amino acid deletion in the aryl hydrocarbon receptor 2 (AHR2), a transcription factor that mediates toxicity and activation of numerous genes in the AHR2 pathway, including CYP1A. A standing variant of AHR2 was nearly fixed in the Hudson River population and almost absent elsewhere (Wirgin et al. 2011).

Several factors may have contributed to resistance evolution in Hudson River tomcod. The selective pressure experienced by the population was strong due to the chronic exposure of critical early life stages to high concentrations of these sediment-borne contaminants. Also, although the population was panmictic within the Hudson River, they were reproductively isolated from other sensitive populations, preventing re-introduction of sensitive alleles via gene flow and facilitating local adaptation (Wirgin et al. 2011). Finally, the fact that resistance was attributable to a single standing variant in one gene likely accelerated the evolutionary process. The year when resistance would have been first detectable is not known because the population was not tracked through time, so we only know that it occurred sometime within the 70 years after contamination began. The generation time of tomcod (approximately 1-2 years) is less than that of sea lamprey, but this example does demonstrate that fish populations are capable of rapidly evolving resistance to toxic chemicals in the environment.

Meeting the Pre-requisites for Resistance Evolution in Sea Lamprey

Resistance has been observed to evolve within a matter of weeks, months, years, or decades depending on the organism in question and the strength and mechanism of selection (Palumbi

2001). Below, we evaluate whether the selective environment has been conducive for resistance evolution by estimating the number of generations of sea lamprey which have been exposed to lampricides and the strength of the selective pressure from TFM.

Has there been enough time for evolutionary change to be observed?

Assuming a generation time of 5 years, Great Lakes sea lamprey have been exposed to lampricides for between 6 and 11.6 generations (Table 1), placing them within the range of other fish populations where evolution has been documented (Table 2). A 5-year generation time for sea lamprey is an approximation based on observations of time between recruitment and metamorphosis. The age at metamorphosis typically ranges from about 3 to 5 years, depending on the lake (generally, ages 3-4 for Michigan, Erie and Ontario; age 4 for Huron; age 5 for Superior; M. Steeves, personal observation), followed by a parasitic juvenile phase of 12 – 18 months prior to spawning (Bergstedt and Swink 1995). We therefore chose a generation time toward the lower end of the range because we wanted to evaluate the possibility that enough time has passed for resistance evolution. Even a more conservative estimate for generation time in Lake Superior of 7 years would mean exposure to lampricides for 8.3 generations, still well within the range reported for rapid genetic change in other fish populations (Table 2).

There is growing awareness of the rapid pace of evolution that is possible in vertebrate populations, most often as a result of human-induced selective pressures (Hendry et al. 2017). In fish populations, where generation times are comparable to sea lamprey, evolution has occurred on surprisingly short timescales in response to selection from a variety of sources

including harvesting, toxins, predation, and introduction into novel environments (Table 2). Based on a review of published accounts of evolution in fishes, we see that evolution has been documented in as few as 3 generations and spanning as many as 140 generations (Table 2). The generation times of fish in these studies ranged from months (e.g., 2-3 months in guppies *Poecilia reticulata*; van Wijk et al. 2013) to several years (e.g., about 4 years in sockeye salmon *Oncorhynchus nerka*; Hendry et al. 1998). The types of selective pressure were diverse, ranging from strong harvesting imposed in a laboratory, such as 90% harvest rate in Atlantic silversides *Menidia menidia* (Conover and Munch 2002) to natural selection acting on a population introduced to a novel environment, as observed for mosquitofish (*Gambusia affinis*) (Stockwell and Weeks 1999).

Rapid changes in fish life history attributed to evolution have also occurred in several wild stocks experiencing selective pressure from commercial fisheries (Jørgensen et al. 2007). The generation time of these commercial species are relatively long; for example close to a decade for Atlantic cod (*Gadus morhua*) (Eikeset et al. 2013). Trends in maturation-related traits attributed to fisheries-induced evolution have been reported in as few as 1.8 to 12 generations (Devine et al. 2012), although identifying the genetic contribution to the changes is difficult (Hansen et al. 2012).

What is the selective pressure from lampricides?

At this time, we don't have a direct measure of the strength of selection from lampricides on sea lamprey in the Great Lakes. Various approaches have been used to estimate selection pressure (Kinnison and Hendry 2001; Siepielski et al. 2009; Laugen et al. 2014), generally

requiring data on responses to selection for a particular trait, behaviour, or mutation. The selection gradient is a measure of direct selection on a trait, and can be measured as the covariation between the trait and fitness (Lande and Arnold 1983). In the case of sea lamprey resistance, where the source of selection is direct mortality from lampricides, the selection gradient is generally expected to increase as mortality increases because there is predicted to be a larger difference between the genotypes of those individuals that survive (the resistance genotypes) and those that are susceptible to the pesticide. The selection gradient together with the genetic variation of the trait in question will affect the magnitude or rate of evolution. For sea lamprey resistance, we lack a specific trait, mechanism, mutation, or behaviour for which to estimate selection gradients. For the purposes of this paper and until such data are available and analyzed, we use the mortality rate from lampricides as an indicator of the strength of selection but it must be emphasized that this is not a direct measure.

To quantify lampricide-induced mortality, we estimated the fraction of the sea lamprey population killed from lampricides since implementation of the control program. Making use of an operating model and existing data we estimated that lampricide-induced mortality in 2012 ranged from a high of 90% in Lake Huron and Ontario to a low of 82% in Lake Superior (Table 3; Appendix 1). For Lake Superior, the mortality estimate derived from the operating model (82%) was only slightly lower than an estimate (89%) obtained from control program data on larval abundances (Appendix 1). Furthermore, these estimates are close to previous suggestions that sea lamprey abundances in the Upper Great Lakes were reduced by 90% of pre-control levels (Smith and Tibbles 1980; Heinrich et al. 2003), although factors other than TFM application likely contributed to the declines in abundance (M. Siefkes pers. comm.). From these various

estimates, we conclude that the strength of selection from lampricides could have been substantial. To provide some context, these mortality rates are comparable to those used in harvest selection experiments on Atlantic silversides where life history evolution arose within 4 generations (Conover and Munch 2002).

Estimates of lampricide-induced mortality might underestimate the overall strength of selection from lampricides if there is selection in response to sub-lethal effects. Lampricide exposure has the potential to cause both mortality and sub-lethal effects in the sea lamprey that survive a treatment (Clifford et al. 2012). Many species exhibit sub-lethal responses to chemicals if exposed to concentrations insufficient to cause mortality (Relyea and Hoverman 2006). Examples of sub-lethal effects include changes in physiology, immune response, behaviour, and reproductive success (Weis et al. 2001; Relyea and Hoverman 2006). Sub-lethal responses to TFM could impact fitness and, if related to genetic variation among individuals, could evolve in response to lampricide treatment. In future, more direct measures of the strength of selection would enable more accurate predictions as to the likelihood and rate of resistance evolution in sea lamprey.

Evidence for Resistance in Sea Lamprey

Several aspects of sea lamprey biology and control offer clues about the likelihood of resistance evolution (Table 4). However, weighing these factors against the potential evolutionary costs of evolving strategies to evade control is challenging. Ultimately, experimental data are required to demonstrate that resistance has evolved or that the number of resistant individuals in the population is increasing.

Evidence for physiological resistance can be obtained by tracking changes in the lethal concentrations of a pesticide required to kill fixed percentages of test organisms. Taking this approach, we analyzed the outcomes of historical TFM toxicity tests that have regularly been conducted on sea lamprey larvae since 1956 (Appendix 2). Over the 57 year period of toxicity testing, no increase in the TFM concentration at which 50% (the LC₅₀ over 19-24 h) of the sample lamprey died (Fig. 2) was detected as predicted if resistance had developed. An earlier analysis covering a shorter period of time (30 years) also found no evidence of changes in TFM toxicity in larval sea lamprey (Scholefield and Seelye 1990).

Our findings of no change in the toxicity to TFM over time should be interpreted cautiously when drawing conclusions about the evidence for resistance. Our analyses relied on historical data that were collected for other purposes, not with the intent of specifically investigating the evolution of resistance (see Appendix 2 for more details about the tests). Sample sizes were low for some years and key variables known to influence toxicity (e.g., pH, source stream of the larvae, temperature, time of source larvae collection) were not always measured or controlled. Furthermore, the cause of higher estimates of LC₅₀s from the archival data relative to the other sources (Fig. 2) is unknown and could indicate differences in testing conditions. Appropriately designed experiments are needed to rule out the possibility of lampricide resistance in sea lamprey.

Looking to the Future: Strategies to Reduce the Risk of Resistance

Management actions aimed at reducing resistance to pesticides include (1) diversifying control measures, (2) allowing survival or immigration of susceptible individuals, (3) monitoring to

detect resistance, and (4) modeling to predict resistance development (Mota-Sanchez et al. 2002). A few of these strategies could be integrated into current sea lamprey control program procedures.

Diversifying control measures using integrated pest management (IPM) could be a key tool to reduce the likelihood of resistance evolution in sea lamprey. IPM includes using alternative control measures to reduce the ecological costs of chemical pesticides (reviewed in Kogan 1998); these alternative controls often involve behavioural manipulation of pests such as push-pull strategies (Cook et al. 2007) or biological control strategies (Lacey et al. 2001). IPM has been adopted in many agricultural systems to reduce reliance on chemical pesticides and in turn reduce contaminant levels in the environment, improve control, and reduce the incidence of resistance (Kogan 1998). The expectation is that it would be rare for an individual in a pest population to possess resistance genes for multiple control tactics that combine different aspects of an organism's life history, behaviour, or physiology. The selective environment becomes more heterogeneous in the presence of multiple control tactics, delaying the evolution of resistance (Bourguet et al. 2013).

As a concept, IPM has already been embraced by the GLFC and the sea lamprey control program (Sawyer 1980; Christie and Goddard 2003; GLFC 2011). The GLFC first endorsed IPM for sea lamprey control during the 1970s and adopted it officially as part of its policy in the 1980s (Sawyer 1980; Christie and Goddard 2003; Siefkes et al. 2013). This paved the way to exploring the development of alternative control measures including barriers, trapping, pheromones, and repellants (Siefkes et al. 2013). Besides barriers, however, these alternative control tactics are still under development and do not contribute to the current suppression of

sea lamprey. Barriers to migration do limit sea lamprey access to suitable spawning and nursery habitats, but are not present on all streams, rely in part on a network of deteriorating structures constructed for purposes other than sea lamprey control, and face constant pressure to restore watershed connectivity through barrier removal (Lavis et al. 2003a; McLaughlin et al. 2013). Thus, although the development of lampricide alternatives remains a top priority for the GLFC (Sawyer 1980; Christie and Goddard 2003; GLFC 2011), the ecology and economy of the Great Lakes fisheries currently depends on TFM to control sea lamprey.

This reliance on a single chemical to control a pest increases the chances of resistance development. Even considering that niclosamide is sometimes used, it is believed to have a similar mode of action. The use of an alternative lampricide that exerts its toxicity through a different mechanism, if discovered, could slow resistance if used alongside TFM. For this reason, the application of two or more toxins with different modes of action has been used to reduce resistance in insect pests (Comins 1986; Roush and Daly 1990; Carrière et al. 2016). The same logic applies when combining chemical pesticides with other control tactics such as barriers – the use of multiple control tactics, each exploiting a different aspect of sea lamprey biology, will reduce the likelihood of TFM resistance because selection would favor individuals based on a different suite of traits and underlying genotypes. In effect, selection for chemical resistance would be diluted in the presence of effective alternative control tactics.

Research on pheromones used by sea lamprey for reproduction and migration (Li et al. 2002; Meckley et al. 2014; Buchinger et al. 2015) and the sequencing of the sea lamprey genome (Smith et al. 2013) could pave the way for the development of alternative control techniques. Next generation (i.e., new) lampricides could be discovered that have a different

mode of action, or are more selective, environmentally friendly, and cost effective. For example, identifying circumstances under which a gene is expressed could be used to develop a lampricide that exploits a unique aspect of sea lamprey development (Buchinger et al. 2015; McCauley et al. 2015). In this respect, greater incorporation of high throughput methods, such as RNA sequencing to generate transcriptomes of potential target tissues or organs, could be very fruitful. An emerging genetic approach to control undesirable species and reverse resistance is the engineered gene drive method based on the CRISPR-Cas9 technology, a gene editing technology using guided RNA that can target and modify DNA and RNA sequences (Champer et al. 2016). The gene drive elements are inherited by offspring and result in the spreading of sterility within the population.

Design and implementation of effective barriers, potentially in association with the use of pheromones (Johnson et al. 2009; Buchinger et al. 2015), offers another potential option for reducing resistance to TFM. Effective barriers have the potential to alter which sea lamprey individuals successfully reproduce, thereby altering the fitness landscape and potentially mitigating selection for pesticide resistance. When upstream migrating sea lamprey encounter a barrier, several responses have been documented including swimming around to find ways to circumvent the barrier or to evade capture within traps, fallback and spawning further downstream, or leaving the tributary to find an alternative spawning site (e.g., Bravener and McLaughlin 2013; McLean et al. 2015; Holbrook et al. 2016). These responses require time and energy that could provide a fitness advantage to individuals with certain heritable characteristics; as with other alternative control techniques, it becomes increasingly unlikely

that individuals would have the suite of genetic characteristics that confer advantages in an environment with both barriers and chemical treatment.

As mentioned above, barriers do contribute to control of sea lamprey but there is growing pressure to remove them in many locations to restore connectivity and reduce habitat fragmentation for other organisms (McLaughlin et al. 2013). Often interests conflict between keeping (or repairing) a barrier to prevent sea lamprey from accessing upstream spawning habitat versus removing a barrier to improve accessibility to upstream habitat for desirable fishes. The management decisions in these cases are not straightforward. Furthermore, barriers are not present on every stream with sea lamprey spawning habitat and many of those that exist are in disrepair. Thus, there is scope for increasing the effectiveness of barriers as a control tactic. Improvements to barrier design that limit sea lamprey passage but permit movement of other fish species (Sherburne and Reinhardt 2016) could provide an alternative control measure used to slow TFM resistance that also serves additional management objectives.

Facilitating the survival and emigration of susceptible individuals is another strategy used to reduce pesticide resistance. This strategy is often employed in agricultural systems to preserve susceptible genes in the population by creating refuges where treatment is not applied, but permitting some gene flow between treated and untreated areas (Gould 1998). At first glance, current control of sea lamprey might appear to be a type of refuge strategy because many stream reaches are untreated. However, the annual, high degree of reproductive mixing of adults makes sea lamprey control different from a true refuge strategy. The absence of natal homing weakens the formation of discrete stocks, making it difficult to

have refuge populations that have never been exposed to lampricides (i.e., because unexposed individuals would mate with exposed individuals at a high rate). A true refuge strategy would require a higher level of reproductive isolation over a number of years between the refuge population and the populations that are exposed – this is not the case with sea lamprey where the reproductive mixing reduces the reproductive isolation of unexposed individuals. However, although different from a refuge strategy in the strict spatial sense, current sea lamprey control does provide opportunities for sea lamprey individuals to survive and reproduce even if they might be susceptible to TFM if exposed. Furthermore, the variety of ways a sea lamprey could evade mortality from TFM in a given year might weaken the strength of selection for any one type of physiological resistance (Table 3; see Behavioral Resistance section), although it could also increase the overall odds of a mutation developing or promote behavioural resistance evolution. Looking to the future, there might be ways of further taking advantage of those sea lamprey already surviving despite possessing susceptible genes. A more direct type of refuge strategy for sea lamprey could include artificially supplementing the wild population with individuals from a population that has not been exposed to lampricides (e.g., an ancestral source population or a population maintained in the laboratory). However, a refuge approach would require careful consideration and weighing of the potential risks associated with increasing the number of spawning adults and enhancing genetic variation in the population versus the benefits of potentially reducing the potential for resistance.

Monitoring for resistance is an integral component to resistance prevention. This is the case whether lampricide resistance arises slowly, with time to act once initial low numbers of resistant individuals are detected, or spread quickly with less opportunity but more urgency to

604 modify operations. In insect pests, the presence of low numbers of resistant individuals
605 provides a warning sign that resistance could become a serious problem if strong selection
606 continues. Early detection has proven to be important in agricultural systems because it is
607 often too late to implement strategies for managing resistance once pesticide failure occurs
608 (Mota-Sanchez et al. 2002; Barres et al. 2016).

609 Routine post-treatment larval assessments already conducted could be used or
610 augmented for the purpose of monitoring for resistance. For example, changes in residual
611 larvae abundances or life history could be tracked and studied. In addition, the control
612 program should develop an annual, standardized toxicity test specifically designed to monitor
613 physiological resistance development. This test could be integrated into existing procedures to
614 conduct quality assurance of new lampricide batches. Residual lamprey that survive treatment
615 in the field or in a laboratory toxicity test could be collected, stored, and later used to
616 investigate mechanisms of resistance development. One potential test for resistance would be
617 a comparison of toxicity between sea lamprey from the Great Lakes and those from a source
618 population outside the Great Lakes basin. Monitoring for physiological resistance would be
619 more straightforward than monitoring for behavioural resistance because of the added
620 challenge of replicating field conditions and inducing natural behavioural responses in a
621 laboratory environment.

622 Screening for resistance alleles, in particular before resistance becomes a problem, can
623 provide valuable information. The initial frequency of resistant alleles in a population critically
624 affects the risk of resistance development, as does the mode of inheritance, i.e., whether the
625 trait is recessive or dominant (Gould et al. 1997). Laboratory crosses where subsequent

generations were followed in the lab (Gould et al. 1997) and DNA-based screening to identify recessive alleles (Gahan et al. 2001) have been used for monitoring the critical early phases of resistance development in insects. Furthermore, data on the frequency of resistant alleles in a population are critical for models which themselves can be a useful tool for predicting, understanding, and managing resistance (Bourguet et al. 2010; Renton et al. 2014).

Beyond Sea Lamprey: Implications for Invasive Species Control and Ecosystem Management

The possibility of sea lamprey developing resistance to chemical control is relevant more broadly in the context of invasive species management. Invasive species possess the capacity to evolve a diversity of physiological and behavioural strategies in response to control and management measures. Many of these strategies better known for bacteria or insects appear relevant for other taxa, including vertebrates. For example, the routes of potential resistance evolution we outlined for sea lamprey (Fig. 1), have basic similarities to the mechanisms insects have naturally evolved to resist plant toxins (Despres et al. 2007).

In many cases, the methods used to control invasive species are likely to cause the strong, directional selection needed to facilitate resistance evolution. Assessing the magnitude of selection from control poses challenges because of compensatory demographic processes and because it can be difficult to tease apart environmental effects when examining phenotypic data collected in the wild. However, in the fight against an invasive species, target mortality rates from a given control measure could be high, and when implemented could produce selection differentials comparable to those that have caused rapid evolution in experimental

settings and in simulations (Conover and Munch 2002; Reznick and Ghalambor 2005; Dunlop et al. 2009).

Awareness that evolution in response to human-induced selection pressure can occur quickly in animal populations is becoming more common. We did not find any evidence that physiological resistance to TFM has developed thus far in sea lamprey, possibly because not enough generations have passed or for some other reason related to sea lamprey biology or control (Table 3). However, numerous other aquatic invasive species have shorter generation times, increasing the speed over which evolution could arise. For example, two prominent invasive fish species, round goby (*Neogobius melanostomus*) and bighead carp (*Hypophthalmichthys nobilis*), can mature at ages of 1-2 years and 2-4 years, respectively (Gutowsky and Fox 2012; Cooke 2016). Furthermore, rapid evolution can occur within timescales relevant to management even in species with later ages of maturity, including fishes (Fraser 2013).

The perception that fish populations will not evolve over time scales relevant to management is outdated. The development of resistance is just one example of how the speed and capacity of populations to evolve is integral to many major global challenges (Carroll et al. 2014). Species targeted for control or harvest can evolve strategies for evasion more quickly than desired, while species targeted for conservation efforts may evolve too slowly in response to climate change and other stressors. Control programs like that for sea lamprey in the Great Lakes can serve as quasi-experiments that help us understand how the risks of resistance evolution extend to a broader range of taxa, ecological circumstances, and management challenges.

670

671 **Acknowledgments**

672 We are grateful to Karen Slaght at the Hammond Bay Biological Station of the United States
673 Geological Survey (USGS) for her assistance with the gathering, organizing, and input of data
674 used for our toxicity analyses. Two anonymous reviewers provided suggestions that improved
675 our paper. This is a contribution from the lampricide resistance workshop funded by the Great
676 Lakes Fishery Commission on 13-15 January 2015. We are grateful to the presenters and
677 workshop audience who provided valuable discussion.

Draft

References

- Adair, R., and Sullivan, P. 2014a. Sea lamprey control in the Great Lakes. Annual Report to the Great Lakes Fishery Commission, Ann Arbor, M.I. Available from http://www.glfc.org/sealamp/ANNUAL_REPORT_2014.pdf [accessed 10 November 2016].
- Adair, R.A., and Sullivan, W.P. 2014b. Standard operating procedures for application of Lampricides in the Great Lakes Fishery Commission integrated management of sea lamprey (*Petromyzon marinus*) control program, Marquette Biological Station, U.S. Fish and Wildlife Service, Marquette, M.I. Available from <http://www.glfc.org/sealamp/sop.php> [accessed 10 November 2016]
- Ahmad, M., and Arif, M.I. 2009. Resistance of Pakistani field populations of spotted bollworm *Earias vittella* (Lepidoptera: Noctuidae) to pyrethroid, organophosphorus and new chemical insecticides. *Pest Manag. Sci.* **65**(4): 433-439.
- Alsan, M., Schoemaker, L., Eggleston, K., Kammili, N., Kolli, P., and Bhattacharya, J. 2015. Out-of-pocket health expenditures and antimicrobial resistance in low-income and middle-income countries: an economic analysis. *Lancet Infect. Dis.* **15**(10): 1203-1210.
- Applegate, V.C., Howell, J.H., Miffed, J.W., Johnson, B.G.H., and Smith, M.A. 1961. Use of 3-trifluoro- methyl-4-nitrophenol as a selective sea lamprey larvicide. Great Lakes Fishery Commission Technical Report 1, Ann Arbor, M.I.
- Applegate, V.C., Howell, J.H., and Smith, M.A. 1957. Toxicity of 4,346 chemicals to larval lampreys and fishes. United States Fish and Wildlife Service Special Science Report Fisheries No. 207. Washington, D.C.

- 700 Applegate, V.C., Howell, J.H., and Smith, M.A. 1958. Use of mononitrophenols containing
701 halogens as selective sea lamprey larvicides. *Science* **127**(3294): 336-338.
- 702 Aykanat, T., Thrower, F.P., and Heath, D.D. 2011. Rapid evolution of osmoregulatory function by
703 modification of gene transcription in steelhead trout. *Genetica* **139**(2): 233-242.
- 704 Barres, B., Micoud, A., Corio-Costet, M.F., Debieu, D., Fillinger, S., Walker, A.S., Delye, C.,
705 Grosman, J., Siegwart, M., and Network, R.P. 2016. Trends and challenges in pesticide
706 resistance detection. *Trends Plant Sci.* **21**(10): 834-853.
- 707 Bergstedt, R.A., and Seelye, J.G. 1995. Evidence for lack of homing by sea lampreys. *Trans. Am.*
708 *Fish. Soc.* **124**(2): 235-239.
- 709 Bergstedt, R.A., and Swink, W.D. 1995. Seasonal growth and duration of the parasitic life stage
710 of the landlocked sea lamprey (*Petromyzon marinus*). *Can. J. Fish. Aquat. Sci.* **52**(6):
711 1257-1264.
- 712 Bills, T.D., Boogaard, M.A., Johnson, D.A., Brege, D.C., Scholefield, R.J., Westman, R.W., and
713 Stephens, B.E. 2003. Development of a pH/alkalinity treatment model for applications of
714 the lampricide TFM to streams tributary to the Great Lakes. *J. Gt. Lakes Res.* **29**: 510-
715 520.
- 716 Birceanu, O. 2009. Mechanism(s) of 3-trifluoromethyl-4-nitrophenol (TFM) toxicity in sea
717 lamprey (*Petromyzon marinus*) & rainbow trout (*Oncorhynchus mykiss*). M.Sc. thesis,
718 Department of Biology, Wilfrid Laurier University, Waterloo, O.N.
- 719 Birceanu, O., McClelland, G.B., Wang, Y.X.S., Brown, J.C.L., and Wilkie, M.P. 2011. The
720 lampricide 3-trifluoromethyl-4-nitrophenol (TFM) uncouples mitochondrial oxidative
721 phosphorylation in both sea lamprey (*Petromyzon marinus*) and TFM-tolerant rainbow

- 722 trout (*Oncorhynchus mykiss*). Comp. Biochem. Physiol. C-Toxicol. Pharmacol. **153**(3):
723 342-349.
- 724 Birceanu, O., Sorensen, L.A., Henry, M., McClelland, G.B., Wang, Y.X.S., and Wilkie, M.P. 2014.
725 The effects of the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) on fuel stores and
726 ion balance in a non-target fish, the rainbow trout (*Oncorhynchus mykiss*). Comp.
727 Biochem. Physiol. C-Toxicol. Pharmacol. **160**: 30-41.
- 728 Bourguet, D., Delmotte, F., Franck, P., Guillemaud, T., Reboud, X., Vacher, C., and Walker, A.S.
729 2010. The skill and style to model the evolution of resistance to pesticides and drugs.
730 Evol. Appl. **3**(4): 375-390.
- 731 Bourguet, D., Delmotte, F., Franck, P., Guillemaud, T., Reboud, X., Vacher, C., Walker, A.S. 2013.
732 Heterogeneity of selection and the evolution of resistance. Trends Ecol. Evol. **28**(2): 110-
733 118.
- 734 Boyle, C.M. 1960. Case of apparent resistance of *Rattus norvegicus*. Nature **188**: 517.
- 735 Bravener, G.A., and McLaughlin, R.L. 2013. A behavioural framework for trapping success and
736 its application to invasive sea lamprey. Can. J. Fish. Aquat. Sci. **70**(10): 1438-1446.
- 737 Breckels, R.D., and Neff, B.D. 2014. Rapid evolution of sperm length in response to increased
738 temperature in an ectothermic fish. Evol. Ecol. **28**(3): 521-533.
- 739 Buchinger, T.J., Siefkes, M.J., Zielinski, B.S., Brant, C.O., and Li, W. 2015. Chemical cues and
740 pheromones in the sea lamprey (*Petromyzon marinus*). Front. Zool. **12**: 32.
- 741 Buchinger, T.J., Wang, H., Li, W., and Johnson, N.S. 2013. Evidence for a receiver bias underlying
742 female preference for a male mating pheromone in sea lamprey. Proc. R. Soc. B-Biol. Sci.
743 **280**(1771).

- 744 Carrière, Y., Fabrick, J.A., and Tabashnik, B.E. 2016. Advances in managing pest resistance to Bt
745 crops: pyramids and seed mixtures. *In* Advances in insect control and resistance
746 management. *Edited by* A.R. Horowitz and I. Ishaaya. Springer, New York. pp. 263-286.
- 747 Carroll, S.P., Jorgensen, P.S., Kinnison, M.T., Bergstrom, C.T., Denison, R.F., Gluckman, P., Smith,
748 T.B., Strauss, S.Y., and Tabashnik, B.E. 2014. Applying evolutionary biology to address
749 global challenges. *Science* **346**(6207): 313-+.
- 750 Champer, J., Buchman, A., and Akbari, O.S. 2016. Cheating evolution: engineering gene drives to
751 manipulate the fate of wild populations. *Nat. Rev. Genet.* **17**(3): 146-159.
- 752 Christie, G.C., and Goddard, C.I. 2003. Sea lamprey international symposium (SLIS II): advances
753 in the integrated management of sea lamprey in the Great Lakes. *J. Gt. Lakes Res.* **29**: 1-
754 14.
- 755 Clifford, A.M., Henry, M., Bergstedt, R., McDonald, D.G., Smits, A.S., and Wilkie, M.P. 2012.
756 Recovery of larval sea lampreys from short-term exposure to the pesticide 3-
757 trifluoromethyl-4-nitrophenol: implications for sea lamprey control in the Great Lakes.
758 *Trans. Am. Fish. Soc.* **141**(6): 1697-1710.
- 759 Comins, H.N. 1986. Tactics for resistance management using multiple pesticides. *Agric. Ecosyst.*
760 *Environ.* **16**(2): 129-148.
- 761 Conover, D.O., and Munch, S.B. 2002. Sustaining fisheries yields over evolutionary time scales.
762 *Science* **297**(5578): 94-96.
- 763 Conover, D.O., Munch, S.B., and Arnott, S.A. 2009. Reversal of evolutionary downsizing caused
764 by selective harvest of large fish. *Proc. R. Soc. B-Biol. Sci.* **276**(1664): 2015-2020.
- 765 Cook, S.M., Khan, Z.R., and Pickett, J.A. 2007. The use of push-pull strategies in integrated pest

- 766 management. Annu. Rev. Entomol. **52**: 375-400.
- 767 Cooke, S.L. 2016. Anticipating the spread and ecological effects of invasive bigheaded carps
768 (*Hypophthalmichthys* spp.) in North America: a review of modeling and other predictive
769 studies. Biol. Invasions **18**(2): 315-344.
- 770 Davies, J., and Davies, D. 2010. Origins and evolution of antibiotic resistance. Microbiol. Mol.
771 Biol. Rev. **74**(3): 417-433.
- 772 Dawson, H.A., and Jones, M.L. 2009. Factors affecting recruitment dynamics of Great Lakes sea
773 lamprey (*Petromyzon marinus*) populations. J. Gt. Lakes Res. **35**(3): 353-360.
- 774 Dawson, H.A., Quintella, B.R., Almeida, P.R., Treble, A.J., and Jolley, J.C. 2015. The ecology of
775 larval and metamorphosing lampreys. *In* Lampreys: biology, conservation, and control.
776 Edited by M.F. Docker. Springer, Dordrecht. pp. 75-138.
- 777 Despres, L., David, J.P., and Gallet, C. 2007. The evolutionary ecology of insect resistance to
778 plant chemicals. Trends Ecol. Evol. **22**(6): 298-307.
- 779 Devine, J.A., Wright, P.J., Pardoe, H.E., and Heino, M. 2012. Comparing rates of contemporary
780 evolution in life-history traits for exploited fish stocks. Can. J. Fish. Aquat. Sci. **69**(6):
781 1105-1120.
- 782 Dlugosch, K.M., and Parker, I.M. 2008. Founding events in species invasions: genetic variation,
783 adaptive evolution, and the role of multiple introductions. Mol. Ecol. **17**(1): 431-449.
- 784 Du, X., Crawford, D.L., Nacci, D.E., and Oleksiak, M.F. 2016. Heritable oxidative phosphorylation
785 differences in a pollutant resistant *Fundulus heteroclitus* population. Aquat. Toxicol. **177**:
786 44-50.
- 787 Dunlop, E.S., Heino, M., and Dieckmann, U. 2009. Eco-genetic modeling of contemporary life-

- 788 history evolution. Ecol. Appl. **19**(7): 1815-1834.
- 789 Eikeset, A.M., Richter, A., Dunlop, E.S., Dieckmann, U., and Stenseth, N.C. 2013. The economic
790 repercussions of fisheries-induced evolution. Proc. Natl. Acad. Sci. U. S. A. **30**: 12259-
791 12264.
- 792 Enayati, A., and Hemingway, J. 2010. Malaria management: past, present, and future. Annu.
793 Rev. Entomol. **55**: 569-591.
- 794 Enberg, K., Jorgensen, C., Dunlop, E.S., Heino, M., and Dieckmann, U. 2009. Implications of
795 fisheries-induced evolution for stock rebuilding and recovery. Evol. Appl. **2**(3): 394-414.
- 796 Eshenroder, R.L. 2014. The role of the Champlain Canal and Erie Canal as putative corridors for
797 colonization of Lake Champlain and Lake Ontario by sea lampreys. Trans. Am. Fish. Soc.
798 (3): 634-649.
- 799 French-Constant, R.H., Daborn, P.J., and Le Goff, G. 2004. The genetics and genomics of
800 insecticide resistance. Trends Genet. **20**(3): 163-170.
- 801 Fine, J.M., Vrieze, L.A., and Sorensen, P.W. 2004. Evidence that petromyzontid lampreys employ
802 a common migratory pheromone that is partially comprised of bile acids. J. Chem. Ecol.
803 **30**(11): 2091-2110.
- 804 Fodale, M.F., Bronte, C.R., Bergstedt, R.A., Cuddy, D.W., and Adams, J.V. 2003. Classification of
805 lentic habitat for sea lamprey (*Petromyzon marinus*) larvae using a remote seabed
806 classification device. J. Gt. Lakes Res. **29**: 190-203.
- 807 Fraser, D.J. 2013. The emerging synthesis of evolution with ecology in fisheries science. Can. J.
808 Fish. Aquat. Sci. **70**(9): 1417-1428.
- 809 Gahan, L.J., Gould, F., and Heckel, D.G. 2001. Identification of a gene associated with Bt

- 810 resistance in *Heliothis virescens*. Science **293**(5531): 857-860.
- 811 Goddard, C. 2013. Forward to Great Lakes Fisheries Policy and Management. *In* Great Lakes
812 fisheries policy and management. *Edited by* W.W. Taylor, A.J. Lynch and N.J. Leonard.
813 Michigan State University Press, East Lansing, M.I.. pp. ix-xi.
- 814 Gordon, S.P., Reznick, D., Arendt, J.D., Roughton, A., Hernandez, M.N.O., Bentzen, P., and
815 Lopez-Sepulcre, A. 2015. Selection analysis on the rapid evolution of a secondary sexual
816 trait. Proc. R. Soc. B-Biol. Sci. **282**(1813): 20151244.
- 817 Gould, F. 1998. Sustainability of transgenic insecticidal cultivars: Integrating pest genetics and
818 ecology. Annu. Rev. Entomol. **43**: 701-726.
- 819 Gould, F., Anderson, A., Jones, A., Sumerford, D., Heckel, D.G., Lopez, J., Micinski, S., Leonard,
820 R., and Laster, M. 1997. Initial frequency of alleles for resistance to *Bacillus thuringiensis*
821 toxins in field populations of *Heliothis virescens*. Proc. Natl. Acad. Sci. U. S. A. **94**(8):
822 3519-3523.
- 823 GLFC (Great Lakes Fishery Commission). 2011. Strategic Vision of the Great Lakes Fishery
824 Commission
825 2011-2020, Ann Arbor, M.I.
- 826 Gutowsky, L.F.G., and Fox, M.G. 2012. Intra-population variability of life-history traits and
827 growth during range expansion of the invasive round goby, *Neogobius melanostomus*.
828 Fisheries Manag. Ecol. **19**(1): 78-88.
- 829 Hansen, M.M., Olivieri, I., Waller, D.M., Nielsen, E.E., and Ge, M.W.G. 2012. Monitoring
830 adaptive genetic responses to environmental change. Mol. Ecol. **21**(6): 1311-1329.
- 831 Heinrich, J.W., Mullett, K.M., Hansen, M.J., Adams, J.V., Klar, G.T., Johnson, D.A., Christie, G.C.,

- 832 and Young, R.J. 2003. Sea lamprey abundance and management in Lake Superior, 1957
833 to 1999. J. Gt. Lakes Res. **29**: 566-583.
- 834 Hendry, A.P., Gotanda, K.M., and Svensson, E.I. 2017. Human influences on evolution, and the
835 ecological and societal consequences Introduction. Philos. Trans. R. Soc. B-Biol. Sci.
836 **372**(1712):13.
- 837 Hendry, A.P., Hensleigh, J.E., and Reisenbichler, R.R. 1998. Incubation temperature,
838 developmental biology, and the divergence of sockeye salmon (*Oncorhynchus nerka*)
839 within Lake Washington. Can. J. Fish. Aquat. Sci. **55**(6): 1387-1394.
- 840 Hendry, A.P., Kinnison, M.T., Heino, M., Day, T., Smith, T.B., Fitt, G., Bergstrom, C.T., Oakeshott,
841 J., Jorgensen, P.S., Zalucki, M.P., Gilchrist, G., Southerton, S., Sih, A., Strauss, S., Denison,
842 R.F., and Carroll, S.P. 2011. Evolutionary principles and their practical application. Evol.
843 Appl. **4**(2): 159-183.
- 844 Hess, J.E., Campbell, N.R., Close, D.A., Docker, M.F., and Narum, S.R. 2013. Population genomics
845 of Pacific lamprey: adaptive variation in a highly dispersive species. Mol. Ecol. **22**(11):
846 2898-2916.
- 847 Holbrook, C.M., Bergstedt, R.A., Barber, J., Bravener, G.A., Jones, M.L., and Krueger, C.C. 2016.
848 Evaluating harvest-based control of invasive fish with telemetry: performance of sea
849 lamprey traps in the Great Lakes. Ecol. Appl. **26**(6): 1595-1609.
- 850 Hollingworth, R.M. 2001. Inhibitors and uncouplers of mitochondrial oxidative
851 phosphorylation. *In* Handbook of Pesticide Toxicology. *Edited by* R.I. Krieger. Academic
852 Press, San Diego, C.A.
- 853 Holmes, J.A., and Youson, J.H. 1994. Fall condition factor and temperature influence the

- 854 incidence of metamorphosis in sea lampreys, *Petromyzon marinus*. Can. J. Zool. **72**(6):
855 1134-1140.
- 856 Howard, D.H., and Scott, R.D. 2005. The economic burden of drug resistance. Clin. Infect. Dis.
857 **41**: S283-S286.
- 858 Howard, D.H., Scott, R.D., Packard, R., and Jones, D. 2003. The global impact of drug resistance.
859 Clin. Infect. Dis. **36**: S4-S10.
- 860 Howell, J.H., King, J.E.L., Smith, A.J., and Hanson, L.H. 1964. Synergism of 5,2'-dichloro-4-
861 nitrosalicylanilide and 3-trifluoromethyl-4-nitrophenol in a selective lamprey larvicide.
- 862 Hua, J., Morehouse, N.I., and Relyea, R. 2013. Pesticide tolerance in amphibians: induced
863 tolerance in susceptible populations, constitutive tolerance in tolerant populations.
864 Evol. Appl. **6**(7): 1028-1040.
- 865 Hubert, J.N., Allal, F., Hervet, C., Ravakarivelo, M., Jeney, Z., Vergnet, A., Guyomard, R., and
866 Vandeputte, M. 2016. How could fully scaled carps appear in natural waters in
867 Madagascar? Proc. R. Soc. B-Biol. Sci. **283**(1837): 20160945.
- 868 Hubert, T.D., Bernardy, J.A., Vue, C., Dawson, V.K., Boogaard, M.A., Schreier, T.M., and
869 Gingerich, W.H. 2005. Residues of the lampricides 3-trifluoromethyl-4-nitrophenol and
870 niclosamide in muscle tissue of rainbow trout. J. Agric. Food Chem. **53**(13): 5342-5346.
- 871 Ishak, M.M., Sharaf, A.A., Mohamed, A.M., and Mousa, A.H. 1970. Studies on the mode of
872 action of some molluscicides on the snail, *Biomphalaria alexandrina*: I. Effect of
873 bayluscide, sodium pentachlorophenate, and copper sulphate on succinate, glutamate,
874 and reduced TMPD oxidation. Comp. Gen. Pharmacol. **1**: 201-208.
- 875 Jablonka, E., and Raz, G. 2009. Transgenerational epigenetic inheritance: Prevalence,

- 876 mechanisms, and implications for the study of heredity and evolution. *Q. Rev. Biol.*
877 **84**(2): 131-176.
- 878 Johnson, N.S., Buchinger, T.J., and Li, W. 2015. Reproductive ecology of lampreys. *In* Lampreys:
879 biology, conservation and control. *Edited by* M.F. Docker. Springer, Dordrecht. pp. 265-
880 303.
- 881 Johnson, N.S., Yun, S.S., Thompson, H.T., Brant, C.O., and Li, W.M. 2009. A synthesized
882 pheromone induces upstream movement in female sea lamprey and summons them
883 into traps. *Proc. Natl. Acad. Sci. U. S. A.* **106**(4): 1021-1026.
- 884 Jones, M.L., Bergstedt, R.A., Tvohey, M.B., Fodale, M.F., Cuddy, D.W., and Slades, J.W. 2003.
885 Compensatory mechanisms in great lakes sea lamprey populations: Implications for
886 alternative control strategies. *J. Gt. Lakes Res.* **29**: 113-129.
- 887 Jones, M.L., Irwin, B.J., Hansen, G.J.A., Dawson, H.A., Treble, A.J., Liu, W., Dai, W., and Bence,
888 J.R. 2009. An operating model for the integrated pest management of Great Lakes sea
889 lampreys. *The Open Fish Science Journal* **2**: 59-73.
- 890 Jørgensen, C., Enberg, K., Dunlop, E.S., Arlinghaus, R., Boukal, D.S., Brander, K., Ernande, B.,
891 Gårdmark, A., Johnston, F., Matsumura, S., Pardoe, H., Raab, K., Silva, A., Vainikka, A.,
892 Dieckmann, U., Heino, M., and Rijnsdorp, A.D. 2007. Managing evolving fish stocks
893 *Science* **318**: 1247-1248.
- 894 Jurgeit, A., McDowell, R., Moese, S., Meldrum, E., Schwendener, R., and Greber, U.F. 2012.
895 Niclosamide is a proton carrier and targets acidic endosomes with broad antiviral
896 effects. *PLoS Pathog.* **8**(10): 1-14.
- 897 Kane, A.S., Kahng, M.W., Reimschuessel, R., Nhamburo, P.T., and Lipsky, M.M. 1994. UDP-

- 898 Glucuronosyl transferase kinetics for 3-trifluoromethyl-4-nitrophenol (TFM) in fish.
899 Trans. Am. Fish. Soc. **123**(2): 217-222.
- 900 Kavanagh, K.D., Haugen, T.O., Gregersen, F., Jernvall, J., and Vollestad, L.A. 2010. Contemporary
901 temperature-driven divergence in a Nordic freshwater fish under conditions commonly
902 thought to hinder adaptation. BMC Evol. Biol. **10**: 12.
- 903 Kinnison, M.T., and Hendry, A.P. 2001. The pace of modern life II: from rates of contemporary
904 microevolution to pattern and process. Genetica **112**: 145-164.
- 905 Kinnison, M.T., Unwin, M.J., Hershberger, W.K., and Quinn, T.P. 1998a. Egg size, fecundity, and
906 development rate of two introduced New Zealand chinook salmon (*Oncorhynchus*
907 *tshawytscha*) populations. Can. J. Fish. Aquat. Sci. **55**(8): 1946-1953.
- 908 Kinnison, M.T., Unwin, M.J., and Quinn, T.P. 1998b. Growth and salinity tolerance of juvenile
909 chinook salmon (*Oncorhynchus tshawytscha*) from two introduced New Zealand
910 populations. Can. J. Zool. **76**(12): 2219-2226.
- 911 Kogan, M. 1998. Integrated pest management: Historical perspectives and contemporary
912 developments. Annu. Rev. Entomol. **43**: 243-270.
- 913 Krabbenhoft, T.J., and Dowling, T.E. 2015. Assessment of population genomics as a tool for
914 discriminating among populations of Great Lakes sea lamprey. 2015 Project Completion
915 Report, Great Lakes Fishery Commission, Ann Arbor, M.I.
- 916 Lanca, M.J., Machado, M., Mateus, C.S., Lourenco, M., Ferreira, A.F., Quintella, B.R., and
917 Almeida, P.R. 2014. Investigating population structure of sea lamprey (*Petromyzon*
918 *marinus*, L.) in Western Iberian Peninsula using morphological characters and heart fatty
919 acid signature analyses. PLoS One **9**(9): 14.

- 920 Lande, R., and Arnold, S.J. 1983. The measurement of selection on correlated characters.
921 Evolution **37**(6): 1210-1226.
- 922 Larson, G.L., Christie, G.C., Johnson, D.A., Koonce, J.F., Mullett, K.M., and Sullivan, W.P. 2003.
923 The history of sea lamprey control in Lake Ontario and updated estimates of
924 suppression targets. J. Gt. Lakes Res. **29**: 637-654.
- 925 Laugen, A.T., Engelhard, G.H., Whitlock, R., Arlinghaus, R., Dankel, D.J., Dunlop, E.S., Eikeset,
926 A.M., Enberg, K., Jorgensen, C., Matsumura, S., Nussle, S., Urbach, D., Baulier, L., Boukal,
927 D.S., Ernande, B., Johnston, F.D., Mollet, F., Pardoe, H., Therkildsen, N.O., Uusi-Heikkila,
928 S., Vainikka, A., Heino, M., Rijnsdorp, A.D., and Dieckmann, U. 2014. Evolutionary impact
929 assessment: accounting for evolutionary consequences of fishing in an ecosystem
930 approach to fisheries management. Fish. Fish. **15**(1): 65-96.
- 931 Lacey, L.A., Frutos, R., Kaya, H.K., and Vail, P. 2001. Insect pathogens as biological control
932 agents: do they have a future? Biol. Control **21**(3): 230-248.
- 933 Lavis, D.S., Hallett, A., Koon, E.M., and McAuley, T.C. 2003a. History of and advances in barriers
934 as an alternative method to suppress sea lampreys in the Great Lakes. J. Gt. Lakes Res.
935 **29**: 362-372.
- 936 Lavis, D.S., Henson, M.P., Johnson, D.A., Koon, E.M., and Ollila, D.J. 2003b. Case history of sea
937 lamprey control in Lake Michigan: 1979 to 1999. J. Gt. Lakes Res. **29**: 584-598.
- 938 Lawrie, A.H. 1970. The sea lamprey in the Great Lakes. Trans. Am. Fish. Soc. **99**: 766-775.
- 939 Lech, J.J., and Statham, C.N. 1975. Role of glucuronide formation in the selective toxicity of 3-

- 940 trifluoromethyl-4-nitrophenol (TFM) for the sea lamprey: comparative aspects of TFM
941 uptake and conjugation in sea lamprey and rainbow trout. *Toxicol. Appl. Pharm.* **31**(1):
942 150-158.
- 943 Leroux, P., and Walker, A.S. 2013. Activity of fungicides and modulators of membrane drug
944 transporters in field strains of *Botrytis cinerea* displaying multidrug resistance. *Eur. J.*
945 *Plant Pathol.* **135**(4): 683-693.
- 946 Lescak, E.A., Bassham, S.L., Catchen, J., Gelmond, O., Sherbick, M.L., von Hippel, F.A., and
947 Cresko, W.A. 2015. Evolution of stickleback in 50 years on earthquake-uplifted islands.
948 *Proc. Natl. Acad. Sci. U. S. A.* **112**(52): E7204-E7212.
- 949 Levinton, J.S., Suatoni, E., Wallace, W., Junkins, R., Kelaher, B., and Allen, B.J. 2003. Rapid loss of
950 genetically based resistance to metals after the cleanup of a Superfund site. *Proc. Natl.*
951 *Acad. Sci. U. S. A.* **100**(17): 9889-9891.
- 952 Li, W.M., Scott, A.P., Siefkes, M.J., Yan, H.G., Liu, Q., Yun, S.S., and Gage, D.A. 2002. Bile acid
953 secreted by male sea lamprey that acts as a sex pheromone. *Science* **296**(5565): 138-
954 141.
- 955 Mackie, J.A., Levinton, J.S., Przeslawski, R., DeLambert, D., and Wallace, W. 2010. Loss of
956 evolutionary resistance by the oligochaete *Limnodrilus hoffmeisteri* to a toxic
957 substance- cost or gene flow? *Evolution* **64**(1): 152-165.
- 958 Manion, P.J., and McLain, A.L. 1971. Biology of larval sea lampreys (*Petromyzon marinus*) of the
959 1960 year class, isolated in the Big Garlic River, Michigan, 1960-65. Great Lakes Fishery
960 Commission Technical Report 15, Ann Arbor, M.I.
- 961 Manion, P.J., and Smith, B.R. 1978. Biology of larval and metamorphosing sea lampreys,

- 962 *Petromyzon marinus*, of the 1960 year class in the Big Garlic River, Michigan, Part II,
963 1966-72. Great Lakes Fishery Commission Technical Report 30, Ann Arbor, M.I.
- 964 Manzon, R.G., Youson, J.H., and Holmes, J.A. 2015. Lamprey metamorphosis. *In* Lampreys:
965 Biology, Conservation and Control. *Edited by* M.F. Docker. Springer, Dordrecht
966 Heidelberg New York London. pp. 139-214.
- 967 Mariath, H.A., Orton, C.J., and Shivas, C.J. 1990. Resistance to oviposition suppressants in *Lucilia*
968 *cuprina*. *In* Resistance management in parasites of sheep. *Edited by* J.A. McKenzie, P.J.
969 Martin and J.H. Arunde. Australian Wool Corp, Melbourne. p. 52.
- 970 McCauley, D.W., Docker, M.F., Whyard, S., and Li, W.M. 2015. Lampreys as diverse model
971 organisms in the genomics era. *BioScience* **65**(11): 1046-1056.
- 972 McLain, A.L., Smith, B.R., and Moore, H.H. 1965. Experimental control of sea lampreys with
973 electricity on the south shore of Lake Superior. Great Lakes Fishery Commission
974 Technical Report 10, Ann Arbor, M.I.
- 975 McLaughlin, R.L., Smyth, E.R.B., Castro-Santos, T., Jones, M.L., Koops, M.A., Pratt, T.C., and
976 Velez-Espino, L.A. 2013. Unintended consequences and trade-offs of fish passage. *Fish.*
977 *Fish.* **14**(4): 580-604.
- 978 McLean, A.R., Barber, J., Bravener, G., Rous, A.M., and McLaughlin, R.L. 2015. Understanding
979 low success trapping invasive sea lampreys: an entry-level analysis. *Can. J. Fish. Aquat.*
980 *Sci.* **72**(12): 1876-1885.
- 981 Meckley, T.D., Wagner, C.M., and Gurarie, E. 2014. Coastal movements of migrating sea

- 982 lamprey (*Petromyzon marinus*) in response to a partial pheromone added to river water:
983 implications for management of invasive populations. Can. J. Fish. Aquat. Sci. **71**(4): 533-
984 544.
- 985 Monteiro, J.P., Martins, A.F., Lucio, M., Reis, S., Geraldés, C., Oliveira, P.J., and Jurado, A.S.
986 2011. Interaction of carbonylcyanide p-trifluoromethoxyphenylhydrazone (FCCP) with
987 lipid membrane systems: a biophysical approach with relevance to mitochondrial
988 uncoupling. J. Bioenerg. Biomembr. **43**(3): 287-298.
- 989 Morales, J.A., Cardoso, D.G., Della Lucia, T.M.C., and Guedes, R.N.C. 2013. Weevil x insecticide:
990 Does 'personality' matter? PLoS One **8**(6).
- 991 Morse, T.J., Ebener, M.P., Koon, E.M., Morkert, S.B., Johnson, D.A., Cuddy, D.W., Weisser, J.W.,
992 Mullett, K.M., and Genovese, J.H. 2003. A case history of sea lamprey control in Lake
993 Huron: 1979 to 1999. J. Great Lakes Res. **29**: 599-614.
- 994 Mota-Sanchez, D., Bills, S.P., and Whalon, M.E. 2002. Arthropod resistance to pesticides: status
995 and overview. In Pesticides in agriculture and the environment. Edited by W. Wheeler
996 and B. Gainesville. CRC Press, New York, N.Y. pp. 241-272.
- 997 Nacci, D., Coiro, L., Champlin, D., Jayaraman, S., McKinney, R., Gleason, T.R., Munns, W.R.,
998 Specker, J.L., and Cooper, K.R. 1999. Adaptations of wild populations of the estuarine
999 fish *Fundulus heteroclitus* to persistent environmental contaminants. Mar. Biol. **134**(1):
1000 9-17.
- 1001 Nacci, D.E., Champlin, D., and Jayaraman, S. 2010. Adaptation of the estuarine fish *Fundulus*
1002 *heteroclitus* (Atlantic Killifish) to polychlorinated biphenyls (PCBs). Estuar Coast **33**(4):
1003 853-864.

- 1004 Nettles, D.C., Staats, N.R., Chipman, B.D., and Nashett, L.J. 2001. Final supplemental
1005 environmental impact statement. A long-term program of sea lamprey control in Lake
1006 Champlain. U.S. Fish & Wildlife, Vermont Department of Fish and Wildlife and New York
1007 State Department of Environmental Conservation.
- 1008 Niblett, P.D., and Ballantyne, J.S. 1976. Uncoupling of oxidative phosphorylation in rat liver
1009 mitochondria by the lamprey larvicide TFM (3-trifluoromethyl-4-nitrophenol). *Pest.*
1010 *Biochem. Physiol.* **6**: 363-366.
- 1011 Oliver, A.J., Wheeler, S.H., and Gooding, C.D. 1982. Field evaluation of 1980 and pindone oat
1012 bait, and the possible decline in effectiveness of poison baiting for the control of the
1013 rabbit, *Oryctolagus cuniculus*. *Aust. Wildlife Res.* **9**(1): 125-134.
- 1014 Palkovacs, E.P., Kinnison, M.T., Correa, C., Dalton, C.M., and Hendry, A.P. 2012. Fates beyond
1015 traits: ecological consequences of human-induced trait change. *Evol. Appl.* **5**(2): 183-
1016 191.
- 1017 Palumbi, S.R. 2001. Evolution - Humans as the world's greatest evolutionary force. *Science*
1018 **293**(5536): 1786-1790.
- 1019 Pearse, D.E., Hayes, S.A., Bond, M.H., Hanson, C.V., Anderson, E.C., Macfarlane, R.B., and Garza,
1020 J.C. 2009. Over the falls? Rapid evolution of ecotypic differentiation in
1021 steelhead/rainbow trout (*Oncorhynchus mykiss*). *J. Hered.* **100**(5): 515-525.
- 1022 Pelz, H.J., Rost, S., Hunerberg, M., Fregin, A., Heiberg, A.C., Baert, K., MacNicoll, A.D., Prescott,
1023 C.V., Walker, A.S., Oldenburg, J., and Muller, C.R. 2005. The genetic basis of resistance to
1024 anticoagulants in rodents. *Genetics* **170**(4): 1839-1847.
- 1025 Phillis, C.C., Moore, J.W., Buoro, M., Hayes, S.A., Garza, J.C., and Pearse, D.E. 2016. Shifting

- 1026 thresholds: rapid evolution of migratory life histories in steelhead/rainbow trout,
1027 *Oncorhynchus mykiss*. J. Hered. **107**(1): 51-60.
- 1028 Posthuma, L., and Vanstraalen, N.M. 1993. Heavy-metal adaptation in terrestrial invertebrates -
1029 a review of occurrence, genetics, physiology and ecological consequences. Comp.
1030 Biochem. Physiol. C-Pharmacol. Toxicol. Endocrinol. **106**(1): 11-38.
- 1031 Potter, I.C. 1980. Ecology of larval and metamorphosing lampreys. Can. J. Fish. Aquat. Sci. **37**:
1032 1595-1615.
- 1033 Potts, D.D., Dawson, H.A., and Jones, M.L. 2015. Validation of a relationship between statolith
1034 size and age of larval Great Lakes sea lamprey (*Petromyzon marinus*). Environ. Biol. Fish.
1035 **98**(8): 1859-1869.
- 1036 Ranson, H., N'Guessan, R., Lines, J., Moiroux, N., Nkuni, Z., and Corbel, V. 2011. Pyrethroid
1037 resistance in African anopheline mosquitoes: what are the implications for malaria
1038 control? Trends Parasitol. **27**(2): 91-98.
- 1039 Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R., Karchner, S.I.,
1040 Hahn, M.E., Nacci, D., Oleksiak, M.F., Crawford, D.L., and Whitehead, A. 2016. The
1041 genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild
1042 fish. Science **354**(6317): 1305-1308.
- 1043 Relyea, R., and Hoverman, J. 2006. Assessing the ecology in ecotoxicology: a review and
1044 synthesis in freshwater systems. Ecol. Lett. **9**(10): 1157-1171.
- 1045 Renaud, C.B., Gill, H.S., and Potter, I.C. 2009. Relationships between the diets and
1046 characteristics of the dentition, buccal glands and velar tentacles of the adults of the
1047 parasitic species of lamprey. J. Zool. **278**(3): 231-242.

- 1048 Renton, M., Busi, R., Neve, P., Thornby, D., and Vila-Aiub, M. 2014. Herbicide resistance
1049 modelling: past, present and future. *Pest Manag Sci* **70**(9): 1394-1404.
- 1050 Reznick, D.A., Bryga, H., and Endler, J.A. 1990. Experimentally induced life-history evolution in a
1051 natural population. *Nature* **346**: 357-359.
- 1052 Reznick, D.N., and Bryga, H. 1987. Life-history evolution in guppies (*Poecilia reticulata*): 1.
1053 Phenotypic and genetic changes in an introduction experiment. *Evolution* **41**(6): 1370-
1054 1385.
- 1055 Reznick, D.N., and Ghalambor, C.K. 2005. Can commercial fishing cause evolution? Answers
1056 from guppies. *Can. J. Fish. Aquat. Sci.* **62**: 791-801.
- 1057 Roush, R.T., and Daly, J.C. 1990. The role of population genetics in resistance research and
1058 management. *In* *Pesticide Resistance in Arthropods. Edited by* R.T. Roush and B.E.
1059 Tabashnik. Chapman & Hall, New York, N.Y.. pp. 97-152.
- 1060 Sawicki, R. 1987. Definition, detection and documentation of insecticide resistance. *In*
1061 *Combating Resistance to Xenobiotics: Biological and Chemical Approaches. Edited by*
1062 M.G. Ford, D.W. Holloman, B.P.S. Khambay and R.M. Sawicki. Ellis Horwood, London,
1063 The United Kingdom. pp. 105-117.
- 1064 Sawyer, A.J. 1980. Prospects for integrated pest management of the sea lamprey (*Petromyzon*
1065 *marinus*). *Can. J. Fish. Aquat. Sci.* **37**: 2081-2092.
- 1066 Scholefield, R.J., and Seelye, J.G. 1990. Resistance to 3-trifluoromethyl-4-nitrophenol (TFM) in
1067 sea lamprey. Great Lakes Fishery Commission Technical Report 56, Ann Arbor, M.I.
- 1068 Scholefield, R.J., Slaght, K.S., and Stephens, B.E. 2008. Seasonal variation in sensitivity of larval

- 1069 sea lampreys to the lampricide 3-trifluoromethyl-4-nitrophenol. N. Am. J. Fish. Manage.
1070 **28**(5): 1609-1617.
- 1071 Sharpe, D.M.T., and Hendry, A.P. 2009. Life history change in commercially exploited fish
1072 stocks: an analysis of trends across studies. *Evol. Appl.* **2**(3): 260-275.
- 1073 Sherburne, S., and Reinhardt, U.G. 2016. First test of a species-selective adult sea lamprey
1074 migration barrier. *J. Gt. Lakes Res.* **42**(4): 893-898.
- 1075 Siefkes, M.J., Steeves, T.B., Sullivan, W.P., Twohey, M.B., and Li, W. 2013. Sea lamprey control:
1076 past, present, and future. *In* Great Lakes Fisheries Policy & Management. *Edited by*
1077 W.W. Taylor, A.J. Lynch and N.J. Leonard. Michigan State University Press, East Lansing
1078 M.I. pp. 651-704.
- 1079 Siepielski, A.M., DiBattista, J.D., and Carlson, S.M. 2009. It's about time: the temporal dynamics
1080 of phenotypic selection in the wild. *Ecol. Lett.* **12**(11): 1261-1276.
- 1081 Smith, B.R., and Tibbles, J.J. 1980. Sea lamprey (*Petromyzon marinus*) in Lakes Huron, Michigan,
1082 and Superior: history of invasion and control, 1936-78. *Can. J. Fish. Aquat. Sci.* **37**: 1780-
1083 1808.
- 1084 Smith, J.J., Kuraku, S., Holt, C., Sauka-Spengler, T., Jiang, N., Campbell, M.S., Yandell, M.D.,
1085 Manousaki, T., Meyer, A., Bloom, O.E., Morgan, J.R., Buxbaum, J.D., Sachidanandam, R.,
1086 Sims, C., Garruss, A.S., Cook, M., Krumlauf, R., Wiedemann, L.M., Sower, S.A., Decatur,
1087 W.A., Hall, J.A., Amemiya, C.T., Saha, N.R., Buckley, K.M., Rast, J.P., Das, S., Hirano, M.,
1088 McCurley, N., Guo, P., Rohner, N., Tabin, C.J., Piccinelli, P., Elgar, G., Ruffier, M., Aken,
1089 B.L., Searle, S.M.J., Muffato, M., Pignatelli, M., Herrero, J., Jones, M., Brown, C.T.,
1090 Chung-Davidson, Y.W., Nanlohy, K.G., Libants, S.V., Yeh, C.Y., McCauley, D.W.,

- 1091 Langeland, J.A., Pancer, Z., Frittsch, B., de Jong, P.J., Zhu, B.L., Fulton, L.L., Theising, B.,
1092 Flicek, P., Bronner, M.E., Warren, W.C., Clifton, S.W., Wilson, R.K., and Li, W.M. 2013.
1093 Sequencing of the sea lamprey (*Petromyzon marinus*) genome provides insights into
1094 vertebrate evolution. *Nature Genet.* **45**(4): 415-421.
- 1095 Smith, S.J., and Marsden, J.E. 2007. Predictive morphometric relationships for estimating
1096 fecundity of sea lampreys from lake Champlain and other landlocked populations. *Trans.*
1097 *Am. Fish. Soc.* **136**(4): 979-987.
- 1098 Sparks, T.C., Lockwood, J.A., Byford, R.L., Graves, J.B., and Leonard, B.R. 1989. The role of
1099 behavior in insecticide resistance. *Pestic. Sci.* **26**(4): 383-399.
- 1100 Stearns, S.C. 1983. The genetic basis of differences in life history traits among 6 populations of
1101 mosquitofish (*Gambusia affinis*) that shared ancestors in 1905. *Evolution* **37**(3): 618-627.
- 1102 Stockwell, C.A., and Weeks, S.C. 1999. Translocations and rapid evolutionary responses in
1103 recently established populations of western mosquitofish (*Gambusia affinis*). *Anim.*
1104 *Conserv.* **2**(2): 103-110.
- 1105 Sullivan, W.P., Christie, G.C., Cornelius, F.C., Fodale, M.F., Johnson, D.A., Koonces, J.F., Larson,
1106 G.L., McDonald, R.B., Mullett, K.M., Murray, C.K., and Ryan, P.A. 2003. The sea lamprey
1107 in Lake Erie: a case history. *J. Gt. Lakes Res.* **29**: 615-636.
- 1108 Swink, W.D., and Johnson, N.S. 2014. Growth and survival of sea lampreys from metamorphosis
1109 to spawning in Lake Huron. *Trans. Am. Fish. Soc.* **143**(2): 380-386.
- 1110 Tabashnik, B.E., Mota-Sanchez, D., Whalon, M.E., Hollingworth, R.M., and Carriere, Y. 2014.
1111 Defining terms for proactive management of resistance to Bt crops and pesticides. *J.*
1112 *Econ. Entomol.* **107**(2): 496-507.

- 1113 Tao, H.L., Zhang, Y., Zeng, X.G., Shulman, G.I., and Jin, S.K. 2014. Niclosamide ethanolamine-
1114 induced mild mitochondrial uncoupling improves diabetic symptoms in mice. *Nature*
1115 *Medicine* **20**(11): 1263-1269.
- 1116 Terada, H. 1990. Uncouplers of oxidative phosphorylation. *Environ. Health Perspect.*
1117 **87**: 213-218.
- 1118 Twigg, L.E., Martin, G.R., and Lowe, T.J. 2002. Evidence of pesticide resistance in medium-sized
1119 mammalian pests: a case study with 1080 poison and Australian rabbits. *J. Appl. Ecol.*
1120 **39**(4): 549-560.
- 1121 Van Leeuwen, T., Van Pottelberge, S., and Tirry, L. 2006. Biochemical analysis of a chlorfenapyr-
1122 selected resistant strain of *Tetranychus urticae* Koch. *Pest Manag. Sci.* **62**(5): 425-433.
- 1123 van Wijk, S.J., Taylor, M.I., Creer, S., Dreyer, C., Rodrigues, F.M., Ramnarine, I.W., van
1124 Oosterhout, C., and Carvalho, G.R. 2013. Experimental harvesting of fish populations
1125 drives genetically based shifts in body size and maturation. *Front. Ecol. Environ.* **11**(4):
1126 181-187.
- 1127 Waldman, J., Grunwald, C., and Wirgin, I. 2008. Sea lamprey *Petromyzon marinus*: an exception
1128 to the rule of homing in anadromous fishes. *Biol. Lett.* **4**(6): 659-662.
- 1129 Walsh, M.R., Broyles, W., Beston, S.M., and Munch, S.B. 2016. Predator-driven brain size
1130 evolution in natural populations of Trinidadian killifish (*Rivulus hartii*). *Proc. R. Soc. B-*
1131 *Biol. Sci.* **283**(1834): 9.
- 1132 Walsh, M.R., and Reznick, D.N. 2008. Interactions between the direct and indirect effects of
1133 predators determine life history evolution in a killifish. *Proc. Natl. Acad. Sci. U. S. A.*
1134 **105**(2): 594-599.

- 1135 Walsh, M.R., and Reznick, D.N. 2011. Experimentally induced life-history evolution in a killifish
1136 in response to the introduction of guppies. *Evolution* **65**(4): 1021-1036.
- 1137 Weis, J.S., Smith, G., Zhou, T., Santiago-Bass, C., and Weis, P. 2001. Effects of contaminants on
1138 behavior: biochemical mechanisms and ecological consequences. *BioScience* **51**(3): 209-
1139 217.
- 1140 Westley, P.A.H., Ward, E.J., and Fleming, I.A. 2012. Fine-scale local adaptation in an invasive
1141 freshwater fish has evolved in contemporary time. *Proc. R. Soc. B-Biol. Sci.* **280**(1751): 9.
- 1142 Wilkie, M.P., Bergstedt, R., and McDonald, D.G. 2014. Establishing physiological indices for
1143 more effective use of TFM to control sea lamprey populations in the Great Lakes. 2014
1144 Project Completion Report, Great Lakes Fishery Commission, Ann Arbor, M.I.
- 1145 Williamson, M.S., MartinezTorres, D., Hick, C.A., and Devonshire, A.L. 1996. Identification of
1146 mutations in the housefly para-type sodium channel gene associated with knockdown
1147 resistance (kdr) to pyrethroid insecticides. *MGG* **252**(1-2): 51-60.
- 1148 Wirgin, I., Roy, N.K., Loftus, M., Chambers, R.C., Franks, D.G., and Hahn, M.E. 2011. Mechanistic
1149 basis of resistance to PCBs in Atlantic tomcod from the Hudson River. *Science*
1150 **331**(6022): 1322-1325.
- 1151 Wirgin, I., and Waldman, J.R. 2004. Resistance to contaminants in North American fish
1152 populations. *Mutat. Res.-Fundam. Mol. Mech. Mutagen.* **552**(1-2): 73-100.
- 1153 Wright, G.D. 2007. The antibiotic resistome: the nexus of chemical and genetic diversity. *Nat*
1154 *Rev Microbiol* **5**(3): 175-186.

1155 **Table 1.** The number of generations of sea lamprey exposed to the lampricide, 3-
 1156 trifluoromethyl-4-nitrophenol (TFM) in the Great Lakes given the timeline of lampricide
 1157 application. Sea lamprey generation time was assumed to be 5 years.

Lake	Year of first application of TFM	Number of generations of sea lamprey exposed to TFM
Superior	1958 ¹	11.6
Huron	1960 ²	11.2
Michigan	1960 ³	11.2
Ontario	1971 ⁴	9.0
Erie	1986 ⁵	6.0

1158 Source: ¹Heinrich et al. 2003; ²Morse et al. 2003; ³Lavis et al. 2003b; ⁴Larson et al. 2003;

1159 ⁵Sullivan et al. 2003.

Draft

Table 2. Published examples of studies in which evolution was documented for fish populations exposed to selection. Examples were restricted to those with strong evidence for evolution as opposed to the sole result of phenotypic plasticity; this included studies where genetic assessment or common garden experiments were conducted to provide evidence of evolution. Note that the number of generations that had passed by the time the study or analysis was done could be greater than the number of generations in which an evolutionary response could have been detected. The list of examples is not exhaustive.

Citation	Species	Trait	Type of study	Agent of selection	Number of generations ¹
van Wijk et al. 2013	Trinidadian guppy (<i>Poecilia reticulata</i>)	Male body size	Artificial selection in laboratory with genetic analysis	Harvest	3
Gordon et al. 2015	Trinidadian guppy (<i>P. reticulata</i>)	Male color (secondary sex characteristic)	Introduction experiment, including monitoring, genetic analysis, and common garden analysis	Introduction to novel environments differing in predation	3
Conover and Munch 2002	Atlantic silverside <i>Menidia menidia</i>	Body size	Artificial selection in laboratory	Harvest	4
Breckels and Neff 2014	Trinidadian guppy (<i>P. reticulata</i>)	Sperm length	Artificial selection in laboratory	Temperature	4
Reznick et al. 1990, Reznick and Bryga 1987	Trinidadian guppy (<i>P. reticulata</i>)	Maturity	Translocation experiment of wild populations	Introduction to novel environments differing in predation	7
Hendry et al. 1998	Sockeye salmon (<i>Onchorhynchus</i>	Developmental rate (time to hatch; time	Common garden experiments of wild populations	Introduction to novel	14

	<i>nerka</i>)	to emerge)		environment	
Aykanat et al. 2011	Rainbow trout (<i>O. mykiss</i>)	Osmoregulatory genes	Common garden experiments and genetic analysis of wild populations	Introduction to novel environment (freshwater)	14
Du et al. 2016, Nacci et al. 1999, 2010, Reid et al. 2016	Killifish (<i>Fundulus heteroclitus</i>)	Hepatocyte oxidative phosphorylation metabolism (PCB resistance)	Genetic and common garden experiments of wild populations	PCBs	15
Kavanagh et al. 2010	Arctic grayling (<i>Thymallus thymallus</i>)	Developmental traits (growth rate; mass development rate; conversion efficiency)	Common garden experiment and genetic analysis of wild populations	Introduction to novel environment (cold vs warm)	22
Pearse et al. 2009, Phillis et al. 2016	Rainbow trout (<i>O. mykiss</i>)	Size-at-age and threshold size for migration tactic (evolution of residency)	Common garden and genetic analysis of wild populations	Translocation above a migration barrier	25
Kinnison et al. 1998a, 1998b	Chinook salmon (<i>O. tshawytscha</i>)	Time to hatch; growth rate	Common garden experiment of wild populations	Introduction to novel environment	26
Westley et al. 2012	Brown trout (<i>Salmo trutta</i>)	Juvenile survival	Common garden and in situ reciprocal transplant	Introduction to novel environment	37
Hubert et al. 2016	Common carp (<i>Cyprinus carpio</i>)	Scale cover	Breeding experiment and genetic analysis of wild and hatchery fish	Introduction to novel environment	40
Wirgin et al. 2011	Atlantic tomcod (<i>Microgadus tomcod</i>)	PCB resistance	Common garden and genetic analysis of wild populations	PCBs	50
Lescak et al. 2015	Threespine	Morphology (mainly	Genetic analysis of wild	Habitat shift	50

	stickleback (<i>Gasterosteus aculeatus</i>)	lateral plate number)	populations		
Stockwell and Weeks 1999	Mosquitofish (<i>Gambusia affinis</i>)	Size at maturity; fat content	Common garden experiment of wild populations	Introduction to novel environment	110
Walsh et al. 2016, Walsh and Reznick 2008, 2011	Trinidadian killifish (<i>Rivulus hartii</i>)	Brain size; size and age at maturity; reproductive investment;	Common garden experiment of natural populations	Predation	114
Stearns 1983	Mosquitofish (<i>G. affinis</i>)	Age and length at maturity; offspring size (weight)	Common garden experiment of wild populations	Introduction to novel environment	140

¹Maximum value in cases where a range of generation times were given.

Table 3. Estimated reductions in parasitic sea lamprey recruitment attributable to lampricide control. Mortality rates were based on a model (Jones et al. 2009) that simulates sea lamprey population dynamics and management at the level of an entire Great Lake and calibrates lake-wide abundance to correspond with recent adult assessments and recent lampricide control expenditures. Model incorporates uncertainty in stream-level lampricide effectiveness and stream-level assessment, and recruitment stochasticity. Results are for 1000 simulations.

Lake	Number of adults	Budget	Median number of parasites with no control	Median number of parasites with control	Proportion killed as a result of control
Superior	760,000	\$1,010,000	2,091,954	370,767	82%
Michigan	1,850,000	\$2,030,000	3,084,211	475,999	85%
Huron	1,910,000	\$2,280,000	5,394,502	521,587	90%
Erie	154,000	\$270,000	199,007	24,796	88%
Ontario	311,000	\$680,000	823,147	78,474	90%

Table 4. Summary of Great Lakes sea lamprey life history and control program attributes relevant to the development of resistance to lampricides.

Life history or control program attribute	What it might mean for resistance
Generation time of about 5 years	Limits the number of generations of sea lamprey that have been exposed to lampricides. Slows the rate of resistance development relative to most other taxa where resistance has been observed. However, does not prevent the possibility of rapid evolution as numerous studies of fish populations have demonstrated.
Absence of natal homing	Sea lamprey born from different tributaries mix during reproduction, potentially diluting selection because mating occurs between larvae exposed to TFM and those not exposed. Similarly, random mating between resistant and non-resistant genotypes might slow the evolution of resistance. Little to no local adaptation would also reduce the chances of a population of resistant (or susceptible) individuals immigrating and spreading resistant (or susceptible) genotypes. Alternatively, this mixing might mean resistance will spread rapidly if it does arise.
Invasion history	The source population(s) of sea lamprey may have possessed limited genetic variability, potentially limiting the number of resistant genotypes present in the founding population. However, some research has suggested that rapid rates of adaptation are still possible in invading populations even though genetic diversity might be low (Dlugosch and Parker 2008).
Mortality rate from lampricides	Lampricide-induced mortality rate estimates are high (>80%), increasing the selective pressure and thus increasing the chances of resistance evolution.
High fecundity	Based on studies of pest resistance in arthropods, high fecundity increases the chances of resistance evolution because the survival and mating of just a few resistant individuals can potentially produce large numbers of resistant offspring. Furthermore, high fecundity increases the chances of novel mutations conferring resistance.

Residual sea lamprey

The many ways in which sea lamprey could survive or avoid treatment will weaken selection and reduce the likelihood of physiological resistance – e.g., lamprey could potentially survive treatment not only by having a resistant genotype, but also because not all streams or stream reaches are treated in a given year (i.e., subject to variable selection). Other examples by which residual sea lamprey might survive include residing in backwaters and eddies where lethal concentrations of TFM are more difficult to achieve or where treatment isn't applied because densities are relatively low. However, the survival of these residual sea lamprey could increase the chances of behavioural resistance evolution and raise the overall probability of a mutation arising by increasing the pathways for resistance evolution.

Reliance on 3-trifluoromethyl-4-nitrophenol (TFM) as the primary method of sea lamprey control

Although integrated pest management is a goal of the sea lamprey control program, chemical treatment of streams with TFM is the main method of control used. This increases the chances of resistance development.

Cyclical nature of treatment

Could increase the chances of certain types of behavioural resistance, for example earlier transformation or outmigration.

Figure Captions

Figure 1. Examples of potential routes of lampricide resistance evolution in sea lamprey. In susceptible lamprey, lampricide is taken from the external environment into the body (uptake) and transported to cell mitochondria where it uncouples mitochondrial oxidative phosphorylation and inhibits ATP production, resulting in death to the individual. Routes of resistance evolution are indicated in red italics, and include *avoidance* of uptake (via behavioural or physiological strategies) or *detoxification*, *excretion*, and *target modification* (via physiological strategies). Example evolutionary changes that could bring about resistance are given in black text. Potential routes and examples given here are not meant to be exhaustive. Modified in part from Despres et al. (2007).

Figure 2. Results of historical toxicity tests showing concentrations of 3-trifluoromethyl-4-nitrophenol (TFM) that killed 50% (LC50) of larval sea lamprey. Results are for static toxicity tests run for 19-24 hours in Lake Huron water conducted at the Hammond Bay Biological Station. Data were from archival records, historical reports (Applegate et al. 1961, Scholefield and Seelye 1990), and quality assurance tests (QA/QC) of TFM supplies (Adair and Sullivan 2014b). Linear regression fit to all data (solid line) and to all but the archival data (dashed line). A small random offset was added to sample from the same year to distinguish overlapping points. The estimated slope for LC50 from all data was -0.009 (95% confidence interval: -0.011 to -0.007). Without the archival data, the LC50 slope was -0.002, (95% confidence interval: -0.004 to 0.001). Six outliers (LC50>3) from the archival records were removed. As these high values occurred early in the time series, their removal does not impact the sign of the slope.

Figure 1.

LC50 TFM (mg/L)

Year

○ Archive
△ Applegate et al. 1961
+ Scholefield and Seelye 1990
× QA/QC

Appendix 1: Sea Lamprey Mortality from Lampricide Treatment

We used two approaches to estimate the mortality rate induced by lampricide treatments. In the absence of direct estimates of selection differentials, we use these estimates as an indication of the potential strength of selection from lampricides. Note that these estimates are different from measurements of treatment effectiveness on a tributary. Lampricide actions are taken at the scale of individual streams, but the sea lamprey they target are from a lake-wide population that does not demonstrate natal homing. Measuring the strength of selection from lampricides requires integrating estimates of effectiveness across all potential lotic and lentic sources of juveniles to a lake.

The first approach involves an empirical approximation of lampricide mortality from data that control agents use to decide where to apply treatments. Potential sea lamprey source streams are surveyed regularly for the presence of larvae potentially large enough to metamorphose into parasitic juveniles within a year. These survey data are used to rank streams for lampricide treatment each year, based on the estimated abundance of large larvae relative to the costs of treating that stream. These data are maintained in a database that can also be used to quantify the expected effectiveness of a multi-year lampricide treatment strategy, based on assumptions about the effectiveness of individual stream treatments. Focusing solely on Lake Superior, we used median estimates of larval abundance by stream for 1996 to 2012, and then used the treatment schedule in 2012-2014 and that proposed for 2015, to apply TFM mortality to the individual streams. Several assumptions were made to derive estimates of mortality from the control data: (i) larval sea lamprey remain in Lake Superior streams for at least four years, (ii) 98% of the 2012 year class of larvae present in Lake Superior streams known to contain larval sea lamprey would have been exposed to lampricide treatments, and (iii) average TFM treatment effectiveness for a stream was 90%. From this, we estimated the mortality rate for the 2012 year class as 89%.

There are several uncertainties associated with this first approach for estimating mortality from lampricides. The assumptions of 90% treatment effectiveness and 98% exposure to lampricides are simplifications that likely are not always reflective of true control outcomes, thus contributing error to the mortality estimates. Another assumption contributing error is that all source populations of sea lamprey have been assessed.

Given the uncertainties of the first approach, we also estimated mortality rates using a second approach based on an existing operating model (Jones et al. 2009). The model simulates sea lamprey population dynamics and management at the level of an entire Great Lake and calibrates lake-wide abundance to correspond with recent adult assessments and recent lampricide control expenditures. The model also explicitly incorporates uncertainty in stream-level lampricide effectiveness and in stream-level assessment, as well as stochasticity in recruitment dynamics. Thus, the model addresses several of the uncertainties associated with the first approach for estimating mortality.

Simulations using the sea lamprey operating model were used to quantify mortality experienced by a lake-wide cohort of sea lamprey due to lampricide treatment (these are the mortality rates presented in the main text of the paper). First, we calibrated the model for each Great Lake by adjusting larval natural mortality rates such that recent observed adult sea lamprey abundance levels were forecasted, on average, when recent levels of lampricide control were applied (*c.f.* Irwin et al. 2012). Second, we ran simulations for each lake with no lampricide control applied, to estimate population abundances associated with no control. Third, we fixed abundance of adults at these “no control” abundances and estimated abundance of parasitic juvenile sea lamprey that would be produced, on average, in the absence of control (averaged across stochastic variation in recruitment dynamics). Fourth, we repeated these simulations with fixed adult abundance but with recent levels of lampricide control and again estimated average abundance of parasitic juvenile sea lamprey produced (in this case averaged across uncertainty in recruitment, assessment, and lampricide effectiveness). Finally, we calculated the ratio of forecasted juvenile abundance with control to that with no control and subtracted this ratio from one to obtain our estimate of lake-wide lampricide-induced mortality. The resulting lake-wide estimate of mortality ranged from 82-90% (Table 3 main text; Fig. A1).

Draft

Figure A1. Estimated reductions in juvenile parasitic sea lamprey recruitment to each of the Great Lakes, attributable to the effect of lampricide control, as indicated by the results of 1000 simulations of a sea lamprey operating model (Jones et al. 2009). Total heights of the bars are forecasted mean numbers of juveniles with no control; heights of the black segments are forecasted mean numbers of juveniles controlled at recent budget levels for each lake; the grey area represents the magnitude of reduction in recruitment due to control.

Appendix 2: Historical Toxicity Test Analysis

We investigated the results of historical toxicity tests for evidence of physiological resistance to TFM in sea lamprey. The toxicity tests used in our analysis were conducted in the field in conjunction with stream treatments and in laboratories as part of ongoing research or chemical quality assurance tests. Information on the testing conditions (e.g., the pH, alkalinity, and temperature of the water) was not regularly recorded. To control for some of the potential variation in testing conditions known to influence toxicity tests (Bills et al. 2003), we focused our analysis on tests conducted at the US Geological Survey Hammond Bay Biological Station in Millersburg, Michigan. This laboratory uses Lake Huron water for its toxicity tests, which has relatively consistent pH and alkalinity (Karen Slaght Pers. Comm). Toxicity tests at this facility are routinely run at 12 degrees C and occur between April and August. Seasonal differences in the lethal concentrations are not generally observed in the lab when conducting the toxicity tests because larvae are typically collected in the spring and held through the testing period (Karen Slaght Pers. Comm). However, it is possible that these conditions were not always met in the earlier tests. Analysis was limited to static tests (between 19-24 hours) because flow-through tests were not common prior to 1980.

We used historic toxicity test results from four sources (Table A1): (1) TFM concentrations and mortality counts from archival records; (2) estimated lethal concentrations from Applegate et al. (1961); (3) estimated lethal concentrations from Scholefield and Seelye (1990); and (4) quality assurance tests of TFM supplies (Adair and Sullivan 2014) (Table 3). Applegate et al. (Applegate et al. 1961) did not report the year their toxicity tests were conducted and we assumed all of their tests were conducted in 1961. The Litchfield Wilcoxon (1949) approach was used to estimate the TFM concentrations at which 50% and 99.9% of the sample individuals died (LC50 and LC99.9, respectively).

Our analysis of historical toxicity data provided no evidence of resistance. The LC50 of TFM did not increase over the 57 year time frame, nor did the relative ratio of LC99.9/LC50 increase (Figure A2). An increase in LC50 over time would have been the most obvious evidence that resistance has developed in sea lamprey exposed to TFM. If only a subset of the sea lamprey population were resistant to TFM, an increase in the relative ratio of LC50 to LC99.9 might be evident before an increase in the LC50 was observed.

There are several potential uncertainties with these toxicity data that limit the applicability of our findings as a test of physiological resistance to TFM. Our analyses relied on historical data that were collected for other purposes, not with the intent of specifically investigating the evolution of resistance. Sample sizes were low for some years and key variables known to influence toxicity (e.g., pH, source stream of the larvae, temperature, timing of larval collection) were not always measured or controlled. Instruments used over time have also changed; for example, concentrations reported by Applegate et al. (1961) were measured using a Klett colorimeter, providing less precision than current methods (Adair and Sullivan 2014). Furthermore, the higher LC50s estimated from the archival data raise concerns (Fig. A2). While this pattern could indicate bias in the historical methods used to record TFM concentration, the

cause is unknown. Thus, although we didn't detect any overt signs that physiological resistance has developed, these results must be interpreted cautiously and further tests are needed.

Draft

Table A1. Historical toxicity tests used to examine temporal trends in the lethal concentration of 3-trifluoromethyl-4-nitrophenol (TFM) required to kill 50% (LC50) and 99.9% (LC99.9) of Sea Lamprey larvae. Range of years and number of toxicity tests with estimates of the LC50 and LC99.9 are indicated.

Source	Year range	No. LC50	No. LC99.9
Archive	1958-1970	38	38
Applegate et al. (1961)	1961-1961	3	3
Scholefield and Seelye (1990)	1963-1987	19	19
Quality assurance and control tests (QA/QC)	1988-2014	181	210

Draft

Figure A2. Toxicity of 3-trifluoromethyl-4-nitrophenol (TFM) to larval sea lamprey exposed for 19-24 hours in static toxicity tests in Lake Huron water conducted at the Hammond Bay Biological Station. Data were from archival records, historical reports (Applegate et al. 1961; Scholefield and Seelye 1990), and quality assurance tests of TFM supplies (Adair and Sullivan 2014). Linear regression fit to all data (solid line) and to all but the archival data (dashed line). A small random offset was added to sample from the same year to distinguish overlapping points. The estimated slope for LC50 from all data was -0.009 (95% confidence interval: -0.011 to -0.007). Without the archival data, the LC50 slope was -0.002, (95% confidence interval: -0.004 to 0.001). The estimated slope for the relative ratio from all the data was -0.00023 (95% confidence interval: -0.0020 to 0.0015). Without the archival data, the relative ratio slope was -0.0046, (-0.0067 to -0.0025). Six outliers (LC50 or relative ratio >3) from the archival records were removed; as these high values occurred early in the time series, their removal does not impact the sign of the slope.

References

- Adair, R.A., and Sullivan, W.P. 2014b. Standard operating procedures for application of Lampricides in the Great Lakes Fishery Commission integrated management of sea lamprey (*Petromyzon marinus*) control program, Marquette Biological Station, U.S. Fish and Wildlife Service, Marquette, M.I. Available from <http://www.glfc.org/sealamp/sop.php> [accessed 10 November 2016]
- Applegate, V.C., Howell, J.H., Miffed, J.W., Johnson, B.G.H., and Smith, M.A. 1961. Use of 3-trifluoro- methyl-4-nitrophenol as a selective sea lamprey larvicide. Great Lakes Fishery Commission Technical Report 1, Ann Arbor, M.I.
- Bills, T.D., Boogaard, M.A., Johnson, D.A., Brege, D.C., Scholefield, R.J., Westman, R.W., and Stephens, B.E. 2003. Development of a pH/alkalinity treatment model for applications of the lampricide TFM to streams tributary to the Great Lakes. J. Gt. Lakes Res. **29**: 510-520.
- Irwin, B.J., Liu, W.H., Bence, J.R., and Jones, M.L. 2012. Defining economic injury levels for sea lamprey control in the Great Lakes basin. N. Amer. J. Fish. Manage. **32**(4): 760-771.
- Jones, M.L., Irwin, B.J., Hansen, G.J.A., Dawson, H.A., Treble, A.J., Liu, W., Dai, W., and Bence, J.R. 2009. An operating model for the integrated pest management of Great Lakes sea lampreys. The Open Fish Science Journal **2**: 59-73.
- Litchfield, J.T.J., and Wilcoxon, F. 1949. A simplified method of evaluating dose-effect experiments. J. Pharmacol. Exp. Ther. **99**(2): 99-113.
- Scholefield, R.J., and Seelye, J.G. 1990. Resistance to 3-trifluoromethyl-4-nitrophenol (TFM) in sea lamprey. Great Lakes Fishery Commission Technical Report 56, Ann Arbor, M.I.