Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

**Predatory behaviour of an invasive amphipod in response to
varying conspecific densities under higher-order predation
risk**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0417.R1
Manuscript Type:	Article
Date Submitted by the Author:	23-Feb-2017
Complete List of Authors:	Iacarella, Josephine; University of Victoria, Biology Hudgins, Emma; McGill University, Biology Dick, Jaimie; The Queen`s University of Belfast Ricciardi, Anthony; McGill University
Please Select from this Special Issues list if applicable:	
Keyword:	BEHAVIOR < General, PREDATOR-PREY INTERACTION < General, INVASIVE SPECIES < Organisms, ZOOBENTHOS < Organisms, CRUSTACEANS < Organisms

SCHOLARONE™
Manuscripts

Predatory behaviour of an invasive amphipod in response to varying conspecific densities under higher-order predation risk

Josephine C. Iacarella^{1,a}, Emma J. Hudgins², Jaimie. T. A. Dick³, Anthony Ricciardi^{1*}

¹*Redpath Museum, McGill University, 859 Sherbrooke Street West, Montreal, Quebec, Canada H3A 0C4*

²*Department of Biology, McGill University, 1205 Avenue Docteur Penfield, Montreal, Quebec Canada H3A 1B1*

³*Institute for Global Food Security, School of Biological Sciences, M.B.C., Queen's University Belfast, 97 Lisburn Road, Belfast, Northern Ireland, UK BT9 7BL*

^a*Current Address: Ecosystem Sciences Division, Fisheries and Oceans Canada, Institute of Ocean Sciences, 9860 W Saanich Rd, Sidney, Canada V8L 5T5*

Email addresses: josie.iacarella@gmail.com, emma.hudgins@mail.mcgill.ca, j.dick@qub.ac.uk, tony.ricciardi@mcgill.ca

*Corresponding author: tony.ricciardi@mcgill.ca, tel: 514-398-4089, fax: 514-398-3185

Abstract

Behavioural responses of invasive animals to biotic interactions can inform predictions of their consumptive impacts; however, such biotic contexts are often overlooked. Here, we assessed the interacting effect of conspecific and higher-order predation risk on the per capita consumption and behaviours of the invasive freshwater amphipod *Gammarus pulex*, using field microcosm and video-recorded lab experiments in Northern Ireland. *G. pulex* exhibited higher per capita consumption in the presence of conspecifics, owing to reduced handling time of prey, regardless of fish presence and despite reduced swimming time and increased time spent physically interacting with each other. Consumption was lower in the presence of fish in the field, and handling time decreased with greater amphipod densities in the presence of fish cue in the lab. Our results show that impacts of *G. pulex* are independently influenced by conspecifics and predation risk, whereas handling time revealed an interacting effect of conspecific density and predation risk. Further assessments of the responses of invasive animals to biotic interactions could help explain variability in their impacts at local spatial scales.

Keywords: biotic interaction, Gammarus pulex, interference competition, invasive species, predation risk

Introduction

Analysis of the behavioural responses of invasive animals under different contexts is an underused but potentially valuable approach to explaining and predicting their ecological impacts. Behavioural responses are a determinant of invasion success across all stages of the invasion process: i.e. transport, introduction, establishment and spread (Chapple *et al.* 2012). Ecological impacts of invasive animals may occur throughout the invasion process following introduction (Ricciardi *et al.* 2013). Higher impacts have been associated with particular behaviours; for example, invaders exhibiting higher levels of aggression are more likely to competitively displace natives (Holway and Suarez 1999). Such invaders may also engage in enhanced foraging activity that maintains high densities despite aggressive conspecific interactions (Pintor *et al.* 2009). Biotic interactions are the primary determinants of invasion success and impact at local scales, whereas the abiotic context of invasion may be a more important predictor at the regional scale (Pintor and Sih 2011).

Invader impact is a function of the per capita effect and abundance of the species (Parker *et al.* 1999; Dick *et al.* 2017). Measurements of per capita effects (e.g. resource consumption rate of individuals) provide a mechanistic understanding of impact that enables the development of context-dependent predictions for risk-assessment (Dick *et al.* 2014). Thus far, most tests of per capita effect are performed in simple systems involving the measurement of one-way interactions, such as predation by an individual invader (Bollache *et al.* 2008; Haddaway *et al.* 2012). However, simple-system experiments may over- or under-estimate per capita effects in more complex biotic contexts (Médoc *et al.* 2013).

The functional response of an invader – the relationship between predation rate and prey supply – is a useful tool to mechanistically measure and predict predatory impacts under various

abiotic (Iacarella and Ricciardi 2015; Iacarella *et al.* 2015a) and biotic conditions (Barrios-O'Neill *et al.* 2014; Paterson *et al.* 2014). Prey-dependent models involving a single predator (Holling 1959) are widely used to model the functional response, with predation rate as a function of prey density. Conversely, ratio-dependent models are a function of the ratio of predators to prey (Arditi and Ginzburg 1989), and have been found to characterize the predatory behaviours of both native and invasive amphipods under varying prey and predator densities (Médoc *et al.* 2013; Médoc *et al.* 2015). Furthermore, amphipod species that aggregate more than others have higher handling times (model estimate of time to handle and digest a prey), likely from interference competition (Médoc *et al.* 2015). The factors that determine the level of interference between predators are not well understood (DeLong and Vasseur 2011) and may be influenced by other biotic interactions such as higher-order predation risk. For instance, the presence of fish cue has been suggested to induce aggregation of the amphipod *Gammarus pulex* (Kullmann *et al.* 2008), potentially also leading to increased conspecific interference. As ratio-dependence has been well established for amphipod species (Médoc *et al.* 2013; Médoc *et al.* 2015), here we evaluate the predatory and behavioural responses of an invasive amphipod in the presence of varying conspecific densities while also under higher-order predation risk.

We assessed several aspects of predation by *G. pulex*, an invasive freshwater amphipod. In field and lab experiments in Northern Ireland, we varied prey density, amphipod density and the presence/absence of fish. Field experiments were used to determine whether previously identified sources of variation in predatory behaviours of other freshwater invaders – specifically, abiotic conditions (Iacarella and Ricciardi 2015; Iacarella *et al.* 2015a) and population source (Iacarella *et al.* 2015b) – altered the amphipods' response to conspecific and fish presence. Reciprocal transplant field experiments improved our ability to make inferences

from the results by testing two source populations of amphipods at both of the two sites. We measured per capita consumption in the field and lab, and quantified behaviour using video in the lab. In addition, we measured handling time of prey (from capture to complete consumption) using video-recordings, in contrast to studies that infer handling times from functional response model estimates (Jeschke *et al.* 2002). *G. pulex* is non-native in Northern Ireland, where it has formed high density aggregations (Dick and Platvoet 1996; MacNeil *et al.* 2003), largely replaced the native amphipod *Gammarus duebeni celticus* (Dick *et al.* 1993; MacNeil *et al.* 2004), and reduced local macroinvertebrate diversity through predation and competition (Kelly and Dick 2005; Kelly *et al.* 2006). It is prey to numerous fishes, including brown trout *Salmo trutta* L. (review by MacNeil *et al.* 1999), and shows behavioural responses to brown trout cues (Åbjörnsson *et al.* 2000). We predicted that an interaction between conspecific and fish presence would lead to the lowest per capita consumption (number of prey eaten per amphipod) in larger conspecific groups while in the presence of fish, owing to increased handling time of prey, reduced time spent swimming (e.g. search effort) and more time spent interacting (e.g. heightened interference). We hypothesized that we would not observe a difference in predation rates between the field sites, nor between the two populations of amphipods. We also assessed the pupation and aggregation response of the larval prey in field and lab settings. Finally, we tested the predatory response of solitary amphipods to increasing prey densities to determine whether observed changes in per capita consumption with amphipod density (in the previous experiments) were a response to prey or amphipod densities.

Methods

88 *Field microcosm experiment: data collection*

89 Reciprocal transplant microcosm experiments were performed in June 2013 at two sites
90 near Lough Neagh, Northern Ireland. *G. pulex* predation was compared at a stream connected to
91 the Salterstown River, 1.6 km (geodesic distance) from a site of introduction in the late 1950s
92 (“Salterstown R. site”, 54°40'52.61"N, 6°31'51.22"W) (Strange and Glass 1979; Dick and
93 Platvoet 1996), and a stream connected to the Bann River approximately 48 km north of the
94 Salterstown R. site (“Bann R. site”, 55°6'4.41"N, 6°37'38.64"W). Only *G. pulex* were found at
95 the Salterstown R. site, whereas the native *G. duebeni* was also found at the Bann R. site.
96 Experiments were run using size-matched *G. pulex* individuals collected by kick-net from
97 Salterstown and Bann R. sites (total length, from the tip of the rostrum to the base of the telson:
98 15.8 ± 0.1 mm ($\pm$ 1SE; $n = 57$) and 16.0 ± 0.2 mm ($n = 56$), respectively). Only male *G. pulex*
99 without visible acanthocephalan parasites, such as *Echinorhynchus truttae*, were used in
100 experiments to avoid differences in feeding rates owing to reproductive or parasitic effects (Dick
101 *et al.* 2010; Paterson *et al.* 2014). Size, sex and absence of parasites were all selected for in the
102 field, thus only individuals used in the experiments were kept in the holding tanks.

103 Throughout the experiments, animals were held in aerated river water at 12 °C on a 12:12
104 hr light:dark cycle. *G. pulex* was collected two days prior to both sets of field experiments at
105 Salterstown and Bann R. sites and kept in source water from their respective sites (Salterstown
106 R. site conductivity: $340 \mu\text{S cm}^{-1}$, Bann R. site: $371 \mu\text{S cm}^{-1}$) with a food resource of leaf litter.
107 Amphipods were held as a group without food 51-52 h prior to the trials in filtered Lagan River
108 (“Minnowburn”, 54°32'54.36"N, 5°57'8.53"W) water ($420 \mu\text{S cm}^{-1}$) to standardize hunger levels.
109 Commercially raised brown trout (fork length: 109.9 ± 5.1 mm, $n=16$; sex not determined) were
110 kept in filtered Minnowburn water and fed juvenile *G. pulex* daily *ad libitum* (collected from

Minnowburn), beginning four days prior to the first experiment. Dipteran larvae *Simulium* spp. (6.9 ± 0.1 mm) were collected at stream sites separate from experimental sites for use as prey on the same day. Experimental treatments had one or four *G. pulex* with 15 dipteran larvae initially provided per amphipod and an individual fish either present or absent. Prey depletion was allowed in all experiments in order to not disturb the behaviours of the animals. Dipteran larvae are commonly used in predation experiments with *G. pulex* with similar densities provided (Paterson et al. 2014; Médoc et al. 2015). These larvae are consumed in high amounts by *G. pulex* relative to other macroinvertebrate prey (Krisp and Maier 2005), and overlap in habitat with *G. pulex* in Northern Ireland (Kelly et al. 2006).

Microcosms consisted of two attached black opaque PVC tubes 100 mm in diameter: the tube containing amphipods and dipteran larvae was 100 mm long, contained four, 1.5 cm flat glass beads attached to the bottom, and was enclosed on either side with 1 mm nitex mesh lined with mosquito netting (approximately $0.2 \text{ holes mm}^{-2}$) to further reduce the mesh size; the tube containing a brown trout was 200 mm long and covered at the end with 1 mm mesh (Fig. 1). Dipteran larvae, amphipods and brown trout were added consecutively and as quickly as possible to microcosms held in tubs with water; dipteran larvae were observed to attach immediately to surfaces as soon as contact was made (J.C.I., pers. obs.). Microcosms were then haphazardly placed along the river bed edges in places where there was a steady current, and were weighted down at 45° angles to the flow of water (MacNeil et al. 2004). Water passed through the tube containing fish to the tube containing *G. pulex* and dipteran larvae so that the amphipods could potentially sense both the movement and chemical cue of the fish. Experiments were run on 4-5 consecutive days at each site for 20 h from 9 pm – 5 pm for a total of six replicates. Past predation experiments with amphipods and dipteran prey have found interference competition

(i.e. ratio-dependent predation) in experiments run for 8 h (Médoc et al. 2015) and 12 h (Médoc et al. 2013) in stable lab conditions, and other similar experiments have run for up to 40 h (Bollache *et al.* 2008, Paterson *et al.* 2014). We ran these experiments for 20 h to allow adequate time to be able to observe predation differences between treatments, particularly in more challenging feeding conditions with current flow and with a live fish present and possibly causing disturbance.

The field experiment used two levels of amphipod densities (1 or 4), two levels of fish presence (present or absent), two sites (Salterstown R. or Bann R.), two amphipod sources (Salterstown R. or Bann R.) and six replicates of each trial for a total of 96 trials. Controls with neither *G. pulex* nor fish present were run simultaneously at the highest dipteran larvae density of 60 individuals per microcosm, with $93.3 \pm 6.7\%$ larvae remaining in microcosms at the Salterstown R. site and $91.3 \pm 3.3\%$ remaining at the Bann R. site. Some dipteran larvae escaped through the mesh and netting during the trials (but escape rate did not differ between sites, binomial Generalized Linear Model, $z=-0.784$, $df_{\text{residual}}=6$, $p=0.43$), and an additional $6.7 \pm 1.8\%$ pupated (see below for analysis of pupation). No *G. pulex* or brown trout escaped during the experiment.

Field microcosm experiment: data analysis

All remaining dipterans (larvae and pupa) were counted at the end of the trial and subtracted from the amount provided to calculate the total amount eaten by *G. pulex*; this amount was then multiplied by the average percent remaining in controls without amphipods (93.3% for trials at Salterstown R. and 91.3% for Bann R.) to account for some prey escape. Replicates in which an individual *G. pulex* moulted (3 cases) were excluded from the analysis as this can

affect predation rates (Bovy *et al.* 2015). Per capita consumption (number of prey eaten divided by number of amphipods present) of *G. pulex* was compared across treatments, with one per capita consumption response for each treatment replicate representing the group response as a whole. A generalized linear mixed-effects model (GLMM) with a binomial distribution (“lme4” in R; Bates *et al.* 2014) was used to assess the effects of amphipod density, fish presence, experimental site and *G. pulex* origin. A partially-crossed random factor of experimental night nested within experimental site was included to account for any effect of time (Bates 2010). Full models with interactions between all factors were reduced using Akaike information criterion for small sample sizes (AICc; Bolker *et al.* 2009). In addition, the additive effects of amphipod density (including controls with no amphipods), fish presence and experimental site were evaluated for the proportion of dipteran larvae that pupated (for 0, 1 and 4 amphipods with 60, 15 and 60 larvae, respectively) using a binomial GLMM and the same random effect structure (interaction terms were not tested in this case as the additional amphipod factor level over-parameterized the model). Best-fit models were determined by the lowest AICc value and were run to determine *post-hoc* contrasts between retained factor levels.

Behavioural lab experiment: data collection

Experiments were performed in the lab to further assess the effect of conspecific presence and fish presence on the predatory response of *G. pulex* with controlled conditions and the ability to observe movement and social behaviours. These experiments differed from the field experiments, in part, with the inclusion of a larger amphipod density and the use of fish cue instead of a live fish; fewer prey were provided per amphipod to accommodate the larger amphipod density while maintaining adequate water conditions. Predation was measured in

groups of one, four and eight *G. pulex*, with eight dipteran larvae provided per amphipod (8, 32 and 64 prey, respectively). *G. pulex* was used from both Salterstown and Bann R. sites, as field experiments showed no differences in their per capita consumption (see Results), but were kept separate in order to verify lack of differences based on origin, with the exception of four cases in which they were mixed to complete the trials. As in the field experiments, amphipods were held as a group without food 51-52 h prior to the trials in Minnowburn water to standardize hunger levels. Aerated, filtered Minnowburn water was also used in the experimental containers. Fish cue water was prepared by separating two brown trout into 4 L of water and feeding them ten juvenile *G. pulex* daily for two days, followed by a two-day period without food in 2 L of clean water (Alexander *et al.* 2013). Experiments were run in 75 mL of water (with fish cue or without) in freshly-washed containers (8 cm diameter) with five, 1.5 cm flat glass marbles attached to the bottom (with 0.5 cm spacing between marbles and 1 cm spacing from the edge of the container). Dipteran larvae were added first and observed to attach to the marbles, bottoms and sides of the containers; amphipods were added immediately thereafter. Experiments were run for 2 h in artificially-lit, 19°C ambient conditions with webcams recording each trial. An effort was made to run each level of amphipod density simultaneously, and fish cue treatments were alternated until 11 replicates were obtained for each treatment. Trials in which amphipods moulted (1 case) or were dead or listless at the end of the experiment (3 cases in group trials where amphipods were observed to attack an individual) were repeated. Some *G. pulex* were re-used in seven of the replicates (out of 66) spread out among treatments. Re-used amphipods were put back into holding tanks for at least four days between trials and those that were used in fish cue treatments were only re-used in no cue treatments, and vice versa. Controls with dipteran larvae in water with and without fish cue were run three times at each prey density (8, 32 and 64)

with $99.4 \pm 0.4\%$ survival across the three densities. No dipteran larvae pupated during the lab experiments.

Behavioural lab experiment: video analysis

From video recordings, the handling time for a single dipteran larva was measured by tracking the consumption of the first prey selected by the last amphipod added to the container. Handling time was determined as the time spanning the initial grasping of the prey to full consumption of the prey. Trials were not included if the prey was not immediately fully consumed (e.g. prey were abandoned or no prey were captured), leaving 6-9 replicates per treatment; the number of trials in which the focal amphipod abandoned its first prey or did not capture any prey were distributed evenly across amphipod and fish cue treatments (Fisher's exact test, $p=0.43$ and $p=0.18$, respectively).

The time focal amphipods spent swimming, resting or interacting was quantified using JWatcher (Blumstein *et al.* 2010). Recordings were first divided into eight, 14 min sections beginning 4 min after the start of each trial; four of the eight sections were randomly selected for analysis using a random number generator. The selected time sections for each treatment were evenly distributed across the experimental duration (chi-squared tests, $p=0.99$). In four and eight amphipod trials, the individual to be observed for each time section was chosen by numbering each amphipod and randomly generating the number to be selected. Swimming was categorized as any motion by the focal amphipod while it was not in contact with conspecifics; this was used to approximate search effort for prey. Though amphipods may swim for other reasons including to search for shelter or mates, the amphipods were able to discover more prey as they swam across the experimental arena. Resting was quantified as any time the focal amphipod was

stationary and was not in contact with conspecifics, whereas interacting (four and eight amphipod trials only) was determined to be any stationary or moving contact between the focal amphipod and one or more conspecifics. We considered any contact between the focal *G. pulex* and its conspecifics as an interaction, rather than limiting it to times when amphipods were directly interfering with each other (e.g. combative behaviour), as we assumed that they would be aware of all contact and this may have an effect on responses to prey and predation risk.

Aggregation of dipteran larvae in response to the presence of amphipods and fish cue was evaluated using three randomly selected 0- (control) and 8-amphipod trials – each initially containing 64 larvae – both with and without fish cue (n=12). Video snapshots were taken at 0 and 120 minutes, and the distances between all larvae were determined in ImageJ (Schneider et al. 2012). The change in distance was measured as the mean of all distances at the end of the experiment minus the mean at the beginning.

Behavioural lab experiment: data analysis

The effects of amphipod density, fish cue and amphipod source (Salterstown R. site, Bann R. site, mixed or re-used) on per capita consumption of *G. pulex* in the lab were analyzed using a generalized linear model (GLM) with a binomial distribution. Tukey's *post hoc* tests were then used to compare per capita consumption between amphipod densities. The effect of amphipod density and fish cue on handling time (i.e. the time taken to fully consume a dipteran larva once captured) was assessed using multiple linear regression followed by Tukey's *post hoc* tests. Handling time was $\log_{10}$ transformed to improve normality (Shapiro-Wilk, $p=0.30$), and variances remained homogenous (Fligner-Killeen, $p=0.42$). Separate binomial GLMMs were run to assess the effect of amphipod density and fish cue on the time spent performing each of the

three identified behaviours. Repeat measures of the four time samples per trial were included as a random effect in these models to account for potential variation in behaviour across different time sections. All full models with interaction terms were reduced to best-fit models for *post-hoc* analyses using AICc, as described previously. Finally, multiple linear regression was used to compare the change in mean distances in the presence/absence of amphipods and fish cue.

Prey-dependent functional response experiment: data collection and analysis

A functional response experiment with solitary *G. pulex* individuals was run to assess the prey density at which per capita consumption no longer rises in response to increasing prey (i.e. the asymptote of the curve). *G. pulex* were collected from a Lagan R. site (54°21'44.42"N, 6°1'26.22"W) within the same invaded waterway as the previous collection sites, and dipteran larvae were collected from Minnowburn. *G. pulex* were kept in source water and starved for 50 h in a 50:50 mix of Lagan R. and Minnowburn water prior to experiments. Experiments were run in 100 mL of mixed water in pots (9 cm diameter) for 24 h at 12 °C beginning at 10:45am, with artificial light from 9am to 9pm. *G. pulex* individuals were provided with prey densities of 2, 4, 8, 16, 30 and 40. Each prey density was replicated six times (except 8 prey, which was replicated 5 times). Trials were redone if *G. pulex* moulted up to 48 h after the experiment. Controls of prey without amphipods were run four times at the highest prey density with $96.3 \pm 3.2\%$ survival.

We determined whether the functional response was of Type II or Type III, following the methods of Alexander et al. (2012) and Dick et al. (2013). Type I responses are generally found for filter feeders (Jeschke *et al.* 2004), so we restricted our analysis to Type II and III. We derived functional response types using logistic regressions of the proportion of prey consumed as a function of prey density. The functional response curve was then modelled using maximum

likelihood estimation (“bbmle” in R; Bolker 2016) with Rogers’ random predator equation (Rogers 1972) for Type II curves with non-replacement of prey (Juliano 2001). Bootstrapping with replacement was then used to generate multiple curve fits ($n = 1500$) with 95% confidence intervals around the mean curve (“frair” in R; Pritchard *et al.* 2016).

Amphipod and prey densities across experiments

We estimated the comparable amount of amphipod and prey densities provided across the field, behavioural lab and functional response experiments based on the basal surface area of the experimental arenas (bottom half of cylindrical surface area with no base = 0.016 m^2 , bottom of round container = 0.005 m^2 , bottom of round container = 0.006 m^2 , respectively) as well as the volume (0.785 L, 0.075 L, 0.1 L, respectively). We provide basal surface area as *G. pulex* is a benthic predator and generally does not utilize the upper extent of the experimental containers (J.C.I., pers. obs); furthermore, field observations of *G. pulex* densities are based on area (MacNeil *et al.* 2003). Our experimental densities of amphipods are well within the range of densities observed in the field (Table 1). *G. pulex* densities in Northern Ireland are generally 200 – 1000 m^{-2} within the invaded range, and can reach up to 2000 m^{-2} (MacNeil *et al.* 2003). Finally, the density of prey provided to solitary amphipods in the field and in behavioural lab experiments was equivalent to six and ten prey provided, respectively, in the functional response experiment (Table 1). Dipteran larvae have been found at densities around $100,000 \text{ m}^{-2}$ (Dudley *et al.* 1986), which exceeds our highest prey density of $12,800 \text{ m}^{-2}$ with eight amphipods in the behavioural lab experiments.

Results

295

296 *Field microcosm experiment*

297 Per capita consumption by *G. pulex* was significantly higher in the presence of
298 conspecifics and lower in the presence of fish ($z=2.123$, $p=0.034$ and $z=2.314$, $p=0.021$,
299 respectively) (Fig. 2a and b; see Table 2 for all model results). The site of the experiments also
300 significantly influenced per capita consumption for amphipods of both origins ($z=2.643$,
301 $p=0.008$), whereas amphipods had similar per capita consumption within a site regardless of their
302 origin ($p>0.05$) (Fig. 2c). The feeding rate of solitary amphipods without fish present in the field
303 was 0.25 ± 0.03 prey h^{-1} (see below for comparable feeding rates in lab experiments).

304

305 *Behavioural lab experiment*

306 In the lab, per capita consumption was affected by conspecific amphipod density, but not
307 by fish cue presence nor amphipod source (Table 2). Groups of four and eight amphipods tended
308 to have higher per capita consumption than amphipods alone ($z=2.175$, $p=0.075$ and $z=2.217$,
309 $p=0.068$, respectively), but per capita consumption was similar between the two higher
310 amphipod densities ($z=0.044$, $p=0.999$) (Fig. 3a). The feeding rate of solitary amphipods without
311 fish cue present in the shorter lab experiments was 0.95 ± 0.22 prey h^{-1} .

312 Amphipod density and fish cue presence had an interacting effect on the handling time
313 of *G. pulex* ($F_{5,38}=3.335$, $p=0.014$) (Fig. 3b). Handling times were not significantly different
314 across amphipod densities with no fish cue, whereas they declined with density in the presence
315 of fish cue (1 v. 8 amphipods: $t=-3.720$, $p=0.008$; Table 2).

316 The importance of amphipod density and fish cue treatments for predicting movement and
317 social behaviours varied across measures (Table 2). Swimming time of *G. pulex* significantly

declined with larger groups (all group level comparisons: $p \leq 0.003$), and tended to decline in the presence of fish cue ($z=1.722$, $p=0.085$) (Fig. 4). The amount of time amphipods spent resting was similar across all treatments, whereas the time spent interacting with each other significantly increased between the four and eight amphipod groups ($z=7.657$, $p<0.001$). Fish cue had no effect on interaction time (Table 2).

The proportion of dipteran larvae that pupated during the field experiments was lower when four amphipods were present than when one or none were present (4 vs. 0, $z=-3.149$, $p=0.005$; 4 vs. 1, $z=-3.933$, $p<0.001$; 0 vs. 1, $z=0.174$, $p=0.983$) (Fig. 5a). Neither fish presence nor the experimental site had an effect on pupation ($z=0.509$, $p=0.611$ and $z=0.432$, $p=0.666$, respectively). In the lab, the change in mean distance between larvae was greater and more negative (i.e. they became closer together) when amphipods were present ($F_{1,8} = 15.500$, $p = 0.004$) (Fig. 5b). Fish cue had no effect on larval aggregation, nor was there an interaction ($F_{1,8} = 0.102$, $p = 0.758$ and $F_{1,8} = 0.479$, $p = 0.509$, respectively).

Prey-dependent functional response experiment

G. pulex showed a Type II functional response to dipteran larvae with an asymptote in per capita consumption beginning at eight prey provided ($z=-9.390$, $df_{\text{residual}}=33$, $p<0.001$) (Fig. 6). The maximum feeding rate of *G. pulex* obtained from the asymptote of the functional response curve was 0.24 prey h^{-1} .

Discussion

The predatory behaviour of *G. pulex* in our field and lab experiments was affected by the presence of conspecifics and higher-order predation risk, though handling time was the only

response variable for which conspecific and fish presence had interactive effects. Consumptive impacts of invasive animals have often been measured as one-way interactions between the invasive consumer and the food resource, in order to make relative comparisons of per capita effects (Bollache *et al.* 2008; Haddaway *et al.* 2012; Dick *et al.* 2013; Alexander *et al.* 2014). A recent meta-analysis of functional response behaviour experiments shows that simple experimental assessments explain and predict community level impacts of invaders across diverse taxonomic and trophic groups (Dick *et al.* 2017). Our results suggest that further biotic interactions may alter the results of such experiments and inclusion of more complex contexts may be necessary to obtain measurements of per capita effects that predict field impacts at a higher resolution (Médoc *et al.* 2013; Paterson *et al.* 2014; Dick *et al.* 2017). Predatory and risk-avoidant responses of invasive animals to the presence of conspecifics and higher-order predation risk vary with species and experimental design (Ellrott *et al.* 2007; Paterson *et al.* 2013). The potential complexity of predatory behavioural responses highlights the need for consideration of biotic context in impact experiments.

Influence of conspecifics

Contrary to our prediction of reduced consumption in the presence of conspecifics and fish, per capita consumption of *G. pulex* increased with conspecific presence, although not with larger amphipod densities (Fig. 2a & 3a). The prey densities provided to solitary amphipods for field and behavioural lab experiments matched densities at the beginning and end of the functional response curve asymptote, respectively (Fig. 6). This indicates, particularly for the lab experiments, that feeding rates may not have increased because of the higher prey densities provided to the larger groups of amphipods. Rather, the observed increase in per capita

consumption in response to the presence of conspecifics may be an effect of the amphipods themselves. Sura and Mahon (2011) attributed a tendency for higher feeding rates in larger groups of snails to exploitative competition for a limiting food resource, in which case the conspecifics do not affect each other through direct contact (de Villemereuil and López-Sepulcre 2011). Conversely, Médoc et al. (2015) related lower consumption of an amphipod species in larger groups to its propensity for aggregating with conspecifics. In our experiments, the initial presence of *G. pulex* individuals appeared to have stimulated each other's predatory response. Per capita consumption rates increased from one to four amphipods, despite time spent interacting and reduced swimming time. Interestingly, this was offset by lowered handling times (Fig. 3b) – i.e. the amphipods consumed their prey more quickly, presumably allowing them to maintain higher consumption. The lack of per capita consumption difference between four and eight amphipod densities may indicate the onset of interference, as swimming time decreased further, interaction time increased, and handling times either increased or decreased, depending on the presence of fish cue (see “Influence of predation risk” below). These results suggest that per capita consumption in larger groups does not continue to increase; increased feeding rates triggered by conspecific presence may become outweighed by interference competition at higher densities.

Feeding satiation at high prey densities may prevent interference competition despite the presence of conspecifics. However, amphipod satiation was unlikely in the 2 h behavioural lab experiments, as amphipods were observed to commence feeding as late as 1 h 45 min into the trials (E.J.H. pers. obs.), and consumption was higher in the 20 h field experiments. Furthermore, results were consistent between the field and lab experiments, whereas interference at the highest amphipod density was only tested in the shortest duration experiment. Other similar

predation studies on amphipods have run experiments ranging from 8 to 40 h (Bollache *et al.* 2008; Médoc *et al.* 2013; Paterson *et al.* 2014; Médoc *et al.* 2015), and provide evidence for interference competition (Médoc *et al.* 2013; Médoc *et al.* 2015).

The addition of more individuals in higher amphipod density treatments could lead to some exhibiting greater predatory behaviours than others. However, we excluded common factors that lead to discrepancies in feeding by collecting animals and running experiments all within a short timeframe (two months), excluding parasite-infested amphipods (Dick *et al.* 2010; Paterson *et al.* 2014), only including males, matching body size (Médoc *et al.* 2013) and not using during- or post-trial moulting individuals (Bovy *et al.* 2015). Ratio-dependent models of amphipod consumption do not distinguish the behaviour of individuals within groups (Médoc *et al.* 2013; Médoc *et al.* 2015), though this could be an interesting avenue of further research.

A better understanding of predatory responses may be gained with observations of movement and social behaviours. To our knowledge, studies of changes in predatory responses with larger conspecific densities rarely observe the movement and social behaviours of the conspecifics. Here, we found that larger amphipod densities led to reduced time spent swimming (i.e. a proxy for reduced search effort) and increased time spent interacting in larger groups (Fig. 4). Single *G. pulex* spent most of their time swimming, whereas amphipods in groups of eight spent most of their time interacting with each other, suggesting heightened interference. Higher densities of amphipods than those used in our experiments could lead to interference levels that eventually reduce per capita consumption (i.e. gradual interference hypothesis; Ginzburg and Jensen 2008). Our experimental system was not large enough to maintain higher densities of prey and amphipods; however, the highest density of amphipods used (Table 1) is within the

range of maximum densities observed in Northern Ireland (1500-2000 individuals m⁻²) (MacNeil *et al.* 2003).

Influence of predation risk

We had predicted that interference would increase, and per capita consumption decline, when *G. pulex* was in the presence of predation risk owing to increased aggregation (Kullmann *et al.* 2008). We did not observe lower per capita consumption or increased interaction times with amphipod density in the presence of fish cue; instead, we found that handling times of *G. pulex* on individual prey items declined with amphipod density in the presence of fish cue (Fig. 2b). Reduced vigilance to predation risk is often observed in groups of animals (Lima and Dill 1990). Lower handling times (e.g. faster consumption of a prey) may be an indication of more focused feeding by individuals within groups, whereas solitary individuals may be more focused on the predation risk and thus forage more cautiously. Milinski and Heller (1978) demonstrated lower feeding rates by a solitary fish in the presence of a mimic avian predator, though the influence of conspecifics was not examined. Fish feeding at higher rates were also found to be less observant of the avian predator (Milinski 1984). Chemical cues are the dominant mechanism of predator detection in *G. pulex*, as amphipods cannot visually detect predators (Baldauf *et al.* 2007). The use of chemical cues may not necessitate a trade-off with foraging, though our study and others indicate reduced activity (here, a trend of reduced swimming) (Alexander *et al.* 2013) and foraging (here, reduced consumption in the field) (Åbjörnsson *et al.* 2000) when amphipods are in the presence of predation risk. However, once an amphipod captures a prey, they may consume it more quickly as a final effort to increase fitness in the face of mortality risk.

Reduction in consumption and active behaviours of amphipods in response to predation risk have been observed in this study and other lab experiments (Wudkevich *et al.* 1997; Åbjörnsson *et al.* 2000; Pennuto and Keppler 2008; Alexander *et al.* 2013), though responses can vary greatly across experiments (Paterson *et al.* 2013). For instance, intermediate fish predators have been found to spend more time hiding in refuge in the presence of a top fish predator (Romare and Hansson 2003). Åbjörnsson *et al.* (2000) also showed that *G. pulex* predation was temporarily reduced in the presence of European bullhead *Cottus gobio* fish cue; however, Paterson *et al.* (2014) found that *G. pulex* per capita consumption did not change in the presence of brown trout. Animals can habituate to the presence of predator cues, particularly if the signal is homogenous and repeated (Åbjörnsson *et al.* 2000; Mirza *et al.* 2006). Both our field and lab experiments were run for a shorter period of time than those of Åbjörnsson *et al.* (2000) and Paterson *et al.* (2014), and may be more representative of an anti-predatory response in field conditions where predator cues are unpredictable and short-term.

We also found evidence of altered behaviours of the dipteran prey in the presence of amphipods. Dipterans pupated less in the field when in the presence of more amphipods, likely to avoid vulnerability while building the casing (Fig. 5a). Increased aggregation of dipterans at the end of treatments with amphipods present also appeared to be a predator avoidant response (Fig. 5b). Alternatively, *G. pulex* may have selected prey that were more solitary, leaving those that were clustered together, though the larvae were observed to actively move away from open spaces during the experiment (E.J.H. pers. obs.). Dipteran larvae can be found at much greater densities than those we used (Dudley *et al.* 1986), but aggregations are highly patchy in the environment (J.C.I. pers. obs.)

Influence of habitat characteristics

Per capita consumption did not vary with the origin of *G. pulex*, in both field and lab experiments; however, per capita consumption of *G. pulex* from both sites was lower at the Salterstown R. site than the Bann R. site (Fig. 1c). Environmental conditions such as temperature and conductivity influence metabolic rates and physiological responses, and can thus mediate the predatory impact of aquatic invaders, as measured by their functional response (Iacarella and Ricciardi 2015; Iacarella *et al.* 2015a). Though weather conditions and temperature remained similar throughout the duration of the field experiments, conductivity was approximately 30 $\mu\text{S cm}^{-1}$ higher at the Bann R. site and perhaps contributed to higher per capita consumption (Dick and Platvoet 1996; Kestrup *et al.* 2011). In addition, the native *G. d. celticus* was observed at the Bann R. site, but not at the Salterstown R. site. *G. pulex* is a mutual, and superior, intra-guild predator of the native (Dick *et al.* 1993); the presence of the native at Bann R. may have triggered a heightened feeding response. Furthermore, *G. pulex* is more tolerant of low dissolved oxygen and has largely replaced the native in areas of poor water quality (MacNeil *et al.* 2004). Though we did not measure dissolved oxygen, the absence of *G. d. celticus* may indicate that Salterstown R. had lower water quality; consumption rates generally increase in more optimal conditions (Iacarella and Ricciardi 2015; Iacarella *et al.* 2015a). We were unable to determine the exact cause of the difference in per capita consumption between experimental sites, as we had *a priori* believed these sites to be fairly similar; however, these results exemplify the importance of assessing invader impacts across the invasion range.

Furthermore, in our experiments, the functional response of *G. pulex* was measured at a lower temperature than the field and behavioural lab experiments owing to logistical constraints, and this may have led to a lower maximum feeding rate (Iacarella *et al.* 2015a). However, field

predation rates of solitary individuals (per hour) were remarkably similar to the functional response maximum feeding rate (experimental durations 20 and 24 h, respectively), whereas the shorter-duration behavioural lab experiments yielded much higher per capita consumption; it is likely that initially high feeding rates are not sustained in longer experiments. The shorter-term lab results may be less representative of long-term impact, whereas the field results may better reflect *G. pulex* daily feeding rates. Similar responses to the presence of conspecifics and predation risk were observed in the field and lab, regardless of differences in experimental duration.

Conclusions

Our study demonstrates how the impacts of invasive animals are mediated by biotic interactions, as the type of interaction (i.e. interference, trophic) influences behaviour in such a way as to affect predation efficiency. Statistically significant differences in per capita consumption rates between invasive and native species, and among contexts, have been shown to relate to actual differences in field impact (Dick *et al.* 2013; 2017). Predictions of ecological impact must also consider behavioural responses of the invader in different biotic contexts. Experiments measuring per capita impacts may be improved by using natural field densities of the invader, as has been done frequently in mesocosm studies (Lodge *et al.* 1994; Mills *et al.* 2004; Gherardi and Acquistapace 2007), yet less commonly in mechanistic predatory studies (but see Médoc *et al.* 2013; Barrios-O'Neill *et al.* 2014). Simulating the level of predation risk that an invader experiences in the field may be problematic, but knowledge of the trophic structure of an invaded habitat can inform predictions of whether per capita effects and abundances will be reduced by risk avoidance and direct predation, respectively (Pimm 1987).

Such context dependencies present a major challenge to risk assessment and for prioritizing management efforts (Ricciardi *et al.* 2013; Kumschick *et al.* 2015). However, the accuracy of risk assessment may be enhanced by further identification of how behavioural responses and interactions of invasive animals affect their ecological impacts, and whether or not these effects can be generalized.

Acknowledgements

The authors thank A. Coyle, R. Paterson and E. McKnight for assistance with the microcosm experiments. This research was funded by the National Geographic Young Explorer's Grant to JCI. Additional support was provided by a McGill International Travel Award (to JCI), the Group for Interuniversity Research in Limnology and Aquatic Environment (to JCI), the Canadian Aquatic Invasive Species Network (to AR), a Natural Sciences and Engineering Research Council (Canada) Discovery Grant (to AR), the Leverhulme Trust (to JTAD) and a Natural Environment Research Council (UK) grant (to JTAD).

514 **References**

- 515 Åbjörnsson, K., Dahl, J., Nyström, P. and Brönmark, C. 2000. Influence of predator and
 516 dietary chemical cues on the behaviour and shredding efficiency of *Gammarus pulex*.
 517 Aquat. Ecol. **34**: 379–387.
- 518 Alexander, M., Dick, J., O'Connor, N., Haddaway, N. and Farnsworth, K. 2012. Functional
 519 responses of the intertidal amphipod *Echinogammarus marinus*: effects of prey supply,
 520 model selection and habitat complexity. Mar. Ecol. Prog. Ser. **468**: 191–202.
- 521 Alexander, M.E., Dick, J.T., Weyl, O.L., Robinson, T.B. and Richardson, D.M. 2014.
 522 Existing and emerging high impact invasive species are characterized by higher
 523 functional responses than natives. Biol. Lett. **10**: 20130946.
- 524 Alexander, M.E., Dick, J.T.A. and O'Connor, N.E. 2013. Trait-mediated indirect
 525 interactions in a marine intertidal system as quantified by functional responses. Oikos
 526 **122**: 1521–1531.
- 527 Arditi, R. and Ginzburg, L. 1989. Coupling in predator-prey dynamics: ratio-dependence. J.
 528 Theor. Biol. **139**: 311–326.
- 529 Barrios-O'Neill, D., Dick, J.T.A., Emmerson, M.C., Ricciardi, A., MacIsaac, H.J.,
 530 Alexander, M.E., et al. 2014. Fortune favours the bold: a higher predator reduces the
 531 impact of a native but not an invasive intermediate predator. J. Anim. Ecol. **83**: 693–701.
- 532 Bates, D.M. 2010. lme4: Mixed-Effects Modeling with R. Springer, New York.
- 533 Bates, D., Mächler, M., Bolker, B. and Walker, S. 2014. Fitting Linear Mixed-Effects Models
 534 using lme4. J Stat. Softw. **67**: 1-48.
- 535 Blumstein, D.T., Evans, C.S. and Daniel, J.C. 2010. JWatcher Software.
- 536 Bolker, B. 2016. bbmle: Tools for general maximum likelihood estimation. R package version

- 537 1.0.18.
- 538 Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H., et al.
- 539 2009. Generalized linear mixed models: a practical guide for ecology and evolution. Trends
- 540 Ecol. Evol. **24**: 127–135.
- 541 Bollache, L., Dick, J.T.A., Farnsworth, K.D. and Montgomery, W.I. 2008. Comparison of the
- 542 functional responses of invasive and native amphipods. Biol. Lett. **4**: 166–169.
- 543 Bovy, H.C., Barrios-O'Neill, D., Emmerson, M.C., Aldridge, D.C. and Dick, J.T.A. 2015.
- 544 Predicting the predatory impacts of the “demon shrimp” *Dikerogammarus haemobaphes*, on
- 545 native and previously introduced species. Biol. Invasions **17**: 597–607.
- 546 Baldauf, S.A., Thünken, T., Frommen, J.G., Bakker, T.C., Heupel, O. and Kullmann, H. 2007.
- 547 Infection with an acanthocephalan manipulates an amphipod’s reaction to a fish
- 548 predator’s odours. Int. J. Parasitol. **37**: 61–65.
- 549 Chapple, D.G., Simmonds, S.M. and Wong, B.B.M. 2012. Can behavioral and personality traits
- 550 influence the success of unintentional species introductions? Trends Ecol. Evol. **27**: 57–64.
- 551 DeLong, J.P. and Vasseur, D.A. 2011. Mutual interference is common and mostly intermediate
- 552 in magnitude. BMC Ecol., **11**: 1.
- 553 Dick, J.T.A., Alexander, M.E., Jeschke, J.M., Ricciardi, A., MacIsaac, H.J., Robinson, T.B., et
- 554 al. 2014. Advancing impact prediction and hypothesis testing in invasion ecology using a
- 555 comparative functional response approach. Biol. Invasions **16**: 735–753.
- 556 Dick, J.T.A., Armstrong, M., Clarke, H.C., Farnsworth, K.D., Hatcher, M.J., Ennis, M., et al.
- 557 2010. Parasitism may enhance rather than reduce the predatory impact of an invader. Biol.
- 558 Lett. **6**: 636–638.
- 559 Dick, J.T.A., Gallagher, K., Avlijas, S., Clarke, H.C., Lewis, S.E., Leung, S., et al. 2013.

- 560 Ecological impacts of an invasive predator explained and predicted by comparative
561 functional responses. *Biol. Invasions* **15**: 837–846.
- 562 Dick, J.T.A., Lavery, C., Lennon, J.J., Barrios-O'Neill, D., Mensink, P.J., Britton, J.R., et al.
563 2017. Invader relative impact potential: a new metric to understand and predict the
564 ecological impacts of existing, emerging and future invasive alien species. *J. Appl. Ecol.*
565 DOI: 10.1111/1365-2664.12849.
- 566 Dick, J.T.A., Montgomery, I. and Elwood, R.W. 1993. Replacement of the indigenous
567 amphipod *Gammarus duebeni celticus* by the introduced *G. pulex*: Differential cannibalism
568 and mutual predation. *J. Anim. Ecol.* **62**: 79–88.
- 569 Dick, J.T.A. and Platvoet, D. 1996. Intraguild predation and species exclusions in amphipods:
570 the interaction of behaviour, physiology and environment. *Freshw. Biol.* **36**: 375–383.
- 571 Dudley, T. L., Cooper, S. D. and Hemphill, N. 1986. Effects of macroalgae on a stream
572 invertebrate community. *J. N. Am. Benthol. Soc.* **5**: 93–106.
- 573 Ellrott, B.J., Marsden, J.E., Fitzsimons, J.D., Jonas, J.L. and Claramunt, R.M. 2007. Effects of
574 temperature and density on consumption of trout eggs by *Orconectes propinquus* and *O.*
575 *rusticus*. *J. Great Lakes Res.* **33**: 7–14.
- 576 Gherardi, F. and Acquistapace, P. 2007. Invasive crayfish in Europe: the impact of
577 *Procambarus clarkii* on the littoral community of a Mediterranean lake. *Freshw. Biol.* **54**:
578 1249–1259.
- 579 Ginzburg, L.R. and Jensen, C.X.J. 2008. From controversy to consensus: the indirect
580 interference functional response. *Verhandlungen des Int. Verein Limnol.* **30**: 297–301.
- 581 Haddaway, N.R., Wilcox, R.H., Heptonstall, R.E.A., Griffiths, H.M., Mortimer, R.J.G.,
582 Christmas, M., et al. 2012. Predatory functional response and prey choice identify predation

- 583 differences between native/invasive and parasitised/unparasitised crayfish. PLoS One 7:
584 e32229.
- 585 Holling, C.S. 1959. Some characteristics of simple types of predation and parasitism. Can.
586 Entomol. **92**: 385–398.
- 587 Holway, D.A. and Suarez, A.V. 1999. Animal behavior: An essential component of invasion
588 biology. Trends Ecol. Evol. **14**: 328–330.
- 589 Iacarella, J.C., Dick, J.T., Alexander, M.E. and Ricciardi, A. 2015a. Ecological impacts of
590 invasive alien species along temperature gradients : testing the role of environmental
591 matching. Ecol. Appl. **25**: 706–716.
- 592 Iacarella, J.C., Dick, J.T.A. and Ricciardi, A. 2015b. A spatio-temporal contrast of the predatory
593 impact of an invasive freshwater crustacean. Divers. Distrib. **21**: 803-812
- 594 Iacarella, J.C. and Ricciardi, A. 2015. Dissolved ions mediate body mass gain and predatory
595 response of an invasive fish. Biol. Invasions **17**: 3237–3246.
- 596 Jeschke, J.M., Kopp, M. and Tollrian R. 2002. Predator functional responses: discriminating
597 between handling and digesting prey. Ecol. Monogr. **72**: 95-112.
- 598 Jeschke, J.M., Kopp, M. and Tollrian R. 2004. Consumer-food systems: why type 1 functional
599 responses are exclusive to filter feeders. Biol. Rev. Camb. Philos. Soc. **79**: 337-349.
- 600 Juliano, S. 2001. Nonlinear curve fitting. Des. Anal. Ecol. Exp. Chapman and Hall, New York,
601 NY.
- 602 Kelly, D.W. and Dick, J.T.A. 2005. Effects of environment and an introduced invertebrate
603 species on the structure of benthic macroinvertebrate species at the catchment level. Arch. für
604 Hydrobiol. **164**: 69–88.

- 605 Kelly, D.W., Bailey, R.J., MacNeil, C., Dick, J.T. and McDonald, R. A. 2006. Invasion by the
606 amphipod *Gammarus pulex* alters community composition of native freshwater
607 macroinvertebrates. *Divers. Distrib.* **12**: 525-534.
- 608 Kestrup, Å.M., Dick, J.T.A. and Ricciardi, A. 2011. Interactions between invasive and native
609 crustaceans: Differential functional responses of intraguild predators towards juvenile hetero-
610 specifics. *Biol. Invasions* **13**: 731–737.
- 611 Krisp, H. and Maier, G. 2005. Consumption of macroinvertebrates by invasive and native
612 gammarids: a comparison. *J. Limnol.* **64**: 55-59.
- 613 Kullmann, H., Thunken, T., Baldauf, S.A., Bakker, T.C. and Frommen, J.G. 2008. Fish odour
614 triggers conspecific attraction behaviour in an aquatic invertebrate. *Biol. Lett.* **4**: 458–460.
- 615 Kumschick, S., Gaertner, M., Vila, M., Essl, F., Jeschke, J.M., Pysek, P., et al. 2015.
616 Ecological impacts of alien species: quantification, scope, caveats, and recommendations.
617 *Bioscience* **65**: 55–63.
- 618 Lima, S.L. and Dill, L.M. 1990. Behavioral decisions made under the risk of predation: a
619 review and prospectus. *Can. J. Zool.* **68**: 619–640.
- 620 Lodge, D.M., Kershner, M.W., Aloï, J.E. and Covich, A.P. 1994. Effects of an omnivorous
621 crayfish (*Orconectes rusticus*) on a freshwater littoral food web. *Ecology* **75**: 1265–1281.
- 622 MacNeil, C., Dick, J.T.A. and Elwood, R.W. 1999. The dynamics of predation on *Gammarus*
623 spp. (Crustacea: Amphipoda). *Biol. Rev.* **74**: 375–395.
- 624 MacNeil, C., Fielding, N.J., Hume, K.D., Dick, J.T.A., Elwood, R.W., Hatcher, M.J., et al.
625 2003. Parasite altered micro-distribution of *Gammarus pulex* (Crustacea : Amphipoda). *Int. J.*
626 *Parasitol.* **33**: 57–64.
- 627 MacNeil, C., Prenter, J., Briffa, M., Fielding, N.J., Dick, J.T., Riddell, G.E., et al. 2004. The

- 628 replacement of a native freshwater amphipod by an invader: roles for environmental
629 degradation and intraguild predation. *Can. J. Fish. Aquat. Sci.* **61**: 1627–1635.
- 630 Médoc, V., Spataro, T. and Arditi, R. 2013. Prey: predator ratio dependence in the functional
631 response of a freshwater amphipod. *Freshw. Biol.* **58**: 858–865.
- 632 Médoc, V., Albert, H. and Spataro, T. 2015. Functional response comparisons among freshwater
633 amphipods: ratio-dependence and higher predation for *Gammarus pulex* compared to the
634 non-natives *Dikerogammarus villosus* and *Echinogammarus berilloni*. *Biol. Inv.* **17**:
635 3625–3637.
- 636 Milinski, M. 1984. A predator's costs of overcoming the confusion-effect of swarming prey.
637 *Anim. Behav.* **32**: 1157–1162.
- 638 Milinski, M. and Heller, R. 1978. Influence of a predator on the optimal foraging behaviour of
639 sticklebacks (*Gasterosteus aculeatus* L.). *Nature* **275**: 642–644.
- 640 Mills, M.D., Rader, R.B. and Belk, M.C. 2004. Complex interactions between native and
641 invasive fish: the simultaneous effects of multiple negative interactions. *Oecologia* **141**: 713–
642 721.
- 643 Mirza, R.S., Mathis, A. and Chivers, D.P. 2006. Does temporal variation in predation risk
644 influence the intensity of antipredator responses? A test of the risk allocation hypothesis.
645 *Ethology* **112**: 44–51.
- 646 Parker, I.M., Simberloff, D., Lonsdale, W.M., Goodell, K., Wonham, M., Kareiva, P.M., et al.
647 1999. Impact: toward a framework for understanding the ecological effects of invader. *Biol.*
648 *Invasions* **1**: 3–19.
- 649 Paterson, R.A., Dick, J.T.A., Pritchard, D.W., Ennis, M., Hatcher, M.J. and Dunn, A.M. 2014.
650 Predicting invasive species impacts: a community module functional response approach

- 651 reveals context dependencies. *J. Anim. Ecol.* **84**: 453–463.
- 652 Paterson, R.A., Pritchard, D.W., Dick, J.T.A., Alexander, M.E., Hatcher, M.J. and Dunn, A.M.
653 2013. Predator cue studies reveal strong trait-mediated effects in communities despite
654 variation in experimental designs. *Anim. Behav.* **86**: 1301–1313.
- 655 Pennuto, C. and Keppler, D. 2008. Short-term predator avoidance behavior by invasive and
656 native amphipods in the Great Lakes. *Aquat. Ecol.* **42**: 629–641.
- 657 Pimm, S.L. 1987. Determining the effects of introduced species. *Trends Ecol. Evol.* **2**: 106–
658 108.
- 659 Pintor, L.M. and Sih, A. 2011. Scale dependent effects of native prey diversity, prey biomass
660 and natural disturbance on the invasion success of an exotic predator. *Biol. Invasions* **13**:
661 1357–1366.
- 662 Pintor, L.M., Sih, A. and Kerby, J.L. 2009. Behavioral correlations provide a mechanism for
663 explaining high invader densities and increased impacts on native prey. *Ecology* **90**: 581–
664 587.
- 665 Pritchard, D., Barrios-O'Neill, D., Bovy, H. and Paterson, R. 2016. frair: Tools for functional
666 response analysis. R package version 0.5.
- 667 Ricciardi, A., Hoopes, M.F., Marchetti, M.P. and Lockwood, J.L. 2013. Progress toward
668 understanding the ecological impacts of nonnative species. *Ecol. Monogr.* **83**: 263–282.
- 669 Rogers, D. 1972. Random search and insect population models. *J. Anim. Ecol.* **41**: 369–383.
- 670 Romare, P. and Hansson, L. A. 2003. A behavioral cascade: top-predator induced behavioral
671 shifts in planktivorous fish and zooplankton. *Limnol. Oceanogr.* **48**: 1956–1964.
- 672 Schneider, C.A., Rasband, W.S., Eliceiri, K.W. 2012. NIH Image to ImageJ: 25 years of image
673 analysis. *Nature Methods* **9**: 671–675.

- 674 Strange, C.D. and Glass, G.B. 1979. The distribution of freshwater Gammarids in Northern
675 Ireland. Proc. R. Irish Acad. **79B**: 145–153.
- 676 Sura, S.A. and Mahon, H.K. 2011. Effects of competition and predation on the feeding rate of
677 the freshwater snail, *Helisoma trivolvis*. Am. Midl. Nat. **166**: 358–368.
- 678 de Villemereuil, P.B. and López-Sepulcre, A. 2011. Consumer functional responses under
679 intra-and inter-specific interference competition. Ecol. Model. **222**: 419–426.
- 680 Wudkevich, K., Wisenden, B.D., Chivers, D.P. and Smith, R.J.F. 1997. Reactions of
681 *Gammarus lacustris* to chemical stimuli from natural predators and injured conspecifics.
682 J. Chem. Ecol. **23**: 1163–1173.

Draft

Table 1. Densities of amphipod predators and dipteran larvae prey used in field, behavioural lab and functional response lab experiments. Densities are based on basal surface area and water volume of the experimental arenas.

Animal #	Field	Behavioural lab	Functional response lab
1 amph.	62.5 m ⁻² ; 1.3 L ⁻¹	200.0 m ⁻² ; 13.3 L ⁻¹	166.7 m ⁻² ; 10.0 L ⁻¹
4 amph.	250.0 m ⁻² ; 5.1 L ⁻¹	800.0 m ⁻² ; 53.3 L ⁻¹	NA
8 amph.	NA	1600.0 m ⁻² ; 106.7 L ⁻¹	NA
Prey per amph.	937.5 m ⁻² ; 19.1 L ⁻¹	1600.0 m ⁻² ; 106.7 L ⁻¹	6 prey: 1000.0 m ⁻² ; 60.0 L ⁻¹ 10 prey: 1666.7 m ⁻² ; 100.0 L ⁻¹ 40 prey: 6666.7 m ⁻² ; 400.0 L ⁻¹

Draft

Table 2. Field and laboratory analysis results for per capita consumption (number of dipteran larvae consumed per amphipod), handling time (time to fully consume a prey item once captured) and social/movement behaviours (swimming, resting, interacting) of *Gammarus pulex*. The effects of amphipod density (Amph #), fish presence (live or cue), experimental site, amphipod origin and source for experiments were analyzed using generalized linear mixed-models (GLMMs), generalized linear models (GLMs) and multiple linear regressions (LMs); residual degrees of freedom (df) are provided for GLMMs and GLMs, and total df are provided for LMs. Best-fit models were reduced from full models containing all interaction terms using Akaike information criterion for small sample sizes (AICc; Δ AICc = full - best-fit model). In the best-fit models, interaction terms (“*”) and/or additive main effects (“+”) were retained.

Experiment, response, test	Model, df	Fixed-effects	Δ AICc	Best-fit term comparisons	Test statistic, p-value
Field, Consumption, GLMM	Full, df=76 Best-fit, df=88	Amph # * Fish * Site * Origin Amph # + Fish + Site	26.90	1. Amph: 1 v. 4 2. Fish: yes v. no 3. Site: Salt. v. Bann	1. $z=2.123$, p=0.034 2. $z=2.314$, p=0.021 3. $z=2.643$, p=0.008
Lab, Consumption, GLM	Full, df=49 Best-fit, df=63	Fish * Amph # * Source Amph #	24.44	1. Amph: 1 v. 4 2. Amph: 1 v. 8 3. Amph: 4 v. 8	1. $z=2.175$, $p=0.075$ 2. $z=2.217$, $p=0.068$ 3. $z=0.044$, $p=0.999$
Lab, Handling time, LM	Full & Best-fit, df=38	Amph # * Fish	0	Fish cue present 1. Amph: 1 v. 4 2. Amph: 1 v. 8 3. Amph: 4 v. 8 Fish cue absent 4. Amph: 1 v. 4 5. Amph: 1 v. 8 6. Amph: 4 v. 8	Fish cue present 1. $t=-1.178$, $p=0.844$ 2. $t=-3.720$, p=0.008 3. $t=-2.430$, $p=0.171$ Fish cue absent 4. $t=-1.238$, $p=0.814$ 5. $t=-0.018$, $p=1.000$ 6. $t=1.013$, $p=0.910$
Lab, Swimming time, GLMM	Full, df=257 Best-fit, df=259	Amph # * Fish Amph # + Fish	1.85	1. Amph: 1 v. 4 2. Amph: 1 v. 8 3. Amph: 4 v. 8 4. Fish: yes v. no	1. $z=-0.790$, p=0.003 2. $z=-6.528$, p<0.001 3. $z=-3.262$, p=0.003 4. $z=-1.722$, $p=0.085$
Lab, Resting time, GLMM	Full, df=257 Best-fit, df=262	Amph # * Fish None retained	6.25	NA	NA
Lab, Interaction time, GLMM	Full, df=171 Best-fit, df=173	Amph # * Fish Amph #	4.02	1. Amph: 4 v. 8	1. $z=7.657$, p<0.001

Figure Legends

Figure 1. Diagram of field microcosms used to test amphipod predation in the presence/absence of conspecifics and fish (number of animals in the diagram do not represent the experimental densities).

Figure 2. Per capita consumption (no. prey eaten divided by no. amphipods present) of *Gammarus pulex* increased in the presence of conspecifics (a) and decreased in the presence of fish (b) in the field. Lower overall per capita consumption was also observed at the Salterstown R. site than at the Bann R. site (c). Different letters indicate significant differences ($p < 0.05$), and error bars are ± 1 SE.

Figure 3. *Gammarus pulex* per capita consumption (no. prey eaten divided by no. amphipods present) tended to be higher in the presence of conspecifics, but was similar between greater amphipod densities (4 vs. 8) in the lab independent of fish cue presence (a). The handling time (time to fully consume a prey item once captured) of *G. pulex* declined with amphipod density in the presence of fish cue (b). Different letters indicate significant differences ($p < 0.05$), and error bars are ± 1 SE.

Figure 4. With increasing amphipod density, the time *Gammarus pulex* spent swimming declined and time spent interacting increased independent of fish cue. Amphipod density had no effect on resting time. Different letters indicate significant differences ($p < 0.05$), and error bars are ± 1 SE.

Figure 5. Dipteran larvae (*Simulium* spp.) pupated proportionally less in the presence of four amphipods than with one or none in the field (a), and were more aggregated by the end of

behavioural lab experiments when in the presence of eight amphipods compared to none (b). Different letters indicate significant differences ($p < 0.05$), and error bars are ± 1 SE.

Figure 6. Functional response of solitary *Gammarus pulex* on dipteran larvae. Dashed lines indicate comparable prey densities provided to solitary amphipods in the field (equivalent to 6 prey) and in the lab (10 prey) based on bottom surface area of the experimental containers. The mean curve (black line), 95% confidence intervals (shaded area) and raw data (gray points, jittered) are shown.

Draft

Figure 1.

Draft

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6