**Linking spawning ground extent to environmental factors -
patterns and dispersal during the egg phase of four North
Sea fishes.**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2016-0310.R2
Manuscript Type:	Article
Date Submitted by the Author:	03-Feb-2017
Complete List of Authors:	Höffle, Hannes; Queen's University Belfast, School of Biological Sciences van Damme, Cindy; Wageningen IMARES Fox, Clive; Scottish Association for Marine Science Lelièvre, Stéphanie; Ifremer, Laboratoire Ressources Halieutiques Loots, Christophe; Ifremer, Laboratoire Ressources Halieutiques Nash, Richard; Institute for Marine Research, Vaz, Sandrine; IFREMER, UMR MARBEC Wright, Peter; Marine Scotland, Munk, Peter; Technical University of Denmark, DTU Aqua, National Institute of Aquatic Resources, Section of Oceanography and Climate
Please Select from this Special Issues list if applicable:	39th Larval Fish Conference
Keyword:	Spawning distribution, DISPERSAL < General, Seismic surveys, CLIMATE CHANGE < General, Demersal fish

SCHOLARONE™
Manuscripts

1 **Running title:** Fish egg distribution in the North Sea

2

3 **Linking spawning ground extent to environmental factors - patterns and**
4 **dispersal during the egg phase of four North Sea fishes.**

5

6 HANNES HÖFFLE^{1*}, CINDY J. G. VAN DAMME², CLIVE FOX³, STÉPHANIE LELIÈVRE⁴,
7 CHRISTOPHE LOOTS⁴, RICHARD D. M. NASH⁵, SANDRINE VAZ⁶, PETER J. WRIGHT⁷
8 AND PETER MUNK¹

9

10 ¹DTU Aqua, National Institute of Aquatic Resources, Technical University of Denmark,
11 Kavalergården 6, DK-2920 Charlottenlund, DENMARK
12 ²IMARES, Institute for Marine Resources and Ecosystem Studies, Haringkade 1, 1976 CP,
13 IJmuiden, The Netherlands
14 ³Scottish Association for Marine Science, Scottish Marine Institute, Oban, PA37 1QA, UK
15 ⁴IFREMER, Laboratoire Ressources Halieutiques, 150 quai Gambetta, BP699, F-62321,
16 Boulogne sur Mer, France
17 ⁵Institute of Marine Research and Hjort Centre for Marine Ecosystem Dynamics, PO Box
18 1870 Nordnes, 5817 Bergen, Norway.
19 ⁶IFREMER, UMR MARBEC, Avenue Jean Monnet, CS 30171, F- 34203, Sète Cedex, France
20 ⁷Marine Scotland Science, Marine Laboratory, 375 Victoria Road, Aberdeen, Scotland, AB11
21 9DB

22

23 *Correspondence. Queen's University Belfast, School of Biological Sciences, Medical
24 Biology Centre, 97 Lisburn Road, Belfast BT9 7BL, Northern Ireland, Tel. +44 (0)28 9024
25 2464, h.hoffle@qub.ac.uk

Abstract

Previous studies have shown that four commercially important demersal species, namely cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), whiting (*Merlangius merlangus*) and plaice (*Pleuronectes platessa*) spawn in distinct areas across the North Sea. Based on two comprehensive ichthyoplankton surveys in 2004 and 2009, the present study uses Generalized Additive Mixed Models (GAMMs) to delimit these spawning grounds using the distribution of recently spawned eggs, investigates their relationship to specific environmental conditions and examines egg dispersal during their development. Results indicate that presence/absence of early stage eggs is more related to temporal and topographic variables, while egg densities are closely linked with hydrography. Egg distribution patterns were relatively consistent during development and only changed near hatching. Compared to historic observations, the location of the spawning grounds appeared stable on the broad scale but centres of egg abundance varied between the surveyed years. Potential effects of long-term climate change and anthropogenic short-term disturbances, such as seismic surveys, on fish reproduction are discussed, pointing out the demand for multi-species studies on these issues.

Keywords: Spawning distribution, dispersal, seismic surveys, climate change, demersal fish

47 Introduction

48 Annually visited spawning grounds are a common feature in the life cycle of many fish
49 populations (Daan et al. 1981; Heath et al. 1994), and have been demonstrated for a number
50 of commercially important fish species in the North Sea (e.g. Brander 1994; González-Irusta
51 and Wright 2016a; 2016b). Such stability in reproductive strategy is thought to have evolved
52 to ensure that, during the average lifetime enough offspring are placed into suitable
53 environments to ensure the population's survival. Suitable conditions for survival may be
54 found in the vicinity of the spawning ground or in nurseries with good connectivity to the
55 spawning grounds. Bakun (1996) coined the term 'ocean triads' for areas where zooplankton
56 is plentiful due to enrichment from nutrients, and where fish offspring are retained in a given
57 water mass. In the North Sea, such conditions could be found at tidal and salinity fronts
58 (Simpson et al. 1993). For fish spawning in late winter and spring salinity fronts may be more
59 important (Munk et al. 2002; Munk et al. 2009) as increased freshwater runoff during fall and
60 winter widens the region of freshwater influence (ROFI), particularly in the southern North
61 Sea (Simpson et al. 1991). Previous studies have suggested that the four commercially
62 important species cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), whiting
63 (*Merlangius merlangus*) and plaice (*Pleuronectes platessa*) all make use of recurrent
64 spawning grounds in the North Sea which may be linked to such hydrographic features.
65 In the southern North Sea cod spawns in relatively cool water (5-7°C) beyond the 32
66 isohaline (Aurich 1941). North Sea cod appears to have four distinct spawning grounds, the
67 Southern and German Bights, the southern and eastern flanks of Dogger Bank and off
68 Flamborough (Aurich 1941; Lelièvre et al. 2010; Daan 1978). The recurrent use of the latter
69 area could, however, not be confirmed from findings of a more recent survey in 2004 (Fox et
70 al. 2008). Further north, ripe and running cod have been observed on the Great Fisher and
71 Ling Banks as well as in the coastal areas (<100 m depth) of Scotland (Graham 1934;

72 González-Irusta and Wright 2016a). In the 2000s the main cod spawning grounds in the
73 northern North Sea were found off the north and east Scottish coasts and east of Shetland
74 towards Viking Bank (González-Irusta and Wright 2016a). Cod in the North Sea do not
75 appear to be a single stock as recent genetic evidence indicates that fish in the deeper
76 northern waters east of Shetland are genetically distinct from other North Sea populations
77 (Heath et al. 2014). In the northernmost North Sea, haddock spawning predominates in the
78 Atlantic influenced water of the deeper outer shelf, (González-Irusta and Wright 2016b;
79 Heath et al. 1994). Haddock rarely spawns in the southern North Sea, spawning has only been
80 extending as far south as the German Exclusive Economic Zone (EEZ) during a pulsed
81 recruitment (Aurich 1941). In the North Sea, haddock egg production appears highest outside
82 the Moray Firth and east of Orkney, though the eggs and larvae spawned to the west of
83 Scotland are advected into the North Sea (Heath and Gallego 1998) and the two regions are
84 treated as one stock for management purposes (ICES 2015).

85 North Sea whiting is divided into two genetically distinct northern and southern spawning
86 populations (Charrier et al. 2007). This species is among the most prolific spawners in the
87 North Sea, and it has both individually and collectively, an extended spawning period
88 (Aurich 1941). Compared to cod it appears to spawn in warmer water, having an optimum of
89 5–8°C, and eggs can be found in the southeastern inshore areas (Lelièvre et al. 2010). The
90 main centres for egg production of the southern population are on the western side, off
91 Flamborough, south of the Dogger Bank and in the Southern Bight, while the northern
92 population has its centre south of Shetland (Loots et al. 2011b).

93 North Sea plaice spawns in a large area of the southeastern North Sea, from the Southern
94 Bight northwards through the German Bight and along the coast of Jutland (Taylor et al.
95 2007). On the western side of the North Sea, plaice eggs have been found from Flamborough
96 up to the Moray Firth and further northeast off Shetland. The southwestern populations of

97 plaice appear to spawn mainly in the Southern Bight and partially in the eastern English
98 Channel (Houghton and Harding 1976; Hunter et al. 2004). Generally plaice spawns further
99 offshore than most other flatfish species, with the main egg production observed 50-200 km
100 offshore, in salinities >32 (Rijnsdorp 1989). Freshwater from the river plumes apparently has
101 a strong effect on spawning outside the core areas (Lelièvre et al. 2014). The location of the
102 spawning areas is thought to offer good connectivity to inshore, spatially limited, soft
103 sediment nursery grounds (Hufnagl et al. 2013).

104 Life-cycle closure requires a succession of suitable habitats and appropriate mechanisms for
105 transiting between them (Petitgas et al. 2013). There are concerns that predicted climate
106 change may not only affect reproduction directly but also affect early life stage habitats and
107 the connectivity between them (Collins et al. 2013). The changes would have a series of
108 direct and indirect effects, taking place on scales ranging from individuals to populations. It
109 has already been seen that North Sea fish distributions have responded to rising sea
110 temperatures (Perry et al. 2005) whilst changes in wind patterns and restricted transport of
111 *Calanus finmarchicus* into the North Sea have been implicated in cod recruitment (Rindorf
112 and Lewy 2006; Fromentin and Planque 1996). Under climate change the location of salinity
113 fronts, which may direct spawning, could also change (Collins et al. 2013; Perry et al. 2005).
114 Such changes may be especially detrimental for fish species on the southern edge of their
115 range, while other species may be able to extend into the North Sea (Dulvy et al. 2008;
116 Planque and Frédou 1999). Many North Sea stocks have also suffered decades of overfishing,
117 leading to low spawning stock biomasses (SSBs) and truncated age structures. Such stocks
118 may have reduced resilience against changes in the abiotic environment (Ottersen et al.
119 2006). Other human activities may put additional stress on spawners and their offspring,
120 particularly in a heavily industrialized area like the North Sea, containing nearly a thousand
121 active above- and sub-surface structures (De Groot 1996; OSPAR Commission 2014). Except

for specific accidents these structures, may, however, only be local disturbances and may even serve as habitat (Løkkeborg et al. 2002; Fujii 2015). Other disturbances might come from seismic exploration which nowadays uses airguns that cause direct physical damage in a much narrower zone than the explosives used in the past (5 m, McCauley et al. 2000). Effects on adults may rather be indirect, causing avoidance of approaching survey vessels. Since seismic surveys are not supposed to be undertaken during periods of fish spawning, or are supposed to use mitigation measures such as soft-starts (Smithies 2014), knowledge of the extent and timing of spawning is essential for proper Environmental Impact Assessment (EIA). Therefore, evaluating the overlap between intense seismic operations and spawning grounds is a normal requirement for EIAs undertaken before seismic operations commence. In 2004 and 2009, dedicated surveys and additional sampling during routine surveys were coordinated by ICES, affording the possibility to examine fish spawning using egg distribution data separated by 5 years. Here we use the egg distribution of the four most abundant species, cod, haddock, whiting and plaice in order to (i) construct statistical models, relating the spawning habitat inferred from stage I eggs, up to the onset of gastrulation, to the abiotic environment; (ii) examine how distribution patterns changed over time by constructing interpolated distribution maps and by statistically comparing the distribution of the successive egg development stages; (iii) evaluate to which extent spawning distribution overlaps with industrial activities and how these, in concert with climate change, may affect future spawning distribution and success.

Materials and methods

Survey design and field sampling

Sampling was carried out across the North Sea (Fig. 1a) by five participating countries, either during the annual International Bottom Trawl Survey for the first quarter (IBTS 1Q) or in dedicated surveys between the 16th February and 23rd of March 2004 and between 17th of January and 6th of March 2009 (Table 1).

The samples were taken with Gulf III/Nackthai (Gehring 1962; Nellen and Hempel 1969), Gulf VII (Nash et al. 1998), BONGO (Posgay and Marak 1980) and WP2 (Fraser 1968; Tranter 1968) nets, covering the entire North Sea from 51.5°N to 62°N. The Gulf and BONGO nets were deployed in double oblique hauls, while the WP2 was deployed in vertical hauls, in both cases down to five metres above sea bottom or a maximum of 100 m depth. Double oblique hauls lasted at least 10 minutes with multiple hauls ensuring the minimum deployment time in shallow water (ICES 2008). To determine cod-like eggs to species level, single eggs without oil globules and diameters of 1.1-1.75 mm (Fox et al. 2008) or 0.97–1.89 mm (Lelièvre et al. 2010) were fixed for later genetic analysis, either in 96% ethanol (Fox et al. 2008) or in a pigment preserving Battaglia-solution containing buffered 0.864% formalin, BHA (butylated hydroxyanisole), EDTA (ethylenediaminetetraacetic acid) and monopropylene glycol and ascorbic acid (Mastail and Battaglia 1978; Bigot 1979).

The locations of installations for oil extraction, seismic survey areas and the outlines of exclusive economic zones were extracted from public online databases (Fig. 1b, OSPAR Commission 2014; Ministerie van Economische Zaken 2015; Oljedirektoratet 2015). Depth and shoreline data were downloaded from NOAA databases (NOAA (NGDC) 2016a; NOAA (NGDC) 2016b).

Egg and larvae identification

Subsamples of at least 100 eggs in the desired size range (>1.16 mm, Russell 1976) were staged (Ryland et al. 1975; Geffen and Nash 2012) and visually identified to the lowest possible taxonomic level (Munk and Nielsen 2005; Russell 1976). Preserved cod-like eggs were identified either with TaqMan probes (Fox et al. 2008) or cytochrome *b* PCR–RFLP (Lelièvre et al. 2010). Accuracy of TaqMan probes is $>98\%$ for cod, haddock and whiting (Taylor et al. 2002) while cytochrome *b* PCR–RFLP confirmed 98% of visual identifications for late-stage cod eggs and 71% for late-stage whiting eggs (Lelièvre et al. 2010). The identified gadoid eggs were assigned to each species according to the proportion obtained from genetic analyses. For stations where this was not possible, the proportion in 2004 was derived from a multinomial Generalized Additive Model (GAM) over the whole survey area (Fox et al. 2008) and in 2009 from the average proportion for the other stations in the same ICES square. Counts of fish eggs were raised by the fraction of the whole sample examined and then converted to density (eggs m^{-2}), by dividing by the sampled volume and multiplying by the sampled depth.

Geostatistics

Hydrographic data were obtained from CTD hauls during the survey and augmented with data from the ICES oceanographic database (ICES 2009). After pre-screening for erroneous measurements, density anomalies σ_t ($\text{kg m}^{-3}-1000$) were calculated from temperature and salinity according to UNESCO standards (Millero and Poisson 1981) using the R package *oce* (Kelley 2013). To calculate the density gradient, an experimental variogram for σ_t was calculated in Surfer 10 (Golden Software 2011) and a linear theoretical variogram model fitted to the experimental variogram, using least square regression. Using ordinary kriging, the density anomaly was then interpolated onto a regular grid of $0.25^\circ \times 0.25^\circ$ and

subsequently smoothed to half that grid resolution. From the interpolated horizontal density anomaly the density gradient in $\text{g m}^{-3} \text{ n.mi.}^{-1}$ was calculated using the formula (Golden Software 1999):

$$(1) \quad \|\vec{g}\| = \sqrt{\left(\frac{Z_E - Z_W}{2\Delta x}\right)^2 + \left(\frac{Z_N - Z_S}{2\Delta y}\right)^2}$$

Where $\vec{g}$ is the density gradient, Z_E and Z_W are adjacent grid points in East-West direction; Z_N and Z_S are adjacent grid points in North-South direction, Δx and Δy the distances in either direction, respectively.

For visual representation, temperature and salinity were interpolated on a 1 n.mi. $\times$ 1 n.mi. grid using the R package gstat (Pebesma 2004). Analogous to the procedure for σ_t an experimental variogram was calculated and a Gaussian, exponential or spherical theoretical variogram was fitted to the experimental variogram using least-square regression and eye-fit. The chosen model was then used for the interpolation. Average temperature and salinity data for the 2000s was also interpolated following the same procedure. However, the value plotted here was the variation, to visually compare the distribution of stage I fish eggs with the variation of hydrography.

Similarly, data for each egg development stage were split into presence/absence and log-transformed non-zero abundance and then interpolated using an ordinary kriging procedure on the default grid in gstat (Pebesma 2004). The values for abundance were then weighted with the probability of occurrence on a cell-by-cell basis and the final maps created by kriging onto the same 1 n.mi. $\times$ 1 n.mi. grid that was used for hydrography.

Statistical comparisons of egg density distributions between stage I and the later stages were performed using the Syrjala test in the R package *ecspa* (de la Cruz 2008). The Syrjala test computes a modified two-sample Cramér–von Mises test for a difference between the cumulative distribution functions of two populations (Syrjala 1996). As the test is sensitive to where the coordinate system has its origin, it is performed once for each of the four corners of the hypothetical rectangle encompassing all sampling sites and then averaged. Although the test is insensitive to sample size, each sample was transformed to ‰ of the sum of the respective stage before conducting the test.

Modelling spatial autocorrelation

Spatial autocorrelation and average patch size were modelled with Moran's eigenvector maps (MEMs) using the R package *pcnm* (Legendre et al. 2013). A truncated distance matrix was computed by converting all distances farther than the longest distance between immediate neighbours to four times that value (Borcard and Legendre 2002). Since there are no interactions in the population of fish eggs, only the MEMs with positive eigenvalues were retained. Before running a global test on all MEMs with positive eigenvalues the hellinger-transformed abundances were detrended to remove any spatial trend. Individual Moran's Eigenvectors (MEVs) were selected by regression-based forward selection with the R package *packfor* (Blanchet et al. 2008; Dray et al. 2013) using p-value and adjusted r^2 as stopping criteria. Similar to Loots et al (2010a) the threshold p-value was set to a low 0.002, to avoid building too complex spatial models. The spatial scale was then determined by first computing a Gaussian variogram for each MEM (Bellier et al. 2007). Then the selected MEMs' eigenvalues were plotted against these ranges and subdivided into sub-models for a fine, medium, medium-broad and broad scale.

Explanatory models

To examine the relationships between the distributions of stage I eggs and the abiotic environment, data from both years were used to construct a generalized additive mixed model (GAMM) with the R package gamm4 (Wood and Scheipl 2014; Wood 2006). As for the kriged distribution maps, separate models were constructed for presence/absence and non-zero abundance, resulting in a hurdle-model. Potential independent variables were temporal (year, day of the year), bathygraphic (bottom depth), hydrographic (temperature, salinity, and σ_t -gradient; from the surface and bottom) and the selected MEVs. These variables were examined for collinearity with pair plots and variance inflation factors (AED, Zuur 2010), entailing the exclusion of the year. Hydrographic values were taken from the CTD casts taken at the locations of the individual samples or, if not available or erroneous, were extracted from the interpolated surfaces. The measurements from 5 m depth and the bottom were strongly correlated. Instead of excluding one or the other, separate models using either surface or bottom measurements were constructed and compared. MEVs were treated as random effects, while the day of the year (=year day), bottom depth and the hydrographic covariates were the fixed-effects. Using either presence/absence in a binomial model with a logit-link or log-transformed non-zero egg densities in a Gaussian model with identity-link as response, the general form of the GAMM was (Wood 2006; Höffle et al. 2014):

$$(2) \quad y_i = \beta + f_1(x_{1i}) + f_2(x_{2i}) + \dots + f_n(x_{ni}) + Z_i b + \varepsilon_i$$

y_i being the response, β the intercept, f_i the smooth functions for the fixed effects x_k , b a random effects coefficient vector from the matrix Z_i and the residual error vector ε_i .

The risk of over-fitting the individual models was reduced by restricting the degrees of freedom to four and by using the Bayesian Information Criterion (BIC, Schwarz 1978) as an indicator of improvement during the backward selection of the optimal models. Initial models contained hydrographic measurements from either the surface (5 m) or the bottom depth. Following the optimization of these models the decisions between either surface or bottom models were based on BIC, Akaike's Information Criterion (AIC, Akaike 1974) and in case of disagreement between the two on the predictive power of the models. Formal assessment of the goodness of fit, since not possible from the GAMM itself (Wood 2006; Zuur et al. 2009), was done by estimating the deviance explained from a generalized additive model (GAM) using the same fixed-effects covariates (Gilman et al. 2012). Post-hoc testing included an evaluation of the random effects' contribution, using the residuals from the GAM as response (Wood 2006). Hereby, linear fixed-effects and mixed models are compared, by using a constant parameter and in the mixed model the random effects. The two models are then compared by AIC, BIC and the log-likelihood ratio test.

The effect of the random covariates on the spatial autocorrelation was evaluated by Moran's I correlograms on the models' residuals, setting a separation distance of 0.7° for non-zero egg densities of cod and plaice, 2.1° for all other models, corresponding to the maximum distance between nearest neighbours (Diniz-Filho et al. 2003; Borcard et al. 2011). The number of lags to cover the entire study area was then 15 and 5, respectively. P -values were adjusted using Holm's correction (Borcard et al. 2011).

The performance of individual fixed covariates was evaluated using three different indicators. Firstly, by dropping each covariate individually and then calculating the log-likelihood ratio between the reduced model and the model selected in the fitting process. Secondly, by computing the ROC AUC (Receiver Operating Characteristic Area Under the Curve) for binomial and the r^2 between response values predicted from individual covariates and the

observations for Gaussian models using the R package MuMIn (Barton 2013). Lastly, partial linear regression and adjusted r^2 were used to partition the explained variation explained by the individual covariates using the R package vegan (Oksanen et al. 2013; Peres-Neto et al. 2006).

Results

Hydrography

In both years hydrography was characterized by a pool of relatively warm ($>7^{\circ}\text{C}$) and saline (>34.5) water in the northern and central North Sea and cooler, fresher water along the margins, which was more pronounced in the East (Fig. 2). Salinity (Figs 2b, d) as well as density gradients (max. $4.3 \text{ g m}^{-3} \text{ n.mi.}^{-1}$ off Norway) were steeper on the eastern side of the North Sea. Surface hydrography was more structured than at the bottom. The main difference between 2004 (Fig. S1) and 2009 (Fig. S2) was that in the southern North Sea water $\leq 6^{\circ}\text{C}$ extended further westwards in 2009. The water in the German Bight was more than 1°C colder than in 2004 and salinity gradients from coastal to offshore waters were shallower, particularly north of the Frisian Islands where a prominent plume of relatively fresh (≤ 34.25) water extended far offshore. North of 57°N the situation was the opposite with the salinity gradient steeper in 2009 (Fig. S2) than 2004 (Fig. S1).

Spatial structuring

The threshold for truncating the distance matrix was 2° . Of the MEMs, 355 were found to have positive eigenvalues. Two of these were significant for cod, four each for haddock and plaice and one for whiting. When plotting eigenvalues against the variogram range, there were four distinct groups (Fig. 3), on ranges of 0-40 n.mi. (fine), 40-75 n.mi. (medium), 75-100 n.mi. (medium-broad) and above 100 n.mi. (broad).

310

311 **Egg distribution**

312 Stage I egg densities on the scale of the entire North Sea were in both years in the same
313 general areas (Fig. 4). High egg densities of cod (Fig. 4a) and plaice (Fig. 4p) were found in
314 the Southeast, along the continental coast, extending as far north as Viking Bank. Whiting
315 and haddock were more common in the western part. The centre of egg densities for haddock
316 was limited to the Northwest (Fig. 4f), two patches on either side of the Fladen Ground and a
317 third, smaller, patch outside the Firth of Forth. Peak densities of whiting eggs also occurred in
318 three locations but they were farther apart. One centre of abundance was east of the Shetland
319 Isles (Fig. 4k) around the Forties, the second, larger, area was between Flamborough and the
320 Dogger Bank and a third area was in the Southern Bight.

321 The overall pattern in the development over time was one of decline and dispersal. For cod
322 and whiting (Figs 4a-e, 4k-o) this process appeared to be quicker compared with haddock and
323 plaice with a sharp drop in egg densities between stages I and II and a lower rate of decline
324 thereafter. For haddock and plaice the peak densities did not decline as quickly (Figs 4f-j, 4p-
325 t), densities of the same magnitude as in stage I were still observed in stages IV and V (Figs
326 4i, 4t). However, the area covered by high densities shrank and in the case of haddock
327 divided into several smaller well-separated patches.

328 The main differences between the years (Figs S3, S4) were that the centres of abundance for
329 cod and plaice were further north and offshore in 2004 than in 2009 and that the rate of
330 dispersal differed between the years for each individual species. In 2004 high cod and plaice
331 densities covered a larger area (Figs S3a-e, S3p-t), from the transition area between the
332 Bights up to the southern edge of the Norwegian trench. For cod in 2009 this area was
333 smaller, more inshore, off the Frisian Islands, while egg densities were higher in the northern
334 patch around Viking Bank (Fig. S4a-e). For plaice the overall location was similarly changed,

extending along an east-west axis rather than a north-south axis, but the extent of the area was similar to 2004. Peak densities of cod eggs and the area covered with such densities, quickly declined in 2009, while in 2004 high densities were still found in stage III (Fig. S3c). There was no such difference for plaice, although in later stages (Figs S3s and S3t) the location of peak densities was similar to 2009. High egg densities of haddock covered much larger areas in 2004 (Figs S3f-j) than in 2009 (Figs S4f-j) and the distribution pattern did not disperse as quickly and completely. Whiting egg densities were generally lower in 2009 than in 2004 (Fig. S4k-o) but there was little difference in how quickly the eggs dispersed. However, the overall pattern was different (Fig. S3k-o) with high densities in the Southern Bight in 2009 while in the earlier year the centres of abundance were off Flamborough and east of Shetland.

The Syrjala-test did not fully confirm the patterns suggested by the interpolated maps (Table 2). The distribution patterns of stage IV and V were more often dissimilar from stage I than those of earlier stages, but that pattern was not consistent across species or years. Although stage IV was significantly dissimilar for all species when comparing mean distributions, it never was for whiting in any of the single years. Similarly, the distribution of stage V was significantly different for all species in 2004, but not in 2009 or for mean distributions. Stage III never exhibited significant dissimilarity to stage I, although there were three cases of significantly different distributions between stages I and II.

Explanatory models

The overall trends in the explanatory models were twofold. Firstly, the best fitting models for non-zero egg density tended to be more complex than those for the probability of occurrence (Table 3, Figs 5, 6). Secondly, although the difference was marginal, three out of four models exhibited a better fit when using hydrographic measurements at the bottom for the probability

of occurrence, while for egg densities it was the opposite. Deviations explained by the corresponding GAMs ranged between 15% for the density of stage I cod eggs and 57% for the presence/absence of haddock eggs. The models for these two species also consistently retained the least fixed-effects covariates (3) for both cod models and the most (5) for both haddock models (Table 3).

Bottom depth and year day were the only fixed-effects covariates retained in every model. They also exhibited similarly shaped relationships for three of the four species (Figs 5a, b). Cod, Haddock and Whiting exhibited an increasing trend towards the end of the period. Plaice differed from the other species, exhibiting a dome shaped relationship between days 30 and 70. The relationships of cod and plaice egg densities with the year day were of similar shape and peaked earlier than their probabilities of occurrence (days 45 and 35; Fig. 6a). Whiting egg densities exhibited an early and a late peak around days 20 and 80, respectively. Haddock egg densities were linearly and positively related to the year day.

Probabilities of occurrence for cod and plaice exhibited overall decreasing trends of similar shape with peak probabilities at ca. 50 m (Fig. 5b). Haddock exhibited a similar pattern, but shifted towards greater depths, with a peak at around 100 m and an increasing trend towards the end of the observed depth range. The probability of finding stage I eggs of whiting declined with increasing depth, becoming zero at ≥ 150 m (Fig. 5b). Relationships between egg densities and bottom depth were of similar shape for cod, whiting and plaice, with peaks at shallower depth than peak probabilities of occurrence (Fig. 6b). Haddock egg densities exhibited a Gaussian distribution around 100 m.

Other than for time and depth, the relationships between the probability of presence and the hydrographic covariates were more species specific with no consistent patterns between any two species. Temperature did not contribute to the model for the occurrence of cod eggs and exhibited negative and positive trends for plaice and whiting, respectively (Fig. 5c). For

haddock the probability of finding eggs was high across the temperature range, except for a local decline around 6°C. Unlike for probability of occurrence, the smooth terms for egg densities related to hydrographic measurements exhibited similarities between two or more species. Haddock consistently differed from the other species, exhibiting linear or near-linear increases in egg densities in relation to year day, salinity and density gradient. Temperature contributed to all models for egg density with the peak occurring as low as ca. 5°C for plaice and as high as 9°C for whiting. The smooth terms for cod and plaice were similar in shape and both egg densities peaked at nearly the same temperatures. Observations at either end of the observed temperature range were few and therefore have wide confidence intervals. Salinity contributed to the model for the probability of occurrence of cod eggs (Fig. 5d) but did not contribute to the model for egg densities (Fig. 6d). Cod probability of occurrence exhibited a weakly declining trend towards higher surface salinities, while haddock was strongly positive related to bottom salinity. Salinity was only retained in the models for egg densities of whiting and plaice, exhibiting similar shapes of the smooth terms, with peaks at the lower end of the range and at salinities of 34. Haddock eggs were only observed in a narrow salinity range and the trends of probability of occurrence as well as egg densities were strongly positive. The density gradient was retained in the presence/absence models for haddock and whiting (Fig. 5e) and the egg density models for these two species as well as for plaice (Fig. 6e). Egg densities of all three species were linearly related to density gradients. Across all species the relationships to the MEVs describing spatial autocorrelation were linear or dome shaped (Figs S5 – S8). Spatial autocorrelation in the residuals was by a factor of 1.5 to 7.5 lower than in the data used for fitting. However, on the first lag it remained significant (Table 3). Predictive power of the selected models was highest for the presence/absence of haddock, the egg density model of whiting and lowest for both models for cod (Table 4). Individual

covariates with high predictive power strongly affected the log-likelihood ratios as well (Table 5). Concerning explained variation this consistency in performance of individual covariates was limited to cod (Table 6). The variation explained by the fixed-effects covariates was lowest for cod and highest for haddock. The highest percentage of explained variation in presence/absence was related to bottom depth for plaice and year day for all other species. For the egg densities the highest percentage of explained variation was related to temperature for cod, salinity for haddock, and density gradient for whiting and plaice, respectively. This increased importance of hydrographic variables in the egg density models was not reflected in the predictive power or the log-likelihood ratios where these covariates were usually weaker than either year day or bottom depth.

The comparison between the GAMMs and their corresponding GAMs using only the fixed-effects covariates revealed that the MEVs only contributed significantly to the models for whiting (Table 7). However, even for whiting the GAM was more parsimonious, as the AIC and BIC was for all GAMs lower.

Comparison with decadal hydrography and oil installations

Averaged over the 2000s the variation in bottom temperature exhibited a range of interannual variability between $<1^{\circ}\text{C}$ and $>1.6^{\circ}\text{C}$ (Fig. 7a). The lower part of this range was found in large areas of the central and southern North Sea, while the rest of the North Sea proper exhibited intermediate variation. High variability was primarily indicated on the northern margins of the North Sea and between the Shetland and Orkney Isles as far south as the Moray Firth. High egg densities across all four species were primarily found at the boundaries between the areas of low, intermediate and high variability. The mean density of stage I eggs across all species, compared to the distribution of oil extraction and seismic survey areas in the eastern North Sea showed that at this time of the year areas of high egg

densities primarily overlapped with the oil industry in the southeastern North Sea in the Dutch and German EEZs (Fig. 7b).

Discussion

The results of the present study indicate that presence/absence and non-zero abundance of cod, haddock, whiting and plaice eggs are primarily governed by different environmental variables. Presence/absence was mainly linked with the day of the year as well as bottom depth whilst non-zero abundance was linked to one or more of the hydrographic covariates. Overall, the spawning of cod and plaice appears more common in the southeastern North Sea, while spawning of haddock and whiting were concentrated in the western areas, and in the case of haddock further limited to the northern North Sea. The overall spawning areas seem to have remained stable, compared to historical observations. However, the, smaller, actual spawning grounds used within those areas varied between the two years surveyed depending on prevailing conditions.

Egg distribution

Where spawning locations persist over many years their use must allow sufficient offspring to survive to sustain the population size (see Hufnagl et al. 2013). Our observations of geographically relatively stable spawning areas contrast findings for other species such as, e.g. mackerel (*Scomber scombrus*, Beare and Reid 2002). However, even for mackerel the spawning occurs mainly along the shelf-break and this location may be linked to patterns in secondary production driven by internal waves (Sharples et al. 2009). Areas of the North Sea where high abundances of cod, haddock, whiting and plaice eggs were found are consistent with earlier studies (Coull et al. 1998; Heath et al. 1994). However, their extent at higher population levels might be greater and at the present time the maximal extent of the spawning

areas may be better indicated by egg occurrence (Brander 1994). The overall better fit for the presence/absence models when including hydrographic measurements from the bottom rather than from the surface may be due to the generally demersal behaviour of the adult fish. On the other hand the better fits for egg densities using surface hydrography may be based in the distribution of pelagic eggs in the upper part of the water column (Coombs et al. 1990; Fritsch 2005). After rising to the surface layer, eggs will be advected from the spawning site. However, in the observed temperature range the eggs develop to the end of stage I within a few days (Thompson and Riley 1981; Geffen and Nash 2012). At common winter current velocities (Sündermann and Pohlmann 2011), dispersal during that time should have been on about the same scale as the survey. This is supported by recent studies on the distribution of mature adults which largely corroborate the finding from the egg surveys (González-Irusta and Wright 2016a; González-Irusta and Wright 2016b; Loots et al. 2011b). The only major difference was that the northern core spawning grounds of whiting extended further south than was observed in the egg surveys (Loots et al. 2011b). Therefore, the distribution of stage I eggs observed in 2004 and 2009 may be considered a reliable source of information on spawning distribution.

Relationship with the abiotic environment

Compared to Otto and Zimmermann (1990) and the studies underlying that review (Goedecke et al. 1967; Tomczak and Goedecke 1962), the overall pattern of hydrography in the North Sea in 2004 and 2009 was not much changed since the 1960s. Patterns were also similar between the sampled years with the main differences in temperature and salinity occurring south of 57°N. A major difference between the two survey years were the higher egg densities of whiting in the Southern Bight in 2009. The difference in egg densities in the Southern Bight correlates with the hydrographic differences, as whiting may have spawned

485 across the boundary between the cool, low salinity coastal water and the warmer water from
486 the South, corresponding to the species preferences (Aurich 1941). In the southern North Sea
487 a comparable data set is available from a series of surveys using the Continuous Underway
488 Fish Egg Sampler (CUFES, Lelièvre 2010). In this study high egg densities of all species
489 were found in the Southern Bight in 2006 and 2008, but to a lesser extent in 2007 (Lelièvre et
490 al. 2012). In 2006 low salinity was not restricted to the German Bight and the interface
491 between fresh and saline water was further offshore, resembling the situation in 2009. From
492 the comparison of the years we deduce that a more seaward position of the boundary between
493 freshwater influenced areas and marine water may attract more spawning fish to the area.
494 The present observations on cod egg distribution are consistent with Aurich (1941) and
495 Brander (1994). The latter author identified the western-central North Sea (2°W - 2°E and
496 52.3 - 56°N) as a minor contributor to the spawning of cod, and the German and Southern
497 Bights as major contributors in alternating periods, 1948-1956 and 1958-1972, respectively.
498 In 2004 and 2009, the distribution of cod eggs in the Southern Bight was limited to a few
499 patches, while in the German Bight cod eggs were widespread, similar to the pattern
500 described by Brander (1994).

501 Densities of cod eggs peaked in areas shallower than 100 m, corresponding to results of
502 previous ichthyoplankton studies (Munk et al. 2002) and in line with González-Irusta and
503 Wright (2016a) who reported that the abundance of cod in spawning condition declined at
504 more than 125 m water depth (González-Irusta and Wright 2016a). The temperature (ca. 4°C)
505 at these locations corresponded to the distribution of spawners which declined when
506 temperatures increased to above 5°C (González-Irusta and Wright 2016a). Cod larvae tend to
507 aggregate at or slightly inshore of a frontal system (Munk et al. 1999) but for the
508 presence/absence and abundance of cod eggs the gradient in water density did not appear to
509 be an important covariate in our analyses. Hence, the aggregation around fronts may only

510 become established by selective processes or behaviours taking place in the larval stage.
511 Patterns of egg production were not entirely consistent with predictions based on the catches
512 of mature female cod in research surveys (Fox et al., 2008). This was particularly the case in
513 the northern North Sea where the bulk of the cod stock is presently located and where high
514 egg production would be expected. Cod in the northern population centred around Viking
515 Bank spawn several weeks later than those in the southern North Sea (González-Irusta and
516 Wright 2016a), which may call for changes in the timing of future ichthyoplankton surveys.
517 The greatest spawning concentrations of haddock have historically been reported from the
518 plateau between the Orkney-Shetland Isles and Norway in water deeper than 100 m
519 (Thompson 1928). A statistical model developed for spawning haddock west of Scotland also
520 indicated a decline at greater depths, which corroborates the egg-based model in the present
521 study (González-Irusta and Wright 2016b). Spawning aggregations have also been reported
522 east of the Moray Firth, at the Buchan and Farne deeps, the Gut and the Fisher Banks. Saville
523 (1959) found haddock eggs to be widely distributed across the northern North Sea from the
524 Scottish coast to the Norwegian deeps except for the Moray Firth and an area around the
525 Fladen Ground. The low egg densities around the Fladen Ground appear a consistent feature,
526 as they were also noted by Damas (1909), González-Irusta and Wright (2016b) and in the
527 present study. The extent of haddock eggs found by Heath et al (1994) in 1992 was
528 comparable with that of Saville's study, although eggs were concentrated further north and
529 west. Heath and Gallego's (1998) study estimated high egg production from mature fish for
530 the Moray Firth, the Scottish east coast and Shetland Isles, which is reflected in the results of
531 the two surveys. However, their model did not predict high abundances to the immediate east
532 of the Shetland Isles, which were present in 2004, 2009 and earlier egg surveys. High
533 abundances of haddock eggs south of the Moray Firth were only found in 2004. In earlier
534 plankton surveys Heath et al. (1994) found a spatial segregation of late stage cod and

haddock eggs, in colder North Sea water and on the warmer outer shelf, respectively. These findings support the suggested preferences for different zones of a frontal system between these gadoids (Munk et al. 1999). More common occurrence and higher abundance of haddock eggs on the offshore side of a frontal system, was also seen in the relationships to the environmental covariates. Observations of haddock eggs were limited to a narrow part of the species' salinity range (30-36, Laurence and Rogers 1976). Within the observed range, the probability of occurrence as well as egg densities increased. Likewise the warmer offshore water, where a local peak was found at 7°C in the present study, falls within the species' temperature range (4-10°C, Laurence and Rogers 1976). Similarly, spawning haddock reached peak probability of presence and abundance at 7°C and exhibited peaks at salinities >34 and >35, respectively (González-Irusta and Wright 2016b). Density gradients in the northern North Sea were lower than in the South. Nevertheless, they positively affected the abundance and to a lesser extent the occurrence of haddock eggs. This may be related to the observed high numbers of eggs on the oceanic side of the front between the Atlantic and the North Sea water in the northwestern North Sea (Heath et al. 1994).

It has been suggested that the whiting population consist of two stocks in the North Sea, one east of England and one east of the Shetland Isles (Charrier et al. 2007). Such separation is consistent with our findings. The preferred habitat of adult whiting can be characterized by shallow depths and relatively high winter temperatures of 6-9°C (Damas 1909). This preference was apparently reflected in the higher probability to find whiting eggs as well as higher egg densities in the upper temperature range. Egg abundance peaked not only at temperatures >7°C but also at sub-optimal temperatures of 4°C, but still within the thermal range for egg survival (Povoa et al. 2011). Probability of occurrence and abundance peaked in the 20-40 m range identified as preferred depth range for spawning by Aurich (1941). The likelihood to find whiting eggs increased at high density gradients, while the egg densities

declined, partially corroborating the findings of Munk et al (1999), who found that whiting eggs were uncommon at central parts of fronts. Whiting has a long spawning period, from February to June with increasingly later spawning at higher latitudes (Hislop 1984). Hence, the earlier sampling period in 2009 may partially explain the difference to 2004, particularly in the northern spawning grounds east off Shetland. In 2009 spawning east of Shetland may not have begun at the time of the survey. However, in 2004 stage IV and V eggs were abundant in the North, while early stages occurred more in the southern North Sea, contrary to the observations by Hislop (1984) and Aurich (1941). As individual whiting spawn many batches over a period between 1.5 and 2 months into late spring (Hislop 1984), the 2004 survey may have sampled the early stages of a later spawning batch in the South, while in the North the previous batch was sampled in the later stages.

The relatively homogenous distribution of plaice eggs in the German Bight and along the Danish coast, particularly in 2004, is a good representation of this species' large spawning area in the southern North Sea and English Channel (Aurich 1941; Metcalfe 2006). Moray Firth, Firth of Forth and the area off Flamborough Head are further suitable for plaice spawning (Loots et al. 2010b) and some eggs were found there, in earlier surveys (Aurich 1941) and in the present study. To develop successfully, plaice eggs need a temperature range of 2-8°C (Wegner et al. 2003), which was reflected in the above average abundance of eggs at 4-6°C. The probability of finding plaice eggs was similarly highest at the lowest observed temperatures and declined linearly over the observed range, indicating that within the species' thermal range the lower part is preferred (Wegner et al. 2003). The high densities of plaice eggs at salinities ≥ 34 corroborated earlier studies indicating high salinities as optimal for plaice eggs (Aurich 1941; Cushing 1990). The linear relationship between egg densities and density gradient was negative and had a similar slope to that for whiting. This result

584 supports earlier studies indicating more offshore spawning grounds than other flatfish
585 (Lelièvre et al. 2014).

586 Overall, egg densities declined quickly between stages I and II and at a lesser rate towards
587 later development stages. However, distribution patterns remained largely the same up to
588 stage IV. Therefore, one can assume that the decline was primarily driven by the high
589 mortality of fish eggs (Bunn et al. 2000), and that mortality rates were similar across the
590 North Sea. Contrary, dispersal may be rather slow. In the southern North Sea, eggs are
591 advected at about 5 km d⁻¹ (Brander 1994), resulting in a transport of around a 100 km from
592 spawning to hatching, based on temperature dependent egg development (Thompson and
593 Riley 1981; Geffen et al. 2006). This slow dispersal may explain why distribution patterns
594 remained similar until the later stages. As indicated by *p*-values in the Syrjala-test,
595 distribution patterns of stages I and III were often more similar than those of stage I and II.
596 This may be due to the difference in duration of development stages. It is less likely to find
597 stage II eggs during a survey, as it is comparatively short (Fox et al. 2003; Geffen et al.
598 2006).

599 Areas with high densities of haddock eggs, similar to densities in stage I, were maintained
600 longer than for cod and whiting. Partially this might be due to a larger contiguous area with
601 high densities to start with. Another factor may be that the distribution of adult haddock in
602 the first quarter changes little (Hedger et al. 2004), suggesting that predators were satiated by
603 a continuous supply of freshly spawned eggs rather than feeding on the less numerous later
604 stages. A similar mechanism may be at work in case of plaice, which also started out with a
605 large area of high densities that shrank but did not disappear over time.

606

607

608

609 Individual covariates

610 The greater importance of the hydrographic variables concerning explained variation, and to
611 a lesser extent predictive power and the log-likelihood ratios, in the egg density models
612 contrasts with the findings of earlier studies. In the present study the year day was the most
613 influential covariate in determining presence/absence. However, earlier studies indicated that
614 environmental covariates are more important for defining the boundaries of a population's
615 habitat, rather than the distribution within (Loots et al. 2011a; Loots et al. 2010a). However,
616 it is not without precedent. Although they used a different measure, González-Irusta and
617 Wright (2016a) came to a similar result for the abundance of North Sea cod in spawning
618 condition. In their study, temporal covariates were likewise the most important ones,
619 followed by salinity as the most important hydrographic covariate. Still, the importance of the
620 year day must be interpreted with care, due to the time series being short and the survey
621 periods differing by about three weeks. Longer time series with several samples on the same
622 year day may reduce its influence (e.g. Höffle et al. 2014; Lelièvre et al. 2014). The
623 distribution of populations on the edge of their environmental range may experience a
624 stronger influence of the environment, which may apply to cod and haddock. Haddock is
625 environmentally restricted to the northern North Sea and while distributed across the North
626 Sea, cod still experiences a strong influence of temperature, because the North Sea stock is
627 one of the most southernmost ones (Hedger et al. 2004; Drinkwater 2005).

628 MEMs are a powerful tool to deal with spatial autocorrelation. However, the additional
629 covariates may be penalized in model selection (Loots et al. 2011a). This may be one reason
630 why only the models for whiting exhibited a significant difference in the log-likelihood ratio
631 test, as they only contained a single MEV on the broad scale. Broad scale structures were also
632 those with the most discernible patterns when plotting observations against predictions,

indicating that these structures were likely more important for characterising the spatial distribution of spawning than those on smaller scales.

Long-term change and short-term disturbances

The four species in the present study may be affected in different ways by long-term changes in environmental conditions of the North Sea, potentially compounded by short-term disturbances (Collins et al. 2013; Petitgas et al. 2013). The debate whether recent changes in the distribution of some North Sea species are related to climate change or to changes in local populations is still ongoing (Neat and Righton 2007; Holmes et al. 2014). The present study indicates that effects would be species specific, with cod and haddock experiencing stronger direct effects of rising temperatures (Hedger et al. 2004; Drinkwater 2005). However, Petitgas et al. (2013) highlighted the potential impact of indirect effects on species with specific habitat requirements for spawning grounds or nurseries as well as on the connectivity between these areas. Plaice appears to be particularly vulnerable as spawning grounds are offshore and the, narrowly defined, nursery areas inshore (Hufnagl et al. 2013). North Sea plaice is one of the more southern populations and suitable spawning grounds and nurseries may decrease in size (Petitgas et al. 2013). Connectivity may be affected by accelerated development and changing current patterns through changes in wind and runoff patterns. For the period 1975 to 2006 Hufnagl et al. (2013) observed a decrease in connectivity related to an increase in temperature, for the spawning grounds in the German Bight and in the English Channel, while connectivity for other areas increased. For cod, connectivity may be affected in the opposite way, as climate change may entail more natal retention (Heath et al. 2008). However, other studies suggested a displacement to the North (Rindorf and Lewy 2006). Whiting may also be affected by changes in salinity and subsequently stronger density gradients in its, previously more homogenous, spawning areas (Loots et al. 2011a).

While the effects of changes in climate are long-term, short-term disturbances may interact with them. Noise producing activities, such as seismic surveys as well as construction work and the ensuing avoidance reactions, may lead to delays in spawning or avoidance of optimal spawning grounds and therefore interfere with successful connectivity between spawning and nursery areas. Together with minor injuries and their effects on long-term survival such indirect effects of noise pollution are understudied and call for further research (Wahlberg and Westerberg 2005; Payne 2004). Such impacts are considered in the EIA that must be undertaken before seismic surveys are conducted. Where potential impacts are judged serious then mitigation measures must be taken. Mitigation can include restricting the timing of work so it is outside of the spawning season or other measures such as soft-starts for pile-driving and air gun surveys. It is therefore important that up-to-date information is available regarding spawning locations and timing so that offshore exploration and construction activities can be properly controlled. Of the four species in the present study, cod and plaice overlap with areas for seismic surveys, whether for oil exploration or construction work, in the southeastern North Sea (Ministerie van Economische Zaken 2015). Cod may be vulnerable due to its SSB being below B_{trigger} , the biomass triggering management actions, while plaice is vulnerable due to its dependence on successful connectivity between early life stages.

Conclusion

Spawning areas appear to be defined by few, time-invariant variables, while centres of abundance within these areas are influenced by the interannual variability of the hydrographic conditions. Explanatory power of the statistical models varied between 15% explained deviation for cod egg density and 57% for haddock egg presence/absence. Models for cod spawning distribution, whether they use eggs or ripe and running adults as response, appear

to be difficult to define as even a far larger number of covariates barely yielded double the explained deviation (González-Irusta and Wright 2016a). Considering that the cod stock is currently the most vulnerable one in the North Sea, the investigation of further drivers governing the spawning distribution should concentrate on this species. Performance of individual covariates indicates that the occurrence of spawning in a given location of the North Sea is more dependent on covariates that do not vary over time, as the most important covariates for these models were year day and bottom depth. Contrary, egg densities were to a greater extent governed by covariates that change over time. This suggests that changes in the ocean climate, not exceeding the fishes' physiological limits, may cause centres of abundance to shift within spawning areas that remain the same. Such centres of abundance should be avoided by potentially damaging activities, and aggregate maps of spawning habitats and egg distributions, such as in the present study and Lelièvre et al. (2014), would be valuable tools in the management of marine areas, protecting the successful reproduction of fishes.

Supplements

Surface and bottom hydrography as well as the distribution of development stages for individual years are available as online supplements (Figs S1-S4). Supplementary figures presenting predicted versus observed values illustrate the information on predictive power of individual covariates given in Table 4 (Figs S5-S8).

Acknowledgements

For the coordination of the surveys, to the Planning Group on North Sea Egg Surveys (PGEGBS), ICES. Captains, crews and scientific staff of the participating research vessels from Denmark, France, Germany, the Netherlands, Norway, Scotland and England. Stéphanie

708 Lelièvre, Christophe Loots and Sandrine Vaz received additional financial support from the
709 INTERREG IVA France (Channel) – England cross-border European cooperation
710 programme, cofinanced by the European Regional Development Fund as part of the Channel
711 integrated Approach for marine Resource Management (CHARM) Phase 3 project. Hannes
712 Höffle was funded by the projects; Sustainable fisheries, climate change and the North Sea
713 ecosystem (SUNFISH, Danish Research Council Project 38135) and High-Resolution and
714 Dynamic Mapping of North Sea Spawning and Larvae Areas (KINO, Statoil and IMR,
715 project 14507). Richard Nash was supported by IMR's North Sea Programme, Early life
716 history dynamics of North Sea fishes (14387) and High-Resolution and Dynamic Mapping of
717 North Sea Spawning and Larvae Areas (KINO, Statoil and IMR, project 14507). The authors
718 would also like to thank the editor and two anonymous reviewers for their helpful
719 suggestions.

720 **References**

- 721 Akaike, H. 1974. A New Look at the Statistical Model Identification. IEEE Trans. Autom. Control
722 **AC19**: 716-723.
- 723 Aurich, H.J. 1941. Die Verbreitung der pelagischen Fischbrut in der südlichen Nordsee während der
724 Frühjahrsfahrten, 1926-1937. DBB Z. Lebensr. **2**: 183-225.
- 725 Bakun, A. 1996. Patterns in the ocean: Ocean processes and marine population dynamics. University
726 of California, San Diego.
- 727 Barton, K. 2013. MuMIn: Multi-model inference. R package version 1.9.5.
- 728 Beare, D.J., and Reid, D.G. 2002. Investigating spatio-temporal change in spawning activity by
729 Atlantic mackerel between 1977 and 1998 using generalized additive models. ICES J. Mar. Sci. **59**:
730 711-724.
- 731 Bellier, E., Monestiez, P., Durbec, J.P., and Candau, J.N. 2007. Identifying spatial relationships at
732 multiple scales: principal coordinates of neighbour matrices (PCNM) and geostatistical approaches.
733 *Ecography* **30**: 385-399.
- 734 Bigot, J.L. 1979. Identification des zoés de tourteau (*Cancer pagurus* L.) et d'étrille (*Macropipus*
735 *puber* L.), comparaison avec d'autres zoés de morphologie très voisine. ICES CM 1979/L:17,
736 International Council for the Exploration of the Sea, Copenhagen, DK.
- 737 Blanchet, F.G., Legendre, P., and Borcard, D. 2008. Forward selection of explanatory variables.
738 *Ecology* **89**: 2623-2632.
- 739 Borcard, D., and Legendre, P. 2002. All-scale spatial analysis of ecological data by means of principal
740 coordinates of neighbour matrices. *Ecol. Model.* **153**: 51-68.
- 741 Borcard, D., Gillet, F., and Legendre, P. 2011. Numerical ecology with R. Springer, New York.

- 742 Brander, K.M. 1994. The Location and Timing of Cod Spawning Around the British Isles. ICES J.
743 Mar. Sci. **51**: 71-89.
- 744 Bunn, N.A., Fox, C.J., and Webb, T. 2000. A literature review of studies on fish egg mortality:
745 Implications for the estimation of spawning stock biomass by the annual egg production method. Sci.
746 Ser., Tech. Rep. (111), CEFAS, Lowestoft, UK.
- 747 Charrier, G., Coombs, S.H., McQuinn, I.H., and Laroche, J. 2007. Genetic structure of whiting
748 *Merlangius merlangus* in the northeast Atlantic and adjacent waters. Mar. Ecol. Prog. Ser. **330**: 201-
749 211.
- 750 Collins, M., Knutti, R., Arblaster, J., Dufresne, J., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski,
751 W., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J., and Michael, W. 2013. Long-
752 term climate change: Projections, commitments and irreversibility. In Climate change 2013: The
753 physical science basis. contribution of working group I to the fifth assessment report of the
754 intergovernmental panel on climate change. Edited by T.F. Stocker, D. Qin, G.-. Plattner, M. Tignor,
755 S.K. Allen, J. Boschung, A. Naules, Y. Xia, V. Bex and P.M. Midgley. Cambridge University Press,
756 Cambridge, United Kingdom and New York NY, USA.
- 757 Coombs, S.H., Nichols, J.H., and Fosh, C.A. 1990. Plaice eggs (*Pleuronectes platessa* L.) in the
758 southern North Sea: abundance, spawning area, vertical distribution, and buoyancy. J. Cons. Int.
759 Explor. Mer. **47**: 133-139.
- 760 Coull, K.A., Johnstone, R., and Rogers, S.I. 1998. Fisheries sensitivity maps in british waters.
761 UKOOA Ltd, 1998, Aberdeen, Scotland.
- 762 Cushing, D.H. 1990. Hydrographic containment of a spawning group of plaice in the Southern Bight
763 of the North Sea. Mar. Ecol. Prog. Ser. **58**: 287-297.
- 764 Daan, N. 1978. Changes in cod stocks and cod fisheries in the North Sea. Rapp. P. -V. Réun. Cons.
765 Int. Explor. Mer **172**: 39-57.

- 766 Daan, N., Heessen, H., Hislop, J.R.G., and Parnell, W. 1981. Results of the international 0-group
767 gadoid survey in the north sea, 1981.
- 768 Damas, D. 1909. Contribution à la biologie des gadides. Rapp. P. -V. Réun. Cons. Int. Explor. Mer
769 **10**: 1-227.
- 770 De Groot, S.d. 1996. Quantitative assessment of the development of the offshore oil and gas industry
771 in the North Sea. ICES J. Mar. Sci. **53**: 1045-1050.
- 772 de la Cruz, M. 2008. Metodos para analizar datos puntuales. *In* Introducción al análisis espacial de
773 datos en ecología y ciencias ambientales: Métodos y aplicaciones. *Edited by* Maestre, F. T., Escudero,
774 A. y Bonet, A. Asociacion Espanola de Ecologia Terrestre, Universidad Rey Juan Carlos y Caja de
775 Ahorros del Mediterraneo Asociacion Espanola de Ecologia Terrestre, Universidad Rey Juan Carlos y
776 Caja de Ahorros del Mediterraneo, Madrid, ES. pp. 75-127.
- 777 Diniz-Filho, J.A.F., Bini, L.M., and Hawkins, B.A. 2003. Spatial autocorrelation and red herrings in
778 geographical ecology. Global Ecol. Biogeogr. **12**(1): 53-64.
- 779 Dray, S., with contributions of Legendre, P., and Blanchet, G. 2013. packfor: Forward Selection with
780 permutation (Canoco p. 46) [online]. Available from <https://r-forge.r-project.org/> [cited 1 2016].
- 781 Drinkwater, K.F. 2005. The response of Atlantic cod (*Gadus morhua*) to future climate change. ICES
782 J. Mar. Sci. **62**: 1327-1337.
- 783 Dulvy, N.K., Rogers, S.I., Jennings, S., Stelzenmüller, V., Dye, S.R., and Skjoldal, H.R. 2008.
784 Climate change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. J.
785 Appl. Ecol. **45**: 1029-1039.
- 786 Fox, C.J., Geffen, A.J., Blyth, R., and Nash, R.D.M. 2003. Temperature-dependent development rates
787 of plaice (*Pleuronectes platessa* L.) eggs from the Irish Sea. J. Plankton Res. **25**: 1319-1329.

- 788 Fox, C.J., Taylor, M., Dickey-Collas, M., Fossum, P., Kraus, G., Rohlf, N., Munk, P., van Damme, C.
789 J. G., Bolle, L.J., Maxwell, D.L., and Wright, P.J. 2008. Mapping the spawning grounds of North Sea
790 cod (*Gadus morhua*) by direct and indirect means. Proc. R. Soc. Lond. B **275**: 1543-1548.
- 791 Fraser, J.H. 1968. Zooplankton sampling. 211, UNESCO, France.
- 792 Fritsch, M. 2005. Traits biologiques et exploitation du bar commun *Dicentrarchus labrax* (L.) dans le
793 pêcheries françaises de la manche et du golfe de gascogne. mémoire de thèse. Université de Bretagne
794 occidentale, Brest, FR.
- 795 Fromentin, J.M., and Planque, B. 1996. Calanus and environment in the eastern North Atlantic II.
796 Influence of the North Atlantic Oscillation on *C. finmarchicus* and *C. helgolandicus*. Mar. Ecol. Prog.
797 Ser. **134**: 111-118.
- 798 Fujii, T. 2015. Temporal variation in environmental conditions and the structure of fish assemblages
799 around an offshore oil platform in the North Sea. **108**: 69-82.
- 800 Geffen, A.J., Fox, C.J., and Nash, R.D.M. 2006. Temperature-dependent development rates of cod
801 *Gadus morhua* eggs. J. Fish Biol. **69**: 1060-1080.
- 802 Geffen, A.J., and Nash, R.D.M. 2012. Egg development rates for use in egg production methods
803 (EPMs) and beyond. Fish. Res. **117-118**: 48-62.
- 804 Gehringer, J.W. 1962. The Gulf III and other modern high-speed plankton samplers. Rapp. P. -V.
805 Réun. Cons. Int. Explor. Mer **153**: 19-22.
- 806 Gilman, E., Chaloupka, M., Read, A., Dalzell, P., Holetschek, J., and Curtice, C. 2012. Hawaii
807 longline tuna fishery temporal trends in standardized catch rates and length distributions and effects
808 on pelagic and seamount ecosystems. Aquat. Conserv. : Mar. Freshwat. Ecosyst. **22**(4): 446-488.
- 809 Goedecke, E., Smed, J., and Tomczak, G. 1967. Monatskarten des salzgehaltes der nordsee dargestellt
810 für verschiedene Tiefenhorizonte. Deutsches hydrographisches Institut, Hamburg.

- 811 Golden Software. 2011. Surfer. Golden Software, Inc, Golden, CO.
- 812 González-Irusta, J., and Wright, P. 2016a. Spawning grounds of Atlantic cod (*Gadus morhua*) in the
813 North Sea. ICES J. Mar. Sci. **73**(2): 304-315.
- 814 González-Irusta, J., and Wright, P. 2016b. Spawning grounds of haddock (*Melanogrammus*
815 *aeglefinus*) in the North Sea and West of Scotland. Fish. Res. **183**: 180-191.
- 816 Graham, M. 1934. Report on the north sea cod. HM Stationery Office, London, UK.
- 817 Heath, M.R., Culling, M.A., Crozier, W.W., Fox, C.J., Gurney, W.S., Hutchinson, W.F., Nielsen,
818 E.E., O'Sullivan, M., Preedy, K.F., and Righton, D.A. 2014. Combination of genetics and spatial
819 modelling highlights the sensitivity of cod (*Gadus morhua*) population diversity in the North Sea to
820 distributions of fishing. ICES J. Mar. Sci. **71**(4): 794-807.
- 821 Heath, M., Rankine, P., and Cargill, L. 1994. Distribution of cod and haddock eggs in the North Sea
822 in 1992 in relation to oceanographic features and compared with distributions in 1952-1957
823 (Abstract). ICES Mar. Sci. Symp. **198**: 438-439.
- 824 Heath, M.R., and Gallego, A. 1998. Bio-physical modelling of the early life stages of haddock,
825 *Melanogrammus aeglefinus*, in the North Sea. Fish. Oceanogr. **7**: 110-125.
- 826 Heath, M.R., Kunzlik, P.A., Gallego, A., Holmes, S., and Wright, P.J. 2008. A model of meta-
827 population dynamics for North Sea and West of Scotland cod - The dynamic consequences of natal
828 fidelity. **93**: 92-116.
- 829 Hedger, R., McKenzie, E., Heath, M., Wright, P., Scott, B., Gallego, A., and Andrews, J. 2004.
830 Analysis of the spatial distributions of mature cod (*Gadus morhua*) and haddock (*Melanogrammus*
831 *aeglefinus*) abundance in the North Sea (1980-1999) using generalised additive models. Fish. Res. **70**:
832 17-25.

- 833 Hislop, J.R.G. 1984. A comparison of the reproductive tactics and strategies of cod, haddock, whiting
834 and norway pout in the north sea. *In* Fish reproduction: Strategies and tactics. *Edited by* G.W. Potts
835 and R.J. Wootton. Academic Press, Oxford. pp. 311-330.
- 836 Höffle, H., Solemdal, P., Korsbrekke, K., Johannessen, M., Bakkeplass, K., and Kjesbu, O.S. 2014.
837 Variability of northeast Arctic cod (*Gadus morhua*) distribution on the main spawning grounds in
838 relation to biophysical factors. *ICES J. Mar. Sci.* **71**: 1317-1331.
- 839 Holmes, S.J., Millar, C.P., Fryer, R.J., and Wright, P.J. 2014. Gadoid dynamics: differing perceptions
840 when contrasting stock vs. population trends and its implications to management. *ICES J. Mar. Sci.*
841 **71**(6): 1433-1442.
- 842 Houghton, R., and Harding, D. 1976. The plaice of the English Channel: spawning and migration. *J.*
843 *Cons. Int. Explor. Mer.* **36**: 229-239.
- 844 Hufnagl, M., Peck, M.A., Nash, R.D., Pohlmann, T., and Rijnsdorp, A.D. 2013. Changes in potential
845 North Sea spawning grounds of plaice (*Pleuronectes platessa* L.) based on early life stage
846 connectivity to nursery habitats. *J. Sea Res.* **84**: 26-39.
- 847 Hunter, E., Metcalfe, J., Arnold, G., and Reynolds, J. 2004. Impacts of migratory behaviour on
848 population structure in North Sea plaice. *J. Anim. Behav.* **73**: 377-385.
- 849 ICES. 2009. ICES dataset on ocean hydrography.
- 850 ICES. 2008. Report of the planning group on north sea cod and plaice egg surveys (PGEGGS), 13-14
851 may 2008, Lowestoft, UK.
- 852 Kelley, D. 2013. *Oce: Analysis of oceanographic data.*
- 853 Laurence, G.C., and Rogers, C.A. 1976. Effects of temperature and salinity on comparative embryo
854 development and mortality of Atlantic cod (*Gadus morhua* L.) and haddock
855 (*Melanogrammus aeglefinus* L.). *J. Cons. Int. Explor. Mer.* **36**: 220-228.

- 856 Legendre, P., Borcard, D., Blanchet, G., and Dray, S. 2013. PCNM: MEM spatial eigenfunction and
857 principal coordinate analyses.
- 858 Lelièvre, S. 2010. Identification and characterisation of winter spawning ground in the english
859 channel and southern north sea. PhD, Université de Nantes, Nantes, FR.
- 860 Lelièvre, S., Verrez-Bagnis, V., Jerome, M., and Vaz, S. 2010. PCR-RFLP analyses of formalin-fixed
861 fish eggs for the mapping of spawning areas in the Eastern Channel and Southern North Sea. J.
862 Plankton Res. **32**: 1527-1539.
- 863 Lelièvre, S., Vaz, S., and Fox, C. 2012. The effectiveness of the continuous underway fish egg
864 sampler compared with standard vertical plankton hauls for sampling the eggs of demersal species in
865 the southern North Sea. Fish. Oceanogr. **21**: 109-124.
- 866 Lelièvre, S., Vaz, S., Martin, C., and Loots, C. 2014. Delineating recurrent fish spawning habitats in
867 the North Sea. J. Sea Res. **91**: 1-14.
- 868 Løkkeborg, S., Humborstad, O., Jørgensen, T., and Soldal, A.V. 2002. Spatio-temporal variations in
869 gillnet catch rates in the vicinity of North Sea oil platforms. ICES Journal of Marine Science: Journal
870 Du Conseil **59**(suppl): S294-S299.
- 871 Loots, C., Vaz, S., Planque, B., and Koubbi, P. 2011a. Understanding what controls the spawning
872 distribution of North Sea whiting (*Merlangius merlangus*) using a multi-model approach. Fish.
873 Oceanogr. **20**: 18-31.
- 874 Loots, C., Sandrine, V., Planque, B., and Koubbi, P. 2011b. Spawning distribution of North Sea plaice
875 and whiting from 1980 to 2007. J. Oceanogr. **3**: 77-95.
- 876 Loots, C., Vaz, S., Planque, B., and Koubbi, P. 2010a. What controls the spatial distribution of the
877 North Sea plaice spawning population? Confronting ecological hypotheses through a model selection
878 framework. ICES J. Mar. Sci. **67**: 244-257.

- 879 Loots, C., Vaz, S., Koubbi, P., Planque, B., Coppin, F., and Verin, Y. 2010b. Inter-annual variability
880 of North Sea plaice spawning habitat. *J. Sea Res.* **64**: 427-435.
- 881 Mastail, M., and Battaglia, A. 1978. Amélioration de la conservation des pigments du zooplancton.
882 CM 1978/L:20, Conseil international pour l'Exploration de la Mer, Roscoff, FR.
- 883 McCauley, R.D., Production, A.P., and Association, E. 2000. Marine seismic surveys: a study of
884 environmental implications. *Appea J.* **40**: 692-708.
- 885 Metcalfe, J. 2006. Fish population structuring in the North Sea: understanding processes and
886 mechanisms from studies of the movements of adults. *J. Fish Biol.* **69**: 48-65.
- 887 Millero, F.J., and Poisson, A. 1981. International One-Atmosphere Equation of State of Seawater.
888 *Deep Sea Res. Pt. A* **28**: 625-629.
- 889 Ministerie van Economische Zaken. 2015. NL Olie- en Gasportaal [online]. Available from
890 http://www.nlog.nl/nl/pubs/maps/other_maps/other_maps.html [cited 06 2015].
- 891 Munk, P., Wright, P.J., and Pihl, N.J. 2002. Distribution of the early larval stages of cod, plaice and
892 lesser sandeel across haline fronts in the North Sea. *Coast. Shelf Sci.* **55**: 139-149.
- 893 Munk, P., Larsson, P.O., Danielssen, D.S., and Moksness, E. 1999. Variability in frontal zone
894 formation and distribution of gadoid fish larvae at the shelf break in the northeastern North Sea. *Mar.*
895 *Ecol. Prog. Ser.* **177**: 221-233.
- 896 Munk, P., Fox, C.J., Bolle, L.J., van Damme, C.J.G., Fossum, P., and Kraus, G. 2009. Spawning of
897 North Sea fishes linked to hydrographic features. **18**: 458-469.
- 898 Munk, P., and Nielsen, J. 2005. Eggs and larvae of north sea fishes. Biofolia, Frederiksberg, DK.
- 899 Nash, R.D.M., Dickey-Collas, M., and Milligan, S.P. 1998. Descriptions of the Gulf VII/PRO-NET
900 and MAFF/Guildline unencased high-speed plankton samplers. *J. Plankton Res.* **20**: 1915-1926.

- 901 Neat, F., and Righton, D. 2007. Warm water occupancy by North Sea cod. *Proc. R. Soc. Lond. B* **274**:
902 789-798.
- 903 Nellen, V.M., and Hempel, G. 1969. Versuche zur Fängigkeit des 'Hai' und des modifizierten Gulf-V-
904 Plankton-Samplers 'Nackthai'/comparisons of the fishing efficiency of the Gulf III sampler 'Hai' and a
905 modified sampler 'Nackthai'. *Meeresforsch.* **20**: 141-154.
- 906 NOAA (NGDC). 2016a. Bathymetry and Global Relief [online]. Available from
907 <https://www.ngdc.noaa.gov/mgg/bathymetry/relief.html> [cited 6 2016].
- 908 NOAA (NGDC). 2016b. Shoreline / Coastline Resources [online]. Available from
909 <https://www.ngdc.noaa.gov/mgg/shorelines/> [cited 6 2016].
- 910 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., O'Hara, R.B., Simpson, G.L., Solymos, P.,
911 Stevens, M.H.H., and Wagner, H. 2013. *Vegan*: Community ecology package.
- 912 Oljedirektoratet. 2015. Seismic surveys [online]. Available from
913 http://gis.npd.no/arcgis/rest/services/FactMaps/2_0/MapServer/7 [cited 6 2015].
- 914 OSPAR Commission. 2014. Inventory of Offshore Installations [online]. Available from
915 <http://www.ospar.org/data> [cited 7 2016].
- 916 Ottersen, G., Hjermann, D.O., and Stenseth, N.C. 2006. Changes in spawning stock structure
917 strengthen the link between climate and recruitment in a heavily fished cod (*Gadus morhua*) stock.
918 *Fish. Oceanogr.* **15**: 230-243.
- 919 Otto, L., and Zimmerman, J.T.F. 1990. Review of the physical oceanography of the North Sea Part 1.
920 General review. *Neth. J. Sea Res.* **26**: 163-193.
- 921 Payne, J.F. 2004. Potential effect of seismic surveys on fish eggs, larvae and zooplankton. Canadian
922 Science Advisory Secretariat, St. John's, CA.

- 923 Pebesma, E.J. 2004. Multivariate geostatistics in S: The gstat package.
- 924 Peres-Neto, P.R., Legendre, P., Dray, S., and Borcard, D. 2006. Variation partitioning of species data
925 matrices: Estimation and comparison of fractions. *Ecology* **87**: 2614-2625.
- 926 Perry, A.L., Low, P.J., Ellis, J.R., and Reynolds, J.D. 2005. Climate change and distribution shifts in
927 marine fishes. *Science* **308**: 1912-1915.
- 928 Petitgas, P., Rijnsdorp, A.D., Dickey-Collas, M., Engelhard, G.H., Peck, M.A., Pinnegar, J.K.,
929 Drinkwater, K., Huret, M., and Nash, R.D. 2013. Impacts of climate change on the complex life
930 cycles of fish. *Fish. Oceanogr.* **22**(2): 121-139.
- 931 Planque, B., and Frédou, T. 1999. Temperature and the recruitment of Atlantic cod (*Gadus morhua*).
932 *Can. J. Fish. Aquat. Sci.* **56**: 2069-2077.
- 933 Posgay, J., and Marak, R.R. 1980. The MARMAP bongo zooplankton samplers. *J. Northw. Atl. Fish.*
934 *Sci* **1**: 91-99.
- 935 Pova, I., Davie, A., Treasurer, J., and Migaud, H. 2011. Broodstock spawning and larviculture of
936 whiting (*Merlangius merlangus* L.) reared in captivity. *Aquacult. Res.* **42**: 386-398.
- 937 Rijnsdorp, A. 1989. Maturation of male and female North Sea plaice (*Pleuronectes platessa* L.). *J.*
938 *Cons. Int. Explor. Mer.* **46**: 35-51.
- 939 Rindorf, A., and Lewy, P. 2006. Warm, windy winters drive cod north and homing of spawners keeps
940 them there. *J. Appl. Ecol.* **43**: 445-453.
- 941 Russell, F.S. 1976. The eggs and planktonic stages of British marine fishes. Academic Press, London,
942 UK.
- 943 Ryland, J.S., Nichols, J.H., and Sykes, A.M. 1975. Effect of temperature on embryonic development
944 of plaice, *Pleuronectes platessa* L. (Teleostei). *J. Exp. Mar. Biol. Ecol.* **18**: 121-137.

- 945 Schwarz, G. 1978. Estimating the dimension of a model. *The Annals of Statistics* **6**(2): 461-464.
- 946 Sharples, J., Moore, C.M., Hickman, A.E., Holligan, P.M., Tweddle, J.F., Palmer, M.R., and Simpson,
947 J.H. 2009. Internal tidal mixing as a control on continental margin ecosystems. *Geophys. Res. Lett.*
948 **36**(23).
- 949 Simpson, J., Sharples, J., and Rippeth, T. 1991. A prescriptive model of stratification induced by
950 freshwater runoff. *Estuar. Coast. Shelf Sci.* **33**(1): 23-35.
- 951 Simpson, J.H., Bos, W.G., Schirmer, F., Souza, A.J., Rippeth, T.P., Jones, S.E., and Hydes, D. 1993.
952 Periodic stratification in the Rhine Rofi in the North Sea. *Oceanol. Acta* **16**: 23-32.
- 953 Smithies, M. 2014. A Review-Worldwide Guidelines to Mitigate the Effect of Seismic on Marine
954 Mammals. 76th EAGE Conference and Exhibition 2014, Amsterdam, NL,.
- 955 Software, G. 1999. Grid operations. *In* Surfer 7 user's guide. *Edited by* Golden Software, Inc, Golden.
956 pp. 361-414.
- 957 Sündermann, J., and Pohlmann, T. 2011. A brief analysis of North Sea physics. *Oceanologia* **53**(3):
958 663-689.
- 959 Syrjala, S.E. 1996. A statistical test for a difference between the spatial distributions of two
960 populations. *Ecology* **77**: 75-80.
- 961 Taylor, M.I., Fox, C., Rico, I., and Rico, C. 2002. Species-specific TaqMan probes for simultaneous
962 identification of cod (*Gadus morhua* L.), haddock (*Melanogrammus aeglefinus* L.) and whiting
963 (*Merlangius merlangus* L.). *Mol. Ecol. Notes* **2**: 599-601.
- 964 Taylor, N., Fox, C.J., Bolle, L.J., Dickey-Collas, M., Fossum, P., Kraus, G., Munk, P., Rohlf, N., van
965 Damme, C.J.G., and Vorbach, M. 2007. Results of the spring 2004 north sea ichthyoplankton
966 surveys: The distribution of fish eggs and larvae from the international ichthyoplankton survey. ICES

- 967 Cooperative Research Report, 285, International Council for the Exploration of the Sea, Copenhagen,
968 DK.
- 969 Thompson, B.M., and Riley, J.D. 1981. Egg and larval development studies in the North Sea cod
970 (*Gadus morhua* L.). Rapp. P. -V. Réun. Cons. Int. Explor. Mer **178**: 553-559.
- 971 Thompson, H. 1928. Haddock biology. International Council for the Exploration of the Sea,
972 Copenhagen, DK.
- 973 Tomczak, G., and Goedecke, E. 1962. Monatskarten der Temperatur der Nordsee dargestellt für
974 verschiedene Tiefenhorizonte. Ergaenzung. Deutsch. Hydrogr. Zeitschr. **7**: 1-112.
- 975 Tranter, D.J. 1968. Zooplankton sampling. UNESCO Report. Monographs on Oceanographic
976 Methodology No. 2, Paris, FR.
- 977 Wahlberg, M., and Westerberg, H. 2005. Hearing in fish and their reactions to sounds from offshore
978 wind farms. Mar. Ecol. Prog. Ser. **288**: 295-309.
- 979 Wegner, G., Damm, U., and Purps, M. 2003. Physical influences on the stock dynamics of plaice and
980 sole in the North Sea. Scient. Mar. **67**: 219-234.
- 981 Wood, S.N. 2006. Generalized additive models: An introduction with R. Chapman & Hall, London,
982 UK.
- 983 Wood, S., and Scheipl, F. 2014. Gamm4: Generalized additive mixed models using mgcv and lme4. R
984 package version 0.2-3.
- 985 Zuur, A.F. 2010. AED: Data files used in mixed effects models and extensions in ecology with R.
986 (2009).
- 987 Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., and Smith, G.M. 2009. Mixed effects models
988 and extension in ecology with R. Springer Science and Business Media, New York.

Table 1: Survey cruises in this study. For the BONGO nets only the mesh size for the samples used in the analysis is given.

Year	Country	Survey ^a	Ship	Start Date	End Date	Gear	Mesh size (µm)	Nos. Hauls
2004	NL	PLACES	Tridens II	1.3.04	4.3.04	Gulf VII	270	64
	DE	PLACES	Heinke	16.2.04	23.2.04	BONGO	500	51
	UK	PLACES	Corystes	18.2.04	8.3.04	Gulf VII	270	137
	DK	PLACES	Dana	25.2.04	6.3.04	BONGO	330	100
	NO	PLACES	Haakon Mosby	8.3.04	23.3.04	Gulf III	330	86
2009	SC	PGEGGS	Scotia	20.2.09	6.3.09	BONGO	350	53
	FR	add. IBTS	Thalassa	17.1.09	13.2.09	WP2	500	136
	NO	add. IBTS	G. O. Sars	8.2.09	21.2.09	Gulf VII	280	57
	NL	IHLS	Tridens II	19.1.09	22.1.09	Gulf VII	280	69
	DK	add. IBTS	Dana	31.1.09	17.2.09	BONGO	330	68

^a) **Abbreviations:** add. IBTS = Additional IBTS-Sampling, IHLS = International Herring Larvae Surveys, PLACES = Plaice and Cod Egg Surveys, PCEGGS = Planning Group on North Sea Cod and Plaice Egg Surveys

994 **Table 2:** Syrjala-test between Stage I and subsequent development stages, for each year and both years combined. P <0.5 indicates
 995 that the distribution patters are significantly dissimilar.
 996

	Stage II				Stage III				Stage IV				Stage V			
	Cramér–von		Kolmogorov		Cramér–von		Kolmogorov		Cramér–von		Kolmogorov		Cramér–von		Kolmogorov	
	Mises		- Smirnov		Mises		- Smirnov		Mises		- Smirnov		Mises		- Smirnov	
	Ψ	p	Ψ	p	Ψ	p	Ψ	p	Ψ	p	Ψ	p	Ψ	p	Ψ	p
2004																
Cod	1.58	0.072	0.18	0.067	0.70	0.546	0.13	0.532	3.92	0.181	0.28	0.139	4.40	0.114	0.29	0.032
Haddock	2.85	0.004	0.22	0.018	1.36	0.061	0.16	0.098	3.78	0.004	0.26	0.012	4.91	0.003	0.33	0.001
Whiting	4.14	0.014	0.22	0.038	2.21	0.221	0.23	0.491	7.27	0.073	0.28	0.274	22.5	0.032	0.47	0.024
Plaice	0.44	0.254	0.11	0.362	1.01	0.061	0.12	0.345	3.39	0.001	0.26	0.002	4.67	0.001	0.31	0.002
2009																
Cod	5.62	0.097	0.33	0.073	5.42	0.062	0.35	0.062	7.09	0.016	0.38	0.011	5.12	0.076	0.35	0.049
Haddock	1.25	0.170	0.27	0.179	1.02	0.202	0.27	0.091	0.67	0.309	0.17	0.488	2.04	0.107	0.33	0.115
Whiting	0.84	0.018	0.25	0.203	0.67	0.075	0.25	0.133	1.59	0.06	0.36	0.305	2.52	0.112	0.54	0.101

Plaice	4.65	0.058	0.36	0.061	1.49	0.144	0.22	0.135	0.92	0.142	0.19	0.091	2.38	0.098	0.30	0.051
<hr/>																
Mean																
<hr/>																
Cod	3.33	0.089	0.16	0.159	2.06	0.219	0.17	0.146	6.69	0.019	0.30	0.016	4.41	0.082	0.21	0.113
Haddock	3.10	0.01	0.17	0.042	1.56	0.116	0.14	0.153	4.64	0.012	0.24	0.022	6.98	0.006	0.32	0.005
Whiting	2.43	0.066	0.20	0.070	3.40	0.104	0.24	0.069	7.99	0.048	0.31	0.13	15.72	0.072	0.37	0.111
Plaice	1.43	0.144	0.15	0.138	0.44	0.521	0.08	0.702	2.10	0.049	0.19	0.009	3.27	0.062	0.21	0.052

Draft

998 **Table 3:** Model formulae, explained deviation, AIC, BIC and Moran's I in the residuals for the best fitting GAMMs fit to presence/absence
 999 and density of fish eggs. Of the Moran's I only the values for the first lag are presented.

1000

Species	Type	N	Model Formula ^a	GAM - Dev. Expl. (%)	AIC	BIC	Moran's I ^b
Cod	Presence/absence	696	YD + s(b_depth, 4) + s5, (1 MEV2) + (1 MEV5)	18.5	688.1	720	0.04**
	Egg density	510	s(YD, 4) + s(b_depth, 4) + s(t5, 4), (1 MEV2) + (1 MEV5)	14.6	1131.2	1173.6	0.23**
Haddock	Presence/absence	696	s(YD, 4) + s(b_depth, 4) + s(tb, 4) + cs.(sb) + σ_{tgb} , (1 MEV3) + (1 MEV10) + (1 MEV28) + (1 MEV83)	57.3	478.5	537.6	0.07**
	Egg density	336	YD + s(b_depth, 3) + s(tb, 3) + s(sb, 3) + σ_{tgb} , (1 MEV3) + (1 MEV10) + (1 MEV28) + (1 MEV83)	42.6	725.3	778.7	0.05**
Whiting	Presence/absence	696	s(YD, 4) + b_depth + tb + s(σ_{tgb} , 4), (1 MEV2)	29.8	672.8	709.2	0.02*
	Egg density	448	s(YD, 4) + s(b_depth, 4) + s(t5, 4) + s(s5, 4) + σ_{tgb} 5, (1 MEV2)	34.8	1081.8	1131.0	0.12**
Plaice	Presence/absence	696	s(YD, 4) + s(b_depth, 4) + tb, (1 MEV2) + (1 MEV8) + (1 MEV12) + (1 MEV15)	21.4	808	853.5	0.02**
	Egg density	363	s(YD, 4) + s(b_depth, 4) + s(t5, 4) + s(s5, 4) + σ_{tgb} 5, (1 MEV2) + (1 MEV8) + (1 MEV12) + (1 MEV15)	29.3	655.1	713.5	0.18**

1001

1002 ^{a)} Model formulae show the fixed-effects covariates, either as linear or smooth (s()) terms with degrees of freedom. The prefix cs. indicates
1003 a scaling function applied to a linear term. Explained devation was calculated with a GAM using the same fixed effects. MEVs were
1004 included as random covariates, e.g. (1|MEV1). YD, day of the year; b_depth, bottom depth; t5, temperature at 5 m depth; s5, salinity at 5 m
1005 depth; σ_{t5} , gradient of σ_t at 5 depth; tb, bottom temperature; sb, bottom salinity; σ_{tgb} , gradient of σ_t at the bottom. MEVs
1006 ^{b)} Significance levels: **: ≤ 0.001 , *: > 0.001

Draft

Table 4: Predictive performance of the selected models and single covariates. The performance for presence/absence models is given as Receiver Operating Characteristic Area Under the Curve (ROC AUC) while for the egg densities it is the r^2 between predictions and observations. N is 636 for presence/absence of all species, for the non-zero abundance it is: $N_{\text{Cod}}=510$, $N_{\text{Haddock}}=336$, $N_{\text{Whiting}}=448$, $N_{\text{Plaice}}=363$.

Response	Covariates ^a	Cod	Haddock	Whiting	Plaice
Presence/Absence (ROC AUC)	Selected Model	0.71	0.85	0.79	0.74
	Year Day	0.64	0.78	0.72	0.63
	Bottom Depth	0.54	0.77	0.50	0.71
	Temperature ₅	-	-	-	-
	Salinity ₅	0.50	-	-	-
	σ_t -Gradient ₅	-	-	-	-
	Temperature _b	-	0.77	0.60	0.63
	Salinity _b	-	0.81	-	-
	σ_t -Gradient _b	-	0.63	0.62	-
	Random	0.50	0.50	0.50	0.50
Egg Density (r^2)	Selected Model	0.45	0.62	0.72	0.51
	Year Day	0.38	0.38	0.59	0.35
	Bottom Depth	0.42	0.56	0.61	0.39
	Temperature ₅	0.42	-	0.57	0.43
	Salinity ₅	-	-	0.58	0.38

σ_t -Gradient5	-	-	0.59	0.31
Temperatureb	-	0.50	-	-
Salinityb	-	0.53	-	-
σ_t -Gradientb	-	0.35	-	-
Random	0.36	0.34	0.56	0.30

1014

1015 ^{a)} Suffixes 5 and b indicate whether a value was measured at the surface (5 m) or at

1016 the bottom.

Draft

Table 5: Log-likelihood ratio test between the selected models and models with each of the covariates removed in turn. Bold font indicates significant p-values.

Species	Dropped Covariate ^a	Presence/Absence			Egg density		
		$\chi^2_{0.05}$	df	p	$\chi^2_{0.05}$	df	p
Cod	Year Day	124.0	1	<0.001	2.2	1	0.13
	Bottom Depth	37.1	2	<0.001	10.2	1	0.001
	Temperature5	-	-	-	13.8	1	<0.001
	Salinity5	5.0	1	0.026	-	-	-
	σ_t -Gradient5	-	-	-	-	-	-
Haddock	Year Day	181.0	2	<0.001	4.8	1	0.029
	Bottom Depth	12.0	2	0.003	14.6	2	0.001
	Temperatureb	35.3	2	<0.001	0.9	2	0.622
	Salinityb	40.4	1	<0.001	15.9	2	<0.001
	σ_t -Gradientb	6.7	1	0.097	14.3	1	<0.001
Whiting	Year Day	134.9	2	<0.001	32.6	2	<0.001
	Bottom Depth	82.4	1	<0.001	79.6	2	<0.001
	Temperature5	-	-	-	24.1	2	<0.001
	Salinity5	-	-	-	0.8	2	0.658
	σ_t -Gradient5	-	-	-	5.4	1	0.020
	Temperatureb	34.1	1	<0.001	-	-	-
	Salinityb	-	-	-	-	-	-
	σ_t -Gradientb	2.0	2	0.374	-	-	-

Plaice	Year Day	35.5	2	<0.001	3.5	2	0.172
	Bottom Depth	78.2	2	<0.001	3.4	2	0.181
	Temperature5	-	-	-	19.1	2	<0.001
	Salinity5	-	-	-	1.8	2	0.408
	σ_t -Gradient5	-	-	-	0.6	1	0.426
	Temperatureb	9.5	1	0.002	-	-	-
	Salinityb	-	-	-	-	-	-
	σ_t -Gradientb	-	1	-	-	-	-

1020

1021 ^{a)} Suffixes 5 and b indicate whether a value was measured at the surface (5 m) or at
1022 the bottom.

Table 6: Variation explained by the fixed effects.

Covariates^a	Cod (%)	Haddock (%)	Whiting (%)	Plaice (%)
Presence/Absence				
Year Day	16.1	33.2	25.7	1.3
Bottom Depth	3.1	27.2	10.8	8.6
Temperature	-	33.0	15.3	8.3
Salinity and σ_{tg}	0.9	22.0	5.1	-
Total	15.9	47.1	29.1	9.6
Egg density				
Year Day	1.7	7.0	1.7	4.4
Bottom Depth	2.1	17.4	4.5	2.9
Temperature	5.0	21.4	6.8	1.1
Salinity and σ_{tg}	-	28.9	9.2	7.9
Total	5.0	30.5	9.9	13.2

1025

1026 ^a) Variation partitioning only allows for four components. Therefore salinity and density
 1027 gradient were, if necessary, grouped. Values for single covariates contain a pure
 1028 component and a component shared with other covariates. Hence, the terms sum up higher
 1029 than the total variation explained. σ_{tg} = density gradient

1030 **Table 7:** Log-likelihood ration test between GAMM and corresponding GAM models for
1031 presence/absence as well as density of eggs for the four species.

Species	Type	AIC		BIC		Log-likelihood ratio test		
		GAMM	GAM	GAMM	GAM	$\chi^2_{0.05}$	df	p
Cod	Presence/Absence	1938.8	1930.0	1957.0	1939.1	4.8	2	0.09
	Egg density	1079.6	1076.2	1096.6	1084.7	0.6	2	0.75
Haddock	Presence/Absence	1626.6	1613.4	1653.9	1622.5	5.2	4	0.26
	Egg density	672.5	659.7	695.4	667.3	4.8	4	0.31
Whiting	Presence/Absence	1921.2	1914.4	1934.8	1923.5	4.7	1	0.03
	Egg density	1005.6	998.7	1017.9	1006.9	4.9	1	0.03
Plaice	Presence/Absence	2046.1	2037.9	2073.3	2046.9	0.2	4	1.0
	Egg density	584.8	574.2	608.1	582.0	2.6	4	0.6

1032

Figure 1: Panel (a) shows the ichthyoplankton stations for 2004 and 2009 over a bathymetric map of the North Sea. Panel (b) exhibits the operational installations related to oil extraction; color-coded depending on the operating country on the same bathymetric map. Red lines depict Exclusive Economic Zones (EEZs). Areas shaded pink; show the areas with planned or ongoing seismic surveys in the Norwegian and Dutch sectors. Germany is not conducting seismic surveys and for the UK and Denmark no information is publicly available.

Figure 2: Mean Temperature and Salinity at the surface (5 m; panels (a, b) and at bottom depth (c, d), based on data from the first quarter of 2004 and 2009. Black dots indicate individual CTD hauls.

Figure 3: Range of structures of spatial autocorrelation (a) for the individual species. Broken lines indicate where the structures were grouped into the scales depicted, in the order fine to broad, on the maps on panels (b) to (e). The plotted structures are those for haddock, except on medium-broad scale (d) where there was only a single structure, for cod.

Figure 4: Mean interpolated surfaces for the distribution of eggs in each stage, drawing from both surveys. Interpolated areas beyond 200 m bottom depth are blanked out. Black dots on panel (a) indicate the sampling stations for both years.

Figure 5: Predicted probability of occurrence for stage I eggs of all four species, on the range of the covariates in the best fitting models. Broken lines indicate models using hydrographic values measured at the surface. Hatches at the bottom of each panel indicate the values observed at samples. As cod was related to surface salinity, while haddock was related to bottom salinity on a narrower range, the corresponding observations are depicted in colours corresponding to the smooth terms. Shaded areas indicate the 95% confidence interval.

Figure 6: Predicted log-transformed stage I egg density for stage I eggs of all four species, related to the covariates in the best fitting models. Broken lines indicate models using hydrographic values measured at the surface. Observations with non-zero egg densities for the individual species are indicated by hatches, in the same colour as the smooth term, at the bottom of each panel. Shaded areas indicate the 95% confidence interval.

Figure 7: Panel (a) shows the average stage I egg density for each ICES rectangle on the interpolated surface of bottom temperature variability, based on CTD data for the 1st quarter of the entire 2000s. Panel (b) depicts installations of the oil industry, categorized as above-surface (red dots) or subsurface structures (white squares) the position of oil industry installation and the seismic survey areas for Norway and the Netherlands, shaded blue, on top of the average distribution of stage I eggs for all four species combined.

Haddock

Whiting

Plaice