Precipitation Input and Antecedent Soil Moisture Effects on Mercury Mobility in Soil – Laboratory Experiments with an Enriched Stable Isotope Tracer

Kristine M. Haynes & Carl PJ Mitchell

Version Post-print/accepted manuscript

Publisher's statement This is the peer reviewed version of the following article:

Haynes, KM, and Mitchell, CPJ (2015), Precipitation input and antecedent soil moisture effects on mercury mobility in soil—laboratory experiments with an enriched stable isotope tracer. *Hydrol. Process.*, 29, 4161–4174. doi: [10.1002/hyp.10442](https://doi.org/10.1002/hyp.10442).

This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

How to cite TSpace items

Always cite the published version, so the author(s) will receive recognition through services that track citation counts, e.g. Scopus. If you need to cite the page number of the **author manuscript from TSpace** because you cannot access the published version, then cite the TSpace version **in addition to** the published version using the permanent URI (handle) found on the record page.

This article was made openly accessible by U of T Faculty.
Please [tell us](#) how this access benefits you. Your story matters.

Precipitation Input and Antecedent Soil Moisture Effects on Mercury Mobility in Soil - Laboratory Experiments with an Enriched Stable Isotope Tracer

Journal:	<i>Hydrological Processes</i>
Manuscript ID:	HYP-14-0717.R1
Wiley - Manuscript type:	Special Issue Canadian Geophysical Union 2015
Date Submitted by the Author:	08-Jan-2015
Complete List of Authors:	Haynes, Kristine; University of Toronto Scarborough, Physical and Environmental Sciences; University of Toronto, Department of Geography Mitchell, Carl; University of Toronto Scarborough, Physical and Environmental Sciences; University of Toronto, Department of Geography
Keywords:	mercury, stable isotope tracer, soils, transport, soil moisture, total organic carbon

SCHOLARONE™
Manuscripts

Review

**Precipitation Input and Antecedent Soil Moisture Effects on Mercury Mobility in Soil –
Laboratory Experiments with an Enriched Stable Isotope Tracer**

Kristine M. Haynes^{a,b*} and Carl P.J. Mitchell^{a,b}

a. University of Toronto Scarborough Department of Physical and Environmental Sciences, 1265
Military Trail, Toronto, Ontario, M1C 1A4, Canada

b. University of Toronto Department of Geography, 100 St. George Street, Toronto, Ontario,
M5S 3G3, Canada

Corresponding author: K.M. Haynes, University of Toronto Scarborough Department of
Physical and Environmental Sciences, 1265 Military Trail, Toronto, Ontario, M1C 1A4, Canada.
(k.haynes@utoronto.ca), Phone: 416-208-2745, Fax: 416-287-7279

Running Title: Hydrological Impacts on Mercury Mobility

Abstract

Terrestrial soils are net mercury (Hg) sinks, but leaching of Hg from upland environments constitutes an important source to downstream wetlands and water bodies. Broadly, hydrology is instrumental in facilitating Hg transport within, and export from watersheds but the relative influences of specific hydrological factors such as antecedent soil moisture and precipitation in controlling the transport of Hg through upland soils are not well understood. The purpose of this research was to elucidate the relative controls of these hydrological factors using a full factorial laboratory experiment involving the application of an enriched stable Hg isotope tracer to intact soil cores. Antecedent soil moisture and precipitation input depth were statistically significant, mutually exclusive controls on tracer Hg mobility. Neither factor however had a strongly significant influence on the mobility of ambient Hg. Tracer Hg mobility was enhanced with larger precipitation events as well as from initially drier soils and appeared to move via simple piston flow. The majority (>99.5%) of added tracer Hg was sorbed to soil organic matter in the surface 3 cm, regardless of the hydrological treatment combinations. Overall, these results suggest that tracer and ambient Hg are differentially controlled by hydrological conditions. Changes in hydrology may have little impact on ambient Hg mobilization in sandy loam soils. If tracer Hg is broadly representative of contemporary soil Hg stocks, extreme precipitation events among otherwise drier conditions could enhance the export of contemporary Hg from upland systems.

Keywords: mercury; stable isotope tracer; soils; transport; soil moisture; total organic carbon

1
2
3
4 46 **1 Introduction**
5

6 47 Soils store an enormous pool of mercury (Hg), containing approximately 1,000,000 Mg of
7
8 48 the global stock (Sunderland and Mason, 2007). In most regions atmospheric deposition is the
9
10 49 principal process delivering Hg from local, regional and global sources (Driscoll *et al.*, 2007;
11
12 50 Fitzgerald *et al.*, 1998). Although soils are large net sinks for atmospherically-deposited Hg
13
14 51 (Gabriel and Williamson, 2004; Grigal, 2003), the accumulated Hg also slowly leaches from
15
16 52 soils and moves via runoff to aquatic ecosystems where its toxic and organic form,
17
18 53 methylmercury (MeHg), may be produced (Branfireun *et al.*, 1996; Harris *et al.*, 2007). High-
19
20 54 flow events such as spring snowmelt appear to amplify the export of both inorganic and organic
21
22 55 Hg species from soils and watersheds, although the mechanisms for increased fluxes vary
23
24 56 (Bishop *et al.*, 1995; Demers *et al.*, 2010; Haynes and Mitchell, 2012; Mitchell *et al.*, 2008a).
25
26 57 Given the toxicological threats of Hg to human and wildlife populations (Mergler *et al.*, 2007;
27
28 58 Scheuhammer *et al.*, 2007), it is important to understand how mercury mobility may be affected
29
30 59 under different and/or changing hydrological conditions through time.
31
32
33
34
35

36
37 60 In field studies it is difficult to discern the relative control of individual hydrological
38
39 61 properties, such as the antecedent soil moisture condition, on Hg transport due to confounding
40
41 62 effects from precipitation inputs and subsequent runoff production as well as groundwater
42
43 63 contributions. At the hillslope scale, Haynes and Mitchell (2012) observed that Hg yield from
44
45 64 sandy loam soils with a shallow surficial O-horizon was flow-driven; increasing proportionally
46
47 65 with snowmelt runoff with no conclusive evidence that antecedent soil moisture was an
48
49 66 important control on the magnitude of runoff Hg fluxes. Previous research in upland
50
51 67 environments (e.g. Bushey *et al.* 2008; Dittman *et al.* 2010) has suggested that high-flow event
52
53 68 runoff enhances Hg mobility. However, few studies consider the potential influence of
54
55
56
57
58
59
60

1
2
3 69 antecedent soil moisture conditions in controlling Hg fluxes through terrestrial environments. A
4
5
6 70 recent study by Oswald and Branfireun (2014) illustrated that antecedent soil moisture conditions
7
8 71 aid in accounting for differences observed in the Hg concentration-discharge relationships that
9
10 72 occur following the wet snowmelt period as compared to the dry late summer period. Higher
11
12 73 THg and soil water dissolved organic carbon concentrations were observed following warmer,
13
14 74 drier periods in the boreal Shield watershed (Oswald and Branfireun 2014). These studies
15
16 75 suggest that a systematic and controlled testing of the relative controls of antecedent soil
17
18 76 moisture and the magnitude of precipitation inputs on Hg mobility is required.
19

20
21
22 77 Due to the myriad variables influencing natural systems it is often difficult to accurately
23
24 78 identify individual hydrological controls on contaminant mobility. Microcosm-scale laboratory
25
26 79 studies reduce the natural variability inherent in field-scale studies and allow for the controlled
27
28 80 manipulation of hydrological variables. Experiments of this nature commonly have been used to
29
30 81 investigate the role of different hydrological factors such as rainfall intensity or soil moisture on
31
32 82 nutrient and pesticide leaching and transport in soil (Phillips, 2001; Phillips and Burton, 2005;
33
34 83 Sigua *et al.*, 1993; Vendelboe *et al.*, 2011). In a recent study Reinmann *et al.* (2012) simulated
35
36 84 soil frost development and spring snowmelt conditions using intact soil columns to determine the
37
38 85 potential effects of frost on carbon and nitrogen losses from forested upland soils. Undisturbed
39
40 86 soil columns have also been used to study the leaching of heavy metals such as lead, copper and
41
42 87 cadmium under different rainfall intensities (Kim *et al.*, 2008). Only a limited number of studies
43
44 88 (e.g. Schlüter *et al.*, 1995; Schlüter 1996) have employed the use of intact soil columns to
45
46 89 investigate the impact of hydrological factors on Hg mobility, despite numerous advances in Hg
47
48 90 analytical techniques including the increasing use of stable isotopes as environmental tracers
49
50
51 91 (Björn *et al.*, 2007).
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Given that Hg in the environment is transported in a highly non-conservative manner, the use of stable isotopes as tracers is an invaluable tool yielding information not only about the transport of Hg but also its speciation under various conditions. The METAALICUS study applied multiple stable Hg isotopes at the landscape scale in order to investigate the potential impacts of altering the atmospheric deposition of Hg and the subsequent downstream effects on Hg bioaccumulation in fish (Harris *et al.*, 2007). A pilot study under the METAALICUS project added an enriched stable Hg isotope to a small, zero-order forested subcatchment as a means of assessing the reactivity and mobility dynamics of newly-deposited Hg as opposed to the large legacy soil pools of Hg at the plot scale (Hintelmann *et al.*, 2002). This field study found that less than 1% of the added Hg appeared in runoff, suggesting that the majority of newly-deposited Hg is incorporated into the organic matter-bound Hg pool in soil and vegetation. Oswald *et al.* (2014) also traced the applied isotope in the watershed upland to assess its distribution (vertical and spatial) and transport, as well as that of ambient Hg through the soil one year following the end of application; with approximately 22% of the spike mass remaining in the upper 15 cm of mineral soil. Despite the increasing application of isotope tracers in field settings, minimal research has been conducted using Hg isotopes in small-scale laboratory experiments to investigate Hg mobility throughout the soil profile. Semu *et al.* (1985) used the radioisotope ²⁰³Hg in soil column experiments to investigate the potential for groundwater contamination by Hg leaching through the soil, observing that minimal vertical translocation of Hg occurred with the majority of added Hg being immobilized in the surface soil horizons. Schlüter (1996) similarly used ²⁰³Hg to assess the translocation of Hg through soil using small-scale column experiments in order to address the fate of Hg in soils under different rainfall intensities and volumes; concluding that practically all of the added ²⁰³Hg is sorbed in the upper organic soil

horizons and that the small amount of Hg translocation is facilitated by complexation with dissolved organic matter. Some more recent work has used column experiments to examine applied non-labelled Hg sorption in soils (e.g. Miretzky *et al.* 2005; Yin *et al.* 1997; Zhu *et al.* 2014). However, such studies did not use intact, undisturbed soil cores working rather with re-packed soil columns (Miretzky *et al.* 2005; Zhu *et al.* 2014) or soil suspensions (Yin *et al.* 1997).

Our work builds on these previous studies by adding antecedent moisture as an experimental variable possibly affecting the mobility and transport of Hg throughout the undisturbed soil profile, and by using new core encasing methods to avoid wall effects and pore blockage that often result from polyvinyl chloride- (PVC) (e.g. Bagarello *et al.*, 2006; Bagarello and Sgroi 2008), resin-encased (e.g. Vanderborght *et al.* 2002) or wax-encased cores.

The overall objective of this study was to determine the relative influences of both antecedent soil moisture and precipitation on the mobility and transport of Hg throughout the soil profile. In a full factorial design we applied an enriched, stable Hg isotope to intact, foam-encased soil cores that had been altered for soil moisture and to which variable amounts of precipitation were added. The experiment was conducted to simulate the broad range in real soil moisture and precipitation input conditions observed in previous field research (Haynes and Mitchell, 2012). A pilot study, described in the Supplementary Information, was also conducted in order to ensure the feasibility of using polyurethane expandable spray foam as a soil-encasing material (e.g., Bagarello and Sgroi 2008) for the purpose of performing Hg leaching experiments while minimizing wall effects.

2 Materials and Methods

2.1 Field Sampling

Eighteen intact soil cores were collected with a 15 cm deep and 10 cm internal diameter stainless steel soil corer in the forested upland of the S7 watershed in the Marcell Experimental Forest (MEF) in north-central Minnesota. The soil corer is equipped with a lever mechanism which allows for extrusion of the soil core from the base of the corer tube without disturbing the soil structure. Characteristic of watersheds in this area, the upper sandy loam soils of the watershed's hillslope are underlain by a low-permeability B-horizon, which acts as a confining layer, directing subsurface runoff as shallow interflow during high-flow events with minimal seepage loss to groundwater (Kolka *et al.*, 2011; Timmons *et al.*, 1977). The soil profile above the clay-rich B-horizon consists of a shallow O-horizon below which there is a well-defined sandy loam A-horizon (mean depth to base of A-horizon 50 ± 27 cm). Hydraulic conductivities in the sandy loam of this area range from 4×10^{-5} to 4×10^{-6} m s⁻¹ (Mitchell *et al.* 2009). The surface 15 cm of soil in this hillslope have a mean bulk density of 1.2 g cm⁻³ and mean porosity of approximately 37%. The upland overstory vegetation in the S7 watershed is predominately comprised of sugar maple (*Acer saccharum*), quaking aspen (*Populus tremuloides*) and balsam poplar (*Populus balsamifera*). We sampled only the upper sandy loam horizon that also included a shallow (< 3 cm) organic horizon. Soil cores were collected from outside of experimental hillslope plots previously described in Haynes and Mitchell (2012), all within a 5 m² area in order to minimize variability among cores and to allow for the most accurate assessment of the influences of only the two hydrological factors in question (antecedent soil moisture and depth of precipitation input). Each core was extruded in the field onto a flat wooden board. We noted no significant compaction or destruction of the soil cores during the collection and extrusion of each

core. In comparing core depth to the cavity from which it was extracted, approximately 2% (0.3 cm) to a maximum of 5% (< 1 cm) compaction was observed. With the core upright on the wooden board, a 20 cm long cardboard mold was placed around it and the cavity between the soil and mold was filled with low-expansion polyurethane spray foam insulation (MONO brand); similar to the methods of Bagarello and Sgroi (2008). A second board was then placed on top of each filled mold to contain the expansion of the foam. The foam was allowed to cure overnight. Once fully cured, each foam-encased core was removed from the wood base and the exposed bottom of each core was also covered with spray foam insulation. No compaction of any soil core was observed as a result of the expansion of the spray foam. The purpose of this elaborate core encasing methodology was to hypothetically remove much of the wall effect, smear and other issues affecting water movement that often occur in intact core studies conducted in rigid tubes and other encasing materials (e.g. Cameron *et al.*, 1990; Hoag and Price, 1997; Bagarello *et al.*, 2006). With other encasing methods, preferential flow directed along the point of contact between the soil and surrounding casing often occurs rather than flow through the soil matrix. A separate study was conducted in order to test the feasibility of using foam-encased soil cores for the study of hydrological and Hg biogeochemical processes. This is detailed in the Supplementary Information.

2.2 Experimental Design

The soil core experiments were conducted in a full factorial design in which the two manipulated variables (antecedent soil moisture and precipitation input depth) incorporated both a low and high treatment level with three replicate cores per treatment combination (see Table 1). High and low treatment levels were quantified based on field data from the study area. For

precipitation input, the low and high levels were 50 mm and 100 mm, respectively. These values were selected in order to mimic the variability in snowpack snow water equivalent (SWE) levels observed in a previous study (Haynes and Mitchell, 2012). The low antecedent soil moisture treatment level was designed to simulate the 65 mm of available soil water in the surface 45 cm of soil recorded in fall 2009 prior to the 2010 winter in the study area. The high antecedent soil moisture treatment level was represented by field capacity as a means of presenting a distinct contrast to the low moisture treatment. All experiments were conducted at room temperature.

To adjust moisture levels in the low antecedent soil moisture treatments, a known volume of tap water (determined based on the volume of the soil column and the physical characteristics of the soil to achieve the target low moisture level) was slowly added to the top of relatively dry cores over the course of approximately one hour with no ponding. For the high antecedent moisture treatments, cores were placed in a basin of tap water overnight to saturate the soil from the base of each core and then drained by gravity the following day; effectively reaching field capacity. To prevent loss of soil during the wetting process, Nytex mesh (50 μm) was secured around the base of each foam-encased soil core prior to manipulation. Once the moisture manipulations were complete, 90 μg of enriched stable ^{199}Hg isotope (approximately three times ambient soil Hg mass per core) was applied evenly to the top of each core in a 10 mL solution of $^{199}\text{HgCl}$ (91.09% ^{199}Hg) via a hydrochloric acid (HCl)-washed Teflon sieve (mean blank ^{199}Hg concentration between uses = $0.6 \pm 0.5 \text{ ng L}^{-1}$; see Figure 1a). The Teflon sieve was used to ensure uniform application of the isotopic solution across the top of each soil core. A 5 mL water rinse of the Teflon sieve followed the isotope addition to ensure complete delivery of the isotope application. Although significantly greater than natural rates of atmospheric deposition at the Marcell Experimental Forest (2004-2013 mean annual deposition of $6.8 \pm 1.4 \mu\text{g m}^{-2}$;

205 National Atmospheric Deposition Program Mercury Deposition Network, 2014), this high level
206 of tracer addition was necessary in order to ensure that the isotope could be analytically detected
207 in the collected outflow waters given the working hypothesis that the majority of the applied
208 tracer would be sorbed in the soil. Given the laboratory setup of this experiment, it would not be
209 feasible to apply the tracer to each of the soil cores at natural deposition rates over the course of
210 multiple years to achieve detectable isotope levels in the core outflow. Immediately following
211 the tracer addition each core was secured to a ring stand balanced on top of a wire mesh above an
212 HCl-washed glass funnel (see Figure 1b). Once secured (approximately five minutes following
213 tracer addition), a rainfall simulator delivered the prescribed level of precipitation input via a
214 metered peristaltic pump. The rainfall simulator was modelled after that used by Weed *et al.*
215 (1998); an evenly-distributed array of hypodermic needles secured through a piece of acrylic
216 tightly bolted to a second piece of acrylic (fitted with a water inlet nut) with a water-tight gasket
217 between the two sheets of acrylic providing a water reservoir. The two levels of precipitation
218 input (low = 50 mm, high = 100 mm) were comprised of tap water (to minimize ionic
219 imbalances), acidified with trace metal grade nitric acid to simulate the pH of natural
220 precipitation (pH~5-6). This simulated precipitation was delivered to each core at a rate of
221 approximately 30-40 mm per hour. The low input experiments were completed over
222 approximately 1.5 hours per core, while the high input experiments took approximately 3 hours
223 to complete. Although designed to simulate two contrasting snowmelt events, the selected levels
224 of input could also be representative of large rain storm events in the study area. There is recent
225 precedent for this intensity of precipitation in the area of north-central Minnesota given the large,
226 record-breaking rainstorms of June 19-20, 2012, in which approximately 185 mm of
227 precipitation (with up to 250 mm reported locally) fell onto already saturated ground and from

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

228 which severe flooding resulted; particularly in the Duluth area. Instantaneous rainfall intensities
229 for this storm were observed to be as high as 50-100 mm per hour during the evening of June 19,
230 2012 (Czuba *et al.*, 2012). In the context of the present study, the timing of the outflow from
231 each of the soil cores may be different than would be observed in the natural environment as a
232 result of the applied precipitation intensity. For each antecedent moisture treatment level a
233 control experiment, in which no further precipitation input was added to each core following
234 isotope addition, was performed to determine the level to which the isotope was transported as a
235 baseline prior to simulated meltwater addition.

236 Outflow from each core was collected in 50 mL intervals in new polyethylene
237 terephthalate glycol (PETG) bottles. A sub-sample for total organic carbon (TOC) analysis was
238 transferred into a 20 mL high density polyethylene (HDPE) scintillation vial and refrigerated in
239 darkness until analysis (approximately 2-3 weeks following collection). Comparisons of filtered
240 and unfiltered samples were analyzed for TOC to determine if filtration significantly alters TOC
241 concentrations. No significant difference was observed between the filtered and unfiltered
242 samples, most likely due to the Nytex mesh acting to filter particulates from the collected
243 outflow. Therefore, outflow samples were not filtered. Outflow samples for Hg analysis were
244 preserved with 0.5% trace metal grade HCl, stored in darkness and refrigerated until analysis,
245 which was within one month of collection. Once outflow from the core had ceased, each core
246 was cut vertically into two equal parts and the soil from each half was sampled in five 3 cm
247 depth intervals (0-3, 3-6, 6-9, 9-12, 12-15 cm). Soil samples were immediately frozen and then
248 lyophilized prior to analysis.

2.3 Analytical Methods

Unfiltered total Hg (for both tracer ^{199}Hg and ambient Hg analysis) water samples were oxidized by the addition of concentrated bromine monochloride (BrCl) and concentrations were determined following stannous chloride reduction and dual gold-trap amalgamation using a Tekran model 2600 total mercury analyzer that was hyphenated to an Agilent 7700 Inductively Coupled Plasma – Mass Spectrometer (ICP-MS) for mercury isotope detection. The freeze-dried soil samples were nitric acid-digested using a microwave digester and diluted prior to THg analysis as for waters. The ^{199}Hg concentrations of the added tracer in both the water and soil samples were determined as the excess total ^{199}Hg concentrations above the naturally-occurring ^{199}Hg concentrations as measured by the ICP-MS. These excess total ^{199}Hg concentrations were calculated based on the magnitude of change from the natural ratio of ^{199}Hg above that of ambient ^{202}Hg ($^{199}\text{Hg}/^{202}\text{Hg} = 0.569$ in nature) as a result of the addition of the labelled ^{199}Hg tracer (Hintelmann and Ogrinc, 2003). Excess total ^{199}Hg will be herein referred to as ‘tracer Hg’. For water samples, recovery of an ambient total mercury spike was $97 \pm 9\%$ (average $\pm$ standard deviation, $n = 11$), replication of duplicates was $6.6 \pm 6.8\%$ ($n = 10$), and the detection limit, calculated as three standard deviations of matrix water blanks, was 0.04 ng L^{-1} ($n = 32$). Replication of ^{199}Hg duplicates in water samples was $6.3 \pm 6.9\%$ ($n = 10$) and the ^{199}Hg detection limit was 0.007 ng L^{-1} ($n = 32$). For digested soil samples, replication of ambient THg duplicates was $5.4 \pm 4.0\%$ ($n = 11$), recovery of an ambient THg spike was $101 \pm 5\%$ ($n = 11$) and the ambient THg detection limit was calculated to be 0.09 ng g^{-1} . Replication of ^{199}Hg duplicates of digested soil samples was $9 \pm 8\%$ ($n = 9$) and the ^{199}Hg detection limit was 0.08 ng g^{-1} . Recovery of standard reference material of known ambient Hg concentration, digested and analyzed with the soil samples, was $97 \pm 4\%$ ($n = 12$). Total organic carbon (TOC) was analyzed using a

1
2
3 273 Shimadzu TOC-V_{CSH} Total Organic Carbon analyzer with an in-line total nitrogen unit (TNM-1)
4
5 274 and a high sensitivity combustion catalyst (720°C). Ultraviolet absorbance of the outflow
6
7
8 275 samples was measured at both 254 and 365 nm using matched quartz cuvettes on a Spectronic
9
10 276 Genesys 5 UV-VIS Spectrophotometer. We used these two absorbance values measured at these
11
12 277 wavelengths to calculate the ratio A₂₅₄/A₃₆₅, where the ratio serves as a proxy for dissolved
13
14 278 organic carbon quality because of its negative relationship with molecular weight of dissolved
15
16 279 organic substances (Berggren *et al.*, 2007). Specific UV absorbance (SUVA) at 254 nm was also
17
18 280 calculated to assess aromaticity of dissolved organic substances in outflow samples by dividing
19
20 281 the absorbance at 254 nm by the TOC concentration while taking into account the cuvette path
21
22 282 length (Weishaar *et al.*, 2003). Loss on ignition (LOI, 550°C for 4 hours) was performed on the
23
24 283 soil samples to determine organic matter content.
25
26
27
28
29
30
31

32 284
33
34 285 **2.4 Statistical Analyses**
35

36 286 Prior to all parametric statistical analyses, data were tested for normality (Shapiro-Wilk
37
38 287 *W* test) and heteroscedasticity and were successfully log-transformed or inverse-transformed to
39
40 288 achieve normality when parametric assumptions were not met. All statistical tests were
41
42 289 performed using SAS Version 9.2 with $\alpha < 0.05$.
43

44 290 A two-way analysis of variance (ANOVA) was performed to determine any significant
45
46 291 influences of, and any significant interaction between both hydrological variables (antecedent
47
48 292 soil moisture and precipitation input depth) on both tracer Hg and ambient THg outflow loads as
49
50 293 well as the proportion of tracer Hg in relation to ambient THg. A one-way ANOVA (with post-
51
52 294 hoc analysis using the Tukey adjustment) was conducted on the overall tracer Hg and ambient
53
54 295 THg outflow loads to assess significant differences among treatment combinations. A one-way
55
56
57
58
59
60

ANOVA with post-hoc Tukey test was performed in order to determine any significant differences in soil tracer ^{199}Hg concentrations among treatments (including controls) for each of the five depths. Correlation matrices were created using Pearson correlation coefficients in order to assess any significant relationships between the measured outflow variables, as well as between soil tracer Hg concentrations and organic matter. The data for one replicate core in the low moisture-high input treatment and one replicate core in the low moisture-low input treatment were omitted from correlation matrices due to exceedingly high organic matter content in the surface layers resulting from an abundance of decomposing woody debris that could not be seen until cores were sectioned at the end of the study. One replicate core in the high moisture-low input treatment was omitted from all analyses due to a lack of outflow production (from a blockage as a result of a large rock within the core).

3 Results

3.1 Tracer ^{199}Hg and Ambient Hg Outflow Loads by Treatment Combinations

Overall, less than 0.5% (range of 0.03 – 0.38%) of the added 90 μg of the ^{199}Hg tracer was present in the total outflow collected across all treatments combinations. The highest tracer Hg outflow load occurred in the low soil moisture – high precipitation input experiments (mean total load = 341 ng; Table 2). The low moisture-high precipitation outflow tracer Hg load was significantly greater than that of any of the other treatment combinations. The second largest tracer Hg outflow load was observed in the low moisture-low precipitation experiments. There was not a statistically significant difference between the tracer outflow loads resulting from the two high soil moisture treatments (high moisture-high precipitation and high moisture-low precipitation). Both of these treatments were significantly less in outflow tracer Hg than those

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

319 resulting from the low moisture treatments ($p < 0.0001$) regardless of the level of precipitation
320 input (Table 2). Similarly, the outflow tracer Hg loads resulting from the high input treatment
321 significantly exceeded the corresponding low input treatment at each level of soil moisture ($p =$
322 0.001). The total ambient THg loads followed the same general pattern by treatment as the tracer
323 Hg loads (Table 2). However, the statistical significance of the outflow ambient THg differences
324 among the treatment combinations was marginal ($p = 0.07$).

325 According to the two-way ANOVA results, both antecedent soil moisture ($p < 0.0001$)
326 and precipitation input depth ($p = 0.001$) were significant controls on tracer Hg transport and
327 mobility through soil. The lack of interaction ($p = 0.16$) between these hydrological variables
328 suggested that these controls on tracer Hg mobility acted independent of one another.
329 Precipitation input depth did not exert a strong influence ($p = 0.18$) on the mobility of ambient
330 THg. The influence of antecedent soil moisture was marginal (close to statistical significance at
331 $p = 0.06$).

332 The time course plots in Figure 2 present the data for a single representative core in each
333 of the experimental combinations. An average could not be adequately depicted due to slight
334 differences in the timing of the outflow for each core, which temporally staggered the data, but
335 the overall pattern in outflow was similar between replicates despite differences in the timing.
336 The time course plots of the tracer Hg and ambient THg concentrations for each of the treatment
337 combinations (Figure 2) illustrated a slight delay in the tracer Hg peak for the high soil moisture
338 treatments (Figure 2*b* and 2*d* each showing one replicate per treatment), with tracer Hg
339 concentrations peaking during the second to fourth 50 mL-interval outflow sample (15-45
340 minutes lag) at a mean maximum concentration of $106 \pm 27 \text{ ng L}^{-1}$ (average $\pm$ standard
341 deviation). However, for the low moisture treatments, tracer Hg concentrations were highest in

the first collected 50 mL outflow sample (mean maximum ^{199}Hg concentration = $939 \pm 664 \text{ ng L}^{-1}$), decreasing with subsequent samples regardless of the precipitation depth (Figure 2a and 2c each showing one replicate per treatment). Tracer isotope concentrations in outflow were considerably higher for the low moisture treatments than the high moisture treatments.

The first 50 mL sample was collected after 10 ± 3 minutes (average $\pm$ standard deviation) on average from the high antecedent moisture experiments and after 17 ± 13 minutes for the low moisture experiments. The rate of outflow was relatively consistent throughout the course of the experiments, with an average overall outflow rate of $5.6 \pm 1.0 \text{ mL min}^{-1}$ for the high moisture soils and $5.0 \pm 1.1 \text{ mL min}^{-1}$ for the low moisture soils (Figure 2). Overall, of the 393 mL of added precipitation for the low input treatments, $380 \pm 39 \text{ mL}$ for the low moisture-low precipitation experiments and $404 \pm 23 \text{ mL}$ for high moisture-low precipitation was recovered in outflow. For the high input treatments, $745 \pm 23 \text{ mL}$ for the low moisture-high precipitation treatment and $787 \pm 16 \text{ mL}$ for high moisture-high precipitation of the 785 mL added as precipitation were collected.

3.2 Proportion of Ambient vs. Tracer Hg in Outflow

Average ratios of tracer Hg to ambient THg outflow loads for each of the treatment combinations were calculated to assess the proportion of tracer Hg vs. ambient THg in outflow. All ratios were greater than one with the mean tracer Hg/ambient THg ratios for both the high moisture-high precipitation and high moisture-low precipitation treatments approximately 4 ± 1 and 4 ± 0.2 , respectively. The ratio for low moisture-low precipitation was approximately 9 ± 4 and the low moisture-high precipitation treatment had the highest ratio at 16 ± 6 . The tracer Hg/ambient THg ratio for the low moisture-high precipitation treatment was significantly greater

1
2
3 365 ($p = 0.0069$) than both of the high moisture treatments (high moisture-low precipitation and high
4
5 366 moisture- high precipitation). However, the ratio for this treatment was not statistically different
6
7
8 367 from the low moisture-low precipitation ratio.
9

10 368 A two-way ANOVA of the tracer Hg/ambient THg ratio across treatment combinations
11
12
13 369 revealed soil moisture to be a significant control on the proportion of tracer Hg mobilized
14
15 370 expressed as a ratio over ambient Hg ($p = 0.002$). Precipitation input depth exerted no
16
17 371 significant influence ($p = 0.26$) on the proportion of tracer Hg present in the outflow runoff.
18
19
20 372

21
22 373 **3.3 Tracer and Ambient Hg Loads by Hydrological Factor**
23

24 374 Under high soil moisture conditions the amount of both tracer and ambient Hg flushed
25
26
27 375 from the soil was nearly doubled ($\sim 1.7 - 1.8X$) however, only marginally significant for tracer
28
29 376 Hg ($p = 0.06$) when the precipitation depth was doubled (Table 2). With low moisture conditions
30
31 377 nearly three times ($\sim 2.7X$, $p < 0.01$) as much tracer Hg was mobilized in runoff with two times
32
33 378 as much precipitation input whereas the amount of liberated ambient Hg remained nearly
34
35
36 379 unchanged ($\sim 1.3X$ increase, $p = 0.35$).
37

38
39 380 With the high level of precipitation input, significantly more tracer Hg ($\sim 6X$, $p < 0.01$)
40
41 381 and approximately 1.5 times as much ambient Hg ($p = 0.02$) were flushed from the low moisture
42
43 382 soil as compared to the soil with high antecedent moisture (Table 2). Low soil moisture soil
44
45 383 cores subjected to low levels of precipitation input mobilized significantly more tracer Hg ($\sim 4X$,
46
47 384 $p < 0.001$) and highly variable, but on average twice as much ambient Hg ($p = 0.25$) as compared
48
49
50 385 to those under high antecedent moisture conditions.
51
52
53 386

3.4 Relationships among Outflow Solutes – Tracer ^{199}Hg , Ambient Hg, TOC

A significant correlation ($p < 0.0001$) existed between the tracer Hg and ambient THg outflow concentrations with all samples included from all replicates within all experimental combinations (Figure 3a). However, only the low soil moisture treatments, low moisture-low precipitation and low moisture-high precipitation, displayed significant correlations ($p < 0.01$ and 0.0001 , respectively) individually between the tracer Hg and ambient THg outflow concentrations (Figure 3b). Neither of the high moisture treatments exhibited significant relationships between these two variables.

A similar trend was observed for the relationship between tracer Hg and TOC concentrations in outflow. A significant, but weak correlation ($p < 0.0001$, $R^2 = 0.11$) existed between these variables when considering all outflow samples across all treatments (Figure 3c). The strength of the relationships improved for each of the low moisture treatments individually ($p < 0.01$ for both low moisture-high precipitation ($R^2 = 0.23$) and low moisture-low precipitation ($R^2 = 0.34$)) (Figure 3d). No significant relationship existed however between tracer Hg and TOC for either of the high soil moisture treatments. For all treatments (overall (Figure 3e) and individually (Figure 3f)), significant correlations were observed between the outflow concentrations of ambient THg and TOC (all $p \leq 0.01$; see Figure 3 for Pearson correlation coefficients).

Significant positive relationships ($p < 0.0001$) between the ratio of A254/A365 and both the outflow ambient THg concentrations and tracer Hg concentrations were observed when considering all treatment combinations (Supplementary Figure A-4). However, only the high moisture-high input treatment exhibited a significant correlation individually. A significant correlation between tracer Hg and A254/A365 was also observed for the high moisture-low input

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

treatment, but not for ambient THg and A254/A365. No significant correlations were observed between either of the tracer or ambient Hg concentrations and SUVA at 254 nm.

3.5 Soil Depth Profiles

The majority of the added tracer Hg was sorbed within the surface 0-3 cm sections of soil across all experimental combinations. A similar trend in tracer Hg concentrations was observed across all treatments (Figure 4). There were no significant differences in tracer Hg concentrations when comparing the same soil depths (0-3, 3-6, 6-9, 9-12 and 12-15 cm) across all treatments including controls. The soil tracer Hg concentrations were significantly positively correlated ($p < 0.0001$; $R^2 = 0.53$) with the percentage of organic matter in the soil, as determined by loss on ignition.

4 Discussion

4.1 Hydrological Controls on Soil Mercury Mobility

Antecedent soil moisture and precipitation input depth were both instrumental in controlling tracer Hg mobility within the soil columns. As there was no observed interaction between these factors, it can be concluded that these hydrological variables act independently of one another. Ambient Hg was not significantly influenced by precipitation input. However, soil moisture did play a marginally statistically significant role; one that we expect may be strengthened with further experimental replication. The mobility of tracer Hg through soil and its presence in outflow was greatly enhanced by the depth of precipitation applied to the soil due to greater flushing. Field studies investigating hydrological transport of Hg at the watershed and hillslope scales have similarly observed increased Hg fluxes as a result of high-flow rain events

and spring snowmelt but differences in Hg concentrations have often been small (e.g. Bishop *et al.*, 1995; Haynes and Mitchell, 2012; Lee *et al.*, 2000; Scherbatskoy *et al.*, 1998). Given the observation of flow-controlled Hg export from terrestrial environments, we investigated only the influence of the amount of precipitation a soil is subjected to rather than the intensity with which precipitation is delivered. The effect of precipitation intensity on both tracer and ambient Hg mobility would be an important avenue for further study.

Although the use of intact soil cores allowed for the investigation of only vertical translocation of Hg through the soil profile, the results may also be used to infer hydrological controls on lateral transport of Hg for the study region. The upland environment in the area is underlain by a low permeability B horizon which acts as a confining layer by directing subsurface runoff as shallow interflow, effectively creating a perched water table (Mitchell *et al.* 2009). During precipitation events, vertical infiltration to the perched water table is the first step in the process of runoff generation before subsurface flow is directed downslope by the confining layer. Therefore, vertical translocation of Hg upon wetting up of the soil profile is important when considering Hg fluxes from upland environments to downstream systems.

The observed timing of tracer and ambient Hg release from these sandy loam soils is likely the result of simple piston flow (Rawlins *et al.* 1997). The chemistry at the onset of outflow from the higher moisture soils would most likely be proportionally dominated by the movement of stored moisture and then replaced proportionally by the tracer added above it over time as the wetting front progressed downward through the core. The fact that outflow ambient THg concentrations were highest initially and subsequently decreased upon further input and that outflow tracer THg concentrations were initially suppressed (Figure 2) support this. Since soil moisture was already low in the low-moisture treatments, the movement of the wetting front

1
2
3 456 through the core length and the chemistry of its outflow were dominated by the tracer.
4
5 457 Additionally, lag times between inputs and outflow were approximately twice as long for low
6
7
8 458 moisture treatments but much more variable than for the higher moisture treatments. Due to this
9
10 459 higher variability in lag time and early timing in the tracer Hg outflow peak it is possible that
11
12 460 preferential flowpaths directed the applied tracer through the soil core in some of our replicates.
13
14
15 461 When sampling the soil of the low moisture treatments, preferential flow pathways (e.g., cracks)
16
17 462 were observed based on the wetting patterns through the low moisture soil cores. The observed
18
19
20 463 piston flow mechanism responsible for the mobility of Hg through these sandy loam soil
21
22 464 columns may not be the governing mechanism of Hg mobility in other soil types such as
23
24 465 predominantly clay soil. This may influence the manner and rate by which Hg is flushed through
25
26
27 466 different soil types as preferential transport may be the more dominant mechanism in fine
28
29 467 textured soils (Weiler and Naef 2003), for example, particularly under low moisture conditions
30
31
32 468 when they are more prone to cracking.

33
34 469 The influence of antecedent soil moisture on the amount of ^{199}Hg tracer present in
35
36 470 outflow does not appear to be the result of dilution. Of the low input treatments, the low
37
38
39 471 moisture-low precipitation experiments yielded approximately four times as much ^{199}Hg tracer in
40
41 472 outflow than the high moisture-low precipitation experiments while for the high input treatments
42
43 473 approximately six times as much tracer Hg was flushed from the low moisture-high precipitation
44
45
46 474 than the high moisture-high precipitation experimental soils (Table 2). Considering that the
47
48 475 volume of water in the high moisture soils was approximately 1.8 times greater than that of the
49
50
51 476 low moisture soils, there is no evidence that the tracer was diluted by soil water.

52
53 477 Given the caveat that the one-time ^{199}Hg application in this experiment was much greater
54
55 478 than natural rates of atmospheric deposition, the applied tracer in this experiment may broadly
56
57
58
59
60

1
2
3 479 represent contemporary Hg stocks deposited in soil or potentially a point source spill or leak
4
5
6 480 from a concentrated source. The measured ambient soil and outflow water Hg may comparably
7
8 481 be indicative of legacy Hg stocks. If the applied tracer is representative of contemporary Hg, our
9
10 482 results may suggest that together antecedent soil moisture and precipitation, whether it is rain or
11
12 483 snowmelt, may be instrumental in controlling the provision of newly-deposited Hg to
13
14 484 downstream wetlands and water bodies; known to be sites of Hg methylation (Mitchell *et al.*,
15
16 485 2008b). Research under the METAALICUS project suggested that newly-deposited Hg was
17
18 486 more readily volatilized and methylated than old, legacy Hg as well as initially more mobile
19
20 487 following deposition (Hintelmann *et al.*, 2002). In the present study the proportion of tracer Hg
21
22 488 as compared to ambient THg present in outflow runoff was significantly affected by soil
23
24 489 moisture with more tracer Hg being flushed from low moisture soils. However, precipitation
25
26 490 input depth had no significant effect on the proportion of tracer vs. ambient Hg in runoff.
27
28 491 Hydrological conditions such as antecedent soil moisture content are key to controlling the
29
30 492 magnitude of tracer Hg in runoff, which if representative of contemporary Hg, may be more
31
32 493 bioavailable for uptake by aquatic organisms. Therefore, alterations to these hydrological factors
33
34 494 may have deleterious downstream consequences on MeHg production and biotic uptake under
35
36 495 potential climatic and other environmental changes.
37
38
39
40
41
42

43 496 Increased frequency of water table fluctuations may affect Hg mobility and transport by
44
45 497 both hydrological and biogeochemical processes. Our results suggest that greater flushing of soil
46
47 498 pores by high-flow events, particularly following periods of water table drawdown, may liberate
48
49 499 more Hg to downstream environments. The antecedent moisture condition of the soil may affect
50
51 500 the partitioning of the Hg between the colloidal and dissolved fractions (although not explicitly
52
53 501 investigated in this study) and therefore impact the phase of Hg mobilized following dry periods
54
55
56
57
58
59
60

1
2
3 502 as opposed to predominantly wet periods (Babiarz *et al.* 2003; Zhu *et al.* 2014). We recognize
4
5 503 that the amount of tracer Hg liberated may be minute in the field as compared to the amount of
6
7
8 504 ambient Hg mobilized given our results following the addition of a high concentration of ¹⁹⁹Hg
9
10 505 stable isotope to the soil cores. However, given the result of significant precipitation and
11
12 506 particularly soil moisture controls on the mobility of tracer Hg, as well as marginally for ambient
13
14
15 507 Hg, changes in hydrological conditions causing drier soils may have implications for the
16
17 508 enhanced mobilization and transport of Hg to downstream aquatic ecosystems. One potential
18
19 509 limitation of the high level of tracer addition in this study may involve the exceedance of typical
20
21 510 binding sites in the soil organic matter. Reduced sulphur binding sites are the dominant sites of
22
23 511 Hg sorption in organic matter and typically far exceed the amount of Hg available in natural
24
25 512 systems (Ravichandran 2004). These sites may have been exceeded, although the concentration
26
27 513 of high quality binding sites in these soils is not known, due to the high concentration of tracer
28
29 514 Hg applied to each core and therefore may have affected the phase partitioning of the mobilized
30
31
32
33
34 515 Hg.
35
36
37
38

39 517 **4.2 Total Organic Carbon and Mercury Transport**

40
41 518 Numerous field studies have observed Hg transport associated with that of dissolved
42
43 519 organic carbon (e.g. Bishop *et al.*, 1995; Dittman *et al.*, 2010; Schuster *et al.*, 2008). The
44
45 520 tendency of Hg to form complexes with organic matter (Driscoll *et al.*, 1995; Schuster, 1991)
46
47 521 may act in facilitating the preferential mobilization of Hg during high-flow events as a result of
48
49 522 phase partitioning of the Hg into a more-readily mobilized phase. The outflow concentrations of
50
51 523 ambient THg were significantly correlated to those of TOC for all four experimental
52
53
54 524 combinations; both individually and collectively. Ambient Hg which has been in the soil for a
55
56
57
58
59
60

longer duration than the applied ^{199}Hg tracer may have equilibrated with the dissolved organic matter over this period. Through the formation of complexes with TOC, the mobility of ambient Hg may be enhanced regardless of the hydrological conditions of the soil. The formation of TOC-Hg complexes may account for the similarity in the ambient THg-TOC relationships as well as the overall ambient Hg outflow loads across the four experimental combinations.

Complexation with TOC in soil may also be important when considering the mobility of newly-applied, tracer Hg. Hintelmann *et al.* (2002) proposed that the initial mobile fraction of newly-deposited Hg flushed from the system during rain events was bound to DOC. Similar laboratory studies have also found that the mobility of added Hg is facilitated by the complexation of Hg with dissolved organic matter (Schlüter, 1996). In the present study we observed the outflow concentrations of tracer Hg to be significantly, albeit weakly, correlated to those of TOC and ambient THg only under low moisture conditions (Figure 3). These observations suggest that tracer Hg is being mobilized from drier soils in a similar manner to that of ambient Hg. Therefore tracer Hg may be mobilized as a result of complex formation with TOC. The drier conditions of the low moisture soils may allow for more of the tracer to interact with the surface area of the organic matter, thereby facilitating greater TOC-tracer Hg complex formation. The relationship between tracer Hg and TOC in outflow runoff may also account for the greater proportion of tracer Hg being mobilized under low moisture conditions as more tracer ^{199}Hg may be partitioned into the mobile, TOC-bound phase. However, there was no significant relationship between tracer Hg and either of ambient THg or TOC for the soils under high moisture conditions. Both high soil moisture experiments (high moisture-low precipitation and high moisture-high precipitation) exhibited significant positive correlations between the tracer Hg concentrations and the absorbance ratios (A_{254}/A_{365}), from which it can be inferred that the

1
2
3 548 tracer Hg was potentially associated with TOC of lower molecular weight. It is therefore
4
5 549 possible that the tracer Hg being flushed from wet soils may be in a different form than that from
6
7
8 550 dry soils. The tracer Hg mobilized from low moisture soil may be found in the colloidal phase
9
10 551 bound to TOC while that from high moisture soils may instead be in the true dissolved fraction.
11
12 552 A study by Babiarz *et al.* (2003) provides evidence to suggest that the phase in which newly
13
14 553 deposited Hg as compared to ambient Hg is transported may differ due to the quality of the
15
16 554 carbon and its associated ligands. This differential phase partitioning according to carbon quality
17
18 555 in the colloidal and dissolved fractions may account for the observed differences in tracer and
19
20 556 ambient Hg mobility. Further study is required for a more detailed assessment of the molecular
21
22 557 makeup of the carbon present in the dissolved and colloidal fractions of outflow (as no
23
24 558 distinction between phases was made in this study) and its potential influences on the phase
25
26 559 partitioning of mobilized Hg. Future research could also focus upon the role of organic matter
27
28 560 decomposition on the mobility of tracer and ambient Hg as this may also influence the
29
30 561 partitioning of Hg into the mobile phase and the mobility of Hg through the soil profile as well as
31
32 562 laterally to downstream ecosystems.
33
34
35
36
37
38
39
40

41 564 **4.3 Distribution of ¹⁹⁹Hg Tracer in Soil Column**

42
43 565 Overall, across all treatments, less than 0.5% of the added Hg tracer was observed in
44
45 566 runoff with the majority being retained in the surface 3 cm section of soil. This result is similar
46
47 567 to that of the field study by Hintelmann *et al.* (2002), in which less than 1% of the added tracer
48
49 568 appeared in runoff over the course of one season. A significant portion of the newly-deposited
50
51 569 Hg was rapidly immobilized by equilibrating with the native pool in both soil and vegetation
52
53
54 570 (Hintelmann *et al.*, 2002). Following six years of tracer application at the same site as
55
56
57
58
59
60

Hintelmann *et al.* (2002) and one year after loading was ended, Oswald *et al.* (2014) were still able to observe approximately 22% of the mass of applied tracer in the top 15 cm of mineral soil, where concentrations decreased with increasing soil depth. Similarly, in the present study, the vast majority of the ^{199}Hg tracer was immobilized in the surface, organic horizons of the soil regardless of both the soil moisture and precipitation conditions. The tracer Hg concentrations decreased rapidly downward with only negligible ^{199}Hg tracer reaching the lower 9 cm of soil across all treatments. The pattern of tracer Hg distribution throughout the soil profiles may be attributed to sorption to soil organic matter as these variables were significantly correlated. Similar soil column studies using an applied Hg tracer concluded that the majority of the added Hg was sorbed near the soil surface with no considerable vertical translocation through the soil (Schlüter, 1996; Semu *et al.*, 1985). It has been proposed that a large proportion of newly-deposited Hg is not immediately mobile rather being initially bound to soil and vegetation (Hintelmann *et al.*, 2002). However, upon repeated flushing with multiple precipitation events following deposition, it is possible that the mobility of this new Hg pool may be altered as the hydrological conditions of the soil change. A similar field investigation by Munthe *et al.* (2001) using an applied ^{199}Hg tracer at the plot scale observed the isotope ratio of $^{202}\text{Hg}/^{199}\text{Hg}$ stabilized after a period of approximately three months. During an initial rainstorm following the application of the tracer, Hintelmann *et al.* (2002) observed a significant decrease in runoff tracer concentrations which was not observed for native Hg and was sustained throughout the rest of the season. Future research could focus on the potential effects of repeated precipitation events of varying magnitudes over soils of different antecedent moisture in a controlled manner in order to assess the mobility of an applied Hg tracer to such changes in soil hydrological conditions

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

593 over time. The role of temperature in influencing the mobility and cycling of Hg is also an
594 important factor to consider in future study.

595
596 **4.4 Conclusions and Implications**

597 Both the antecedent moisture condition of the soil and the precipitation input depth are
598 important controls on tracer Hg mobility in upland ecosystems. Soil moisture exerts significant
599 influence on the proportion of tracer Hg mobilized. Overall, mercury mobility is enhanced as a
600 result of high-flow events and from soils with low initial moisture content. Piston flow appears
601 to be the dominant process governing the mobilization of both tracer and ambient Hg through
602 these upland sandy loam soil cores. Preferential transport flowpaths and complex formation with
603 solutes such as TOC may also be important in controlling Hg fluxes, particularly when these
604 soils are dry. Despite significant differences in Hg mobility according to hydrological
605 treatments, the majority of added Hg was sorbed in the surface 3 cm of this sandy loam soil with
606 less than 0.5% of ¹⁹⁹Hg tracer appearing in runoff. Given that this result was observed following
607 the addition of a large amount of ¹⁹⁹Hg tracer, we acknowledge that the potential magnitude of
608 contemporary Hg mobilized in the field may be minimal given natural rates of atmospheric
609 deposition. However, this study does broaden our understanding of the influence of hydrological
610 conditions; particularly soil moisture on Hg mobility.

611 The dependence of Hg transport on hydrological factors may be important when
612 considering natural variability in hydrological processes (particularly during high-flow
613 precipitation and spring snowmelt events) as well as the potential impacts of climate change on
614 these processes. Further research on Hg mobility and transport in upland ecosystems is
615 warranted given the potential manifestation of changes in upland Hg export on downstream

processes such as Hg methylation in wetlands (Mitchell *et al.*, 2009) and subsequent biotic uptake of MeHg; particularly during times of greatest Hg export such as spring snowmelt (Bishop *et al.*, 1995; Lee *et al.*, 2000; Mitchell *et al.*, 2008a).

Performing studies of this nature at the scale of a soil column is an invaluable tool given the level of control and manipulation of key hydrological variables afforded by such a technique. The use of expandable spray foam insulation to encase intact soil cores allows for the preservation of soil structure without introducing wall effects or blockage of soil pores for hydrological purposes while also acting as a suitable material for use with an applied Hg tracer. Studies involving the use of undisturbed, intact soil cores also provide a more realistic and natural medium for investigation as opposed to an artificial or repacked soil. Scaling results from the level of a soil core to that of a hillslope or watershed presents limitations and may provide an oversimplification of the processes that could potentially influence Hg mobility in natural systems. However, studies at small scales such as this study provide valuable information and an important first step to a greater understanding of the processes involved at the hillslope or watershed scale. Given the level of heterogeneity at the landscape scale, investigations such as this study at smaller scales facilitate the determination of the relative controls of hydrological processes on contaminant transport in the environment, and broaden our understanding of the level of small-scale spatial variability therein.

5 Acknowledgements

Funding was provided through a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant and Great Lakes Air Deposition (GLAD) program grant to C.P.J.M. and a NSERC Canada Graduate Scholarship to K.M.H. We would like to acknowledge the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

laboratory assistance of P. Huang and S. Dhal. We also thank two anonymous reviewers for their constructive comments.

6 Supporting Information

An overview of a separate study that was conducted in order to test the feasibility of using foam-encased soil cores for the study of hydrological and mercury (Hg) biogeochemical processes is described in the Supplementary Information.

7 References

Babiarz CL, Hurley JP, Krabbenhoft DP, Gilmour C, Branfireun BA. 2003. Application of ultrafiltration and stable isotopic amendments to field studies of mercury partitioning to filterable carbon in lake water and overland runoff. *The Science of the Total Environment* **304** : 295-303.

Bagarello V, Elrick DE, Iovino M, Sgroi A. 2006. A laboratory analysis of falling head infiltration procedures for estimating the hydraulic conductivity of soils. *Geoderma*. **135** : 322-334.

Bagarello V, Sgroi A. 2008. Testing Soil Encasing Materials for Measuring Hydraulic Conductivity of a Sandy-Loam Soil by the Cube Methods. *Soil Sci. Soc. Am. J.* **72** : 1048-1057. DOI: 10.2136/sssaj2007.0022.

Berggren M, Laudon H, Jansson M. 2007. Landscape Regulation of Bacterial Growth Efficiency in Boreal Freshwaters. *Global Biogeochem. Cycles* **21** : GB4002. DOI: 10.1029/2006GB002844.

- 661 Bishop K, Lee Y-H, Pettersson C, Allard B. 1995. Methylmercury Output from the Svartberget
662 Catchment in Northern Sweden during Spring Flood. *Water Air Soil Pollut.* **80** : 445-454.
- 663 Björn E, Larsson T, Lambertsson L, Skyllberg U, Frech W. 2007. Recent Advances in Mercury
664 Speciation Analysis with Focus on Spectrometric Methods and Enriched Stable Isotope
665 Applications. *Ambio* **36** : 443-451.
- 666 Branfireun BA, Heyes A, Roulet NT. 1996. The Hydrology and Methylmercury Dynamics of a
667 Precambrian Shield Headwater Peatland. *Water Resour. Res.* **32** : 1785-1794.
- 668 Bushey JT, Driscoll CT, Mitchell MJ, Selvendiran P, Montesdeoca MR. 2008. Mercury transport
669 in response to storm events from a northern forest landscape, *Hydrol. Processes* **22** : 4813–
670 4826.
- 671 Cameron KC, Harrison DF, Smith NP, McLay CDA. 1990. A method to prevent edge-flow in
672 undisturbed soil cores and lysimeters. *Australian Journal of Soil Research.* **28** : 879-886.
673 DOI: 10.1071/SR9900879.
- 674 Czuba CR, Fallon JD, Kessler EW. 2012. Floods of June 2012 in northeastern Minnesota: U.S.
675 Geological Survey Scientific Investigations Report 2012-5283, 42 p. with 3 app.
- 676 Demers JD, Driscoll CT, Shanley JB. 2010. Mercury Mobilization and Episodic Stream
677 Acidification during Snowmelt: Role of Hydrologic Flow Paths, Source Areas, and Supply
678 of Dissolved Organic Carbon. *Water Resour. Res.* **46** : W01511. DOI:
679 10.1029/2008WR007021.
- 680 Dittman JA, Shanley JB, Driscoll CT, Aiken GR, Chalmers AT, Towse JE, Selvendiran P. 2010.
681 Mercury Dynamics in Relation to Dissolved Organic Carbon Concentration and Quality
682 During High Flow Events in Three Northeastern U.S. Streams. *Water Resour. Res.* **46** :
683 W07522. DOI: 10.1029/2009WR008351.

1
2
3 684 Driscoll CT, Blette V, Yan C, Schofield CL, Munson R, Holsapple J. 1995. The Role of
4
5 685 Dissolved Organic Carbon in the Chemistry and Bioavailability of Mercury in Remote
6
7 686 Adirondack Lakes. *Water Air Soil Pollut.* **80** : 499-508.
8
9
10 687 Driscoll CT, Han Y-J, Chen CY, Evers DC, Lambert KF, Holsen TM, Kamman NC, Munson
11
12 688 RK. 2007. Mercury Contamination in Forest and Freshwater Ecosystems in the
13
14 689 Northeastern United States. *BioScience* **57** : 17-28.
15
16
17 690 Fitzgerald WF, Engstrom DR, Mason RP, Nater EA. 1998. The Case for Atmospheric Mercury
18
19 691 Contamination in Remote Areas. *Environ. Sci. Technol.* **32** : 1-7.
20
21
22 692 Gabriel MC, Williamson DG. 2004. Principal Biogeochemical Factors Affecting the Speciation
23
24 693 and Transport of Mercury through the Terrestrial Environment. *Environ. Geochem. Health*
25
26 694 **26** : 421-434.
27
28
29 695 Grigal DF. 2003. Mercury Sequestration in Forests and Peatlands: A Review. *J. Environ. Qual.*
30
31 696 **32** : 393-405.
32
33
34 697 Harris RC, Rudd JWM, Amyot M, Babiarz CL, Beaty KG, Blanchfield PJ, et al. 2007. Whole-
35
36 698 ecosystem Study Shows Rapid Fish-mercury Response to Changes in Mercury Deposition.
37
38 699 *PNAS* **104** : 16586-16591.
39
40
41 700 Haynes KM, Mitchell CPJ. 2012. Inter-annual and Spatial Variability in Hillslope Runoff and
42
43 701 Mercury Flux during Spring Snowmelt. *J. Environ. Monit.* **14** : 2083-2091. DOI:
44
45 702 10.1039/C2EM30267E.
46
47
48 703 Hintelmann H, Harris R, Heyes A, Hurley JP, Kelly CA, Krabbenhoft DP, Lindberg S, Rudd
49
50 704 JWM, Scott KJ, St. Louis VL. 2002. Reactivity and Mobility of New and Old Mercury
51
52 705 Deposition in a Boreal Forest Ecosystem during the First Year of the METAALICUS
53
54 706 Study. *Environ. Sci. Technol.* **36** : 5034-5040.
55
56
57
58
59
60

- Hintelmann H, Ogrinc N. 2003. Determination of Stable Mercury Isotopes by ICP/MS and Their Application in Environmental Studies. In: Cai Y, Braids CO. (eds) *Biogeochemistry of environmentally important trace elements*. ACS Symp. Ser. Vol. **835**, Washington, DC, p. 321-338.
- Hoag RS, Price JS. 1997. The effects of matrix diffusion on solute transport and retardation in undisturbed peat in laboratory columns. *Journal of Contaminant Hydrology*. **28** : 193-205.
- Kim Y-J, Steenhuis TS, Nam K. 2008. Movement of Heavy Metals in Soil through Preferential Flow Paths under Differential Rainfall Intensities. *Clean* **36** : 984-989.
- Kolka RK, Sebestyen SD, Verry ES, Brooks KN. (eds.) 2011. Peatland Biogeochemistry and Watershed Hydrology at the Marcell Experimental Forest. CRC Press Taylor & Francis Group: Boca Raton.
- Lee YH, Bishop KH, Munthe J. 2000. Do Concepts About Catchment Cycling of Methylmercury and Mercury in Boreal Catchments Stand the Test of Time? Six Years of Atmospheric Inputs and Runoff Export at Svartberget, Northern Sweden. *Sci. Total Environ.* **260** : 11-20.
- Mergler D, Anderson HA, Chan LHM, Mahaffey KR, Murray M, Sakamoto M, Stern AH. 2007. Methylmercury Exposure and Health Effects in Humans: A Worldwide Concern. *Ambio* **36** : 3-11.
- Miretszky P, Bisinoti MC, Jardim WF. 2005. Sorption of mercury (II) in Amazon soils from column studies. *Chemosphere* **60** : 1583-1589.
- Mitchell CPJ, Branfireun BA, Kolka RK. 2008a. Total Mercury and Methylmercury Dynamics in Upland-Peatland Watersheds during Snowmelt. *Biogeochemistry* **90** : 225-241. DOI: 10.1007/s10533-008-9246-z.

1
2
3 730 Mitchell CPJ, Branfireun BA, Kolka RK. 2008b. Spatial Characteristics of Net Methylmercury
4
5 731 Production Hot Spots in Peatlands. *Environ. Sci. Technol.* **42** : 1010-1016.
6
7
8 732 Mitchell CPJ, Branfireun BA, Kolka RK. 2009. Methylmercury Dynamics at the Upland-
9
10 733 Peatland Interface: Topographic and Hydrogeochemical Controls. *Water Resour. Res.* **45** :
11
12 734 W02406. DOI: 10.1029/2008WR006832.
13
14
15 735 Munthe J, Lyvén B, Parkman H, Lee Y-H, Iverfeldt Å, Haraldsson C, Verta M, Porvari P. 2001.
16
17 736 Mobility and Methylation of Mercury in Forest Soils: Development of an In-Situ Stable
18
19 737 Isotope Tracer Technique and Initial Results. *Water Air Soil Pollut.: Focus* **1** : 385-393.
20
21
22 738 National Atmospheric Deposition Program Mercury Deposition Network, 2014,
23
24 739 <http://nadp.sws.uiuc.edu/data/MDN/annual.aspx>, accessed 18 December 2014.
25
26
27 740 Oswald CJ, Branfireun BA. 2014. Antecedent moisture conditions control mercury and dissolved
28
29 741 organic carbon concentration dynamics in a boreal headwater catchment. *Water Resour.*
30
31 742 *Res.* **50** : 6610-6627. DOI:10.1002/2013WR014736.
32
33
34 743 Oswald CJ, Heyes A, Branfireun BA. 2014. Fate and transport of ambient mercury and applied
35
36 744 mercury isotope in terrestrial upland soils: Insights from the METAALICUS watershed.
37
38 745 *Environ. Sci. Technol.* **48** : 1023–1031, DOI:10.1021/es404260f.
39
40
41 746 Phillips IR. 2001. Nitrogen and Phosphorus Transport in Soil Using Simulated Waterlogged
42
43 747 Conditions. *Commun. Soil Sci. Plant Anal.* **32** : 821-842.
44
45
46 748 Phillips I, Burton E. 2005. Nutrient Leaching in Undisturbed Cores of an Acidic Sandy Podzol
47
48 749 Following Simultaneous Potassium Chloride and Di-ammonium Phosphate Application.
49
50 750 *Nutrient Cycling in Agroecosystems* **73** : 1-14. DOI: 10.1007/s10705-005-6080-8.
51
52
53 751 Ravichandran M. 2004. Interactions between mercury and dissolved organic matter – a review.
54
55 752 *Chemosphere* **55** : 319-331.
56
57
58
59
60

- Rawlins BG, Baird AJ, Trudgill ST, Hornung M. 1997. Absence of preferential flow in the percolating waters of a coniferous forest soil. *Hydrological Processes* **11** : 575-585.
- Reinmann AB, Templer PH, Campbell JL. 2012. Severe Soil Frost Reduce Losses of Carbon and Nitrogen from the Forest Floor during Simulated Snowmelt: A Laboratory Experiment. *Soil Biol. Biochem.* **44** : 65-74.
- Scherbatskoy T, Shanley JB, Keeler GJ. 1998. Factors Controlling Mercury Transport in an Upland Forested Catchment. *Water Air Soil Pollut.* **105** : 427-438.
- Scheuhammer AM, Meyer MW, Sandheinrich MB, Murray MW. 2007. Effects of Environmental Methylmercury on the Health of Wild Birds, Mammals and Fish. *Ambio* **36** : 12-18.
- Schlüter K. 1996. Translocation of ^{203}Hg Labelled HgCl_2 and CH_3HgCl in an Iron-Humus Podzol Studied by Radio-analytical Techniques. *Z. Pflanzenernähr. Bodenk.* **159** : 215-226.
- Schlüter, K., Alstad, J., Seip, H.M. 1995. Mercury Translocation in and Evaporation from Soil. I. Soil Lysimeter Experiments with ^{203}Hg -Radiolabeled Compounds. *Journal of Soil Contamination*, 4, 327-353.
- Schuster E. 1991. The Behaviour of Mercury in the Soil with Special Emphasis on Complexation and Adsorption Processes – A Review of the Literature. *Water Air Soil Pollut.* **56** : 667-680.
- Schuster PF, Shanley JB, Marvin-Dipasquale M, Reddy MM, Aiken GR, Roth DA, Taylor HE, Krabbenhoft DP, DeWild JF. 2008. Mercury and Organic Carbon Dynamics during Runoff Episodes from a Northeastern USA Watershed. *Water Air Soil Pollut.* **187** : 89-108. DOI: 10.1007/s11270-007-9500-3.

1
2
3 775 Semu E, Singh BR, Steenberg K, Selmer-Olsen AR. 1985. Mobility of ²⁰³Hg-Labelled Mercuric
4
5 776 Chloride in Soil Columns. *Acta. Agric. Scand.* **35** : 129-135.
6
7
8 777 Sigua GC, Isensee AR, Sadeghi AM. 1993. Influence of rainfall intensity and crop residue on
9
10 778 leaching of atrazine through intact no-till soil cores. *Soil Sci.* **156** : 225-232.
11
12
13 779 Sunderland EM, Mason RP. 2007. Human impacts on open ocean mercury concentrations.
14
15 780 *Global Biogeochemical Cycles.* **21** : GB4022. DOI: 10.1029/2006GB002876.
16
17
18 781 Timmons DR, Verry ES, Burwell RE, Holt RF. 1977. Nutrient transport in surface runoff and
19
20 782 interflow from an aspen-birch forest. *J. Environ. Qual.* **6** : 188–192.
21
22
23 783 Vanderborght J, Ghäwiler P, Flühler H. 2002. Identification of Transport Processes in
24
25 784 Soil Cores using Fluorescent Tracers. *Soil Science Society of America Journal* **66** : 774-
26
27 785 787.
28
29
30 786 Vendelboe AL, Moldrup P, Heckrath G, Jin Y, Wollesen de Jonge L. 2011. Colloid and
31
32 787 Phosphorus Leaching from Undisturbed Soil Cores Sampled Along a Natural Clay
33
34 788 Gradient. *Soil Sci.* **176** : 399-406. DOI: 10.1097/SS.0b013e31822391bc.
35
36
37 789 Weed DAJ, Kanwar RS, Cambardella C, Moorman TB. 1998. Alachlor Dissipation in Shallow
38
39 790 Cropland Soil. *Journal of Environmental Quality* **27** : 767-776.
40
41
42 791 Weiler M, Naef F. 2003. An experimental tracer study of the role of macropores in infiltration in
43
44 792 grassland soils. *Hydrological Processes* **17** : 477-493.
45
46
47 793 Weishaar JL, Aiken GR, Bergamaschi BA, Fram MS, Fujii R, Mopper K. 2003. Evaluation of
48
49 794 Specific Ultraviolet Absorbance as an Indicator of the Chemical Composition and
50
51 795 Reactivity of Dissolved Organic Carbon. *Environ. Sci. Technol.* **37** : 4702-4708.
52
53
54 796 Yin Y, Allen HE, Huang CP, Sanders PF. 1997. Adsorption/Desorption Isotherms of Hg(II) by
55
56 797 Soil. *Soil Science* **162** : 35-45.
57
58
59
60

1
2
3 798 Zhu Y, Ma LQ, Dong X, Harris WG, Bonzongo JC, Han F. 2014. Ionic strength reduction and
4
5 799 flow interruption enhanced colloid-facilitated Hg transport in contaminated soils. *Journal*
6
7
8 800 *of Hazardous Materials* **264** : 286-292.
9
10 801
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Full factorial design with snowmelt input and antecedent soil moisture as the experimental variables; 3 replicate soil cores per treatment combination. Low and high treatment levels are summarized below.

		SNOWMELT INPUT		
		LOW	HIGH	NO INPUT/CONTROL
ANTECEDENT SOIL MOISTURE	LOW	n = 3	n = 3	n = 3
	HIGH	n = 3	n = 3	n = 3
Antecedent Soil Moisture:		Snowmelt Input:		
Low: similar to 65 mm available soil water in surface 45 cm soil, observed prior to winter 2010		Low: 50 mm (spring 2010 snowpack SWE)		
High: field capacity		High: 100 mm (spring 2011 snowpack SWE)		

Table 2 Tracer total ^{199}Hg and ambient THg outflow loads (ng) for all hydrological treatment combinations. Lower-case letter superscripts represent significant differences for tracer (total ^{199}Hg).

Treatment	Tracer THg Load (ng)	Ambient THg Load (ng) *
Low Moisture – Low Input	127 ± 12^a	17 ± 11
High Moisture – Low Input	31 ± 0.8^b	7.9 ± 0.5
High Moisture – High Input	55 ± 14^b	14 ± 1
Low Moisture – High Input	341 ± 100^c	22 ± 4

^{a,b,c} denote significant differences in tracer THg outflow loads among treatments

* No significant differences in ambient THg outflow loads among treatments

Figure 1 Depiction of procedure for experiments: a.) Following alteration of soil moisture, ¹⁹⁹Hg isotope (as ¹⁹⁹HgCl) applied to soil surface with Teflon sieve. b.) Immediately after isotope addition, foam-encased core secured to ring stand above acid-washed glass funnel. Precipitation input delivered to soil surface via peristaltic pump and rainfall simulator.

343x258mm (72 x 72 DPI)

Figure 2 Time course plots of tracer Hg and ambient THg concentrations as well as outflow rates for each of the four treatment combinations. One replicate, representative core was selected from each treatment. 167x103mm (300 x 300 DPI)

Figure 3 Scatterplots of correlations between outflow variables: a) tracer (total ¹⁹⁹Hg) concentrations vs. ambient THg concentrations including all treatments, b) tracer Hg vs. ambient with treatments designated, c) tracer Hg concentrations vs. DOC concentrations including all treatments, d) tracer Hg vs. DOC with treatments designated, e) ambient THg concentrations vs. DOC concentrations including all treatments, and f) ambient THg vs. DOC with treatments designated. *denote significant correlations.

144x74mm (300 x 300 DPI)

Figure 4 Soil depth profiles of tracer (total ^{199}Hg) concentrations (ng g^{-1}) for all treatment combinations including moisture controls: a) Low Moisture Control, b) High Moisture Control, c) Low Moisture Low Input, d) High Moisture Low Input, e) Low Moisture High Input, and f) High Moisture High Input. Error bars represent standard deviation of the three replicates per treatment.

133x84mm (300 x 300 DPI)