Canadian Journal of Zoology
Revue canadienne de zoologie

**Leeches (Annelida: Hirudinea) from the far north of Ontario:
distribution, diversity, and diagnostics**

Journal:	<i>Canadian Journal of Zoology</i>
Manuscript ID	cjz-2017-0078.R1
Manuscript Type:	Article
Date Submitted by the Author:	23-Aug-2017
Complete List of Authors:	Langer, Sarah; Trent University Vezensyi, Kathryn; Trent University de Carle, Danielle; University of Toronto Beresford, David; Trent University, Biology Kvist, Sebastian; Royal Ontario Museum, Department of Natural History; University of Toronto, Department of Ecology and Evolutionary Biology
Keyword:	HIRUDINEA < Taxon, Leeches, northern Ontario, species distribution, specimen records, checklist

SCHOLARONE™
Manuscripts

**Leeches (Annelida: Hirudinea) from the far north of Ontario: distribution, diversity, and
diagnostics**

Sarah V. Langer¹, Kathryn A. Vezsenyi¹, Danielle de Carle^{2,3}, David V. Beresford⁴, Sebastian
Kvist^{2,3,*}

¹ Environmental and Life Sciences Graduate Program, Trent University, 1600 West Bank Drive,
Peterborough Ontario, Canada, K9J 7B8, Canada

² Department of Natural History, Royal Ontario Museum, 100 Queen’s Park, Toronto, Ontario,
M5S 2C6, Canada

³ Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street,
Toronto, Ontario, M5S 3B2, Canada.

⁴ Biology Department, Trent University, 1600 West Bank Drive, Peterborough Ontario, Canada,
K9J 7B8, Canada.

*Corresponding author: E-mail: sebastian.kvist@utoronto.ca

Abstract

Leeches have a worldwide distribution, yet numerous geographical regions remain to be adequately surveyed. Here, we present leech species records for one of these regions: the far north of Ontario. This region is primarily wetland habitat and includes two of Ontario's three ecozones. Morphological examinations, and a single instance of successful DNA amplification with subsequent molecular identification, allowed us to identify representatives of 12 species from two predatory families (Erpobdellidae and Haemopidae) and one parasitic family (Glossiphoniidae) among samples of 130 individuals. To provide a more inclusive list of species records for this remote region, our data were also augmented by 25 largely unpublished collection records (for 102 individuals) from the Canadian Museum of Nature, which revealed the presence of an additional species. We comment on finds of particular interest in our sampling with comparison to relevant literature, and provide new distribution data for these species.

Key Words: Hirudinea, leeches, northern Ontario, species distribution, specimen records, checklist.

Introduction

Leeches (Annelida: Hirudinea) are a globally distributed, sparsely diverse, but highly abundant group of annelids. They have been recorded from every ocean and every continent, and inhabit freshwater, marine, and terrestrial ecosystems alike. They are defined by the absence of parapodia and chetae, unpaired genital openings on the clitellum, and suckers on both ends of their elongate bodies (Sket and Trontelj 2008). Only half of the roughly 680 known species represent parasitic forms, which feed either on the blood of vertebrates or haemolymph of invertebrates; others are predaceous, scavenging or macrophagous, feeding on smaller invertebrates (Sket and Trontelj 2008; Bielecki et al. 2011). Leeches are a primary food source for numerous freshwater and marine fishes within their habitats (Clady 1974; Davies and Kasserra 1989; Koperski 2006), making them important components of aquatic food webs (Sawyer 1986; Adamiak-Brud et al. 2016). Because predaceous leeches are often found in abundance across ecosystems, they may act as useful gauges of water pollution (Bendell and McNicol 1989; Macova et al. 2009). Beyond their use as bioindicators, leeches are also frequently used in modern authoritative medicine, as relievers of venous congestion following flap or digit replantation surgery (*e.g.* Dabb et al. 1992; Soucacos et al. 1994). One of the key innovations granting them medical utility is their production of highly potent anticoagulants that keep the blood flowing in and around the incision site during the extended feeding period (Kvist et al. 2013, 2014, 2016; Siddall et al. 2016).

Approximately 480 of the 680 described leech species are associated with freshwater, with the remaining species classified as marine or terrestrial (Sket and Trontelj 2008). About 80 freshwater and terrestrial species have been recorded from North America (Moser et al. 2006; Sket and Trontelj 2008; Kubova et al. 2013), and Davies (1973) acknowledged 42 of these

species as inhabiting Canada. Much like other cosmopolitan taxa, leech populations can shift in the face of ecological changes in climate, biota, or landscape. Variance in substrate, for example, can have a large impact on leech communities (Kubova et al. 2013): many species, like those in the family Glossiphoniidae, use inchworm-type locomotion, such that their movement is impeded by softer substrates (Adamiak-Brud et al. 2016). Environmental features such as rocks and submerged logs assist with this type of movement and act as shelter, protecting them from daylight, predators, and wave action (Adamiak-Brud et al. 2016); although, several leeches, *e.g.* members of Hirudinidae and Macrobdellidae, are capable swimmers. Tracking changes that might occur at a community level could be indicative of large-scale landscape alterations. Such endeavours are contingent on comprehensive distribution records covering extended periods of time. Hirudinea is just one of many taxa that have large gaps in knowledge regarding its detailed geographical distribution.

One such gap in our knowledge is the far north of Ontario, which covers 451,808 square kilometres (42% of the province; Marshall and Jones 2011) as defined in Ontario's Far North Act of 2010 (S.O. 2010, c.18 s.2). This area includes two of Ontario's three ecozones: the Hudson Bay Lowlands, and the Ontario Shield (Crins et al. 2009). Ecozones are characterized by physical processes that can be used to explain differences in the flora and fauna present (Crins et al. 2009). The Hudson Bay Lowlands ecozone contains the third largest wetland in the world, is the only area in Ontario influenced by oceanic currents, and has an area of continuous permafrost that extends along the Hudson Bay coastline (Crins et al. 2009). The Ontario Shield ecozone contains tens of thousands of lakes that range in size, depth, transparency, and shoreline complexity (Marshall and Jones 2011). The larger lakes have been surveyed for fish since the Patricia inventory in 1959 (Marshall and Jones 2011). Since then, extensive fieldwork has been

conducted in Ontario’s far north through the Ontario Ministry of Natural Resources and Forestry (OMNRF) Far North Biodiversity Project (FNBP). From 2009 to 2014, selected areas were surveyed using a variety of methods aimed at recording a diverse inventory of flora and fauna. The present study is based on these collection efforts and aims to provide insight into the composition of northern Ontario’s hirudofauna. The first of its kind, our research may serve as a starting point for future studies investigating changes to population distributions as a result of biotic or abiotic influences.

Methods

Leech specimens were collected in the far north of Ontario by the FNBP between 2010 and 2014. Importantly, leeches were not necessarily the target taxa for these collection efforts, which likely resulted in fewer organisms collected, as compared to more leech-centric, targeted approaches. Individuals were most often captured using aquatic net and trap surveys. Aquatic net surveys used D-nets to sample freshwater habitats at scheduled times throughout the day. Aquatic traps were galvanized steel minnow traps baited with dog food and a ball of tinfoil – intended to attract minnows – and left in the water for 24 hours before recovery. Four traps were placed at each locality. For detailed aquatic sampling methods from this project, see Goertz and Phoenix (2015). Two individual leeches were also caught in pitfall and pan trap surveys. Pitfall traps were constructed from 250 mL plastic cups, inserted into the ground and left for three days before recovery. Pan traps of blue, white, and yellow bowls were dug into shallow holes arranged in a cross, and were checked daily. Additionally, leeches were collected when observed outside of scheduled trapping methods, here referred to as incidental. Baited traps are often employed in leech-specific surveys to target sanguivorous and scavenging leech species, but this

may bias catches based on feeding habits (Bendell and McNicol 1991). The surveys conducted during this project were not intended to be leech-specific, allowing us the opportunity to sample leeches using a number of methods, but potentially limiting our catches for some species. For example, no large fish specimens were collected during the current project, which resulted in a complete absence of fish leeches (Piscicolidae) – it is likely that extended fishing efforts would reveal several piscicolid species in the region. All collected specimens were preserved in 70% denatured ethanol. Leeches were sorted from these samples and brought to the Royal Ontario Museum (ROM) for identification and long-term curation. Identification was done according to specialized literature (*e.g.* Davies 1971, 1973; Klemm 1985; Sawyer 1986) and examination of specimens was conducted using a Leica Wild M10 dissecting microscope (Leica Microsystems) fitted with a Spot Flex 15.2 64 megapixel camera (Spot Imaging). Several juvenile individuals proved refractory to morphological identification and, for five of these, molecular identification via DNA barcoding (see Hebert et al. 2003) using cytochrome *c* oxidase subunit I (COI) was attempted; for this purpose, DNA extraction, amplification and sequencing methods followed Phillips et al. (2010). Molecular identification relied both on a BLAST approach (Altschul et al. 1997), and on a phylogenetic approach using the parsimony criterion. All COI sequences pertaining to the genus *Haemopsis* were downloaded from GenBank on August 21, 2017 and these were then jointly aligned using the online version of MAFFT ver. 7 (Katoh and Standley 2013). A phylogenetic tree was then constructed using TNT (Goloboff et al. 2008); the analysis employed 100 iterations, 5 rounds of ratcheting and 3 rounds of fusing. Because the tree is intended to infer specimen identifications rather than relationships, the tree was mid-point rooted. Maps were produced with ArcMap ver. 10.2 (Esri, 2011) using open sources outlines from the OMNRF and an ecozone layer (Commission for Environmental Cooperation 2009).

Our ROM-curated records were augmented by largely unpublished records in the online database of the Canadian Museum of Nature (CMN). To increase the comprehensiveness of the present study, 102 individuals collected at 25 localities, and representing 10 different species, were included from the CMN database, only four of which have been previously published (see Hovingh 2006).

Results

A total of 130 individual leeches were collected between 2010 and 2014, representing three families: Erpobdellidae (n=51) (Fig. 1), Haemopidae (n=33) (Fig. 2), and Glossiphoniidae (n=46) (Fig. 3). Among these, 12 species were recorded, representing both predacious and parasitic taxa. Locations (including GPS coordinates), dates, and methods of capture for all specimens are listed in Table 1. Several specimens were damaged or juvenile, precluding successful morphological identification. DNA extraction and amplification was attempted for five individuals whose specific identities were unknown; however – likely due to long-term preservation in relatively low concentrations of ethanol (70%) – four of these attempts were unsuccessful. As a result, these four individuals are only identified to genus level, as are juvenile individuals and individuals in which the diagnostic features were obscured due to preservation methods. Each species is briefly presented below, with location of capture summarized by ecodistrict; this method of classifying the far north is described by Crins (2009). CMN records are accompanied by the location descriptions from the museum’s database.

Family Erpobdellidae

Erpobdella obscura (Verrill 1872)

Fig. 4A

Specimens possess 4 pairs of anterior eyes (2 anterior pairs and 2 posterior pairs); variable pigmentation, most often with relatively small, scattered black blotches on a lighter background; and 2 annuli between gonopores (in furrows between annuli). One specimen was opened to reveal a spirally-coiled atrial cornua. All of these characters suggest that the specimens belong to *Erpobdella obscura*. This species was one of the most abundant among the samples and individuals were collected from the ecodistricts Sandy Lake, Wunnummin Lake, Albany River, Winisk River, Dickey River, and Lower Sachigo River. In addition, three records (five individuals) of this species were available in the CMN database; these were collected from “between Mattic and Opazatika River”, Moose Factory, and “North Point, 29 km north of Moosonee”, respectively (Table 1).

Erpobdella punctata (Leidy 1870)

Figure 4B

Members of *Erpobdella punctata* exhibit remarkable variation in dorsal pigmentation patterns, but all possess 2 or 4 longitudinal rows of darkly pigmented dots; these dots are sometimes fused into laterally oriented black bars, which can occasionally be interpreted as continuous stripes of pigmentation. This species is moderate to large in size (up to 100 mm), has gonopores separated by 2 annuli (with large male pore), and 3 pairs of eyes. *Erpobdella punctata* is abundant throughout southern Canada (Davies 1973; Klemm 1985), but the northernmost limit of its distribution is still unclear. The three specimens were collected from three different ecodistricts: Lower Sachigo River, Wunnummin Lake, and Winisk River. Two additional records from south of Albany River and Moose Factory were present in the CMN database (Table 1).

170

171

Erpobdella parva (Moore 1912)

172

Figure 4C

173

174

175

176

177

178

Specimens possess gonopores separated by $3\frac{1}{2}$ annuli, with the male gonopore elevated on a ring, 4 pairs of eyes (2 posterior pairs of labial eyes behind the anterior 2 pairs), uniform dorsal colouration and absence of a mid-dorsal stripe. Collectively, these characters identify *Erpobdella parva*. Only a single individual of this species was collected from the Wunnummin Lake ecodistrict and a single record from the Winisk tributary existed in the CMN database (Table 1).

179

180

Erpobdella dubia (Moore and Meyer 1951)

181

Fig. 4D

182

183

184

185

186

187

188

189

Members of *Erpobdella dubia* possess a dark, mid-dorsal, longitudinal stripe that immediately sets them apart from most other species of *Erpobdella*. In addition, the dorsum is often greenish in colour and heavily mottled, gonopores are normally separated by $3\frac{1}{2}$ annuli ($2\frac{1}{2}$ and 3 annuli have also been reported [Klemm 1985]), and 4 pairs of eyespots are present (2 posterior pairs of labial eyes behind the anterior 2 pairs). A single specimen of *Erpobdella dubia* was collected from the Lower Sachigo River ecodistrict; one record was also available from the CMN database, collected from “narrows at source of Shell Brook” (Table 1).

190

Family Haemopidae

191

Haemopsis grandis (Verrill 1874)

192

Fig. 5A

Specimens of *Haemopsis grandis* are relatively large in size, and exhibit a thin margin of the oral sucker, no jaws, 12 pharyngeal folds, and black blotches of dorsal pigmentation (or entirely black background with beige or light green blotches), as is diagnostic for the species (Klemm 1985). A total of four individuals were collected from three different ecodistricts: Wunnummin Lake, Sandy Lake, and Lower Kenogami Lake. Two individuals were recorded from the CMN database, both collected from Fort Albany, 1.5 km west of Albany River (Table 1).

Haemopsis marmorata (Say 1824)

Fig. 5B

Specimens of *Haemopsis marmorata* is diagnosed by the relatively large body size, the presence of jaws with 12-16 teeth, a caudal sucker that was about half the width of the widest part of the body, and a uniformly grey dorsal colouration with moderate to heavy black, irregular blotching. Superficially, this species resembles *H. grandis* but is distinguished therefrom by the presence of jaws, and by the general size of the adult forms: *H. grandis* grows to be almost twice as long as *H. marmorata* (Klemm 1985). *Haemopsis marmorata* was the most abundant species among our samples, and was represented by 24 individuals from five different ecodistricts: Wunnummin Lake, Lower Sachigo River, Albany River, Winisk River, and Dickey River. Another four records for this species existed in the CMN database and these were collected from the Albany River, Moosonee District 3 km from Partridge River, west of Fort Severn and Moose Factory (see Table 1).

Haemopsis cf. lateromaculata Mathers 1963

Fig. 5C

A single individual presents morphological characters that suggest an affinity with *Haemopsis lateromaculata*. The specimen possesses a green dorsum with heavy dark blotching and few scattered yellow blotches, a darker venter with few indistinct dark or yellow pigmentation blotches, a caudal sucker broadly attached by a pedicel, and jaws with 10-12 teeth. This individual was collected from the Wunnummin Lake ecodistrict (Table 1). In addition, four records exist in the CMN database [identity adopted from Hovingh (2006), the database lists one record without a species identification, and the others as *H. marmorata*]. These were collected from “North-northeast of east end of Anderson Island in Albany River”, “between Mattis and Opazatika River Missinabi River”, “North end of lake near outlet Severn Lake” and “between new post M.B.C. and Moose River Abitibi River” (see Table 1). Because of this species’ resemblance to *H. marmorata*, this specimen was subjected to DNA sequencing to enable robust molecular identification via DNA barcoding of the COI locus (GenBank accession number MF706261). The resulting sequence was BLASTed against GenBank’s non-redundant (nr.) nucleotide sequence database using a BLASTn protocol. The closest match recovered was to *Haemopsis terrestris* (Forbes 1890), but these sequences still showed 14% difference. When constraining the BLAST search against *H. lateromaculata* (only a single COI sequence available on GenBank), the sequences showed 18% dissimilarity. Moreover, in the tree resulting from the phylogenetic analysis (Supplementary Figure S1) of the COI region, the new specimen was located on a long branch, and did not nest closely with other species, suggesting that this taxon is not otherwise represented in the dataset. The identification of our specimen as *Haemopsis* cf. *lateromaculata* should therefore be viewed with appropriate caution – it could be that this

specimen represents undescribed diversity or, alternatively, that the COI sequence for *H. lateromaculata* in GenBank is mislabelled.

Family Glossiphoniidae

Helobdella stagnalis (Linnaeus 1758)

Figure 6A

Specimens are small (often less than 1 cm); possess a dorsal, nuchal, chitinous scute on somite VIII; and are light grey or beige in color. Much like *Haemopsis marmorata*, the abundance of *Helobdella stagnalis* was relatively high compared to that of most other species: 23 individuals were collected from four different ecodistricts (Wunnumin Lake, Sandy Lake, Albany River and Lower Sachigo River; and, two records (6 adults and 71 juveniles) for this species are present in the CMN database, both of which were collected from the Attawapiskat River (Table 1). The taxonomic history of this species and its constituents is rather muddled. Historically, individuals bearing a distinct nuchal scute have all been identified as *H. stagnalis* but, concurrently with our growing knowledge of species boundaries, several other nuchal scute-bearing species have been described (*e.g.* Siddall 2001; Ocegüera-Figueroa et al. 2010; Salas-Montiel et al. 2014; Beresic-Perrins et al. 2017), some with geographic distributions overlapping that of *H. stagnalis*. Indeed, the boundaries between nuchal scute-bearing species are obscured, and a thorough revision of the genus is direly needed. All of the specimens examined here show morphological characters that are compatible with those of *H. stagnalis*.

Glossiphonia elegans (Verrill 1872) Castle 1900

Fig. 6B

Our specimens of *Glossiphonia elegans* possess 3 pairs of eyes arranged in parallel (*i.e.*, not in an arch-like manner), a pair of dark paramedial stripes dorsally and ventrally, interrupted by metameric white spots (sometimes lacking from juvenile individuals), and an opaque body with beige to green or gray colouration. Molecular data have suggested that the North American form of the species commonly identified as *Glossiphonia complanata* (Linnaeus 1758) may not be conspecific with its European counterpart (Siddall et al. 2005). As a result, Siddall et al. (2005) resurrected the specific epithet *Glossiphonia elegans* to represent the North American form. The specimens were collected from the ecodistricts of Sandy Lake, Albany River, Dickey River, Wunnummin Lake and Lower Sachigo River; in addition, four records for this species exist in the CMN database, and these were collected from Moose Factory, “Winisk 100 metres from east bank of river, 14 km upstream from village”, “10 km west of Attawapiskat Monument Channel” and at outlet of Stout Lake (Table 1).

Placobdella rugosa (Verrill 1874)

Fig. 6C

The single specimen of *Placobdella rugosa* possesses heavy dorsal papillation with 5 rows of more prominent papillae, lateral margins with alternating light and dark pigmentation (in the order of light, light, dark, light, light, dark, etc.) and scattered dark chromatophores on the ventral and anterodorsal surfaces. The identification of members of *Placobdella*, particularly the papillated forms, is fraught with difficulty, and the taxonomy of *Placobdella rugosa* is no exception. The species was originally described as a subspecies of *Placobdella ornata* (Verrill 1872), subsequently elevated to full species status, reassigned to *P. ornata*, and finally resurrected in 2012 (Moser et al. 2012, and references therein). The species is abundant across

Canada and is associated with varying climates and faunas (Klemm 1982, de Carle et al. 2017).

This particular specimen was collected from the Sandy Lake ecodistrict (Table 1).

Placobdella montifera Moore 1906

Whereas the newly collected material did not include individuals pertaining to *Placobdella montifera*, the collections from CMN do include one record. This species occurs in Ontario and westward across Canada (Davies 1973), and can be distinguished readily by its 3 prominent pointed keels or ridges on the dorsal surface – no other species of *Placobdella* possess this character – and its constricted neck region, resulting in a discoid head. This specimen was collected from Central Patricia Crow River (Table 1).

Theromyzon rude (Baird 1869)

Fig. 6D

A single specimen of *Theromyzon rude* was collected from the Sandy Lake ecodistrict (no species records were available from the CMN database; Table 1). It possesses 2 annuli between gonopores, a gelatinous and slightly transparent body with 2 paramedial rows of brown spots on the dorsum. Members of this genus are typically distinguished from congeners by the number of annuli between gonopores, but Baird (1869) failed to note the number of annuli in his original description of *T. rude*. Based on material collected from the type locality, Moore and Meyer (1951) and Meyer and Moore (1954) re-described the species and noted 3 annuli between gonopores. Later, the original type material was re-examined by Oosthuizen and Davies (1992) who noted that only 2 annuli separated the gonopores; the confused species, with 3 annuli between gonopores, was subsequently described by Davies and Oosthuizen (1993). As a result,

much uncertainty has surrounded the taxonomy of the species. The problem was exacerbated by the resulting difficulties in separation of *T. rude sensu lato* and *Theromyzon biannulatum* Klemm 1977, both of which possess 2 annuli between gonopores. Notwithstanding that our specimen is compatible with the morphology of *T. rude*, we concede that a taxonomic revision of the group, including molecular techniques, is needed before robustly inferring the identity of any specimen of *Theromyzon* with 2 annuli between gonopores.

Theromyzon tessulatum (Müller 1774)

Fig. 6E

Specimens display 4 annuli between gonopores and gelatinous bodies with 2 paramedial lines on the dorsum. Four leech individuals (one adult and a brood of three juveniles) were collected in the Lower Sachigo River ecodistrict (no specimen records were available in the CMN database; Table 1). As opposed to the congeneric *T. rude*, this species is distinct and therefore more easily identified by virtue of the 4 annuli separating the gonopores. This species was thought to be confined to Europe but it is now understood that its distribution is transatlantic (Pawlowski 1948; Herrman 1970; Klemm 1977, 1982; Davies 1973). The origin of this distribution is indeed intriguing, seeing as this species is often encountered attached to, and feeding from, waterfowl, which may explain its spread and current range.

Discussion

Of the 42 species of leeches that were recorded in the last comprehensive study of Canada’s hirudofauna (Davies 1973), 33 are non-piscicolid taxa. Members of Piscicolidae are almost exclusively encountered attached to their fish hosts, but the remaining species can

perhaps be more easily collected from rocks and debris. Of these 33 species, 25 have previously been recorded from Ontario (Davies 1973). In total, 12 species were recovered from our newly collected samples, and one additional species was recorded in the database of the Canadian Museum of Nature. It is remarkable that, despite the harsh winter climates in northern Ontario (including the area of continuous permafrost on the northern coast), 52% of Ontario's known non-piscicolid leech diversity is also recovered from the far north of the province. Taken together, the collection sites cover a substantial part of this region and, as far as the authors can tell, eight of the twelve species records are first reported in the present study (see Hovingh 2006). Some of the species recovered herein are more or less ubiquitous across Canada, but some represent taxa that are only infrequently recovered. Below, we discuss records of particular interest and provide a general discussion on species distributions across North America in general, and Ontario in particular.

Members of Erpobdellidae are considered widespread and some species, *e.g. Erpobdella punctata*, have been described as the most commonly encountered and widely distributed leeches in North America (Sawyer 1970; Klemm 1985; Kutschera and Wirtz 2001), though only three individuals were present among the newly collected samples. In contrast, other species within the genus, such as *Erpobdella parva*, have been recorded relatively infrequently (Klemm 1982); this species is represented by a single specimen among our samples. Given their abundance throughout North America, it is likely that species of *Erpobdella* play important roles in the ecosystems they inhabit, and it is already known that they represent a significant source of food for several freshwater fish species (*e.g.* Sinha and Jones 1967; Clady 1974; Davies and Kasserra 1989; Koperski 2006).

Of the haemopid species recovered, *Haemopsis grandis* and *H. marmorata* are considered common and widely distributed in North America, with *H. grandis* being especially well represented in the Great Lakes region (Klemm 1982). *Haemopsis lateromaculata* has more recently been described as widely distributed in North America (Hovingh 2006). Members of the genus are known to be amphibious (Madill and Hovingh 2007), yet only *Haemopsis terrestris* (Forbes 1890) is described as fully terrestrial. It is worth noting, however, that the distinction between amphibious and terrestrial is sometimes difficult to accurately identify: many primarily aquatic leeches of the order Arhynchobdellida are known to spend time foraging and depositing cocoons on land immediately surrounding freshwater habitats (Siddall and Bureson 1996). Interestingly, two specimens of an unknown species of *Haemopsis* were collected in terrestrial traps, but these specimens do not conform to the morphological description of *Haemopsis terrestris*. The specimens lack a median longitudinal stripe and possess jaws with distichous teeth and a relatively narrow caudal sucker, suggesting that they belong to *Haemopsis marmorata*. However, because of the terrestrial nature of these collecting events, we have refrained from providing a specific epithet for these individuals. Freshly collected material from which DNA can be extracted and successfully amplified will likely shed light on the identity of these unknown leeches.

Both *Erpobdella* and *Haemopsis* are predacious, with prey items ranging from oligochaetes and insect larvae, to snails, small crustaceans, and dead fish (Sawyer 1972, Barton and Metcalfe 1986; Kutschera and Wirtz 2001; Moser et al. 2006). Moreover, previous expeditions have successfully collected representatives of both genera using aquatic traps baited with beef liver and ground beef (de Carle et al. 2017). Predacious leeches swallow their prey either whole or in pieces (Kutschera and Wirtz 2001; Pfeiffer et al. 2005; Sket and Trontelj

2008). For the present study, one specimen of a *Haemopsis* species was caught in an aquatic trap while feeding on an adult predaceous diving beetle (Dytiscidae: *Dytiscus dauricus*) (Fig. 7). A similar observation was made by Bielecki *et al.* (2011), wherein *Haemopsis sanguisuga* was collected in the act of feeding on a predaceous diving beetle larva (*Dytiscus marginalis*). Species of *Haemopsis* are opportunistic feeders, and are frequently found with exoskeletons of larval aquatic insects in their digestive tracts (Moser *et al.* 2006).

All of the newly collected glossiphoniid species are considered common and widespread (Klemm 1982). *Glossiphonia complanata* (the European counterpart to the North American *G. elegans*; Siddall *et al.* 2005) and *Helobdella stagnalis* are considered some of the most widely distributed species in Europe (Kutschera and Wirtz 2001) and are also commonly encountered across North America (Klemm 1985). Although *Theromyzon* is considered holarctic (Mason *et al.* 2005), members of this genus are less frequently encountered than other glossiphoniids (Klemm 1982). This may be a function of their particular host affinity: members of *Theromyzon* have a propensity to feed on birds (Mason *et al.* 2005), an association which may also explain why they are so widely distributed across the landscape (Sket and Trontelj 2008).

Glossiphoniids are unique among leeches, in that they exhibit a high degree of parental care (Siddall *et al.* 2005). Leeches of this family produce membranous cocoons – unlike the hardened or spongy cocoons of other leeches – which are attached to the venter of the parent, and carried until the offspring hatch. After hatching, the juvenile leeches remain attached to their parent until being transported to their first meal (Siddall *et al.* 2005). Members of Glossiphoniidae mainly employ a parasitic lifestyle, feeding on the blood of vertebrates or the haemolymph of invertebrates (Siddall *et al.* 2005), though some may also eat small prey (Sket and Trontelj 2008). Species of *Placobdella* are primarily ectoparasitic on aquatic reptiles and

amphibians, though some are known to feed on fish, and many will feed opportunistically on other vertebrates such as birds and humans (Sawyer 1972; Jones and Woo 1990; Siddall and Gaffney 2004; Siddall and Bowerman 2006; Moser et al. 2014). Oddly, only a single specimen of *Placobdella rugosa* was newly collected and one additional record of *Placobdella montifera* was available in the CMN database. This paucity of *Placobdella* diversity and abundance starkly contrasts that of the southern parts of the province. Indeed, 8 species of *Placobdella* are currently known from Ontario (Davies 1973; Klemm 1985; de Carle et al. 2017), some of which are commonly encountered in high densities (de Carle et al. 2017).

Each of the collected families shows widespread distribution in the far north, and is seemingly not limited by ecozone – including the area of continuous permafrost on Ontario’s northern coast. By and large, the diversity of leeches recorded from the far north of Ontario is lower than the diversity recovered in the southern regions (possibly as a result of more leech-directed research in southern Ontario), and the species composition of natural populations seems to be somewhat different between the two regions. This may reflect biotic or abiotic barriers to dispersal that are difficult to conquer. Future studies of leech species distributions across Canada that incorporate diversity from northern provinces and territories as well as northern regions of southern provinces, will likely evince the mechanisms that have produced distributions among these economically and ecologically influential organisms.

Acknowledgments

The authors thank the Ontario Ministry of Natural Resources and Forestry’s Northeast Science and Information Section and Wildlife Research and Development Section for project coordination and logistics, and Far North Branch for funding of the Far North Biodiversity

Project. We give special thanks to Dean Phoenix and the field crews of the Far North Biodiversity Project. We also thank the First Nations communities of Peawanuck, Marten Falls, Webequie, Nibinamik, Kitchenuhmaykoosib Inninuwug, Keewaywin, Fort Albany, and Fort Severn for their hospitality and support for working on their traditional lands. We thank the field crews who collected and provided the samples used in this paper. Finally we thank Sherri DeGasparro for providing the identification on our dytiscid beetle and Don Stacey and Shannon Asencio for providing technical assistance. Peter Trontelj (University of Ljubljana) provided excellent feedback that greatly improved an earlier version of the manuscript. SK was funded by an NSERC discovery grant and a scholarship from the Olle Engkvist Byggmästare Foundation.

References

- Adamiak-Brud, Ź., Bielecki, A., Kobak, J., and Jabłońska-Barna, I. 2016. Rate of short-term colonization and distribution of leeches (Clitellata: Hirudinida) on artificial substrates. *J. Zool. (Lond.)* 299: 191-201.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25: 3389-3402.
- Baird, W. 1869. Descriptions of some new sectorial Annelides in the collection of the British Museum. *Proc. Zool. Soc. Lond.* 1869: 310-318.
- Barton, D.R., and Metcalfe, J.L. 1986. Life cycles, reproduction, and diets of *Dina dubia* and *Erpobdella punctata* (Hirudinea: Erpobdellidae) in Canagagigue Creek, Ontario. *Can. J. Zool.* 64: 640-648.

- 443 Bendell, B.E., and McNicol, D.K. 1991. An assessment of leeches (Hirudinea) as indicators of
444 lake acidification. *Can. J. Zool.* 69: 130-3.
- 445 Beresic-Perrins, R.K., Govedich, F.R., Banister, K., Bain, B.A., Rose, D., and Shuster, S.M.,
446 2017. *Helobdella blinni* sp. n. (Hirudinida, Glossiphoniidae) a new species inhabiting
447 Montezuma Well, Arizona, USA. *ZooKeys*, 661: 137-155.
- 448 Bielecki, A., Cichocka, J.M., Jeleń, I., Świątek, P., and Adamiak-Brud, Ż. 2011. A checklist of
449 leech species from Poland. *Wiadomości Parazytologiczne*, 57: 11-20.
- 450 Clady, M.D. 1974. Food habits of yellow perch, smallmouth bass and largemouth bass in two
451 unproductive lakes in northern Michigan. *Am. Midl. Nat.* 91: 453-459.
- 452 Commission for Environmental Cooperation. 2009. Ecological regions of North America
453 geospatial data presentation form: vector digital data. Montreal, Qu: Commission for
454 environmental cooperation. Digital Media.
- 455 Crins, W.J., Gray, P.A., Uhlig, P.W.C., and Wester, M.C. 2009. The Ecosystems of Ontario, Part
456 I: Ecozones and Ecoregions. Science and Information Branch, Inventory, Monitoring and
457 Assessment, Ontario Ministry of Natural Resources, Peterborough Ontario.
- 458 Dabb, R.W., Malone, J.M., and Leveret, L.C. 1992. The use of medicinal leeches in the salvage
459 of flaps with venous congestion. *Ann. Plast. Surg.* 29: 250-256.
- 460 Davies, R.W. 1971. A key to the freshwater Hirudinoidea of Canada. *J. Fish. Res. Board Can.*
461 28: 543-552.
- 462 Davies, R.W. 1973. The geographic distribution of freshwater Hirudinoidea in Canada. *Can. J.*
463 *Zool.* 51: 531-545.
- 464 Davies, R.W., and Kasserra, C.E. 1989. Foraging activity of two species of predatory leeches
465 exposed to active and sedentary prey. *Oecologia*, 81: 329-334.

- 466 Davies, R.W., and Oosthuizen, J.H. 1993. A new species of duck leech from North America
467 formerly confused with *Theromyzon rude* (Rhynchobdellida: Glossiphoniidae). Can. J.
468 Zool. 71: 770-775.
- 469 de Carle, D., Ocegüera-Figueroa, A., Tessler, M., Siddall, M.E., and Kvist, S. 2017.
470 Phylogenetic analysis of *Placobdella* (Hirudinea: Rhynchobdellida: Glossiphoniidae)
471 with consideration of COI variation. Mol. Phylogenet. Evol. 114: 234-248.
- 472 Environmental systems research institute (ESRI). (2011). ArcGIS desktop version 10.2. USDA,
473 Redlands, CA.
- 474 Forbes, S.A. 1890. An american terrestrial leech. Am. Nat. 24: 646-649.
- 475 Goertz, D.E., and Phoenix, R.D. 2015. Range extensions for northern redbelly dace (*Chrosomus*
476 *eos*), fathead minnow (*Pimephales promelas*), and Iowa darter (*Etheostoma exile*) in
477 Ontario, Canada. Can. Field-Nat. 129: 237-9.
- 478 Goloboff, P.A., Farris, J.S., and Nixon, K.C. 2008. TNT, a free program for phylogenetic
479 analysis. Cladistics, 24: 774-786.
- 480 Herbert, P.D., Ratnasingham, S., and deWaard, J.R. 2003. Barcoding animal life: cytochrome c
481 oxidase subunit 1 divergences among closely related species. Proc. R. Soc. Lond. B. Biol.
482 Sci. (Supplemental) 270: S96-S99.
- 483 Herrman, S.J. 1970. Systematics, distribution, and ecology of Colorado Hirudinea. Am. Midl.
484 Nat. 83: 1-37.
- 485 Hovingh, P. 2006. The leech *Haemopsis lateromaculata* (Hirudinea: Haemopidae): its North
486 America distribution and additional notes on species description. Can. Field-Nat. 120:
487 443-451.

- 488 Jones, S.R.M., and Woo, P.T.K. 1990. Redescription of the leech *Desserobdella phalera* (Graf,
489 1899) n. comb. (Rhynchobdellida: Glossiphoniidae), with notes on its biology and
490 occurrence on fishes. Can. J. Zool. 68: 1951-1955.
- 491 Katoh, K., and Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7:
492 improvements in performance and usability. Mol. Biol. Evol. 30: 772-780.
- 493 Klemm, D.J. 1982. Leeches (Annelida: Hirudinea) of North America. U.S. Environmental
494 Protection Agency: Office of research and development, Cincinnati Ohio. 1-177.
- 495 Klemm, D.J. 1985. Freshwater leeches (Annelida: Hirudinea). A guide to the freshwater
496 Annelida (Polychaeta, Naidid and Tubificid Oligochaeta, and Hirudinea) of North
497 America. Kendall/Hunt Publishing Co., Dubuque.
- 498 Koperski, P. 2006. Relative importance of factors determining diversity and composition of
499 freshwater leech assemblages (Hirudinea; Clitellata): a meta analysis. Arch. Hydrobiol.
500 166: 325-341.
- 501 Kubová, N., Schenková, J., and Horsák, M. 2013. Environmental determinants of leech
502 assemblage patterns in lotic and lenitic habitats. Limnologica, 43: 516-524.
- 503 Kutschera, U., and Wirtz, P. 2001. The evolution of parental care in freshwater leeches. Theory
504 in Biosciences, 120: 115-37.
- 505 Kvist, S., Gi-Sik, M., and Siddall, M.E. 2013. Diversity and selective pressures of anticoagulants
506 in three medicinal leeches (Hirudinida: Hirdinidae, Macrobdellidae). Ecol. Evol. 3: 918-
507 933.
- 508 Kvist, S., Brugler, M.R., Goh, T.G., Giribet, G., and Siddall, M.E. 2014. Pyrosequencing the
509 salivary transcriptome of *Haemadipsa interrupta* (Annelida: Clitellata: Haemadipsidae):

- 510 anticoagulant diversity and insight into the evolution of anticoagulation capabilities in
511 leeches. *Invertebr. Biol.* 133:74-98.
- 512 Kvist, S., Ocegüera-Figueroa, A., Tessler, M., Jiménez-Armenta, J., Freeman Jr., R.M., Giribet,
513 G., and Siddall, M.E. 2016. When predator becomes prey: investigating the salivary
514 transcriptome of the shark-feeding leech *Pontobdella macrothela* (Hirudinea:
515 Piscicolidae). *Zool. J. Linn. Soc.* 179: 725-737.
- 516 Macova, S., Harustiaková, D., Kolarová, J., Machová, J., Zlabek, V., Vykusová, B., Randák, T.,
517 Velisek, J., Poleszczuk, G., Hajslová, J., Pulkrabová, J., and Svobodová, Z. 2009.
518 Leeches as sensor-bioindicators of river contamination by PCBs. *Sensors*, 9: 1807-1820.
- 519 Madill, J., and Hovingh, P. 2007. Freshwater leech (Annelida: Hirudinida) distribution in the
520 Canadian Province of Newfoundland and Labrador and adjacent regions: check-list, new
521 records, new pigmentation forms, and Pleistocene refugia. *Zootaxa*, 1657: 1-21.
- 522 Marshall, T.R., and Jones, N.E. 2011. Aquatic Ecosystems of the Far North of Ontario: State of
523 Knowledge. Ontario Legislative Library, Toronto, Ontario, Canada.
- 524 Mason, T.A., Sayers, C.W., Paulson, T.L., Coleman, J., and Shain, D.H. 2005. Cocoon
525 deposition and hatching in the aquatic leech, *Theromyzon tessulatum* (Annelida,
526 Hirudinea, Glossiphoniidae). *Am. Midl. Nat.* 154: 78-87.
- 527 Meyer, M.C., and Moore, J.P. 1954. Notes on Canadian leeches (Hirudinea), with the description
528 of a new species. *Wasmann J. Biol.* 12: 63-96.
- 529 Moore, J.P., and Meyer, M.C. 1951. Leeches (Hirudinea) from Alaskan and adjacent waters.
530 *Wasmann J. Biol.* 9: 11-77.

- 531 Moser, W.E., Klemm, D.J., Richardson, D.J., Wheeler, B.A., Trauth, S.E., and Daniels, B.A.
532 2006. Leeches (Annelida: Hirudinida) of Northern Arkansas. J. Ark. Acad. Sci. 60: 84-
533 95.
- 534 Moser, W.E., Richardson, D.J., Hammond, C.I., and Lazo-Wasem, E.A. 2012. Redescription of
535 *Placobdella ornata* (Verrill, 1872) (Hirudinida: Glossiphoniidae). Bull. Peabody Mus.
536 Nat. Hist. 53: 325-330.
- 537 Moser, W.E., Bowerman, J., Hovingh, P., Pearl, C.A., and Ocegüera-Figueroa, A. 2014. New
538 host and distribution records of the leech *Placobdella sophieae* Ocegüera-Figueroa et al.
539 2010 (Hirudinida: Glossiphoniidae). Comp. Parasitol. 81: 199–202.
- 540 Ocegüera-Figueroa, A., León-Règagnon, V., and Siddall, M.E. 2010. DNA barcoding reveals
541 Mexican diversity within the freshwater leech genus *Helobdella* (Annelida:
542 Glossiphoniidae). Mitochondrial DNA, 21: 24-29.
- 543 Oosthuizen, J.H., and Davies, R.W. 1992. Redescription of the duck leech, *Theromyzon rude*
544 (Baird, 1869) (Rhynchobdellida: Glossiphoniidae). Can. J. Zool. 70: 2028-2033.
- 545 Pfeiffer, I., Brenig, B., and Kutschera, U. 2005. Molecular phylogeny of selected predacious
546 leeches with reference to the evolution of body size and terrestrialism. Theory in
547 Biosciences, 124: 55-64.
- 548 Phillips, A.J., Arauco-Brown, R., Ocegüera-Figueroa, A., Gomez, G.P., Beltrán, M., Lai, Y.T.,
549 and Siddall, M.E. 2010. *Tyrannobdella rex* n. gen. n. sp. and the evolutionary origins of
550 mucosal leech infestation. PLoS ONE, 5: e10057.
- 551 Pawlowski, L.K. 1948. Contribution a la systematique des sangsues du genre *Erpobdella* de
552 Blainville. Acta Zoologica et Oecologica, Universitatis Lodziensis, 8: 55.

- 553 Salas-Montiel, R., Phillips, A.J., De Leon, G.P.P., and Ocegüera-Figueroa, A. 2014. Description
554 of a new leech species of *Helobdella* (Clitellata: Glossiphoniidae) from Mexico with a
555 review of Mexican congeners and a taxonomic key. *Zootaxa*, 3900: 77-94.
- 556 Sawyer, R.T. 1970. Observations on the natural history and behavior of *Erpobdella punctata*
557 (Leidy)(Annelida: Hirudinea). *Am. Midl. Nat.* 83: 65-80.
- 558 Sawyer, R.T. 1972. North American freshwater leeches, exclusive of the Piscicolidae, with a key
559 to all species. *Illinois Biological Monographs* 46: University of Illinois press. 1-154.
- 560 Sawyer, R.T. 1986. *Leech biology and behaviour*. Vol. 1. Oxford: Clarendon Press.
- 561 Siddall, M.E., and Burreson, E.M. 1996. Leeches (Oligochaeta?: Euhirudinea), their phylogeny
562 and the evolution of life-history strategies. *Hydrobiologia*, 334: 227-285.
- 563 Siddall, M.E. 2001. Leeches of Laguna Volcán, Bolivia, including a new species of *Helobdella*
564 (Clitellata: Hirudinea). *Am. Mus. Novit.* No. 3313: 1-12.
- 565 Siddall, M.E., and Gaffney, E.S. 2004. Observations on the leech *Placobdella ornata* feeding
566 from bony tissues of turtles. *J. Parasitol.* 90: 1186–1188.
- 567 Siddall, M.E., Budinoff, R.B., and Borda, E. 2005. Phylogenetic evaluation of systematics and
568 biogeography of the leech family Glossiphoniidae. *Invertebr. Syst.* 19:105-112.
- 569 Siddall, M.E., and Bowerman, J. 2006. A new species of glossiphoniid leech from *Rana pretiosa*
570 (Amphibia: Ranidae) in Oregon. *J. Parasitol.* 92: 855–7.
- 571 Siddall, M.E., Brugler, M., and Kvist, S. 2016. Comparative transcriptomic analyses of 3 species
572 of *Placobdella* (Rhynchobdellida: Glossiphoniidae) confirms a single origin of blood
573 feeding in leeches. *J. Parasitol.* 102: 143-150.
- 574 Sinha, V.R.P., and Jones, J.W. 1967. On the food of the freshwater eels and their feeding
575 relationships with salmonids. *J. Zool. (Lond.)* 153: 119-137.

576 Sket, B., and Trontelj, P. 2008. Global diversity of leeches (Hirudinea) in freshwater.
577 Hydrobiologia, 595: 129-137.

578 Soucacos, P.N., Beris, A.E., Malizos, K.N., Kabani, C.T., and Pakos, S. 1994. The use of
579 medicinal leeches, *Hirudo medicinalis*, to restore venous circulation in trauma and
580 reconstructive microsurgery. Int. Angiol. 13: 251-258.

581

582

Draft

Figure legends

Figure 1. Distribution of leech samples of the family Erpobdellidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

Figure 2. Distribution of leech samples of the family Haemopidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

Figure 3. Distribution of leech samples of the family Glossiphoniidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

Figure 4. Dorsal views of select representative specimens of Erpobdellidae from the newly collected material; scale bars represent 0.5 cm. (A) *Erpobdella obscura* (ROMIZ I12051), (B) *Erpobdella punctata* (ROMIZ I12062), (C) *Erpobdella parva* (ROMIZ I12072), (D) *Erpobdella dubia* (ROMIZ I12101).

Figure 5. Dorsal views of select representative specimen of Haemopidae from the newly collected material; scale bars represent 1 cm. (A) *Haemopsis grandis* (ROMIZ I12001), (B) *Haemopsis marmorata* (ROMIZ I12015), (C) *Haemopsis lateromaculata* (ROMIZ I12115).

605 Figure 6. Dorsal views of select representative specimen of Glossiphoniidae from the newly
606 collected material; scale bars represent 0.25 cm. (A) *Helobdella stagnalis* (ROMIZ I12030), (B)
607 *Glossiphonia elegans* (ROMIZ I12095), (C) *Placobdella rugosa* (ROMIZ I12037), (D)
608 *Theromyzon rude* (ROMIZ I12018), (E) *Theromyzon tessulatum* (ROMIZ I12109).

609

610 Figure 7. Preserved *Haemopsis* sp. feeding on a predacious diving beetle, *Dytiscus dauricus*
611 (Holomuzki 1986) (ROMIZ I12122), recovered from an aquatic trap in the Albany River
612 ecodistrict.

Draft

Table 1. Metadata for leeches collected from the far north of Ontario, deposited at the Royal Ontario Museum (ROM) or Canadian Museum of Nature (CMN).

Family: Erpobdellidae						
Species	Survey	Date	Latitude	Longitude	Specimens	Catalogue Number
<i>Erpobdella dubia</i> (Moore and Meyer 1951)	Aquatic Net	20-Jun-2013	53.592	-88.364	1	ROMIZ I12101
	Unknown	24-Jul-1965	55.333	-87.283	1	CMNA 1987-0646
		6-Jun-2010	52.981	-87.926	1	ROMIZ I12119
<i>Erpobdella obscura</i> (Verrill 1872)	Aquatic Trap	13-Jun-2010	52.560	-87.300	5	ROMIZ I10631, I10632, I10633, I10634, I10636
		15-Jun-2010	52.560	-87.298	2	ROMIZ I12110, I12111
		23-Jun-2010	52.340	-86.163	2	ROMIZ I12105, I12106
		30-Jun-2011	52.826	-94.215	1	ROMIZ I12002
		30-Jun-2011	52.827	-94.217	2	ROMIZ I12011, I12012
		16-Jul-2011	53.907	-93.214	1	ROMIZ I10630
		20-Jun-2014	54.752	-89.363	1	ROMIZ I12091
		22-Jun-2014	54.752	-89.364	1	ROMIZ I12092

		23-Jun-2014	54.441	-87.902	1	ROMIZ I12081
		23-Jun-2014	54.441	-87.901	1	ROMIZ I12080
		25-Jun-2014	55.482	-87.880	1	ROMIZ I12086
		5-Jun-2010	52.983	-87.932	1	ROMIZ I12112
		5-Jun-2011	53.198	-89.112	1	ROMIZ I12067
		17-Jun-2011	54.456	-90.365	1	ROMIZ I12051
	Aquatic Net	21-Jun-2011	54.456	-90.365	1	ROMIZ I12055
		11-Jul-2011	52.456	-91.820	1	ROMIZ I12071
		11-Jul-2011	53.602	-93.538	1	ROMIZ I12022
		23-Jun-2014	54.443	-87.898	1	ROMIZ I12083
	Incidental	9-Jul-2011	53.602	-93.537	1	ROMIZ I12006
		24-Jun-1920	50.733	-81.483	1	CMNA 1900-5992
	Unknown	14-15-July-1920	51.267	-80.617	3	CMNA 1900-5993
		22-Aug-1979	51.483	-80.45	1	CMNA 1988-0132
<i>Erpobdella parva</i> (Moore 1912)	Aquatic Net	7-Jul-2011	52.462	-91.820	1	ROMIZ I12072
	Unknown	27-Jul-1965	55.283	-85	1	CMNA 1978-0329

<i>Erpobdella punctata</i> (Leidy 1870)	Aquatic Net	9-Jun-2011	54.164	-92.044	1	ROMIZ I12062
		28-Jun-2014	55.482	-87.881	1	ROMIZ I12087
	Incidental	11-Jun-2011	53.752	-88.919	1	ROMIZ I12013
	Unknown	14-15-July-1920	51.267	-80.617	1	CMNA 1900-5968
		12-Jul-1965	52.217	-81.8	1	CMNA-1978-0328
<i>Erpobdella sp.</i>	Aquatic Trap	26-Jun-2014	55.582	-87.877	1	ROMIZ I12075
	Aquatic Net	5-Jun-2010	52.782	-85.679	1	ROMIZ I12116
		2-Jun-2011	53.196	-89.108	2	ROMIZ I12064, I12065
		5-Jun-2011	54.431	-89.667	1	ROMIZ I12063
		17-Jun-2011	54.455	-90.366	1	ROMIZ I12121
		11-Jul-2011	53.602	-93.538	1	ROMIZ I12024
		15-Jul-2011	53.733	-92.762	6	ROMIZ I12042, I12043, I12044, I12045, I12046, I12047
		23-Jun-2012	52.893	-82.684	1	ROMIZ I12069
		11-Jun-2013	52.586	-89.675	1	ROMIZ I12096
		1-Jul-2013	50.686	-85.790	1	ROMIZ I12094
		20-Jul-2013	51.314	-85.748	1	ROMIZ I12103

		22-Jul-2013	51.314	-85.748	1	ROMIZ I12104
		4-Jul-2014	56.566	-88.592	1	ROMIZ I12089
Family: Haemopidae						
<i>Haemopsis grandis</i> (Verrill 1874)	Aquatic Trap	4-Jun-2011	53.196	-89.112	1	ROMIZ I12001
		21-Jun-2011	54.449	-90.359	1	ROMIZ I12003
	Aquatic Net	12-Jul-2013	50.937	-84.852	1	ROMIZ I12098
	Incidental	12-Jul-2011	53.606	-93.522	1	ROMIZ I12016
	Unknown	10-Jul-1965	52.217	-81.883	1	CMNA 1987-0667
		11-Jul-1965	52.217	-81.883	1	CMNA 1987-0668
<i>Haemopsis lateromaculata</i> Mathers 1963	Incidental	20-Jun-2010	52.163	-87.394	1	ROMIZ I12115
	Unknown	1920	51.05	-80.917	1	CMNA 1985-0259
		24-Jun-1920	50.733	-81.483	1	CMNA 1985-0262
		12-Jul-1965	52.217	-81.65	1	CMNA 1985-0279
		24-Jul-1965	54.083	-90.7	1	CMNA 1987-0670
<i>Haemopsis marmorata</i> (Say 1824)	Aquatic Trap	4-Jun-2010	52.981	-87.926	1	ROMIZ I12113
		6-Jun-2010	52.981	-87.926	1	ROMIZ I12120
		14-Jul-2010	53.219	-86.225	1	ROMIZ I12118
		18-Jun-2011	54.456	-90.365	1	ROMIZ I12014

	18-Jun-2011	54.457	-90.363	1	ROMIZ I12017
	21-Jun-2011	54.449	-90.359	1	ROMIZ I12004
	15-Jun-2012	51.743	-80.650	5	ROMIZ I12057, I12058, I12059, I12060, I12061
	25-Jun-2014	55.582	-87.877	2	ROMIZ I12084, I12085
	26-Jun-2014	55.582	-87.877	5	ROMIZ I12074, I12076, I12077, I12078, I12079
	28-Jun-2014	55.621	-88.838	1	ROMIZ I12093
Aquatic Net	7-Jul-2010	53.064	-87.515	1	ROMIZ I12117
	5-Jun-2011	53.196	-89.108	1	ROMIZ I12066
	28-Jun-2014	55.482	-87.881	1	ROMIZ I12088
Incidental	11-Jun-2011	54.164	-92.043	1	ROMIZ I12005
	11-Jun-2011	53.753	-88.919	1	ROMIZ I12015
Unknown	14-15 July- 1920	51.267	-80.617	1	CMNA 1977-0404
	16-Aug- 1920	52.283	-81.517	1	CMNA 1985-0281
	11-Jun-1975	51.3	-80.3	1	CMNA 1986-0143

		2-Aug-1980	56	-87.717	1	CMNA 1981-0062
<i>Haemopsis</i> sp.	Aquatic Trap	15-Jun-2012	51.743	-80.650	1	ROMIZ I12122
	Incidental	20-Jun-2010	52.163	-87.394	1	ROMIZ I12114
	Pitfall	13-Jul-2012	51.358	-82.057	1	ROMIZ I12073
	Pan Trap (blue)	8-Jun-2013	52.586	-89.677	1	ROMIZ I12082
Family: Glossiphoniidae						
<i>Glossiphonia elegans</i> (Verrill 1872) Castle 1900	Aquatic Trap	23-Jun-2010	52.340	-86.163	2	ROMIZ I12107, I12108
		7-Jul-2011	53.602	-93.537	1	ROMIZ I12040
		12-Jul-2012	52.306	-83.384	1	ROMIZ I12070
	Aquatic Net	10-Jun-2013	52.586	-89.677	1	ROMIZ I12095
		20-Jun-2013	53.592	-88.362	1	ROMIZ I12102
		6-Jul-2013	51.008	-87.146	1	ROMIZ I12100
		10-Jul-2014	56.289	-89.567	1	ROMIZ I12090
		14-Jul-1920	51.267	-80.617	1	CMNA 1900-8243
	Unknown	15-Jul-1965	52.933	-82.6	1	CMNA 1978-0327
		22-Jul-1965	55.283	-85.083	1	CMNA 1978-0326

		11-Aug-1967	52.133	-94.733	1	CMNA 1987-0671
<i>Glossiphonia</i> sp.	Incidental	9-Jul-2011	53.602	-93.537	1	ROMIZ I12007
		7-Jul-2011	53.603	-93.536	1	ROMIZ I12041
		11-Jul-2011	53.602	-93.537	5	ROMIZ I12030, I12031, I12032, I12033, I12148
		11-Jul-2011	53.602	-93.538	8	ROMIZ I12019, I12020, I12021, I12023, I12025, I12026, I12027, I12028
<i>Helobdella stagnalis</i> (Linnaeus 1758)	Aquatic Net	15-Jul-2011	53.733	-92.765	2	ROMIZ I12048, I12049
		17-Jun-2011	54.456	-90.365	3	ROMIZ I12052, I12053, I12054
		13-Jun-2011	53.753	-88.920	1	ROMIZ I12056
		21-Jun-2012	52.893	-82.683	1	ROMIZ I12068
		25-Jun-2013	53.397	-87.800	1	ROMIZ I12097
		6-Jul-2013	51.012	-87.154	1	ROMIZ I12099
		15-Jul-1965	52.967	-82.35	56	CMNA 1987-0669
	Unknown	16-Aug-1965	52.933	-82.4	21	CMNA 1987-0666

<i>Helobdella</i> sp.	Unknown	26-Jul-1963	51.267	-80.617	1	CMNA 1987-0662
<i>Placobdella montifera</i> (Moore 1906)	Unknown	5-Aug-1961	51.5	-90.15	1	CMNA 1987-0659
<i>Placobdella rugosa</i> (Verrill 1874)	Aquatic Net	11-Jul-2011	53.602	-93.537	1	ROMIZ I12037
<i>Theromyzon rude</i> (Baird 1869)	Aquatic Net	11-Jul-2011	53.602	-93.538	1	ROMIZ I12018
<i>Theromyzon tessulatum</i> (Müller 1774)	Aquatic Trap	13-Jun-2010	52.560	-87.298	4	ROMIZ I12109
<i>Theromyzon</i> sp.	Aquatic Trap	11-Jul-2011	53.603	-93.536	1	ROMIZ I12038
		7-Jul-2011	53.603	-93.536	1	ROMIZ I12147
	Aquatic Net	11-Jul-2011	53.602	-93.537	4	ROMIZ I12029, I12034, I12035, I12036
		11-Jul-2011	53.603	-93.536	1	ROMIZ I12039
		15-Jul-2011	53.737	-92.763	1	ROMIZ I12050

Figure 1. Distribution of leech samples of the family Erpobdellidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

170x170mm (300 x 300 DPI)

Figure 2. Distribution of leech samples of the family Haemopidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

170x170mm (300 x 300 DPI)

Figure 3. Distribution of leech samples of the family Glossiphoniidae, collected for the present study across the far north of Ontario's two ecozones; Hudson Bay Lowlands and Ontario Shield. Projection: Canadian Lambert Conformal Conic, Datum: NAD 1983.

170x170mm (300 x 300 DPI)

Figure 4. Dorsal views of select representative specimens of Erpobdellidae from the newly collected material; scale bars represent 0.5 cm. (A) *Erpobdella obscura* (ROMIZ I12051), (B) *Erpobdella punctata* (ROMIZ I12062), (C) *Erpobdella parva* (ROMIZ I12072), (D) *Erpobdella dubia* (ROMIZ I12101).

160x148mm (300 x 300 DPI)

Figure 5. Dorsal views of select representative specimen of Haemopidae from the newly collected material; scale bars represent 1 cm. (A) *Haemopsis grandis* (ROMIZ I12001), (B) *Haemopsis marmorata* (ROMIZ I12015), (C) *Haemopsis lateromaculata* (ROMIZ I12115).

179x198mm (300 x 300 DPI)

Figure 6. Dorsal views of select representative specimen of Glossiphoniidae from the newly collected material; scale bars represent 0.25 cm. (A) *Helobdella stagnalis* (ROMIZ I12030), (B) *Glossiphonia elegans* (ROMIZ I12095), (C) *Placobdella rugosa* (ROMIZ I12037), (D) *Theromyzon rude* (ROMIZ I12018), (E) *Theromyzon tessulatum* (ROMIZ I12109).

213x488mm (300 x 300 DPI)

Figure 7. Preserved *Haemopsis* sp. feeding on a predaceous diving beetle, *Dytiscus dauricus* (Holomuzki 1986) (ROMIZ I12122), recovered from an aquatic trap in the Albany River ecodistrict.

109x84mm (300 x 300 DPI)