Canadian Journal of Plant Science
Revue canadienne de phytotechnie

**Yield and nutritive value of binary legume-grass mixtures
under grazing or frequent cutting**

Journal:	<i>Canadian Journal of Plant Science</i>
Manuscript ID	CJPS-2017-0183.R1
Manuscript Type:	Article
Date Submitted by the Author:	19-Sep-2017
Complete List of Authors:	Bélanger, Gilles; Agriculture and Agri-Food Canada, Tremblay, Gaetan; Agriculture and Agri-Food Canada, Quebec Research and Development Centre Papadopoulos, Yousef; Agriculture and Agri-Food Canada, Duynisveld, John; AAFC, Nappan Research Farm Lajeunesse, Julie; Agriculture and Agri-Food Canada, Lafrenière, Carole; Université du Québec en Abitibi-Témiscamingue Station de Recherche Agroalimentaire, Fillmore, Sherry; Agriculture Agri-food Canada, Kentville RDC
Keywords:	Alfalfa, Clover, Forage, Grass, Grazing

SCHOLARONE™
Manuscripts

Yield and nutritive value of binary legume-grass mixtures under grazing or frequent cutting

Gilles Bélanger¹, Gaëtan F. Tremblay¹, Yousef A. Papadopoulos², John Duynisveld³, Julie Lajeunesse⁴, Carole Lafrenière⁵, and Sherry A. E. Fillmore⁶

¹Agriculture and Agri-Food Canada, Quebec Research and Development Centre, 2560 Hochelaga Blvd., Québec, QC, G1V 2J3 Canada; ²Agriculture and Agri-Food Canada, Faculty of Agriculture, Dalhousie University, PO Box 550, Truro, NS, B2N 5E3 Canada; ³Agriculture and Agri-Food Canada, Kentville Research and Development Centre, Nappan, NS, B0L 1C0 Canada; ⁴Agriculture and Agri-Food Canada, Normandin Research Farm, 1468 Saint-Cyrille St., Normandin, QC, G8M 4K3 Canada; ⁵Université du Québec en Abitibi-Témiscamingue, 79 Côté St., Notre-Dame-du-Nord, QC, J0Z 3B0 Canada; ⁶Agriculture and Agri-Food Canada, Kentville Research and Development Centre, 32 Main St., Kentville, NS, B4N 1J5 Canada.

Corresponding author: gilles.belanger@agr.gc.ca

Abbreviations: **ADF**, acid detergent fiber; **aNDF**, neutral detergent fiber assayed with a heat stable α -amylase; **DM**, dry matter; **IVTD**, *in vitro* true digestibility of dry matter; **NDFd**, *in vitro* aNDF digestibility; **PCA**, principal component analysis; **RPD**, ratio of prediction to deviation; **RSQ**, coefficient of determination for the prediction; **SD**, standard deviation; **SEP**, standard error of prediction; **SEP(C)**, standard error of

prediction corrected for the bias; **TDN**, total digestible nutrients; **NSC**, non-structural carbohydrates

For Review Only

Abstract

Although most forage production in eastern Canada is comprised of species mixtures, little research has identified the best species to include in forage mixtures. Our objective was to identify binary legume-grass mixtures with high forage yield and nutritive value under either simulated grazing with frequent cutting or cattle grazing. The experiment was conducted at three sites in eastern Canada with 18 binary legume-grass mixtures of one of six grass species [Kentucky bluegrass (*Poa pratensis* L.), meadow fescue (*Festuca eliator* auct. Amer.), orchardgrass (*Dactylis glomerata* L), tall fescue (*Schedonorus phoenix* (Scob.) Holub), timothy (*Phleum pratense* L.), and meadow brome (*Bromus biebersteinii* Roem. & Schult.)] seeded in 2010 with a grazing-type alfalfa (*Medicago sativa* L.), white clover (*Trifolium repens* L.), or birdsfoot trefoil (*Lotus corniculatus* L.). The six grass species grown in mixture with alfalfa, birdsfoot trefoil, or white clover persisted well under frequent cutting or rotational grazing at the three sites. White clover grown in a binary mixture with a grass species did not perform well under frequent cutting or rotational grazing. Meadow brome-based binary mixtures were overall the best performing in terms of DM yield; although their nutritive value was average, meadow brome combined with alfalfa or birdsfoot trefoil were among the best legume-grass mixtures for the estimated milk production per hectare. The greatest estimated milk production per hectare was obtained with birdsfoot trefoil mixed with meadow brome followed by the alfalfa-timothy and the alfalfa-meadow brome mixtures.

Introduction

Legume-grass mixtures generally provide more consistent forage yield across a wide range of environments than grass or legume monocultures (Sleugh et al. 2000; Bélanger et al. 2014). Legume-grass mixtures have also been shown to reduce weed invasion compared to monocultures (Tracy and Sanderson 2004; Picasso et al. 2008; Frankow-Lindberg et al. 2009; Sanderson et al. 2012; Finn et al. 2013; Bélanger et al. 2014).

Nutritive value should also be considered because of its impact on animal productivity, and meat and milk quality. Results of a pan-European experiment, which included a Canadian site, have demonstrated that mixing grasses and legumes increases dry matter (DM) yield (Finn et al. 2013) with no negative effects on nutritive value (Sturludóttir et al. 2013). Adding a legume into a grass sward has been shown to increase forage DM yield and crude protein concentration (Barnett and Posler 1983), and improve forage nutritive value (Papadopoulos et al. 2001). Furthermore, mixing timothy (*Phleum pratense* L.) with alfalfa (*Medicago sativa* L.) has been shown to increase the non-structural carbohydrate concentration of forages (Bélanger et al. 2014), potentially resulting in a more efficient use of N by ruminants (Brito et al. 2009).

Kentucky bluegrass (*Poa pratensis* L.), meadow fescue (*Festuca eliator* auct. Amer.), orchardgrass (*Dactylis glomerata* L), tall fescue (*Schedonorus phoenix* (Scob.) Holub), timothy, and meadow brome grass (*Bromus biebersteinii* Roem. & Schult.) are forage grass species that are well adapted to cool seasons and recommended in eastern Canada. Alfalfa, white clover (*Trifolium repens* L.), and birdsfoot trefoil (*Lotus corniculatus* L.) are perennial legume species also recommended in eastern Canada but their performance and nutritive value in mixtures with grasses and under grazing are not well documented.

Although most forage production in eastern Canada is comprised of species mixtures, there is limited research on identifying the best species to include within forage mixtures. Our objective was to identify persistent binary legume-grass mixtures with high forage yield and nutritive value to be used under both simulated grazing with frequent cutting or cattle grazing in eastern Canada.

Materials and Methods

The experiment was conducted at three sites: 1) Chapais Research Farm of Agriculture and Agri-Food Canada, Lévis, QC, Canada, 2) Normandin Research Farm of Agriculture and Agri-Food Canada, Normandin, QC, Canada, and 3) Nappan Research Farm of Agriculture and Agri-Food Canada, Nappan, NS, Canada. Site characteristics are presented in Table 1. Binary legume-grass mixtures (18) of one of six grass species (orchardgrass cv. Killarney, Kentucky bluegrass cv. Troy, meadow brome cv. Fleet, meadow fescue cv. Pradel, tall fescue cv. Courtenay, and timothy cv. Express) were seeded in 2010 with a grazing-type alfalfa cv. CRS1001, birdsfoot trefoil cv. AC Langille, or white clover cv. Milkanova. Seeding rates for each species are presented in Simili da Silva et al. (2013). At each site, binary mixtures were replicated three times in a split-plot layout, with legume species as main plots set out as a Latin square and grass species randomized to the subplots. Phosphorus and K fertilizers were applied before seeding and each year if needed based on provincial recommendations in Québec for the sites at Lévis and Normandin (CRAAQ 2003) and in Nova Scotia for the site at Nappan (A.A.C.C.P.C.F.C. 1991). Lime was applied at Lévis in 2011 at a rate of 3.8 Mg ha⁻¹ after the first and second cutting, and in the spring of 2012 at a rate of 3.2 Mg ha⁻¹. Nitrogen

was applied at seeding at rates of 30 kg N ha⁻¹ at Lévis, 25 kg N ha⁻¹ at Normandin, and 24 kg N ha⁻¹ at Nappan. No N was applied in the post-seeding years, except for 40 kg N ha⁻¹ after the second cutting at Lévis and Normandin in 2014 and 2015, and for 34 kg N ha⁻¹ after the first grazing from 2013 to 2015 at Nappan. Our intent was to rely entirely on legume N₂ fixation to provide N to the legume grass-mixtures. Towards the end of the study, we applied a low rate of fertilizer N to help the forage grasses because the legume component of the mixture was much decreased.

For logistics reasons, all mixtures were grazed or cut at the same time and the timing of those events was based on timothy, the main forage grass species in eastern Canada, reaching 33 cm in height (Table 2). Because of the lack of grazing facilities and cattle at Lévis and Normandin, grazing was simulated by frequent cutting of all plots to a 5-cm height. Dry matter yield was determined by cutting an area of 7.3 m² at Lévis using a self-propelled flail-type CarterTM forage harvester (Carter MGF Co., Inc., Brookston, IN, USA) and an area of 6.0 m² at Normandin using a walk-behind flail harvester (Swift Machine and Welding, Swift Current, SK, Canada). A fresh forage sample of approximately 500 g was taken from each plot, weighed, dried at 55°C in a force-draft oven to determine DM concentration, and then ground using a Wiley mill (Standard model 4, Arthur H. Thomas Co., Philadelphia, PA) to pass through a 1-mm screen. At Nappan, rotational grazing with beef steers was initiated in the spring and concluded in late summer or fall. A block of 8 to 16 growing beef steers, weighing 500 kg on average, was allocated to this trial and animals started on the first grazing paddock (replicate 1) and were moved to the next paddock at the required sward height. Grazing of all four replicates, each 0.3 ha in size, was completed in a maximum of six days from initiation

that was set when timothy reached about 33 cm in height. Sward height was monitored to ensure that cattle would exit the paddocks at the appropriate sward exit height (6–8 cm). Forage samples were taken just prior to the beginning of each grazing cycle using a 0.25 m² quadrat randomly placed within each plot. Samples were dried in an oven at 55°C, and weighed to estimate DM yield. All samples from each grazing cycle were then ground with a Wiley mill to pass a 1-mm screen.

In the first two years, the frequency grid technique (Vogel and Masters 2001) was used approximately two weeks after the first cutting or grazing to determine the presence of each seeded species in each plot. Two grids of 25 squares (5 × 5 cm) were placed in each plot. The presence of at least one seeded plant species and one other species was noted for each square. This information is an estimate of the minimum plant density. In the last three years, seeded grasses, seeded legumes, and weeds were manually separated twice during the season (first and third cutting or grazing) from one sample taken in each plot from a 0.25 m² quadrat; each component was dried at 55°C in a force-draft oven and weighed to determine their proportion assessed as their contribution to DM yield.

Dried and ground forage samples were scanned by visible near infrared reflectance spectrometry (VNIRS) using a NIRSystem 6500 monochromator (Foss, Silver Spring, MD, USA) in the range from 400 to 2500-nm intervals. Out of approximately 1100 forage samples per post-seeding year, the WinISI IV (ver. 4.5.0.14017) software (Infrasoft International LLC, State College, PA, USA) was used to select approximately 75 samples per post-seeding year from 2011 to 2014 with spectra that contributed the most to the variability within all samples. Of these 75 samples per post-seeding year, approximately 60 and 15 were randomly selected for the calibration and validation sets,

respectively. Samples selected for the calibration and validation sets (≈ 75 samples per post-seeding year $\times$ 4 post-seeding years ≈ 300 samples) were chemically analyzed in duplicates that were averaged prior to the development of the calibration equations. The DM and ash concentrations (Leco Corporation, 2009) were determined using a thermogravimetric analyser (model TGA701, Leco Corporation, St. Joseph, MI, USA). Crude fat (ether extract) was determined using Ankom xt15 Extractor Technology Method (AOCS, 2003). Concentrations of water soluble carbohydrates (WSC) and starch were measured according to dos Passos Bernardes et al. (2015) and the concentration of non-structural carbohydrates (NSC) was calculated as the sum of WSC and starch concentrations. Total N concentration was measured using a method adapted from Isaac and Johnson [1976]. Ground samples (100 mg) were digested for 60 min at 380 °C in a 1.5-mL mixture of selenious and sulfuric acid plus 2 mL of 30 % H₂O₂. After cooling, the mixture was diluted to 75 mL with deionized water. An auto analyzer (QuikChem 8000 Lachat Zellweger Analytics Inc., Lachat Instruments, Milwaukee, WI) was used to measure total N with the method 13–107–06–2–D and P with the method 13–115–01–2–A (Lachat, 2013, Zellweger Analytics, Lachat Instruments, Milwaukee, WI). The acid detergent fiber (ADF) was determined according to AOAC (1990). The neutral detergent fiber (aNDF) was analyzed following Mertens (2002) with addition of a heat-stable α -amylase and sodium sulfite. These fiber extractions were done using the Ankom filter bag technique (ANKOM Technology, NY, USA). The IVTD was measured using the method of Goering and Van Soest (1970) based on a 48-h incubation with buffered rumen fluid followed by an aNDF determination of the post digestion residues. The rumen fluid incubation was performed with Ankom F57 filter bags and an Ankom Daisy II incubator,

using the bath incubation procedures outlined by Ankom Technology Corp. (ANKOM Technology, NY, USA). Rumen fluid was obtained from a ruminally fistulated dairy cow that was offered a diet of 37% grass silage, 15% corn silage, 8% hay, 30% corn grain, and 10% concentrate mix formulated to meet the nutritional requirements of a lactating dairy cow expected to produce 10200 kg milk yr⁻¹. The *in vitro* true digestibility of DM (IVTD; g kg⁻¹ DM) and the *in vitro* aNDF digestibility (NDFd; g kg⁻¹ aNDF) were calculated as below:

$$\text{IVTD} = [1 - (\text{post digestion dry weight following aNDF wash/predigestion dry weight})] \times 1000$$

$$\text{NDFd} = [1 - (\text{post digestion dry weight following aNDF wash/predigestion dry weight of aNDF})] \times 1000.$$

The total N, ADF, aNDF, neutral detergent insoluble crude protein (NDICP; Licitra et al. 1996), ash, and ether extract concentrations, along with NDFd were used to calculate the total digestible nutrients (TDN; NRC, 2001) using the University of Wisconsin Alfalfa/Grass Evaluation System and the milk production per hectare with Milk2013 (Undersander et al. 2013).

The VNIRS calibration equations were developed using a modified least squares regression method of the WinISI IV software. Depending on the nutritive attribute and the calibration set of approximately 240 forage samples (≈ 60 samples per post-seeding year $\times 4$ post-seeding years ≈ 240), the number of calibration samples used to develop the final calibration equations varied between 223 and 237. Calibration equations were selected based on Martens and Naes (2001) as follows: Reference data = $f(\text{spectral data}) + \text{SEC}$, where $f()$ means “function of” and SEC is the standard error of calibration. The

best NIRS calibration equations were the ones that minimize the SEC. Cross-validation was performed by using four subgroups from the calibration set to choose the optimal number of terms and to avoid over fitting the calibration model (Shenk and Westerhaus, 1991). Calibration equations were validated using WinISI IV software by comparing predicted against reference values. Statistics on the VNIRS performance to predict nutritive attributes in the validation set ($n = 61$) are presented in Table 3. The ratio of standard error of prediction to standard deviation [RPD = standard deviation (SD) of the reference data used in the validation set divided by the standard error of prediction corrected for bias (SEP(C))] was greater than 3 and the VNIRS predictions were therefore considered successful for all nutritive attributes.

Data were assessed by analyses of variance (ANOVA) using the GENSTAT 17 statistical software (VSN International 2013). Sites, legume species, and grass species were considered fixed effects. Differences were considered significant when $P < 0.05$. Seasonal values of DM yield were calculated as the sum of the DM yield at each cutting or grazing. Seasonal values of the nutritive attributes were calculated as the average of their values at each cutting or grazing. Seasonal values were reported with the objective of presenting and discussing the overall response over the five years of the study. The question of seasonal distribution of DM yield and nutritive attributes will be addressed in future manuscripts. For each variate, extreme high or low values were identified after calculating an upper [overall mean + $(2.81 \times \text{SEM} / 2)$] and a lower [overall mean – $(2.81 \times \text{SEM} / 2)$] limit centered about the overall mean. A principal component analysis (PCA) was used to assess the relationships among variates (DM yield and nutritive attributes) and how variations in these variates were related to legume-grass mixtures.

The PCA was performed on the least squares means of the treatments using the correlation matrix method to give equal weight to all variates.

Results and Discussion

Main effects of sites, legume species, and grass species

The three sites differed significantly for most variates [contrasts: Nappan vs. Lévis and Normandin, Lévis vs. Normandin; Table 4]. Average forage DM yields across mixtures and years were, respectively, 6.67, 5.21, and 4.15 Mg ha⁻¹ at Nappan, Lévis, and Normandin. The forage nutritive value was lower at Nappan than at Lévis and Normandin [contrast: Nappan vs. Lévis and Normandin; Table 4] with greater ADF (345 vs. 300 and 316 g kg⁻¹ DM) and aNDF (538 vs. 440 and 453 g kg⁻¹ DM) concentrations along with lower total N concentration (21.8 vs. 26.5 and 25.5 g kg⁻¹ DM), IVTD (826 vs. 895 and 888 g kg⁻¹ DM), and NDFd (676 vs. 761 and 739 g kg⁻¹ aNDF). Site differences in forage DM yield and nutritive value are expected and can be explained by differences in soil and climatic conditions, and the different management practices (grazing vs. frequent cutting) used at the three sites.

Sites also differed in the composition of the mixtures in the last three post-seeding years. The average proportion of the legume species in each of the last three post-seeding years (2013, 2014, and 2015) was greater at Lévis (23, 20, and 4%) and Normandin (30, 27, and 5%) than at Nappan (4, 3, and 2%) (Table 5). Our visual observations in the first post-seeding year suggest that selective grazing of birdsfoot trefoil and alfalfa might explain the lack of legume persistence under grazing at Nappan. This difference in the proportion of the legume species might explain in part the lower concentrations of ADF and aNDF, and the greater N concentration in forages at Lévis and Normandin than at

Nappan because of the known lower ADF and aNDF concentrations of legume species and their greater N concentration compared to grass species (Pelletier et al., 2010). Along with their differences in species composition, lower forage DM yields at Lévis and Normandin could explain the greater forage TDN concentration and IVTD at those two sites because of the known negative relationship between forage DM yield and digestibility (Bélanger et al. 2001).

The three legume species in binary mixtures with grasses differed significantly for most variates (contrast: C vs. A+B, A vs. B; Table 4). The average seasonal forage DM yield across sites, years, and grass species in the mixtures was the least with the white clover-based mixtures and the greatest with the birdsfoot trefoil-based mixtures (Table 6). The white clover-based mixtures also had the greatest ADF and aNDF concentrations, the greatest NDFd, and the lowest total N concentration (Table 6). Although statistically significant, differences in TDN concentration and IVTD were small. The proportion of white clover was much less than that of birdsfoot trefoil and alfalfa at Lévis and Normandin in the last three post-seeding years (Table 5). This lower proportion of white clover could explain the greater ADF and aNDF concentrations, the greater NDFd, and the lower total N concentrations of the white clover-based mixtures.

White clover was nearly absent in the last three post-seeding years at all three sites (Table 5). The decline of white clover in mixtures with grasses under frequent cutting has been reported previously. In a study conducted in Québec where white clover was grown in a mixture with either meadow fescue or meadow bromegrass under frequent cutting (Drapeau and Bélanger, 2009), the proportion of white clover decreased from the first to the third post-seeding year, reaching values below 10%. In a study of white clover in

mixtures with grasses conducted in Newfoundland, the proportion of white clover also decreased from 40% in the first post-seeding year to 27% in the third post-seeding year (McKenzie et al. 2005). White clover in mixtures with forage grasses is therefore not well adapted to frequent cutting or rotational grazing under the conditions of eastern Canada.

The proportion of alfalfa and birdsfoot trefoil was also poor in the last three post-seeding years of the study at Nappan where cattle grazing was used (Table 5). At Lévis and Normandin, two sites with frequent cutting, the proportion of alfalfa and birdsfoot trefoil ranged between 23 and 43% in the third and fourth post-seeding years. As mentioned above, selective grazing in the first two post-seeding years might explain the poor persistence of alfalfa and birdsfoot trefoil under grazing at Nappan. In the first post-seeding year, all seeded legume species were present at the three sites. Measurements with the frequency grid technique indicated that at least 27 plants m⁻² of each species were present (data not shown).

The six grass species in binary mixtures with a legume species also differed significantly for most variates (Table 4). Average forage DM yields of Kentucky bluegrass- and timothy-based mixtures across sites, years, and legume species in the mixture did not differ from those of the other grass-based mixtures [contrast: Kb vs. (Ti+Tf+Or+Mf+Mb), Ti vs. (Tf+Or+Mf+Mb); Table 4]. Forage DM yields of other grass-based mixtures, however, differed with average DM yields being the lowest with the meadow fescue- and orchardgrass-based mixtures and the greatest with the meadow brome- and tall fescue-based mixtures (Table 6). The timothy-based mixtures had lower average ADF and aNDF concentrations, and greater TDN concentration than the average of all grass species-based mixtures (Table 6). The Kentucky bluegrass-based

mixtures had lower IVTD and NDFd, and TDN concentration than the five other grass species-based mixtures. The meadow fescue-based mixtures had lower concentrations of ADF and aNDF, greater IVTD and NDFd, and greater concentrations of TDN and NSC than the tall fescue-based mixtures.

All grass species persisted well at the three sites. Their proportion ranged between 21 and 86%, and this proportion was relatively stable in the last three post-seeding years (Table 5). The proportion of timothy and Kentucky bluegrass, although significant, tended to be less than that of other forage grasses in the last three post-seeding years, while that of tall fescue and meadow fescue tended to be greater. All six grass species in mixtures with a legume species can therefore tolerate frequent cutting or grazing under the conditions of eastern Canada.

Comparison of the eighteen binary legume-grass mixtures

The effects of both legume and grass species in the mixtures were affected by the sites as indicated by the significant site $\times$ legume and site $\times$ grass interactions for all variates (Table 4). There was also a significant interaction between grass species and legume species in the mixtures for DM yield, NDFd, and concentrations of aNDF, NSC, and N. The 18 binary legume-grass mixtures were therefore compared at each site using PCA with the overall objective of determining the best binary mixture for several forage nutritive attributes and DM yield.

The first principal component explained 55, 58, and 50% whereas the second component explained 30, 20, and 29% of the total variation at Nappan, Lévis, and Normandin, respectively (Fig. 1); the first two principal components therefore explained at least 78% of the total variation. At Nappan, the first component was defined mostly by

the forage ADF concentration, DM yield, aNDF concentration, and total N concentration on the positive side, and by IVTD, NDFd, and concentrations of NSC and TDN on the negative side. At Lévis and Normandin, the first component was mostly defined by concentrations of total N and TDN, DM yield, and IVTD on the positive side, and by ADF and aNDF concentrations, and NDFd on the negative side. Attributes within the same group on each side were positively correlated, while attributes in opposing groups were negatively correlated.

The first component of the PCA mostly defined differences among grass species in the mixtures at Nappan and differences among legume species in the mixtures at Lévis and Normandin (Fig. 1). At Nappan, Kentucky bluegrass-based mixtures with high ADF concentration along with low IVTD, NDFd, and concentrations of NSC and TDN were opposed to mixtures with timothy and meadow fescue with high IVTD, NDFd, concentrations of NSC and TDN, and low ADF concentration. At Lévis and Normandin, mixtures with alfalfa and birdsfoot trefoil with high concentrations of total N and TDN, and low ADF and aNDF concentrations were opposed to white clover-based mixtures with low concentrations of total N and TDN, and high concentrations of ADF and aNDF.

The PCA confirmed differences among sites in their response to the 18 binary legume-grass mixtures. The legume component in the last three post-seeding years at Nappan was very low and this might explain why the grass species was the main driver of the first component of the PCA at that site. At Lévis and Normandin, the legume proportion was greater than at Nappan and the legume species were the main driver of the first component of the PCA. The main drivers of the second component of the PCA also differed among sites. At Nappan, forage aNDF concentration was opposed total N

concentration, while forage DM yield was opposed to IVTD and NSC concentration at Lévis and Normandin.

Forage DM yield was one of the main drivers of the relationship among variates on the first component at Nappan and on the first and second component at Lévis and Normandin. It is an important attribute for ensuring farm profitability and it should be considered along with nutritive attributes in the selection of legume-grass mixtures. At Nappan, the timothy- and meadow fescue-based mixtures had above average nutritive value but they tended to have below average DM yield (Table 7). Conversely, Kentucky bluegrass-based mixtures had below average nutritive value and had an average DM yield. Of the 18 binary legume-grass mixtures at Nappan, the alfalfa-timothy mixture is the only one that combined above average DM yield along with above average IVTD, and concentrations of TDN and total N, and below average ADF and aNDF concentrations. At Lévis and Normandin, white clover-based mixtures had above average ADF and aNDF concentrations and below average DM yield (Fig. 1). Conversely, alfalfa- and birdsfoot-based mixtures had below average ADF and aNDF concentrations, and above average total N and TDN concentrations along with greater DM yield. At these two sites, timothy-based mixtures with either alfalfa or birdsfoot trefoil tended to have above average DM yield, IVTD, and concentrations of total N and TDN along with below average ADF and aNDF concentrations (Tables 8 and 9).

The potential milk production per hectare was estimated for the 18 legume-grass mixtures in an effort to integrate both forage DM yield and nutritive value into one variate. Averaged across the three sites and the five years, birdsfoot trefoil mixed with either timothy, Kentucky bluegrass, tall fescue, orchardgrass, or meadow bromegrass, or

alfalfa mixed with either timothy or meadow brome grass resulted in above average estimated milk production (Table 10). The highest estimated milk production was obtained with birdsfoot trefoil mixed with meadow brome grass followed by the alfalfa-timothy mixture.

Meadow brome grass-based mixtures were overall the best performing in terms of DM yield (data not shown). The meadow brome grass-based mixtures, however, had above average ADF and aNDF concentrations, average IVTD and NDFd, and below average TDN concentration. Although the nutritive value of meadow brome grass-based mixtures was average, it provided one of the best combinations with alfalfa and birdsfoot trefoil for the estimated milk production. There is limited information in eastern Canada on the alfalfa-meadow brome grass mixture. In a study with frequent cutting, meadow brome grass yielded more than meadow fescue in the second and third post-seeding years when they were grown with white clover (Drapeau and Bélanger, 2009).

Tall fescue-based mixtures also performed well in terms of DM yield but they had above average ADF and aNDF concentrations, and below average TDN concentration. Meadow fescue-based mixtures had below average DM yield over the five years of the study but it had lower than average ADF concentration and above average IVTD, NDFd, and NSC concentration. Alfalfa and timothy are known to be not well adapted to grazing. Under the conditions of our study, however, the grazing-type alfalfa cultivar performed very well in mixture with timothy.

Limitations and Perspectives

This study was not specifically designed to compare the effect of cattle grazing and simulated grazing with frequent cutting but our results indicate that the performance of

binary legume-grass mixtures managed under frequent cutting over five post-seeding years seem to differ with cattle grazing. Although it was not measured, our visual observations in the first post-seeding year at Nappan suggest that selective grazing of birdsfoot trefoil and alfalfa might have reduced their persistence and their contribution to forage DM yield in subsequent years. Selective grazing of birdsfoot trefoil over tall fescue has been reported previously (Wen et al., 2004). Our results confirm the importance of evaluating forage mixtures under cattle grazing if those mixtures are to be used mainly for grazing.

Birdsfoot trefoil and alfalfa persisted well for four post-seeding years under frequent cutting, while white clover did not perform well under frequent cutting or rotational grazing. The six grass species persisted well under frequent cutting or rotational grazing. Although the persistence of most legume and grass species was acceptable, except for white clover, the productivity and nutritive value of the binary mixtures varied. Meadow brome-grass-based binary mixtures were overall the best performing in terms of DM yield. Although the nutritive value of meadow brome-grass-based binary mixtures was average, they provided one of the best combinations with alfalfa or birdsfoot trefoil for the estimated milk production per hectare. The greatest estimated milk production per hectare was obtained with birdsfoot trefoil mixed with meadow brome-grass (11.23 Mg ha^{-1}) followed by the alfalfa-timothy (10.56 Mg ha^{-1}) and the alfalfa-meadow brome-grass (10.39 Mg ha^{-1}) mixtures (Table 10). The performance of birdsfoot trefoil, grazing-type alfalfa, meadow brome-grass, and timothy in this experiment conducted over five post-seeding years confirms their potential for grazing and frequent cutting.

Although no or little N fertilizer was applied in this study, the forage mixtures performed well and the forage grasses were relatively productive over the five years of the study. The grasses in the binary legume-grass mixtures depended mostly on soil N and on the transfer of N from the legume species. The importance of N fertilization for binary legume-grass mixtures remains, however, poorly understood, primarily when the legume component is declining with time.

This study, conducted at three sites and over five post-seeding years, provides valuable information on the performance of binary mixtures of legume and grass species that are adapted to the cool and humid climate conditions of eastern Canada. For logistics reasons, all mixtures were grazed or cut at the same time and the timing of those events was based on timothy reaching a certain height. This approach may have introduced a bias in favor of timothy, the main forage grass species in eastern Canada. More research is required to assess some of those binary mixtures with cutting or grazing based on the development and growth of each mixture.

Acknowledgments

The authors are grateful to Andrée-Dominique Baillargeon, Geneviève Bégin, Camille Lambert-Beaudet, Lucie Lévesque, and Danielle Mongrain at the Québec Research and Development Centre in Québec City, Jean-Noël Bouchard at the Québec Research and Development Centre in Normandin, and Matthew Crouse and Carla MacKay at the Kentville Research and Development Centre in Nappan for their technical assistance. The authors also wish to acknowledge the assistance of the Farm Service Division and the livestock crew of the Nappan Research Farm for providing assistance during seeding, fencing, and data collection. This work was supported through a Beef Cluster grant by Agriculture and Agri-Food Canada and Beef Cattle Research Council, a division of the Canadian Cattlemen's Association.

References

- A.A.C.C.P.C.F.C. 1991. Fertility management of established forage stand. Atlantic Provinces Field Crop Guide. Publication 100. Atlantic Advisory Committees on Cereal, Protein, Corn and Forage Crops. p.9
- AOAC. 1990. Method 973.18: Determination of Acid Detergent Fiber by refluxing. Official Method of Analysis, 15th edition. AOAC International, Gaithersburg, MC, USA.
- AOCS, 2003. Method AM 5–04: Rapid determination of oil/fat utilizing high temperature solvent extraction. Official methods and recommended practices of the AOCS. 5th ed. 2nd printing. Firestone D., ed., AOCS, Champaign, IL.
- Bélanger, G., Castonguay, Y., and Lajeunesse, J. 2014. Benefits of mixing timothy with alfalfa for forage yield, nutritive value, and weed suppression in northern environments. *Can. J. Plant Sci.* **94**: 51–60.
- Bélanger, G., Michaud, R., Jefferson, P.G., Tremblay, G.F., and Brégard, A. 2001. Improving the nutritive value of timothy through management and breeding. *Can. J. Plant Sci.* **81**: 577–585.
- Barnett, F.L., and Posler, G.L. 1983. Performance of cool-season perennial grasses in pure stands and in mixtures with legumes. *Agron. J.* **75**: 582–586.
- Brito, A.F., Tremblay, G.F., Lapierre, H., Bertrand, A., Castonguay, Y., Bélanger, G., Michaud, R., Benchaar, C., Ouellet, D.R., and Berthiaume, R. 2009. Alfalfa cut at sundown and harvested as baleage increases bacterial protein synthesis in late-lactation dairy cows. *J. Dairy Sci.* **92**: 1092–1107.

- CRAAQ. 2003. Guide de référence en fertilisation. 1st ed. (In French.) Centre de Référence en Agriculture et Agroalimentaire du Québec, Québec City, QC.
- dos Passos Bernardes, A., Tremblay, G.F., Bélanger, G., Brégard, A., Seguin, P., and Vanasse, A. 2015. Sugar yield of sweet pearl millet and sweet sorghum as influenced by harvest dates and delays between biomass chopping and pressing. *Bioenerg. Res.* **8**: 100–8.
- Drapeau, R., and Bélanger, G. 2009. Comparison of meadow fescue and meadow bromegrass in monoculture and in association with white clover. *Can. J. Plant Sci.* **89**: 1059–1063.
- Finn, J.A., Kirwan, L., Connolly, J., Sebastià, M. T., Helgadóttir, Á., and 27 co-authors. 2013. Ecosystem function enhanced by combining four functional types of plant species in intensively managed grassland mixtures: a 3-yr continental-scale field experiments. *J. Appl. Ecol.* **50**: 365–375.
- Frankow-Lindberg, B.E., Brophy, C., Collins, R.P., and Connolly, J. 2009. Biodiversity effects on yield and unsown species invasion in a temperate forage ecosystem. *Ann. Bot.* **103**: 913–921.
- Goering, H.K., and Van Soest, P.J. 1970. Forage Fiber Analysis (apparatus, reagents, procedures and some applications). USDA Agriculture Handbook 379. U.S. Gov. Print Office, Washington, D.C.
- Isaac, R.A., and Johnson, W.C. 1976. Determination of total nitrogen in plant tissue, using a block digester. *J. Assoc. Off. Anal. Chem.* **59**: 98–100.
- Leco corporation. 2009. Leco Corporation, 2009. Moisture and ash determination in flour. Organic application note.

<http://www.leco.com/index.php/component/edocman/?view=document&id=193>

(accessed 16 May 2017, available after joining the mailing list). Leco Corporation, St. Joseph, MI.

Licitra, G., Hernandez, T.M., and Van Soest, P.J. 1996. Standardization of procedures for nitrogen fractionation of ruminant feeds. *Anim. Feed Sci. Technol.* **57**: 347–358.

Martens, H., and Naes, T. 2001. Multivariate calibration by data compression. p. 59–100.

In: P.C. Williams and K.H. Norris, editors, *Near infrared technology in the agricultural and food industries*. 2nd ed. Am. Assoc. Cereal Chem., St. Paul, MN.

McKenzie, D.B., Papadopoulos, Y.A., McRae, K.B., and Butt, E. 2005. Compositional changes over four years for binary mixtures of grass species grown with white clover. *Can. J. Plant Sci.* **85**: 351–360.

Mertens, D.R. 2002. Gravimetric determination of amylase-treated neutral detergent fibre in feeds with refluxing beakers or crucibles: collaborative study. *J. Assoc. Chem. Int.* **85**: 1217–1240.

NRC. 2001. *Nutrient requirements of dairy cattle*, 7th rev. ed. National Academic Press, Washington, DC.

Papadopoulos, Y.A., Martin, R.C., Fredeen, A.H., McRae, K.B., and Price, M.A. 2001. Grazing and the addition of white clover improve the nutritional quality of orchardgrass cultivars. *Can. J. Anim. Sci.* **81**: 597–600.

Pelletier, S., Tremblay, G.F., Bélanger, G., Bertrand, A., Castonguay, Y., Pageau, D., and Drapeau, R. 2010. Forage nonstructural carbohydrates and nutritive value as affected by time of cutting and species. *Agron. J.* **102**: 1388–1398.

- Picasso, V.D., Brummer, E.C., Liebman, M., Dixon, P.M., and Wilsey, B.J. 2008. Crop species diversity affects productivity and weed suppression in perennial polycultures under two management strategies. *Crop Sci.* **48**: 331–342.
- Sanderson, M.A., Brink, G., Ruth, L., and Stout, R. 2012. Grass-legume mixtures suppress weeds during establishment better than monocultures. *Agron. J.* **104**: 36–42.
- Shenk, J.S., and Westerhaus, M.O. 1991. Population definition, sample selection, and calibration procedures for near infrared reflectance spectroscopy. *Crop Sci.* **31**: 469–474.
- Simili da Silva, M., Tremblay, G.F., Bélanger, G., Lajeunesse, J., Papadopoulos, Y.A., Fillmore, S.A.E., and Jobim, C.C. 2013. Energy to protein ratio of grass-legume binary mixtures under frequent clipping. *Agron. J.* **105**: 482–492.
- Sleugh, B., Moore, K.J., George, J.R., and Brummer, E.C. 2000. Binary legume-grass mixtures improve forage yield, quality, and seasonal distribution. *Agron. J.* **92**: 24–29.
- Sturludóttir, E., Brophy, C., Bélanger, G., Gustavsson, A.-M., Jørgensen, M., Lunnan, T., and Helgadóttir, Á. 2013. Benefits of mixing grasses and legumes for herbage yield and nutritive value in Northern Europe and Canada. *Grass Forage Sci.* **69**: 229–240.
- Tracy, B.F., and Sanderson, M.A. 2004. Forage productivity, species evenness and weed invasion in pasture communities. *Agric. Ecosyst. Environ.* **102**: 175–183.
- Undersander, D., Combs, D., Shaver, R., and Hoffman, P. 2013. University of Wisconsin alfalfa/grass evaluation system – MILK 2013. UW Extension.
- <http://www.uwex.edu/ces/forage/articles.htm#milk2000> (accessed 2017-05-12).
- VSN International (2013). GenStat for Windows 17th Edition. VSN International, Hemel Hempstead, UK. Web page: GenStat.co.uk

- Vogel, K.P., and Masters, R.A. 2001. Frequency grid – a simple tool for measuring grassland establishment. *J. Range Manage.* **54**: 653–655.
- Wen, L., Williams, J. E., Kallenbach, R. L., Roberts, C. A., Beuselinck, P. R., and McGraw, R. L. 2004. Cattle preferentially select birdsfoot trefoil from mixtures of tall fescue and birdsfoot trefoil. Online. *Forage and Grazinglands* doi:10.1094/FG-2004-0924-01-RS.

For Review Only

List of figures

Figure 1. Diagram of the first two principal components (PC) of a principal component analysis to illustrate the relationship among forage nutritive attributes [ADF, acid detergent fiber; aNDF, neutral detergent fiber assayed with a heat-stable α -amylase; IVTD, *in vitro* true digestibility of dry matter; N; NDFd, *in vitro* aNDF digestibility; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch); TDN, total digestible nutrients] and dry matter yield averaged across five post-seeding years for 18 legume-grass binary mixtures (A = alfalfa, B = birdsfoot trefoil, C = white clover, Kb = Kentucky bluegrass, Mb = meadow brome grass, Mf = meadow fescue, Or = orchardgrass, Tf = tall fescue, Ti = timothy.). λ_1 and λ_2 are the contribution of the first and second principal components to the total variation.

Table 1. Site characteristics

Soil/crop information	Lévis (QC)	Normandin (QC)	Nappan (NS)
Latitude	46°48'N	48°51'N	45°46'N
Longitude	71°23'W	72°32'W	64°15'W
Elevation (above sea level)	43	137	20
Annual rainfall ¹ , mm	924	612	916
Annual temperature ¹ , °C	4.0	0.9	5.8
Growing degree-days (5°C basis) ¹	1713	1359	1718
Soil texture	Fine sandy loam	Silty clay	Loam
Soil pH (water) ²	5.2	5.9	7.1
Soil available P ² , mg kg ⁻¹	86	143	133
Soil available K ² , mg kg ⁻¹	199	284	118
Plot size, m ²	12	15	10

¹ 30-yr average (1971-2000); http://climate.weather.gc.ca/climate_normals/index_e.html.

² Values at the start of the experiment (0-20 cm).

Table 2. Cutting or grazing dates in the five post-seeding years at the three sites.

Year	Cutting/grazing	Lévis	Normandin	Nappan
2011	1	3 June	6 June	15 to 20 June
	2	22 June	27 June	9 Aug.
	3	13 July	28 July	7 Oct.
	4	17 Aug.	31 Aug.	-
	5	6 Oct.	-	-
2012	1	24 May	7 June	22 to 31 May
	2	18 June	12 July	4 to 10 July
	3	9 July	23 Aug.	-
	4	21 Aug.	-	-
	5	3 Oct.	-	-
2013	1	28 May	12 June	27 to 31 May
	2	14 June	16 July	4 to 10 July
	3	5 July	10 Sept.	19 to 23 Aug.
	4	21 Aug.	-	10 to 11 Oct.
	5	19 Sept.	-	-
2014	1	29 May	9 June	16 June
	2	16 June	3 July	4 to 10 July
	3	8 July	19 Aug.	19 to 23 Aug.
	4	8 Aug.	-	-
	5	10 Sept.	-	-
2015	1	29 May	15 June	5 to 8 June
	2	17 June	15 July	4 to 10 July
	3	9 July	18 Aug.	19 to 23 Aug.
	4	13 Aug.	-	-
	5	16 Sept.	-	-

Table 3. Statistics² of the performance of near-infrared spectroscopy to predict the nutritive attributes of the validation set of forage samples.

Attribute ¹	<i>n</i>	Slope	Mean	SD	SEP	RSQ	SEP(C)	RPD
ADF, g kg ⁻¹ DM	61	1.03	338	50.4	10.8	0.96	10.8	4.7
aNDF, g kg ⁻¹ DM	61	0.98	509	90.7	14.5	0.97	14.6	6.2
IVTD, g kg ⁻¹ DM	61	1.02	837	61.5	15.8	0.94	15.4	4.0
NDFd, g kg ⁻¹ aNDF	61	0.96	676	93.5	29.8	0.90	30.1	3.1
TDN, g kg ⁻¹ DM	61	0.96	573	77.6	20.0	0.94	19.7	3.9
NSC, g kg ⁻¹ DM	61	0.97	77	24.7	5.9	0.94	5.9	4.2
Total N, g kg ⁻¹ DM	61	0.96	24.4	7.21	1.01	0.98	1.01	7.2

¹ ADF = acid detergent fiber; aNDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* aNDF digestibility; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).

² *n* = number of samples in the validation set; SD = standard deviation; SEP = standard error of prediction; SEP(C) = standard error of prediction corrected for the bias; RSQ = coefficient of determination for the prediction; RPD = ratio of prediction to deviation [SD/SEP(C)].

Table 4. Analysis of variance (probability values) for the effects of sites, and legume species and grass species in the binary mixtures on forage dry matter yield and several nutritive attributes.

Sources of variation	DM yield	ADF ²	aNDF	IVTD	NDFd	TDN	NSC	Total N
Site (S)	0.002	0.001	<0.001	<0.001	<0.001	0.002	<0.001	<0.001
Nappan vs. (Lévis+Normandin)	0.001	ns	<0.001	<0.001	<0.001	<0.001	0.002	<0.001
Lévis vs. Normandin	0.037	<0.001	0.10	0.048	<0.001	ns	<0.001	<0.001
Legume (L)	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
C vs. (A+B) ³	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
A vs. B	<0.001	0.011	ns	0.014	<0.001	<0.001	ns	ns
Grass (G)	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Kb vs. (Ti+Tf+Or+Mf+Mb) ³	ns ¹	0.064	ns	<0.001	<0.001	<0.001	0.002	<0.001
Ti vs. (Tf+Or+Mf+Mb)	ns	<0.001	<0.001	<0.001	<0.001	<0.001	0.007	<0.001
Or vs. (Tf+Mf+Mb)	0.002	ns	0.028	ns	<0.001	<0.001	<0.001	<0.001
Mb vs. (Tf+Mf)	<0.001	<0.001	ns	<0.001	<0.001	<0.001	<0.001	<0.001
Tf vs. Mf	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	ns
S × L	<0.001	<0.001	<0.001	0.006	0.057	<0.001	<0.001	<0.001
S × G	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
L × G	<0.008	ns	0.043	ns	0.001	ns	0.005	0.013
S × L × G	ns	ns	ns	ns	0.083	ns	<0.001	0.003

¹ Not significant at $P < 0.10$.² ADF = acid detergent fiber; aNDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* aNDF digestibility; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).³ A = alfalfa, B = birdsfoot trefoil, C = white clover, Kb = Kentucky bluegrass, Mb = meadow brome grass, Mf = meadow fescue, Or = orchardgrass, Tf = tall fescue, Ti = timothy.

Table 5. Proportion (%) of each seeded species assessed at their contribution to forage DM yield of eighteen binary legume-grass mixtures in years 3, 4, and 5 after seeding. Values are the average of two measurements in each post-seeding year.

Main effects/Years	Lévis			Normandin			Nappan		
	3	4	5	3	4	5	3	4	5
Legume									
White clover	<u>2</u>	<u>8</u>	<u>2</u>	<u>15</u>	<u>7</u>	<u>0</u>	4	3	1
Birdsfoot trefoil	38	29	4	43	43	5	<u>1</u>	<u>1</u>	1
Alfalfa	30	23	6	33	31	11	8	6	4
Mean	23	20	4	31	27	6	4	3	2
SEM	2.4	2.9	0.8	1.5	1.9	0.9	0.7	1.0	0.7
Upper limit ¹	27	24	5	33	30	7	5	5	3
Lower Limit	20	16	3	28	25	4	3	2	1
Grass									
Timothy	<u>28</u>	<u>21</u>	<u>23</u>	<u>35</u>	<u>36</u>	<u>50</u>	<u>48</u>	<u>41</u>	<u>44</u>
Kentucky bluegrass	59	<u>35</u>	<u>43</u>	<u>49</u>	<u>47</u>	<u>51</u>	57	<u>39</u>	49
Tall fescue	74	71	82	64	58	61	86	69	78
Orchardgrass	58	50	62	58	57	60	60	51	49
Meadow fescue	72	72	78	65	<u>47</u>	68	70	64	65
Meadow brome	56	52	65	63	55	74	<u>39</u>	<u>21</u>	<u>23</u>
Mean	58	50	59	56	50	61	60	48	51
SEM	4.8	4.4	3.4	1.9	2.2	2.8	7.6	3.8	4.2
Upper limit ¹	63	54	62	58	52	63	68	51	56
Lower Limit	53	45	55	54	48	58	52	44	47

¹Mixture values that are greater by more than ½ the LSD above the grand mean are in bold type, while those less the same amount are underlined.

Table 6. Means of all nutritive attributes and dry matter (DM) yield across five post-seeding years and three sites for the main effects of one of three legume species and one of six grass species present in binary legume-grass mixtures.

Main effects	DM yield ¹ Mg ha ⁻¹	ADF g kg ⁻¹ DM	aNDF g kg ⁻¹ DM	IVTD g kg ⁻¹ DM	NDFd g kg ⁻¹ aNDF	TDN g kg ⁻¹ DM	NSC g kg ⁻¹ DM	Total N g kg ⁻¹ DM
Legume								
White clover	<u>4.86</u>	311	483	878	750	<u>610</u>	99	<u>23.3</u>
Birdsfoot trefoil	5.76	<u>303</u>	<u>464</u>	877	<u>734</u>	620	94	25.2
Alfalfa	5.45	305	<u>464</u>	<u>874</u>	<u>725</u>	613	<u>93</u>	25.4
SEM	0.059	0.7	1.8	0.7	0.9	1.1	0.6	0.11
Upper limit ²	5.45	307	474	877	737	617	96	24.9
Lower Limit	5.27	305	468	875	735	613	94	24.5
Grass								
Timothy	5.38	<u>290</u>	<u>439</u>	884	734	643	98	25.7
Kentucky bluegrass	5.33	308	472	<u>856</u>	<u>701</u>	<u>605</u>	<u>93</u>	25.2
Tall fescue	5.52	311	495	875	750	<u>595</u>	99	<u>23.3</u>
Orchardgrass	<u>5.12</u>	308	473	879	737	619	<u>90</u>	24.7
Meadow fescue	<u>4.92</u>	<u>304</u>	467	888	758	614	107	<u>23.3</u>
Meadow brome	5.86	316	476	876	<u>737</u>	<u>612</u>	<u>85</u>	25.6
SEM	0.083	1.0	2.5	0.9	1.3	1.6	0.8	0.15
Upper limit ²	5.48	307	475	877	738	617	96	24.9
Lower Limit	5.24	305	467	875	734	613	94	24.5
Overall mean	5.36	306	471	876	736	615	95	24.7

¹ DM = Dry matter; ADF = acid detergent fiber; aNDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* aNDF digestibility; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).

² Legume or grass values that are greater by more than ½ the LSD above the overall mean are in bold type, while those less the same amount are underlined.

Table 7. The first two component scores and variate means¹ for the 18 binary legume-grass mixtures sorted according to the first principal component (PC) score and averaged across five post-seeding years (2011-2015) at Nappan.

Mixture ²	PC 1 ($\lambda_1=55\%$)	PC 2 ($\lambda_2=30\%$)	IVTD ³ g kg ⁻¹ DM	ADF g kg ⁻¹ DM	NDFd g kg ⁻¹ aNDF	NSC g kg ⁻¹ DM	TDN g kg ⁻¹ DM	DM yield Mg ha ⁻¹	aNDF g kg ⁻¹ DM	Total N g kg ⁻¹ DM
B-Kb	4.36	0.85	<u>790</u>	363	<u>616</u>	<u>76.2</u>	<u>538</u>	7.00	562	21.8
A-Kb	3.45	-1.61	<u>795</u>	354	<u>608</u>	<u>73.8</u>	<u>556</u>	7.07	531	24.2
C-Kb	3.08	-0.12	<u>797</u>	356	<u>630</u>	<u>79.8</u>	<u>548</u>	6.69	550	23.2
A-Mb	0.98	<u>-2.06</u>	825	348	<u>658</u>	<u>75.8</u>	580	6.90	525	24.6
B-Or	0.87	0.34	820	351	673	<u>82.8</u>	572	6.66	548	21.5
C-Mb	0.77	-0.31	830	354	681	<u>82.6</u>	574	7.11	537	22.6
C- Tf	0.73	1.84	830	348	701	91.2	<u>552</u>	7.78	559	20.8
B-Tf	0.39	2.23	827	349	699	91.7	<u>553</u>	7.02	562	<u>19.9</u>
A-Tf	0.04	1.38	828	344	691	93.3	562	7.08	555	20.9
A-Or	-0.38	-1.08	830	342	673	<u>82.9</u>	585	6.37	532	22.8
C-Or	-0.45	-0.06	827	346	681	<u>86.9</u>	577	<u>5.99</u>	536	21.6
A-Mf	-0.79	1.36	834	345	696	95.5	568	6.59	547	<u>20.2</u>
A-Ti	<u>-1.40</u>	<u>-2.14</u>	839	<u>332</u>	679	86.8	602	7.28	<u>514</u>	23.4
B-Mb	<u>-1.50</u>	<u>-2.88</u>	837	<u>336</u>	680	84.3	595	6.41	<u>499</u>	24.5
B-Ti	<u>-2.28</u>	-0.36	837	<u>335</u>	690	94.0	596	<u>5.90</u>	528	20.9
C-Mf	<u>-2.36</u>	2.13	841	341	708	109.4	570	6.23	539	<u>18.8</u>
B-Mf	<u>-2.39</u>	1.31	840	340	706	102.7	571	<u>5.70</u>	532	<u>19.6</u>
W-Ti	<u>-3.10</u>	-0.82	845	<u>331</u>	706	96.2	605	6.30	<u>520</u>	21.4
Mean	(0.00)	(0.00)	826	345	676	88.1	572	6.67	538	21.8
SEM	(1.00)	(1.00)	6.4	5.1	8.2	3.63	8.1	0.329	9.8	0.72
LSD (5%)										
upper limit ⁴	1.20	1.21	835	353	688	93.3	584	7.14	552	22.8
Lower limit	-1.21	-1.21	817	338	665	82.9	561	6.20	524	20.8

¹Variates were arranged according to their correlation with the first PC score. λ_1 = The contribution of the first principal component to the total variation; λ_2 = The contribution of the second principal component to the total variation.

²A = alfalfa, B = birdsfoot trefoil, C = white clover, Kb = Kentucky bluegrass, Mb = meadow brome grass, Mf = meadow fescue, Or = orchardgrass, Tf = tall fescue, Ti = timothy.

³ADF = acid detergent fiber; aNDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* digestibility of NDF; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).

⁴Mixture values that are greater by more than ½ the LSD above the overall mean are in bold type, while those less the same amount are underlined.

Table 8. The first two component scores and variate means¹ for the 18 binary legume-grass mixtures sorted according to the first principal component (PC) score and averaged across five post-seeding years (2001-2015) at Lévis.

Mixture ²	PC 1 ($\lambda_1=58\%$)	PC 2 ($\lambda_2=20\%$)	Total N ³ g kg ⁻¹ DM	aNDF g kg ⁻¹ DM	TDN g kg ⁻¹ NDF	ADF g kg ⁻¹ DM	dNDFd g kg ⁻¹ aNDF	DM yield Mg ha ⁻¹	NSC g kg ⁻¹ DM	IVTD g kg ⁻¹ DM
A-Ti	4.08	-0.78	30.5	<u>382</u>	663	<u>278</u>	<u>740</u>	5.68	<u>94</u>	903
B-Ti	3.32	-1.30	29.3	<u>401</u>	664	<u>278</u>	<u>757</u>	6.19	98	905
A Kb	2.11	0.51	29.6	<u>408</u>	637	<u>289</u>	<u>732</u>	5.21	97	<u>888</u>
B Or	1.59	0.24	28.3	<u>427</u>	642	<u>295</u>	<u>760</u>	5.99	<u>92</u>	899
B-Kb	1.54	0.77	28.3	<u>422</u>	637	<u>290</u>	<u>735</u>	5.43	97	<u>885</u>
B-Mb	1.17	1.91	28.5	442	633	306	765	7.01	<u>84</u>	897
A-Or	1.02	0.39	27.8	<u>426</u>	634	299	<u>751</u>	5.03	<u>91</u>	896
A-Mb	0.75	2.19	28.4	441	621	308	<u>753</u>	6.34	<u>85</u>	892
B-Mf	0.57	<u>-1.45</u>	26.6	<u>428</u>	633	<u>292</u>	774	5.77	104	904
C-Ti	-0.30	<u>-1.37</u>	<u>25.2</u>	<u>425</u>	628	297	759	<u>3.93</u>	103	896
A-Mf	-0.43	<u>-1.48</u>	<u>25.1</u>	435	623	296	770	<u>4.88</u>	104	902
C-Or	<u>-1.39</u>	-0.27	<u>25.0</u>	461	624	309	778	<u>4.32</u>	95	897
C-Kb	<u>-1.42</u>	0.01	<u>24.9</u>	453	622	305	<u>754</u>	<u>3.77</u>	102	<u>884</u>
A-Tf	<u>-1.56</u>	1.06	<u>25.1</u>	465	<u>601</u>	305	758	5.32	100	<u>884</u>
B-Tf	<u>-2.24</u>	1.35	<u>24.1</u>	477	<u>596</u>	309	764	5.93	102	<u>883</u>
C-Mb	<u>-2.35</u>	0.56	<u>24.6</u>	477	<u>611</u>	320	787	<u>4.76</u>	<u>92</u>	897
C-Mf	<u>-2.55</u>	<u>-2.29</u>	<u>23.1</u>	461	<u>614</u>	305	788	<u>3.93</u>	110	902
C-Tf	<u>-3.90</u>	-0.03	<u>22.3</u>	492	<u>592</u>	312	780	<u>4.35</u>	107	<u>886</u>
Mean	(0.00)	(0.00)	26.5	440	626	300	761	5.21	98	895
SEM	(1.00)	(1.00)	0.29	5.1	3.3	2.4	3.6	0.216	2.1	1.8
LSD (5%)										
upper limit ⁴	1.20	1.21	26.9	447	631	303	767	5.52	101	897
Lower limit	-1.21	-1.21	26.1	433	621	296	756	4.90	95	892

¹Variates were arranged according to their correlation with the first PC score. λ_1 = The contribution of the first principal component to the total variation; λ_2 = The contribution of the second principal component to the total variation.

² A = alfalfa, B = birdsfoot trefoil, C = white clover, Kb = Kentucky bluegrass, Mb = meadow brome grass, Mf = meadow fescue, Or = orchardgrass, Tf = tall fescue, Ti = timothy.

³ADF = acid detergent fiber; NDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* digestibility of NDF; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).

⁴Mixture values that are greater by more than ½ the LSD above the overall mean are in bold type, while those less the same amount are underlined.

Table 9. The first two component scores and variate means¹ for the 18 legume-grass mixtures sorted according to the first principal component (PC) score and averaged across five post-seeding years (2011-2015) at Normandin.

Mixture ²	PC1 ($\lambda_1=50\%$)	PC2 ($\lambda_2=29\%$)	NDF ³ g kg ⁻¹ DM	N g kg ⁻¹ DM	TDN g kg ⁻¹ DM	ADF g kg ⁻¹ DM	NDFd g kg ⁻¹ aNDF	DM yield Mg DM ha ⁻¹	IVTD g kg ⁻¹ DM	NSC g kg ⁻¹ DM
B-Ti	4.32	-0.26	<u>396</u>	29.0	663	<u>292</u>	<u>726</u>	4.81	895	<u>76</u>
A-Ti	2.99	-0.73	<u>405</u>	27.5	649	<u>298</u>	<u>719</u>	4.07	894	81
C-Ti	1.96	<u>-1.48</u>	<u>420</u>	26.9	651	<u>303</u>	745	<u>3.61</u>	897	<u>77</u>
B-Kb	1.89	<u>1.58</u>	<u>429</u>	27.0	631	<u>308</u>	<u>710</u>	4.89	<u>874</u>	83
B-Mf	1.37	<u>-1.48</u>	<u>429</u>	26.6	634	<u>305</u>	752	4.23	899	84
B-Or	0.78	0.92	450	26.3	629	315	<u>733</u>	4.66	888	<u>72</u>
A-Kb	0.77	1.42	<u>438</u>	26.1	618	315	<u>699</u>	4.13	<u>871</u>	86
B-Mb	0.45	2.78	458	27.0	620	329	<u>730</u>	5.42	<u>881</u>	<u>66</u>
B-Tf	-0.21	-0.17	460	25.3	<u>614</u>	<u>311</u>	755	4.64	890	79
A-Mf	-0.32	<u>-1.81</u>	447	24.9	617	312	746	<u>3.54</u>	895	91
A-Or	-0.77	0.12	461	25.0	<u>615</u>	321	<u>733</u>	<u>3.53</u>	886	<u>76</u>
A-Mb	-0.80	2.03	468	25.8	<u>607</u>	333	<u>727</u>	4.61	<u>881</u>	<u>71</u>
C-Mf	<u>-1.24</u>	<u>-2.63</u>	460	<u>24.1</u>	617	316	771	<u>3.45</u>	901	89
C-Kb	<u>-1.47</u>	0.85	477	<u>24.3</u>	<u>608</u>	324	<u>727</u>	<u>3.79</u>	<u>873</u>	84
A-Tf	<u>-1.86</u>	-0.73	475	<u>23.7</u>	<u>599</u>	318	753	<u>3.80</u>	886	86
C-Or	<u>-1.94</u>	0.32	482	<u>24.2</u>	<u>611</u>	328	750	<u>3.56</u>	885	<u>70</u>
C-Tf	<u>-2.46</u>	<u>-1.98</u>	484	<u>23.0</u>	<u>601</u>	319	777	<u>3.79</u>	895	89
C-Mb	<u>-3.46</u>	1.26	508	<u>23.3</u>	<u>591</u>	342	755	4.14	886	<u>68</u>
Mean	(0.00)	(0.00)	453	25.5	621	316	739	4.15	888	79
SEM	(1.00)	(1.00)	5.5	0.37	3.6	2.7	1.8	0.136	2.3	1.3
LSD (5%)										
upper limit ⁴	1.20	1.21	460	26.1	626	320	742	4.34	891	81
Lower limit	-1.21	-1.21	445	25.0	616	312	737	3.95	884	78

¹Variates were arranged according to their correlation with the first PC score. λ_1 = The contribution of the first principal component to the total variation; λ_2 = The contribution of the second principal component to the total variation.

² A = alfalfa, B = birdsfoot trefoil, C = white clover, Kb = Kentucky bluegrass, Mb = meadow brome grass, Mf = meadow fescue, Or = orchardgrass, Tf = tall fescue, Ti = timothy.

³ ADF = acid detergent fiber; NDF = neutral detergent fiber assayed with a heat-stable α -amylase; IVTD = *in vitro* true digestibility of dry matter; NDFd = *in vitro* digestibility of NDF; TDN = total digestible nutrients; NSC, non-structural carbohydrates (water soluble carbohydrates plus starch).

⁴Mixture values that are greater by more than ½ the LSD above the overall mean are in bold type, while those less the same amount are underlined.

Table 10. Estimated milk production per hectare from the 18 legume-grass mixtures averaged across five post-seeding years (2011-2015) at the three sites.

Legume	Grass	Milk production (Mg ha ⁻¹)			
		Lévis	Normandin	Nappan	Mean
White clover	Timothy	<u>7.26</u>	<u>6.79</u>	11.15	<u>8.52</u>
White clover	Kentucky bluegrass	<u>6.70</u>	<u>6.55</u>	10.57	<u>8.12</u>
White clover	Tall fescue	<u>7.09</u>	<u>6.35</u>	12.20	<u>8.79</u>
White clover	Orchardgrass	<u>7.67</u>	<u>6.19</u>	<u>9.94</u>	<u>8.12</u>
White clover	Meadow fescue	<u>6.79</u>	<u>5.95</u>	<u>10.10</u>	<u>7.91</u>
White clover	Meadow brome	<u>8.23</u>	7.07	11.83	9.28
Birdsfoot trefoil	Timothy	11.87	9.28	<u>10.22</u>	10.30
Birdsfoot trefoil	Kentucky bluegrass	9.94	8.84	10.90	10.05
Birdsfoot trefoil	Tall fescue	9.72	8.02	10.97	9.76
Birdsfoot trefoil	Orchardgrass	11.15	8.41	10.99	10.35
Birdsfoot trefoil	Meadow fescue	10.39	7.59	<u>9.25</u>	9.22
Birdsfoot trefoil	Meadow brome	12.72	9.66	10.90	11.23
Alfalfa	Timothy	10.87	7.61	12.75	10.56
Alfalfa	Kentucky bluegrass	9.50	7.24	11.39	9.51
Alfalfa	Tall fescue	8.85	<u>6.37</u>	11.34	<u>9.01</u>
Alfalfa	Orchardgrass	9.14	<u>6.22</u>	10.70	<u>8.85</u>
Alfalfa	Meadow fescue	<u>8.62</u>	<u>6.16</u>	10.72	<u>8.65</u>
Alfalfa	Meadow brome	11.17	7.95	11.55	10.39
Mean		9.32	7.35	10.97	9.37
SEM		0.384	0.244	0.499	0.245
LSD (5%)					
Upper limit ¹		9.87	7.70	11.69	9.72
Lower limit		8.76	7.00	10.25	9.02

¹Mixture values that are greater by more than ½ the LSD above the overall mean are in bold type, while those less the same amount are underlined.

Figure 1.