Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

Heavy sea louse infection is associated with decreased stomach fullness in wild juvenile sockeye salmon

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0267.R1
Manuscript Type:	Article
Date Submitted by the Author:	29-Nov-2017
Complete List of Authors:	Godwin, Sean; Simon Fraser University, Krkošek, Martin; University of Toronto, Department of Ecology and Evolutionary Biology Reynolds, John D.; Simon Fraser University, Department of Biological Sciences Rogers, Luke; University of Toronto, Department of Ecology and Evolutionary Biology Dill, Lawrence; Simon Fraser University,
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	FORAGING < General, PARASITES < Organisms, SALMON < Organisms, COMPETITION < General

SCHOLARONE™
Manuscripts

Heavy sea louse infection is associated with decreased stomach fullness in wild juvenile sockeye
salmon

Sean C. Godwin^{1,*}, Martin Krkošek^{2,3}, John D. Reynolds¹, Luke A. Rogers², Lawrence M. Dill¹

¹ *Earth to Ocean Research Group, Department of Biological Sciences, Simon Fraser University,
Burnaby, British Columbia, Canada, V5A 1S6*

² *Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario,
Canada, M5S 3B2*

³ *Salmon Coast Field Station, Simoom Sound, British Columbia, Canada, V0P 1S0*

** Corresponding author: E-mail: sgodwin@sfu.ca, Tel: 778-782-3989*

Co-author e-mail addresses:

Martin Krkošek: martin.krkosek@utoronto.ca

John Reynolds: reynolds@sfu.ca

Luke Rogers: luke.rogers@mail.utoronto.ca

Lawrence Dill: ldill@sfu.ca

Abstract

Foraging success can be mediated by parasites, but this is poorly understood for marine fish whose aggregations and patchy prey fields create conditions for intense intraspecific competition. We evaluated whether sea louse infection is associated with decreased stomach fullness of wild juvenile sockeye salmon (*Oncorhynchus nerka*) in Johnstone Strait, BC during their marine migration from the Fraser River. *Caligus clemensi* comprised 98.6% of the pre-adult and adult lice and 86.5% of the copepodites (freshly attached juvenile lice); the rest were *Lepeophtheirus salmonis*. We found that infection status was an important predictor of relative stomach fullness for juvenile sockeye (wet stomach content weight divided by body weight), as indicated by mixed-effects model selection, and that highly infected fish had $17\% \pm 8\%$ lower relative stomach fullness than did lightly infected fish. This louse-associated reduction in relative stomach fullness occurs as the juvenile sockeye migrate through a food-limited environment, and presumably elevated competition. Given that early marine growth for juvenile salmon is often a predictor of survival, our results highlight the importance of understanding sub-lethal effects of parasites on salmonids and possibly other fish species.

Introduction

Foraging success is tightly linked to growth and survival of individuals, both of which are predictors of fitness and population dynamics (Crombie 1947; Sutherland 1996). Fish frequently experience patchy foraging opportunities and aggregate in large groups that intensify competition (see review by Ward et al. (2006)), and consequently survival often depends on these competitive outcomes (e.g., Resetarits 1995; Welker et al. 1994). This is especially the case for juvenile fishes, whose early growth depends on food supply and often determines survival and recruitment (Anderson 1988; Bergenius et al. 2002; Houde and Hoyt 1987). Evidence suggests that parasites affect competitive foraging outcomes of hosts, thereby influencing host population dynamics, community structure, and biodiversity (see Hatcher et al. 2006). However, this evidence is primarily from terrestrial species (e.g., Grosholz 1992; Maksimowich and Mathis 2000; Schall 1992). Despite what is likely a more competitive environment with food patchiness and consumer aggregation for fishes, there is little work in the fisheries literature on how parasites mediate competition in wild fish and whether this translates to decreased survival through reduced foraging success and growth (but see Finley and Forrester 2003).

Pacific salmon (*Oncorhynchus* spp.), like many fishes, experience high juvenile mortality from predation and starvation (Groot and Margolis 1991; Parker 1968). Consequently, the impacts of parasitism may be primarily expressed through the mechanisms by which parasitism affects inter- and intra- specific interactions. Juvenile Pacific salmon migrate in large groups to swamp and evade predators (Eggers 1978; Furey et al. 2016), so competition for food is probable in regions with low prey availability (McKinnell et al. 2014). Foraging success during the marine migration of juvenile salmon likely affects growth, which is often a predictor of survival (Duffy and Beauchamp 2011; Farley et al. 2007; Moss et al. 2005), so competition in environments of

low foraging opportunity is one plausible mechanism through which parasites may affect survival of juvenile salmon (Godwin et al. 2015). Although parasitism can raise energetic requirements and thereby increase foraging rate (e.g., Giles 1987; Shi et al. 2002), parasitism may also interfere with the behavioural process of obtaining food in a food-limited environment and thereby reduce foraging success (e.g., Barber and Ruxton 1998).

During the ocean phase of their juvenile migration, Pacific salmon are susceptible to infection by sea lice (*Lepeophtheirus salmonis* and *Caligus clemensi*), which are native ectoparasites that feed on the surface tissue of their host (Wootten et al. 1982). Juvenile Pacific salmon normally have low infection levels of sea lice, especially the salmonid specialist *L. salmonis*, because these salmon are temporally and spatially separated from adult Pacific salmon (Krkošek et al. 2007). However, in recent decades domesticated Atlantic salmon (*Salmo salar*) farmed in open-net pens in coastal British Columbia (BC) have provided year-round reservoirs for sea lice that allow substantial transmission of *L. salmonis* to juvenile Pacific salmon (Costello 2009b; Groner et al. 2016). For the generalist *C. clemensi*, there also exist other natural host infection reservoirs, such as Pacific herring (*Clupea pallasii*) (Beamish et al. 2009; Morton et al. 2008), which share nearshore coastal waters with juvenile salmon as spawners and larvae (Beamish et al. 2012).

In recent years, the generalist *C. clemensi* has infected over 98% of out-migrating juvenile sockeye salmon (*O. nerka*) from the Fraser River (Godwin et al. 2015; Godwin et al. 2017), an iconic set of Pacific salmon populations that forms Canada's largest sockeye run. Juvenile sockeye salmon that experience high infection intensity by *C. clemensi* exhibit reduced foraging success in a competitive and food-limited experimental setting (Godwin et al. 2015). Whether this translates to reduced foraging success in the wild is still unknown, but is the focus

of this work. Here, we tested whether the relative stomach fullness (wet stomach content weight divided by body weight) of wild juvenile sockeye is lower when sea louse infection intensity is higher, which has implications for our understanding of how parasites mediate competition in fishes and potentially for the management of *C. clemensi* in BC.

Methods

Fish collection

We collected juvenile sockeye salmon in the wild as they migrated through western Johnstone Strait, BC, between May 26 and June 7, 2014 (Fig. 1). At this point in their migration, the sockeye post-smolts have well-developed scales and average 114 mm in fork length. We used a hand-operated purse seine net (bunt: 27 x 9 m with 13 mm mesh, tow: 46 x 9 m with 76 mm mesh) that we set from a small (6 m) open boat. After surrounding the fish, we brought the net next to the boat to form a pocket of sufficient width and depth to allow the fish to swim freely and minimize their contact with the mesh.

We transferred captured fish from the net into an insulated fish tote (0.58 m deep and 0.97 x 0.55 m across) filled with fresh seawater. We moved fish from the net into the tote by dipping them and their surrounding seawater into a 3.79 L container (an inverted milk jug with the top capped and bottom cut off). This transfer method minimized or prevented sea louse detachment as fish were never exposed to air and there was minimal contact between the fish and sampling equipment (Godwin et al. 2015). All subsequent transfers were also performed using the same method. We used ice packs to regulate water temperature in the tote and aquarium bubblers to maintain adequate aeration. We transferred 50–100 sockeye into the tote during each capture event. See Supplementary Table S1 for collection locations, catch sizes, and

oceanographic data.

Infection status assessment

We transferred sockeye individually from the tote into 13.2 L clear plastic aquaria and assessed them for sea louse infection by eye. If a fish appeared to be in one of our two infection categories (see below), we recorded that category and transferred the fish to an individual sterile 532 mL sample bag (Whirl-Pak® Write-On Bags; Nasco, Fort Atkinson, Wisconsin, USA) and euthanized it with an overdose of MS-222 (240 mg L⁻¹). After euthanizing a fish, we performed a full assessment of its infection status using a hand lens (Krkošek et al. 2005a) to confirm its infection category. Fish that were euthanized but found not to meet our infection category criteria were not used; such fish accounted for approximately 10% of euthanized fish and were usually identified initially to be in the lightly infected category by eye but then found to have too many small juvenile lice upon inspection by hand lens. We alternated between processing highly infected and lightly infected fish so as to not confound digestion time with infection status.

Larger, more developed sea lice have greater effects on their hosts (Jakob et al. 2013; Nendick et al. 2011; Wootten et al. 1982). We created infection categories that reflected this differential level of pathogenicity, so that small juvenile lice were not weighted equally to large adult lice (similar to Peacock et al. 2015). Sea lice initially attach to their host as copepodites, then develop through two (*L. salmonis*; Hamre et al. 2013) or four (*C. clemensi*; Kabata 1972) attached chalimus stages of increasing size before molting into their motile pre-adult and adult stages. Here, we consider individuals in their second *L. salmonis* chalimus stage or their third or fourth *C. clemensi* chalimus stages as ‘large chalimus’ sea lice; we also consider pre-adult and adult individuals as ‘motile’ sea lice. To weight the infection statuses of fish according to the

development of lice infecting them, we defined a louse infection scale in which one large chalimus louse was equal to one infection unit, one motile louse was equal to two infection units, and copepodite and small chalimus lice were equal to zero. Because of the high infection prevalence observed ($> 98\%$), we were unable to create a category for uninfected fish. Instead, we created a ‘lightly infected’ category in which all the fish had zero infection units and no more than three copepodite or small chalimus lice (Table 1). In all collections, the ‘lightly infected’ and ‘highly infected’ categories differed by a minimum of three infection units. See Supplementary Table S2 for the detailed sea louse infection data.

Approximately one hour after capture, we released the remaining fish at the collection site. In each collection, we retained five to seven pairs of fish, each consisting of a lightly and highly infected sockeye. In total, we retained 130 juvenile sockeye salmon from across 11 collections to analyze stomach fullness in relation to infection status.

Zooplankton sampling and analysis

Immediately following the infection status assessments and subsequent release of fish, we collected zooplankton samples with a horizontal plankton tow at the fish capture site. Plankton tows were performed with a 0.5 m diameter plankton net with 250 μm mesh. The top of the net was kept 5-10 cm below the ocean surface, on average, and the tow lasted for 30 s. We attached a calibrated flow meter (General Oceanics, Miami, Florida, USA) at the mouth of the net to measure the volume of water sampled. We used horizontal tows instead of vertical ones because plankton samples near the surface are more similar to sockeye diets than those from deeper in the water column (Landingham et al. 1998) and our field observations indicate that juvenile sockeye frequently occupy the surface waters at the collection sites.

Each plankton sample was placed in a glass jar containing 250 mL of 10% formalin-seawater solution. Within three days of the collection, the samples were poured into a 63 μ m sieve and rinsed with distilled water. The samples were then halved repeatedly using a Folsom plankton splitter (Aquatic Research Instruments, Hope, Idaho, USA) until approximately 200-250 individuals remained, after which they were transferred to a Bogorov counting tray (Wildlife Supply Company, Yulee, Florida, USA). From these samples, we identified and enumerated individuals from the high-level taxa previously found in juvenile Fraser River sockeye salmon stomachs (Price et al. 2013); these were termed 'sockeye prey'. These high-level taxa from Price et al. (2013) were: Copepoda, Brachyura, Oikopleura, Euphausiacea, Cladocera, Pteropoda, Decapoda, Amphipoda, Insecta, Cumacea, fish, and eggs. We calculated sockeye prey density by multiplying the number of sockeye prey in the Bogorov tray by the reciprocal of the splitting fraction, then dividing by the volume of water that passed through the plankton net.

Determination of relative stomach fullness

We used wet stomach content weight as our measure of stomach fullness, and divided this by the fish's body weight to calculate relative stomach fullness. Following the field collections, we transported the euthanized fish on ice to a laboratory facility 45 min away by boat. Upon arrival at the laboratory, fish were analyzed in the original order of collection by dissecting them immediately to weigh the wet stomach contents. Stomachs were excised between the lower oesophagus and the pyloric sphincter, and their contents extruded with forceps and weighed, as in Godwin et al. (2015).

Statistical analysis

To determine which biological variables best explained juvenile sockeye salmon relative stomach fullness, we fit a suite of 13 mixed-effects models to our data (see Supplementary Table S3 for the full model set). Our models included the biologically relevant combinations of five fixed effects: infection status, body size, prey density, the two-way interaction between infection status and body size, and the two-way interaction between infection status and prey density. We included infection status as a fixed effect to test our main hypothesis that relative stomach fullness decreases with high intensities of sea louse infection. We included body size and its interaction with infection status to account for any additional benefit of a larger body size to relative stomach fullness, and the potential decreasing effect of infection with body size (Godwin et al. 2015). Finally, we included prey density and its interaction with infection status to account for the probable association between relative stomach fullness and prey density, and the possible decreasing effect of infection with increasing density of prey. Prey density and relative stomach fullness were centered and scaled by one standard deviation to allow the model-fitting optimizer to function correctly, since their variances were different by 10 orders of magnitude. Each model included a random effect of collection number on the intercept, which was determined *a priori* to account for the hierarchical structure of the sampling design, and a variance structure allowing for different variances in each collection to account for heteroscedasticity in the residuals. We performed model selection using Akaike Information Criterion corrected for small sample sizes (Hurvich and Tsai 1989) as our measure of model parsimony. We calculated relative variable importance (RVI) values based on the AICc weights.

We derived our measure of body size from a principal component analysis using three highly correlated body metrics: fork length, body depth, and weight. The first principal component explained 98% of the original variation in these metrics and so we used that as the

variable representing body size in the statistical analyses.

We tested for differences in body size between infection categories using a two-sample t -test, and used linear regression to assess whether motile or overall louse abundance increased with body size for the highly infected fish. We completed all our analysis in R 3.2.1 (R Foundation for Statistical Computing, 2015) using the nlme and MuMIn packages.

Results

The juvenile sockeye salmon in our highly infected category were primarily infected by *C. clemensi* rather than *L. salmonis*, with 98.6% of the motile sea lice and 86.5% of the copepodid lice infecting these fish belonging to the former species. Neither motile abundance ($R^2 = 0.009$, d.f. = 63, $p = 0.455$) nor overall louse abundance ($R^2 = 0.025$, d.f. = 63, $p = 0.207$) increased with body size for fish in the highly infected category. Body size also did not differ between infection categories ($t = 0.282$, d.f. = 128, $p = 0.779$). The mean ($\pm$ SE) zooplankton density across all collections was 941 individuals $L^{-1} \pm 141$ individuals L^{-1} , and the mean sockeye prey density was 772 individuals $L^{-1} \pm 134$ individuals L^{-1} , indicating that potential sockeye prey constituted 82% of the zooplankton community in the surface waters during our collections.

Lightly infected fish had higher relative stomach fullness than highly infected fish (Fig. 2), and this trend held in each of the 11 collections (Fig. 3). Infection status was the most important predictor of relative stomach fullness; of the 13 mixed-effects models, the top nine all included an infection status term, while none of the bottom four did (Supplementary Table S3). The importance of infection status to relative stomach fullness was corroborated by this predictor having the highest RVI (0.97), compared with body size (0.66), prey density (0.60), the

interaction between infection status and prey density (0.16), and the interaction between infection status and body size (0.14). Regardless of their rank, all models with the relevant terms revealed that relative stomach fullness was higher for lightly infected fish, that it increased with body size, and that it increased with prey density. The top-ranked model included only an infection status term, but two other models were within 2 AICc units of the top model and therefore also had substantial support (Burnham and Anderson 2002; Table 2). One of these highly-supported models included a body size term, while the other included a predictor for prey density (Table 2).

The highest-ranked model without an infection status predictor was 7.8 AICc units higher than the top model and accounted for only 0.7% of model support, as judged by AICc weights (Supplementary Table S2). The top-ranked model was 51 times more likely than the highest-ranked model without an infection status predictor, and its coefficients indicated that highly infected fish had $17\% \pm 8\%$ lower relative stomach fullness than lightly infected fish, on average.

Discussion

Theory and empirical evidence suggest that pathogens and parasites can influence host survival and population dynamics by modulating competitive foraging interactions (Hatcher et al. 2006). However, there is little evidence of parasite-mediated intraspecific competition in wild fishes, for whom intraspecific competition may be particularly intense due to fish aggregation and food patchiness. Our results indicate that, for wild juvenile sockeye salmon, high levels of sea louse infection are associated with reduced relative stomach fullness. The juvenile sockeye used in this study were captured during their early marine migration from their natal freshwater

systems, which based on the timing of capture and genetic analyses from previous studies (Godwin et al. 2017; Groot and Cooke 1987; Price et al. 2011) were mostly in the Fraser River watershed of BC.

While our stomach fullness data describe the quantity of prey consumed by sockeye, it should be noted that stomach fullness is not a true measure of foraging success or efficiency. Stomach fullness does not account for the energy densities or digestibility of prey, which vary among the zooplankton prey items that dominate the diet of juvenile sockeye (Foy and Norcross 1999; Lee 1974) as well as spatially and temporally with the availability of those items (Landingham et al. 1998; Mackas et al. 2013; Tanasichuk and Routledge 2011). Parasitized individuals can shift their diet to prey items of lower energy density or digestibility when they struggle to compete with unparasitized conspecifics for higher-quality prey (Milinski 1984). Since juvenile sockeye with heavy sea louse infection have lower competitive foraging abilities (Godwin et al. 2015), it is possible that they too shift toward capturing more prey items of lower quality. If that were the case, then by using stomach fullness data we produced conservative estimates of the differences in foraging success between infection categories.

Animal migration is generally demanding metabolically, so managing energy gain and depletion is vital for most migrating animals to avoid starvation or the sub-lethal effects of depleted energy reserves (Sapir et al. 2011). Unlike some migratory species, juvenile sockeye salmon forage during their migration, but this foraging is temporally variable because feeding opportunities are patchy (McKinnell et al. 2014; Parsons et al. 1970). In regions with relatively high productivity, such as the northern Strait of Georgia, BC (Masson and Peña 2009; Parsons et al. 1970), no evidence of food limitation has been observed (Price et al. 2013). By contrast, the region in which we captured fish for this study, Johnstone Strait, has a sparse prey field due to

strong tidal mixing (B. Hunt et al., Hakai Institute, unpublished data), which may cause high mortality for juvenile sockeye that enter the Strait with insufficient energy reserves (McKinnell et al. 2014). Sea lice levels on juvenile sockeye are also considerably higher in Johnstone Strait than in the more productive southern regions (Godwin et al. 2015; Price et al. 2011). Hence, the potential effects of sea lice on sockeye salmon growth and survival in Johnstone Strait are likely to involve the elevated abundances of the parasite itself, its effects on intraspecific competition, and the intensification of competition due to food limitation and the energy expenditure of migration.

Our results provide equivocal evidence for an association between prey density and the relative stomach fullness of juvenile sockeye. If food competition is indeed higher in Johnstone Strait for sockeye than in nearby regions and that contributed to our finding that relative stomach fullness is associated with heavy sea louse infection, then we might have expected to find a stronger relationship between prey density and relative stomach fullness. Only one of our three models with considerable support (as judged by AICc values; Burnham and Anderson 2002) contained a prey density predictor, which across our model set was our third most important predictor variable ($RVI = 0.60$). It is possible that prey density did not severely limit relative stomach fullness of sockeye in Johnstone Strait in 2014 even if food availability is commonly much lower there than in the Strait of Georgia (McKinnell et al. 2014; B. Hunt et al., Hakai Institute, unpublished data.). However, if the prey field was not limiting sockeye foraging rates then we might not expect to observe the differences in relative stomach fullness between infection categories that we did, since highly infected (and therefore less competitive (Godwin et al. 2015)) individuals would still have the opportunity to feed to satiation. It is also possible that the prey density estimates from our zooplankton collections did not represent the prey field

encountered by the fish when they were feeding, due to patchiness in time or space (Parsons et al. 1970) or the strong tidal currents known to occur in Johnstone Strait (Sutherland et al. 2007). Most of the sockeye in this study must have eaten in the preceding eight hours (see Appendix), which limits the potential effects of patchiness and currents on our results, but these remain possibilities that cannot be excluded. Perhaps the most likely explanation for our equivocal evidence of a relationship between prey density and relative stomach fullness was that the sample size of 11 collection sites was too small to detect an effect. While our study design was suitable for looking at consistent differences in relative stomach fullness within groups, the sample size was limited for investigating variation among collections.

The relationship that we found between sea louse infection and relative stomach fullness is correlative, but the weight of evidence is building that sea lice have sub-lethal effects on important determinants of sockeye salmon survival. Sea lice are associated with Pacific salmon population declines (Connors et al. 2010; Krkošek et al. 2011a), but these declines cannot be explained by direct mortality alone, which may only be significant at small host sizes (Jones et al. 2008). However, sea lice also appear to have sub-lethal effects that influence mortality through their hosts' ecological interactions, for example by reducing swimming endurance (Mages and Dill 2010), increasing risk-taking behaviours (Krkošek et al. 2011b), and elevating predation (Peacock et al. 2015). For juvenile sockeye salmon specifically, individuals that are heavily infected by sea lice, primarily *C. clemensi*, have lower competitive foraging ability (Godwin et al. 2015), reduced body growth (Godwin et al. 2017), and actively attempt to dislodge these parasites by leaping (Atkinson et al. In review). Nonetheless, laboratory studies utilizing experimental infections are needed to help differentiate two alternative interpretations of our results: 1) that sea lice abundance is a consequence rather than a cause of variation in

relative stomach fullness; or 2) that sea louse abundance and relative stomach fullness are both correlated with (signals of) fundamental underlying fitness variation among individuals. We consider these alternative explanations to be unlikely because they both require sustained differences in relative stomach fullness that would have led to a difference in body size between the two infection categories, which was not observed. It is also striking that the observed louse-associated differences in relative stomach fullness occurred in each of the 11 collections (Fig 3), so any correlation between an underlying condition and sea louse infection would have to be very strong indeed.

In addition to our main result that heavy sea louse infection is associated with reduced sockeye stomach fullness, we also found moderate evidence that relative stomach fullness increased with body size. This result runs counter to the negative exponential relationship between relative stomach fullness and body size reported by Brett (1971) for juvenile sockeye, suggesting that larger juvenile sockeye may have a foraging advantage due to their body size. The obvious potential mechanism for this is that smaller sockeye are prevented from foraging on larger (and possibly more abundant) prey due to gape limitation, which often, but not always, determines foraging success in young fish (Bremigan and Stein 1994; Devries et al. 1998; Hargreaves and LeBrasseur 1986; Scharf et al. 2000). Together with the relationship between infection status and relative stomach fullness, this potential effect of body size would be consistent with our previous findings that high infection intensities and smaller body sizes are associated with lower competitive foraging ability (Godwin et al. 2015) and reduced growth (Godwin et al. 2017) for juvenile sockeye. When considered together (Fig. 4), these interconnected relationships suggest that sea louse infection may not just be accompanied by a single, temporary effect for the duration of infection. Infection may instead have long-lasting

effects by initiating and/or intensifying divergent growth among individuals through differences in foraging success, which would create and reinforce intraspecific heterogeneity in body sizes and ultimately differential survival. To detect such differences in body size, the amount of time between sea louse exposure and fish collection would need to be greater than that of this study (3-10 days; Price et al. 2011; Welch et al. 2011), since the small amount of differential body growth that would occur between infection categories in that time (< 0.5 mm; Godwin et al. 2017) would be statistically swamped by the individual-level variation in body size. This potential for lasting effects of sea lice on factors critical to sockeye survival highlights the need to consider the anthropogenic influence on the parasite burdens of these fish, and potentially reduce it through management actions when possible.

Over 98% of the motile sea lice infecting the juvenile sockeye salmon were *C. clemensi*, but although these infections have been linked to open net-pen salmon farms (Price et al. 2011), there are currently no management actions directed at regulating this louse species on farms in BC. *Lepeophtheirus salmonis*, the main species of sea louse that causes fish mortality and financial loss to salmon aquaculture in the northern hemisphere (Costello 2009a; Johnson et al. 2004; Mustafa et al. 2001), is controlled on BC farms through application of in-feed parasiticide (emamectin benzoate; Saksida et al. 2010) when their abundance exceeds three motile lice per fish (Fisheries and Oceans Canada 2016). While emamectin benzoate also reduces *C. clemensi* numbers, *C. clemensi* abundance is not directly managed. Since *Caligus* spp. are considered less pathogenic than *L. salmonis* (Igboeli et al. 2014; Johnson et al. 2004) and most of the research into the effects of sea lice on wild salmon has focussed on *L. salmonis* (e.g., Connors et al. 2010; Johnson et al. 1996; Krkošek et al. 2005b), there has been little reason to target *C. clemensi* with treatments on farms until now. However, given the mounting evidence for sub-lethal effects of

C. clemensi on wild salmon, and the ability for *C. clemensi* to reach extreme abundances on farms without targeted treatment (e.g., 100% prevalence and 47.2 lice per fish (Di Cicco et al. 2017)), it seems prudent to start considering *C. clemensi* in the fish health management plans of farmed salmon in BC.

There is increasing concern over the potential impacts of *C. clemensi* on wild Fraser River sockeye salmon (e.g., Moore et al. 2017), especially in the context of the record-low Fraser sockeye returns in 2009 and 2016 (Pacific Salmon Commission 2016). Unlike *L. salmonis*, which can only infect salmonids, the generalist *C. clemensi* may have multiple sea lice reservoirs along the juvenile sockeye migration route, including Pacific herring, which may amplify extinction risk for salmon by allowing parasite abundances to remain high when an imperilled host population declines (De Castro and Bolker 2005) and perhaps threaten herring stocks as well. Our results shed further light on the sometimes subtle interactions between parasites and wild Pacific salmon (Miller et al. 2014; Peacock et al. 2014), and underscore the need to study not only the direct mortality from parasites but their sub-lethal effects as well. The impacts of parasite-mediated intraspecific competition on host survival and recruitment may be particularly influential for fishes, which often experience intense food competition and growth-dependent mortality, but there has been little to no work in this area. Competition is a fundamental driver of populations, and identifying how and when parasites mediate the competitive interactions of their hosts may be essential to understanding the host-parasite dynamics of many systems.

Acknowledgements

We are grateful to Salmon Coast Field Station and its staff for making this research possible. We thank M. Bartlett, L. Walker, and C. White for their diligent work in the laboratory and field. This work was supported by Simon Fraser University, the University of Toronto, Watershed Watch Salmon Society, V. and D. Bradshaw, a Natural Sciences and Engineering Research Council of Canada (NSERC) Industrial Postgraduate Scholarship (to S. Godwin), an NSERC Alexander Graham Bell Canada Graduate Scholarship (to S. Godwin), NSERC Discovery Grants (to L. Dill, M. Krkošek, and J. Reynolds), an Alfred P. Sloan Foundation Fellowship in Ocean Science and Canada Research Chair (to M. Krkošek), and the Tom Buell Endowment Fund supported by the Pacific Salmon Foundation and the BC Leading Edge Endowment Fund (to J. Reynolds). We performed all fish collection and euthanasia under permits and approval by Fisheries and Oceans Canada (license number XR472014) and the Simon Fraser University Animal Care Committee (protocol number 1088-B13).

References

- Anderson, J.T. 1988. A review of size dependent survival during pre-recruit stages of fishes in relation to recruitment. *Journal of Northwest Atlantic Fishery Science*, **8**: 55-66.
- Atkinson, E., Bateman, A.W., Dill, L.M., Krkošek, M., Reynolds, J.D., and Godwin, S.C. In review. Oust the louse: leaping behaviour removes sea lice from wild juvenile sockeye salmon.
- Barber, I., and Ruxton, G.D. 1998. Temporal prey distribution affects the competitive ability of parasitized sticklebacks. *Animal Behaviour*, **56**(6): 1477-1483.
- Beamish, R., Wade, J., Pennell, W., Gordon, E., Jones, S., Neville, C., Lange, K., and Sweeting, R. 2009. A large, natural infection of sea lice on juvenile Pacific salmon in the Gulf Islands area of British Columbia, Canada. *Aquaculture*, **297**(1): 31-37. doi: 10.1016/j.aquaculture.2009.09.001.
- Beamish, R., Neville, C., Sweeting, R., and Lange, K. 2012. The synchronous failure of juvenile Pacific salmon and herring production in the Strait of Georgia in 2007 and the poor return of sockeye salmon to the Fraser River in 2009. *Marine and Coastal Fisheries*, **4**(1): 403-414.
- Bergenius, M.A., Meekan, M.G., Robertson, R.D., and McCormick, M.I. 2002. Larval growth predicts the recruitment success of a coral reef fish. *Oecologia*, **131**(4): 521-525.
- Bremigan, M.T., and Stein, R.A. 1994. Gape-dependent larval foraging and zooplankton size: implications for fish recruitment across systems. *Canadian Journal of Fisheries and Aquatic Sciences*, **51**(4): 913-922.
- Brett, J. 1971. Satiation time, appetite, and maximum food intake of sockeye salmon (*Oncorhynchus nerka*). *Journal of the Fisheries Board of Canada*, **28**(3): 409-415.

- Burnham, K.P., and Anderson, D.R. 2002. Model selection and multimodel inference: a practical information-theoretic approach. Springer, New York.
- Connors, B.M., Krkošek, M., Ford, J., and Dill, L.M. 2010. Coho salmon productivity in relation to salmon lice from infected prey and salmon farms. *Journal of Applied Ecology*, **47**(6): 1372-1377. doi: 10.1111/j.1365-2664.2010.01889.x.
- Costello, M.J. 2009a. The global economic cost of sea lice to the salmonid farming industry. *Journal of Fish Diseases*, **32**(1): 115-118. doi: 10.1111/j.1365-2761.2008.01011.x.
- Costello, M.J. 2009b. How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proceedings of the Royal Society of London B: Biological Sciences*, **276**(1672): 3385-3394.
- Crombie, A.C. 1947. Interspecific competition. *The Journal of Animal Ecology*, **16**(1): 44-73. doi: 10.2307/1506.
- De Castro, F., and Bolker, B. 2005. Mechanisms of disease-induced extinction. *Ecology Letters*, **8**(1): 117-126. doi: 10.1111/j.1461-0248.2004.00693.x.
- Devries, D.R., Stein, R.A., and Bremigan, M.T. 1998. Prey selection by larval fishes as influenced by available zooplankton and gape limitation. *Transactions of the American Fisheries Society*, **127**(6): 1040-1050.
- Di Cicco, E., Ferguson, H.W., Schulze, A.D., Kaukinen, K.H., Li, S., Vanderstichel, R., Wessel, Ø., Rimstad, E., Gardner, I.A., and Hammell, K.L. 2017. Heart and skeletal muscle inflammation (HSMI) disease diagnosed on a British Columbia salmon farm through a longitudinal farm study. *PLoS One*, **12**(2): e0171471. doi: 10.1371/journal.pone.0171471.
- Duffy, E.J., and Beauchamp, D.A. 2011. Rapid growth in the early marine period improves the marine survival of Chinook salmon (*Oncorhynchus tshawytscha*) in Puget Sound,

- Washington. Canadian Journal of Fisheries and Aquatic Sciences, **68**(2): 232-240. doi:
10.1139/F10-144.
- Eggers, D.M. 1978. Limnetic feeding behavior of juvenile sockeye salmon in Lake Washington
and predator avoidance. Limnology and Oceanography, **23**(6): 1114-1125. doi:
10.4319/lo.1978.23.6.1114.
- Farley, E.V., Murphy, J.M., Adkison, M.D., Eisner, L.B., Helle, J.H., Moss, J.H., and Nielsen, J.
2007. Early marine growth in relation to marine-stage survival rates for Alaska sockeye
salmon (*Oncorhynchus nerka*). Fishery Bulletin, **105**(1): 121-130.
- Finley, R.J., and Forrester, G.E. 2003. Impact of ectoparasites on the demography of a small reef
fish. Marine Ecological Progress Series, **248**: 305-309.
- Fisheries and Oceans Canada. (2016). Marine Finfish Aquaculture Licence under the Fisheries
Act. Available from [http://www.pac.dfo-mpo.gc.ca/aquaculture/licence-permis/docs/licence-
cond-permis-mar/licence-cond-permis-mar-eng.pdf](http://www.pac.dfo-mpo.gc.ca/aquaculture/licence-permis/docs/licence-cond-permis-mar/licence-cond-permis-mar-eng.pdf). Accessed 2017-04-26.
- Foy, R.J., and Norcross, B.L. 1999. Spatial and temporal variability in the diet of juvenile Pacific
herring (*Clupea pallasii*) in Prince William Sound, Alaska. Canadian Journal of Zoology,
77(5): 697-706.
- Furey, N.B., Hinch, S.G., Bass, A.L., Middleton, C.T., Minke-Martin, V., and Lotto, A.G. 2016.
Predator swamping reduces predation risk during nocturnal migration of juvenile salmon in
a high-mortality landscape. Journal of Animal Ecology, **85**(4): 948-959. doi: 10.1111/1365-
2656.12528.
- Giles, N. 1987. Predation risk and reduced foraging activity in fish: experiments with parasitized
and non-parasitized three-spined sticklebacks, *Gasterosteus aculeatus* L. Journal of Fish
Biology, **31**(1): 37-44.

- Godwin, S.C., Dill, L.M., Reynolds, J.D., and Krkošek, M. 2015. Sea lice, sockeye salmon, and foraging competition: lousy fish are lousy competitors. *Canadian Journal of Fisheries and Aquatic Sciences*, **72**(7): 1113-1120. doi: 10.1139/cjfas-2014-0284.
- Godwin, S.C., Dill, L.M., Krkošek, M., Price, M.H.H., and Reynolds, J.D. 2017. Reduced growth in wild juvenile sockeye salmon infected with sea lice. *Journal of Fish Biology*, **91**(1): 41-57. doi: 10.1111/jfb.13325.
- Groner, M.L., Rogers, L.A., Bateman, A.W., Connors, B.M., Frazer, L.N., Godwin, S.C., Krkošek, M., Lewis, M.A., Peacock, S.J., and Rees, E.E. 2016. Lessons from sea louse and salmon epidemiology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**(1689): 20150203.
- Groot, C., and Cooke, K. 1987. Are the migrations of juvenile and adult Fraser River sockeye salmon (*Oncorhynchus nerka*) in near-shore waters related. In *Sockeye Salmon (O. nerka) Population Biology and Future Management*. Edited by H.D. Smith, L. Margolis and C.C. Wood. Canadian Special Publication of Fisheries and Aquatic Sciences. No 96. pp. 53-60.
- Groot, C., and Margolis, L. 1991. Pacific salmon life histories. UBC press, Vancouver, Canada.
- Grosholz, E.D. 1992. Interactions of intraspecific, interspecific, and apparent competition with host-pathogen population dynamics. *Ecology*, **73**(2): 507-514.
- Hamre, L.A., Eichner, C., Caipang, C.M.A., Dalvin, S.T., Bron, J.E., Nilsen, F., Boxshall, G., and Skern-Mauritzen, R. 2013. The salmon louse *Lepeophtheirus salmonis* (Copepoda: Caligidae) life cycle has only two chalimus stages. *PloS one*, **8**(9): e73539.
- Hargreaves, N.B., and LeBrasseur, R.J. 1986. Size selectivity of coho (*Oncorhynchus kisutch*) preying on juvenile chum salmon (*O. keta*). *Canadian Journal of Fisheries and Aquatic Sciences*, **43**(3): 581-586. doi: 10.1139/f86-069.

- 477 Hatcher, M.J., Dick, J.T., and Dunn, A.M. 2006. How parasites affect interactions between
478 competitors and predators. *Ecology Letters*, **9**(11): 1253-1271. doi: 10.1111/j.1461-
479 0248.2006.00964.x
- 480 Houde, E., and Hoyt, R. 1987. Fish early life dynamics and recruitment variability. *Transactions*
481 *of the American Fisheries Society*.
- 482 Hurvich, C.M., and Tsai, C.-L. 1989. Regression and time series model selection in small
483 samples. *Biometrika*, **76**(2): 297-307. doi: 10.1093/biomet/76.2.297.
- 484 Igboeli, O.O., Burka, J.F., and Fast, M.D. 2014. *Lepeophtheirus salmonis*: a persisting challenge
485 for salmon aquaculture. *Animal Frontiers*, **4**(1): 22-32. doi: 10.2527/af.2014-0004.
- 486 Jakob, E., Sweeten, T., Bennett, W., and Jones, S. 2013. Development of the salmon louse
487 *Lepeophtheirus salmonis* and its effects on juvenile sockeye salmon *Oncorhynchus nerka*.
488 *Diseases of Aquatic Organisms*, **106**(3): 217-227. doi: 10.3354/dao02642.
- 489 Johnson, S., Blaylock, R., Elphick, J., and Hyatt, K. 1996. Disease induced by the sea louse
490 (*Lepeophtheirus salmonis*)(Copepoda: Caligidae) in wild sockeye salmon (*Oncorhynchus*
491 *nerka*) stocks of Alberni Inlet, British Columbia. *Canadian Journal of Fisheries and Aquatic*
492 *Sciences*, **53**(12): 2888-2897. doi: 10.1139/f96-226.
- 493 Johnson, S.C., Bravo, S., Nagasawa, K., Kabata, Z., Hwang, J., Ho, J., and Shih, C. 2004. A
494 review of the impact of parasitic copepods on marine aquaculture. *Zoological Studies*, **43**(2):
495 229-243.
- 496 Jones, S., Kim, E., and Bennett, W. 2008. Early development of resistance to the salmon louse,
497 *Lepeophtheirus salmonis* (Krøyer), in juvenile pink salmon, *Oncorhynchus gorbuscha*
498 (Walbaum). *Journal of Fish Diseases*, **31**(8): 591-600.

- 499 Kabata, Z. 1972. Developmental stages of *Caligus clemensi* (Copepoda: Caligidae). Journal of
500 the Fisheries Research Board of Canada, **29**(11): 1571-1593. doi: 10.1139/f72-245.
- 501 Krkošek, M., Morton, A., and Volpe, J.P. 2005a. Nonlethal assessment of juvenile pink and
502 chum salmon for parasitic sea lice infections and fish health. Transactions of the American
503 Fisheries Society, **134**(3): 711-716. doi: 10.1577/t04-133.1.
- 504 Krkošek, M., Lewis, M.A., and Volpe, J.P. 2005b. Transmission dynamics of parasitic sea lice
505 from farm to wild salmon. Proceedings of the Royal Society of London B: Biological
506 Sciences, **272**(1564): 689-696. doi: 10.1098/rspb.2004.3027.
- 507 Krkošek, M., Gottesfeld, A., Proctor, B., Rolston, D., Carr-Harris, C., and Lewis, M.A. 2007.
508 Effects of host migration, diversity and aquaculture on sea lice threats to Pacific salmon
509 populations. Proceedings of the Royal Society of London B: Biological Sciences, **274**(1629):
510 3141-3149. doi: 10.1098/rspb.2007.1122.
- 511 Krkošek, M., Connors, B.M., Morton, A., Lewis, M.A., Dill, L.M., and Hilborn, R. 2011a.
512 Effects of parasites from salmon farms on productivity of wild salmon. Proceedings of the
513 National Academy of Sciences, **108**(35): 14700-14704.
- 514 Krkošek, M., Connors, B.M., Ford, H., Peacock, S., Mages, P., Ford, J.S., Morton, A., Volpe,
515 J.P., Hilborn, R., Dill, L.M., and Lewis, M.A. 2011b. Fish farms, parasites, and predators:
516 implications for salmon population dynamics. Ecological Applications, **21**(3): 897-914. doi:
517 10.1890/09-1861.1.
- 518 Landingham, J.H., Sturdevant, M.V., and Brodeur, R.D. 1998. Feeding habits of juvenile Pacific
519 salmon in marine waters of southeastern Alaska and northern British Columbia. Fishery
520 Bulletin, **96**(2): 285-302.

- 521 Lee, R.F. 1974. Lipids of zooplankton from Bute inlet, British Columbia. *Journal of the Fisheries*
522 *Board of Canada*, **31**(10): 1577-1582.
- 523 Mackas, D., Galbraith, M., Faust, D., Masson, D., Young, K., Shaw, W., Romaine, S., Trudel,
524 M., Dower, J., and Campbell, R. 2013. Zooplankton time series from the Strait of Georgia:
525 Results from year-round sampling at deep water locations, 1990–2010. *Progress in*
526 *Oceanography*, **115**: 129-159.
- 527 Mages, P.A., and Dill, L.M. 2010. The effect of sea lice (*Lepeophtheirus salmonis*) on juvenile
528 pink salmon (*Oncorhynchus gorbuscha*) swimming endurance. *Canadian Journal of*
529 *Fisheries and Aquatic Sciences*, **67**(12): 2045-2051.
- 530 Maksimowich, D.S., and Mathis, A. 2000. Parasitized salamanders are inferior competitors for
531 territories and food resources. *Ethology*, **106**(4): 319-329.
- 532 Masson, D., and Peña, A. 2009. Chlorophyll distribution in a temperate estuary: The Strait of
533 Georgia and Juan de Fuca Strait. *Estuarine, Coastal and Shelf Science*, **82**(1): 19-28.
- 534 McKinnell, S., Curchitser, E., Groot, K., Kaeriyama, M., and Trudel, M. 2014. Oceanic and
535 atmospheric extremes motivate a new hypothesis for variable marine survival of Fraser
536 River sockeye salmon. *Fisheries Oceanography*, **23**(4): 322-341. doi: 10.1111/fog.12063.
- 537 Milinski, M. 1984. Parasites determine a predator's optimal feeding strategy. *Behavioral Ecology*
538 *and Sociobiology*, **15**(1): 35-37.
- 539 Miller, K.M., Teffer, A., Tucker, S., Li, S., Schulze, A.D., Trudel, M., Juanes, F., Tabata, A.,
540 Kaukinen, K.H., and Ginther, N.G. 2014. Infectious disease, shifting climates, and
541 opportunistic predators: cumulative factors potentially impacting wild salmon declines.
542 *Evolutionary Applications*, **7**(7): 812-855.

- Moore, J.W., Mooers, A.O., and Gallagher, P. (2017). Speaking for the salmon. The protection of Pacific wild salmon: understanding potential risk factors and seeking solutions. Available from <http://www.sfu.ca/content/dam/sfu/pwrc/PDFs/SfSJJan2017Recommendations.pdf>. Accessed 2017-04-26.
- Morton, A., Routledge, R., and Krkošek, M. 2008. Sea louse infestation in wild juvenile salmon and Pacific herring associated with fish farms off the east-central coast of Vancouver Island, British Columbia. *North American Journal of Fisheries Management*, **28**(2): 523-532. doi: 10.1577/M07-042.1.
- Moss, J.H., Beauchamp, D.A., Cross, A.D., Myers, K.W., Farley, E.V., Murphy, J.M., and Helle, J.H. 2005. Evidence for size-selective mortality after the first summer of ocean growth by pink salmon. *Transactions of the American Fisheries Society*, **134**(5): 1313-1322. doi: 10.1577/t05-054.1.
- Mustafa, A., Rankaduwa, W., and Campbell, P. 2001. Estimating the cost of sea lice to salmon aquaculture in eastern Canada. *The Canadian Veterinary Journal*, **42**(1): 54.
- Nakagawa, S., and Schielzeth, H. 2013. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, **4**(2): 133-142. doi: 10.1111/j.2041-210x.2012.00261.x
- Nendick, L., Sackville, M., Tang, S., Brauner, C., and Farrell, A. 2011. Sea lice infection of juvenile pink salmon (*Oncorhynchus gorbuscha*): effects on swimming performance and postexercise ion balance. *Canadian Journal of Fisheries and Aquatic Sciences*, **68**(2): 241-249.

- Pacific Salmon Commission. (2016). Fraser river panel weekly report - August 19, 2016.
Available from <http://www.psc.org/download/71/2016/4572/weeklyreport06.pdf>. Accessed
2017-04-26.
- Parker, R.R. 1968. Marine mortality schedules of pink salmon of the Bella Coola River, central
British Columbia. Journal of the Fisheries Board of Canada, **25**(4): 757-794.
- Parsons, T., LeBrasseur, R., and Barraclough, W. 1970. Levels of production in the pelagic
environment of the Strait of Georgia, British Columbia: a review. Journal of the Fisheries
Board of Canada, **27**(7): 1251-1264.
- Peacock, S.J., Connors, B.M., Krkošek, M., Irvine, J.R., and Lewis, M.A. 2014. Can reduced
predation offset negative effects of sea louse parasites on chum salmon? Proceedings of the
Royal Society of London B: Biological Sciences, **281**(1776): 20132913.
- Peacock, S.J., Krkošek, M., Bateman, A.W., and Lewis, M.A. 2015. Parasitism and food web
dynamics of juvenile Pacific salmon. Ecosphere, **6**(12): 1-16.
- Price, M.H.H., Proboyszcz, S.L., Routledge, R.D., Gottesfeld, A.S., Orr, C., and Reynolds, J.D.
2011. Sea louse infection of juvenile sockeye salmon in relation to marine salmon farms on
Canada's west coast. PLoS ONE, **6**(2): e16851. doi: 10.1371/journal.pone.0016851.
- Price, M.H.H., Glickman, B.W., and Reynolds, J.D. 2013. Prey selectivity of Fraser River
sockeye salmon during early marine migration in British Columbia. Transactions of the
American Fisheries Society, **142**(4): 1126-1133. doi: 10.1080/00028487.2013.799517.
- Resetarits, W.J. 1995. Limiting similarity and the intensity of competitive effects on the mottled
sculpin, *Cottus bairdi*, in experimental stream communities. Oecologia, **104**(1): 31-38.
- Saksida, S.M., Morrison, D., and Revie, C.W. 2010. The efficacy of emamectin benzoate against
infestations of sea lice, *Lepeophtheirus salmonis*, on farmed Atlantic salmon, *Salmo salar*

L., in British Columbia. Journal of Fish Diseases, **33**(11): 913-917. doi: 10.1111/j.1365-2761.2010.01192.x.

Sapir, N., Butler, P.J., Hedenström, A., and Wikelski, M. 2011. Energy gain and use during animal migration. In Animal Migration. Edited by E. Milner-Gulland, J.M. Fryxell and A.R. Sinclair. Oxford University Press, New York. pp. 52-67.

Schall, J.J. 1992. Parasite-mediated competition in *Anolis* lizards. Oecologia, **92**(1): 58-64.

Scharf, F.S., Juanes, F., and Rountree, R.A. 2000. Predator size-prey size relationships of marine fish predators: interspecific variation and effects of ontogeny and body size on trophic-niche breadth. Marine Ecology Progress Series, **208**: 229-248.

Shi, Z.-h., Liu, S.-s., and Li, Y.-x. 2002. *Cotesia plutellae* parasitizing *Plutella xylostella*: host-age dependent parasitism and its effect on host development and food consumption. BioControl, **47**(5): 499-511.

Sutherland, G., Foreman, M., and Garrett, C. 2007. Tidal current energy assessment for Johnstone Strait, Vancouver island. Proceedings of the Institution of Mechanical Engineers, Part A: Journal of Power and Energy, **221**(2): 147-157.

Sutherland, W.J. 1996. From individual behaviour to population ecology. Oxford University Press, New York.

Tanasichuk, R., and Routledge, R. 2011. An investigation of the biological basis of return variability for sockeye salmon (*Oncorhynchus nerka*) from Great Central and Sproat lakes, Vancouver Island. Fisheries Oceanography, **20**(6): 462-478.

Ward, A.J., Webster, M.M., and Hart, P.J. 2006. Intraspecific food competition in fishes. Fish and Fisheries, **7**(4): 231-261.

- 609 Welch, D.W., Melnychuk, M.C., Payne, J.C., Rechisky, E.L., Porter, A.D., Jackson, G.D., Ward,
610 B.R., Vincent, S.P., Wood, C.C., and Semmens, J. 2011. *In situ* measurement of coastal
611 ocean movements and survival of juvenile Pacific salmon. Proceedings of the National
612 Academy of Sciences, **108**(21): 8708-8713.
- 613 Welker, M.T., PiERCE, C.L., and Wahl, D.H. 1994. Growth and survival of larval fishes: roles
614 of competition and zooplankton abundance. Transactions of the American Fisheries Society,
615 **123**(5): 703-717.
- 616 Wootten, R., Smith, J.W., and Needham, E. 1982. Aspects of the biology of the parasitic
617 copepods *Lepeophtheirus salmonis* and *Caligus elongatus* on farmed salmonids, and their
618 treatment. Proceedings of the Royal Society of Edinburgh. Section B: Biological Sciences,
619 **81**(03): 185-197. doi: 10.1017/s0269727000003389.
- 620

Tables

Table 1. Infection categories and sample sizes for each fish collection. The infection scale was weighted such that one large chalimus louse was equal to one infection unit, one motile louse was equal to two infection units, and copepodite and small chalimus lice were equal to zero units. For each collection, a single fish pair was comprised of one fish from the highly infected category and one from the lightly infected category.

Collection	Fish pairs	Lightly infected		Highly infected	
		Infection scale	Max. lice	Infection scale	Min. lice
1	5	0	3	4	6
2	6	0	1	3	2
3	7	0	1	4	5
4	7	0	2	4	4
5	6	0	3	4	4
6	7	0	3	4	7
7	6	0	3	4	6
8	5	0	3	5	7
9	5	0	3	5	4
10	6	0	3	5	9
11	5	0	3	5	10
Average	5.9	0.0	2.5	4.3	5.8

Table 2. Model selection results for the six models of relative foraging success that accounted for at least 5% of model support. Relative stomach fullness was calculated as wet stomach content weight divided by body weight. The models included combinations of infection category (infection), body size (size; see description of principal component analysis), and prey density fixed effects. Each model included a random effect on the intercept for collection number. Interaction terms are distinguished with an asterisk symbol. See Supplementary Table S3 for the full model set and selection results.

Rank	Model	$\Delta AICc^*$	$w_i^\dagger$	$R^{2\ddagger}$
1	infection	0	0.358	0.549
2	infection + size	1.24	0.192	0.530
3	infection + prey	1.66	0.156	0.542
4	infection + size + prey	2.66	0.095	0.510
5	infection*size	3.85	0.052	0.530
6	infection*prey	3.91	0.051	0.536

* Difference from the top model AICc ($\Delta AICc$)

† Akaike model weight (w_i)

‡ R^2 for mixed-effects models calculated using the method developed by Nakagawa and Schielzeth (2013)

Figures

Fig. 1. Map of study region. The solid black box indicates the area in which salmon collections occurred.

Fig. 2. Relative stomach fullness of juvenile sockeye salmon for the two categories of sea louse infection. Error bars indicate the 95% confidence intervals around the relative stomach fullness for each infection category.

Fig. 3. Mean relative stomach fullness ($\pm$ SE) for fish in both infection categories of each collection. Each vertical pair of points (one grey, one white) comprises a single collection. The relationship between relative stomach fullness and prey density had equivocal support in our model selection results (see Table S3).

Fig. 4. Relationships among juvenile sockeye salmon traits and sea louse infection. Solid double-ended arrows indicate established correlative sea louse relationships ([a] Godwin et al. 2015, [b] this paper, [c] Godwin et al. 2017). Dashed black arrows indicate established correlative body size relationships, and grey arrows indicate implicit mechanistic relationships.

Appendix. Juvenile sockeye salmon digestion experiment

Methods

To assess the extent to which our prey density estimates were representative of the prey field met by the juvenile sockeye salmon (*O. nerka*) when they had been feeding, we needed to determine how quickly juvenile sockeye digested their prey. To accomplish this, we performed a small feeding experiment at a floating field facility comprised of several floating docks and net pens. We collected fish at the same Johnstone Strait location used in the main study and transported them by boat for 1 h to the experimental facility (see Fig. 1 in Godwin et al. (2015) for map). During transport, the sockeye were again held in insulated fish totes with bubblers and ice packs. We did not collect temperature or salinity data for this experiment, but previous studies using the same experimental facility and juvenile salmon collection sites have indicated that their water temperatures and salinities are very similar (Atkinson et al. In review; Godwin et al. 2015).

Upon arrival at the facility, we transferred the fish to a large (2.8 m deep and 6.1 x 6.1 m across) net pen and weaned them onto frozen brine shrimp (Brine Shrimp Direct, Ogden, Utah, USA) over the next five days. Brine shrimp were thawed in freshwater and were fed to the fish by adding them to the center of the net pen. Medium-sized fish (between 107 and 120 mm) were removed for another study, leaving 31 smaller ($104.1 \text{ mm} \pm 0.4 \text{ mm}$) and 37 larger ($124.9 \pm 0.7 \text{ mm}$) fish. The fork length range of these experimental fish was 97 mm to 132 mm, which was fully within the fork length range of the fish from the main study (88 mm to 133 mm).

One hour after sunrise on their sixth day at the experiment facility, the 68 fish were fed to satiation and ten were immediately sacrificed with a lethal dose of MS-222. At half, one, two,

three, five hours after initial feeding, ten fish were again randomly removed and euthanized, leaving remaining fish. At eight hours, these eight fish were removed and euthanized. We weighed the wet stomach contents of the fish in the same manner as for those sacrificed in the field, and calculated relative stomach fullness by dividing the weight of each fish by the weight of its wet stomach contents.

We fit a Michaelis-Menten curve to the relative stomach fullness data and used these parameter estimates to calculate the amount of digestion that occurred between time points in the experiment.

Results

The digestion rate of the experimental fish began high and gradually slowed over time (Fig. A1). Six fish did not consume any brine shrimp and were therefore removed from the dataset. After one hour, 37% of the initial stomach contents remained, and after the final hour of the experiment (hour eight) only 18% remained. The predicted relative stomach fullness of the experimental fish at hour eight (0.6%) was similar to the mean relative stomach fullness of the main study's fish (0.6%), suggesting that most of the main study's fish must have eaten in the previous eight hours since not all of them would have fed to satiation like the experimental fish.

Fig. A1. Relative stomach fullness of juvenile sockeye salmon over the course of the digestion rate experiment, with a Michaelis-Menten curve fit to the data (black line). Six fish were removed from this dataset because they did not consume any food.

References

- Atkinson, E., Bateman, A.W., Dill, L.M., Krkošek, M., Reynolds, J.D., and Godwin, S.C. In review. Oust the louse: leaping behaviour removes sea lice from wild juvenile sockeye salmon.
- Godwin, S.C., Dill, L.M., Reynolds, J.D., and Krkošek, M. 2015. Sea lice, sockeye salmon, and foraging competition: lousy fish are lousy competitors. *Canadian Journal of Fisheries and Aquatic Sciences*, **72**(7): 1113-1120. doi: 10.1139/cjfas-2014-0284.