Canadian Journal of Fisheries and Aquatic Sciences
 Journal canadien des sciences halieutiques et aquatiques

**Temperature regimes, growth, and food consumption for
 female and male adult walleye in Lake Huron and Lake Erie:
 a bioenergetics analysis**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0280.R1
Manuscript Type:	Article
Date Submitted by the Author:	17-Nov-2017
Complete List of Authors:	Madenjian, Charles; U.S. Geological Survey, Hayden, Todd; US Geological Survey Peat, Tyler; Carleton University Department of Biology, Department of Biology Vandergoot, Christopher; US Geological Survey Fielder, David; Michiganm DNR, Fisheries Gorman, Ann Marie; Fairport Fish Research Station, Ohio Department of Natural Resources Pothoven, Steven; Great Lakes Environmental Research Lab - NOAA Dettmers, John; Great Lakes Fishery Commission Cooke, Steven; Carleton University Zhao, Yingming; Ontario Ministry of Natural Resources Krueger, Charles; Michigan State University, Fisheries and Wildlife
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	bioenergetics modeling, food availability, interbasin variability, interlake variability, temperature regime

SCHOLARONE™
 Manuscripts

**Temperature regimes, growth, and food consumption for female and male
adult walleye in Lake Huron and Lake Erie: a bioenergetics analysis**

**Charles P. Madenjian, Todd A. Hayden, Tyler B. Peat, Christopher S. Vandergoot,
David G. Fielder, Ann Marie Gorman, Steven A. Pothoven, John M. Dettmers,
Steven J. Cooke, Yingming Zhao, and Charles C. Krueger**

C.P. Madenjian. US Geological Survey, Great Lakes Science Center, 1451 Green Road, Ann Arbor, MI 48105, USA.
T.A. Hayden. Michigan State University, Department of Fisheries and Wildlife, Hammond Bay Biological Station, 11188 Ray Road Millersburg, MI 49759, USA.
T.B. Peat and S.J. Cooke. Fish Ecology and Conservation Physiology Laboratory, Department of Biology, Carleton University, Ottawa, ON K1S 5B6, Canada.
C.S. Vandergoot. US Geological Survey, Great Lakes Science Center, Lake Erie Biological Station, Sandusky, OH 44870, USA.
D.G. Fielder. Michigan Department of Natural Resources, Lake Huron Research Station, 160 Fletcher Street, Alpena, MI 49707, USA.
A. Gorman. Ohio Department of Natural Resources, Fairport Harbor Fisheries Research Station, 1190 High Street, Fairport Harbor, OH 44077, USA.
S.A. Pothoven. National Oceanic and Atmospheric Administration, Great Lakes Environmental Research Laboratory, Lake Michigan Field Station, 1431 Beach Street, Muskegon, MI 49441, USA.
J.M. Dettmers. Great Lakes Fishery Commission, 2100 Commonwealth Boulevard, Suite 100, Ann Arbor, MI 48105, USA.
Y. Zhao. Ontario Ministry of Natural Resources and Forestry, Aquatic Research and Monitoring Section, Lake Erie Fisheries Station, 320 Milo Road, Wheatley, ON N0P 2P0, Canada.
C.C. Krueger. Michigan State University, Center for Systems Integration and Sustainability, 1405 South Harrison Road, East Lansing, MI 48823, USA.

Corresponding author: Charles P. Madenjian (e-mail: cmadenjian@usgs.gov).

Abstract. Bioenergetics modeling was used to assess the relative importance of food availability and water temperature in determining walleye (*Sander vitreus*) growth. Temperature regimes experienced by both female and male adult walleye in three basins of Lake Huron and in Lake Erie were determined by use of surgically implanted temperature loggers and acoustic telemetry. Temperatures experienced by walleye were higher in Lake Erie than in Lake Huron. Walleye from Lake Erie grew at nearly double the rate of walleye from Lake Huron, and weight at age for adult females averaged about 50% greater than that for adult males in both lakes. Food consumption rate for an average adult walleye in Lake Erie was nearly twice as high as that in Lake Huron. Interbasin and interlake variability in temperature regimes accounted for a moderate degree of variability in walleye growth. We concluded that the driver for faster growth in Lake Erie compared with Lake Huron was higher food availability in Lake Erie compared with Lake Huron. The sex difference in temperature regimes explained 15% of the sex difference in Lake Erie walleye growth.

Keywords: acoustic telemetry; bioenergetics modeling; food availability; interbasin variability; interlake variability; sex differences; temperature loggers; temperature regime.

Introduction

For a given species of fish, individual-level growth of fish is a function of food availability and the temperature regime experienced by the fish (Hewett and Johnson 1992; Jobling 1994; Quinn and Deriso 1999; Enberg et al. 2008). Food availability is a complex function of several factors, including prey abundance, prey size, and handling times (Werner 1974; Persson and Greenberg 1990; Rennie and Venturelli 2015). Typically, metabolic costs for the fish increase with increasing water temperature, and metabolic costs represent a substantial energy loss for fish (Hewett and Johnson 1992; Jobling 1994; Quinn and Deriso 1999; Enberg et al. 2008). Thus, if the rate of food consumption is identical in two aquatic ecosystems, and other factors affecting growth other than water temperature are also identical across the two ecosystems, then fish growth would be expected to be faster in the aquatic ecosystem with the lower temperatures experienced by fish compared with fish growth in the other aquatic ecosystem, owing to higher metabolic costs in the warmer ecosystem (Hewett and Johnson 1992; Kao et al. 2015a, 2015b).

Bioenergetics modeling has proven invaluable in assessing the effects of various factors, including temperature regime, on fish growth (Chipps and Wahl 2008; Madenjian 2011; Deslauriers et al. 2017). A fish bioenergetics model represents an energy budget for a fish. Energy input via food consumption is equal to metabolic losses, waste losses, and growth:

$$C = (R + A + SDA) + (F + U) + G$$

where C = consumption, R = standard metabolism, A = energy expenditure due to activity, SDA = specific dynamic action (energy required to digest food), F = egestion, U = excretion, and G = growth. Standard metabolism is typically modeled as an exponential function of water temperature. Bioenergetics model applications to evaluate the effects of various factors on fish growth are well illustrated in the study by Hayward and Margraf (1987), who used bioenergetics

71 modeling to show that faster growth of yellow perch (*Perca flavescens*) in the central basin of
72 Lake Erie compared with the western basin of Lake Erie was due to an interbasin difference in
73 food availability rather than an interbasin difference in temperature regime.

74 Coupling archival temperature tag technology with acoustic telemetry has recently enabled
75 researchers to record thermal histories of fish within different basins of the same lake, as well as
76 within lakes connected by waterways (Cooke et al. 2013; Hayden et al. 2014; Peat et al. 2015).
77 By constructing an appropriate acoustic receiver network, interbasin and/or interlake movement
78 of fish can be detected. Implanting both an acoustic transmitter and an archival temperature tag
79 in each fish, and subsequent monitoring of any interbasin and interlake movements of the fish
80 using acoustic telemetry, can be used by researchers to determine temperature regimes
81 experienced by fish while confined to a specific basin or lake.

82 To date, the effects of both interbasin and interlake variability in temperatures experienced by
83 fish on fish growth in a two-lake system, with a connecting waterway between the two lakes,
84 have not been investigated. Moreover, the effect of a difference in temperature regimes between
85 the sexes of a fish population on the difference in growth between the sexes has yet to be
86 assessed for any fish population. Use of archival temperature tagging and acoustic telemetry, in
87 conjunction with fish bioenergetics modeling, affords the opportunity to pursue these new lines
88 of research. In turn, these new lines of research may have fishery management implications,
89 because knowledge of movement and spatial ecology of fishes provides critical information to
90 managers about how fishes are distributed in both space and time (Lucas and Baras 2000;
91 Landsman et al. 2011). For example, bioenergetics analysis of growth of a piscivorous fish
92 within a two-lake system may be useful in accurately characterizing the fish's role as a piscivore
93 in both lakes.

Walleye (*Sander vitreus*) populations support valuable recreational and/or commercial fisheries in many North American lakes, reservoirs, and rivers (Feiner and Höök 2015). Moreover, this species fulfills the ecological role as a top predator in the aquatic ecosystems that it inhabits (Feiner and Höök 2015; He et al. 2015). Female walleye grow substantially faster than male walleye, but reasons for this sexually dimorphic growth are not clear (Bozek et al. 2011; Rennie and Venturelli 2015). In the Laurentian Great Lakes, the most valuable walleye fisheries are situated in Lake Huron and Lake Erie (Roseman et al. 2008; Melstrom and Lupi 2013; Fielder et al. 2014). Despite the importance of the Lake Huron and Lake Erie walleye fisheries, a direct and rigorous comparison of walleye growth in Lake Huron with walleye growth in Lake Erie has yet to be undertaken.

Walleye in Lake Huron exhibit three migration behaviors, as elucidated from an acoustic telemetry study (Hayden et al. 2014). Walleye may reside in Saginaw Bay throughout the year or out-migrate to the main basin of Lake Huron for residence during the summer and fall. Those walleye out-migrating to the main basin either reside in the northern main basin or the southern main basin (Hayden et al. 2014). Walleye remaining in Saginaw Bay are expected to experience warmer temperatures than those walleye out-migrating to the main basin, while food availability for walleye is presumably higher in Saginaw Bay than in the main basin (Fielder et al. 2014; Hayden et al. 2014; Peat et al. 2015).

The primary goals of our study were to compare walleye growth in Lake Huron with walleye growth in Lake Erie and to assess the relative importance of food availability and temperature regime as determinants of walleye growth in the Lake Huron – Lake Erie system. A secondary goal of our study was to evaluate the effect of sex differences in temperature regimes on sex differences in walleye growth. Specific objectives included: (i) use temperature loggers and

acoustic telemetry to determine temperatures experienced by female and male walleye in Saginaw Bay, the northern main basin of Lake Huron, the southern main basin of Lake Huron, and Lake Erie, (ii) develop growth trajectories for female and male walleye in both Lake Huron and Lake Erie, and then compare growth between the two lakes, (iii) use bioenergetics modeling to estimate food consumption by female and male walleye in Saginaw Bay, northern main basin of Lake Huron, southern main basin of Lake Huron, and Lake Erie, (iv) using a bioenergetics modeling framework, exchange temperature regimes experienced by walleye between basins and lakes to assess the effects of interbasin and interlake variability in temperature regimes experienced by walleye on walleye growth, and (v) again using a bioenergetics modeling framework, exchange temperature regimes experienced by walleye between the sexes to evaluate the effect of sex differences in temperature regimes on sex differences in walleye growth.

Materials and methods

Field methods

Walleye in this study were captured and tagged as part of a large project to understand movement of two adfluvial walleye populations in the Great Lakes. A detailed description of walleye capture and tagging is found elsewhere (Hayden et al. 2014; Peat et al. 2015). Walleye in spawning condition were captured using boat-mounted electroshocking gear in the Tittabawassee River (Lake Huron watershed) immediately downstream of Dow Dam (Midland, MI) and near Orleans Park in the Maumee River (tributary to Lake Erie) during March-April in 2011 and 2012. The primary spawning area in the Tittabawassee River is located approximately 2-3 km downstream of the Dow Dam and approximately 100 km upstream of the mouth of the Saginaw River in Saginaw Bay. Primary walleye spawning grounds in the Maumee River are

located approximately 25-30 km upstream of the Maumee River mouth in Lake Erie. After capture, biological measurements (total length in millimeters, sex) and dorsal fin spine clips were collected from each walleye chosen for transmitter implanting and paired t-bar tags (Floy Tag, Inc., Seattle, WA) were inserted below the base of the second dorsal fin. T-bar tags allowed identification of walleye during the implanting process and informed anglers of the presence of an internal acoustic transmitter in the event the fish was caught after release. Dorsal fin spine clips were used for non-lethal age (in years) estimation. Fish were then transferred to streamside holding tanks for surgical implantation of an acoustic transmitter and a temperature logger. Walleye selected for implanting transmitters in the Maumee and Tittabawassee rivers were greater than 381 mm, which represented the minimum length limit for recreational angling limit for Michigan waters of Lake Huron and Ohio waters of Lake Erie during 2011-2012. In total, 199 walleye were implanted with transmitters in the Tittabawassee River and 200 walleye were implanted with transmitters in the Maumee River in 2011. In 2012, 60 walleye were implanted with transmitters in the Tittabawassee River and 33 walleye were implanted with transmitters in the Maumee River. Our research project was reviewed and approved by the Carleton University Animal Care Committee in accordance with the Canadian Council of Animal Care Guidelines for the Use of Fish in Research (Project number 8890; Approval number B10-29).

Each walleye was implanted with an acoustic transmitter programmed to emit a unique code (frequency of 69 kHz) at random intervals between 60 seconds and 180 seconds (mean: 120 seconds) (Vemco, Halifax, Nova Scotia; model V16-4x; 16 mm dia X 86 mm, 24 g; battery life = 1338 days). Each acoustic transmitter was outfitted with an iButton temperature logger (DS1921z; resolution = ± 0.1 °C, precision = 0.2 ± 0.3 °C, accuracy = 0.4 ± 0.3 °C, Maxim Integrated Products, Inc., Sunnyvale, CA) programmed to record temperatures at four-hour

intervals for a period of approximately one year (Donaldson et al. 2009). All iButtons were attached to acoustic transmitters using hot glue, and dipped in an inert waterproof coating (Plasti Dip, Performix Brand, Blaine, MN) prior to implantation.

Walleye were anesthetized prior to surgery using a Smith-Root (Vancouver, WA) portable electroanesthesia system operating at 35 V pulsed direct current. Treatments of 3 seconds induced stage-4 anesthesia for several minutes (Vandergoot et al. 2011). All surgeries were conducted on fish supported in a v-shaped surgical table lined with soft non-slip material. Walleye received a constant supply of fresh river water pumped across gills during surgery and all surgical tools and transmitters were cleaned with povidone iodine prior to surgery. Acoustic transmitters with attached iButtons were inserted into the coelomic cavity through a small incision located along the ventral side of each fish, posterior to the pelvic girdle. Incisions were closed using 2–3 interrupted sutures (absorbable monofilament, PDS-II, 3-0, Ethicon, Somerville, NH). After surgery, fish were placed in holding tanks until fish were able to maintain balance in the water (approximately 30 minutes), and then fish were released near the capture site.

The acoustic receiver array was deployed as part of the Great Lakes Acoustic Telemetry Observation System network (<http://data.glos.us/glatos>) and consisted of approximately 140 omnidirectional receivers operating at 69 kHz (VR2W, Vemco, Halifax, Nova Scotia). Hayden et al. (2014) and Peat et al. (2015) provide complete details of anchoring acoustic receivers and receiver line detection probabilities. All receivers were suspended in the water column and retrieved annually for maintenance and data download. Deployment locations of acoustic receivers during 2011–2012 in Lake Huron consisted of multiple receivers near the Tittabawassee and Saginaw river mouths, double receiver lines located across Saginaw Bay, and

multiple-receiver lines extending perpendicular to the shoreline to depths < 40 m (extending 3-10 km offshore) (Fig. 1). Receiver lines were deployed to monitor walleye movements in and out of the Saginaw Bay, movements in and out of the Tittabawassee and Saginaw rivers, and nearshore movements in Lake Huron. Acoustic receivers were deployed in the St. Clair River flowing out of Lake Huron, and Detroit River to monitor walleye movement between Lake Huron and Lake Erie through the Huron-Erie corridor (Fig. 1). Acoustic receivers were deployed near the mouth of the Maumee River in Lake Erie to monitor walleye movements in and out of the river. Thermal data were obtained via returned transmitter/iButtons from fish caught in recreational and commercial fisheries. A \$100 USD reward was offered to anyone that returned the acoustic transmitter with the attached iButton to the investigators. All acoustic detection data were screened for false positives caused by environmental noise and signal collisions using the short-interval criteria described by Pincock (2009). False detections were detections that were not accompanied by at least one other detection from the same transmitter on the same receiver within 1 hour and were removed from the analysis. Of all of the detections, 99% of them passed the false detection filter.

We used acoustic telemetry detections to classify the walleye recaptured by anglers and commercial fishers into four geographic areas: (1) northern main basin of Lake Huron, (2) southern main basin of Lake Huron, (3) Saginaw Bay, and (4) Lake Erie (Fig. 1). Please refer to Hayden et al. (2014) and Peat et al. (2015) for more details on the classification assignments.

Temperature regimes

We followed the procedure described by Bergstedt et al. (2003), Madenjian et al. (2003, 2006), and He et al. (2015) to develop temperature regimes for both female and male walleye

from each of the above-mentioned four geographic areas. The calendar year was divided into half-month intervals. For each combination of geographic area and sex, temperatures recorded on the iButton temperature loggers were averaged within each half-month interval, and the standard error of the mean was also calculated. The mean temperature for each half-month interval was assigned to the midpoint day, in time, for the half-month interval. Temperatures for days between consecutive midpoints were estimated via linear interpolation. Thus, each day of the calendar year was assigned a temperature.

In 15 cases, the calculated mean temperature for the half-month interval was less than 0.10 °C. To successfully perform the bioenergetics model simulations, all calculated mean temperatures less than 0.10 °C were assigned the value of 0.10 °C. Waste loss components of the bioenergetics model yielded inaccurate estimates of waste losses for temperatures less than 0.10 °C (Kao et al. 2015a, 2015b), and therefore these assignments were necessary to successfully complete the bioenergetics modeling.

We were well justified in developing temperature regimes by sex and geographic area, for four reasons. First, two of the objectives of our study were to assess the effects of interbasin and interlake variability in temperature regimes on walleye growth and to assess the effect of the sex difference in temperature regimes on the sex difference in walleye growth. Development of temperature regimes by sex and geographic area was necessary to accomplish these two objectives. Second, previous research has shown that, on average, (i) surface water temperatures in Saginaw Bay are higher than those in the main basin of Lake Huron, (ii) surface water temperatures in the southern section of Lake Huron's main basin are higher than those in the northern main basin of Lake Huron, and (iii) surface water temperatures in Lake Erie are higher than those in any of the basins of Lake Huron (Moukomla and Blanken 2016). Third,

habitat use may vary with sex both in the Saginaw Bay walleye population and in the Lake Erie walleye population (Madenjian et al. 1998; Wang et al. 2007). Fourth, female walleye grow considerably faster than male walleye (Bozek et al. 2011; Rennie and Venturelli 2015).

Growth trajectories

We estimated growth trajectories for both female and male walleye from Lake Huron based on data from annual gillnet surveys conducted by the Michigan Department of Natural Resources (DNR) each September in Saginaw Bay (Fielder and Thomas 2014). Total length (TL, to the nearest millimeter), weight (to the nearest gram), and age (in years) of each walleye caught in the gill nets were determined. We pooled survey data from years 2010-2012, which closely corresponded with the years of our acoustic telemetry study, and then fitted von Bertalanffy growth curves to the TL at age data for both sexes. In addition, a weight-length regression curve was fitted to TL and weight data for each sex. Total lengths at age for ages 3-13 years were calculated from the fitted von Bertalanffy growth curves for both sexes. Then, these TLs at age were substituted into the fitted weight-length regression equations to estimate weights at age. We focused on ages 3-13 years because this range of ages is used in the population models developed for the Lake Huron walleye population (Fielder and Bence 2014; Fielder and Thomas 2014; He et al. 2015). Walleye growth trajectories were assumed not to vary across the basins (i.e., northern main basin, southern main basin, and Saginaw Bay) of Lake Huron. A similar assumption was made by He et al. (2015) in quantifying piscivory patterns in Lake Huron. Standard errors were calculated for weights at age by first calculating standard errors for TLs at age, using routines described by Ogle (2016), and then substituting the error bounds for TLs at age into the fitted weight-length equations.

Growth trajectories for Lake Erie were based on data from annual gillnet surveys conducted by the Michigan DNR, Ohio DNR, and Ontario Ministry of Natural Resources and Forestry (OMNRF) each October in the western and central basins of Lake Erie (Ohio Division of Wildlife 2011, 2012, 2013). Again, TL (to the nearest millimeter), weight (to the nearest gram), and age (in years) of each walleye caught in the gill nets were determined, and survey data from years 2010-2012 were pooled. Von Bertalanffy growth curves were fitted to TLs at age for both sexes, and weight-length regression curves were fitted to TL and weight data for both sexes. Weights at age were estimated by substituting TLs at age into the fitted weight-length regression equations. Standard errors were calculated for weights at age, using the same procedures described above.

To determine whether walleye growth significantly differed between the two lakes and between the sexes, we applied t tests for paired comparisons to the above-mentioned estimated weights at ages 3-13. Specifically, t tests for paired comparisons were used to: (i) compare weight at age of Lake Huron female walleye with weight at age of Lake Erie female walleye, (ii) compare weight at age of Lake Huron male walleye with weight at age of Lake Erie male walleye, (iii) compare weight at age of Lake Huron female walleye with weight at age of Lake Huron male walleye, and (iv) compare weight at age of Lake Erie female walleye with weight at age of Lake Erie male walleye.

Bioenergetics modeling: fit to observed growth

The Wisconsin walleye bioenergetics model, originally developed by Kitchell et al. (1977) and refined by Hewett and Johnson (1992), was used to estimate food consumption by both female and male walleye in each of the four geographic areas. Bioenergetics model simulations

were performed using the Fish Bioenergetics 4.0 software developed by Deslauriers et al. (2017). The model was fit to observed growth, as represented by the growth trajectory, to estimate food consumption. For each combination of geographic area and sex, the appropriate temperature regime, as derived from our iButton temperature logger data, was used in the bioenergetics model simulations. Each bioenergetics model simulation was one year in duration, and thus 10 simulations were run in order to estimate food consumption by an average walleye from age 3 through age 13 for each combination of geographic area and sex. The bioenergetics model operated on a daily time step (Hewett and Johnson 1992).

With regard to bioenergetics modeling for Lake Huron walleye, inputs for diet composition, prey energy density, and walleye energy density were taken from Pothoven et al. (2017). Because 8 September represented the typical midpoint date for the duration of the Michigan DNR gillnet survey in Saginaw Bay, we assigned 8 September as the starting day for the bioenergetics model simulations. Spawning day was assigned 4 April for Lake Huron walleye (Madenjian et al. 1998). Spawning was simulated by females and males losing 22.1% and 2.9%, respectively, of their body weight on the spawning day (Madenjian et al. 1998). Females and males were assumed to mature by ages 5 and 4 years, respectively (Wang et al. 2009; Fielder 2014; Fielder and Thomas 2014).

With regard to bioenergetics modeling for Lake Erie walleye, diet composition was based on stomach contents data for age-2 and older walleye caught by the Ohio DNR during monthly bottom trawl surveys from May through October and fall gillnet surveys (late September through early November) during 2000-2015 (Ohio Division of Wildlife 2016). Bottom trawls were conducted from sunrise to sunset and gill nets were set overnight on the bottom or suspended approximately 1.5-3 m from the surface. All fish were held on ice in coolers < 12 hours until

being processed. In the laboratory, total length of each walleye was measured to the nearest millimeter, and wet weight was recorded to the nearest gram. For stomach contents analysis, prey items were examined macroscopically, identified to the lowest taxonomic level possible, and measured to the nearest millimeter. Whole prey items were measured for total length, while partially digested items were measured for standard, fork, or backbone length. All prey lengths were converted to total length when necessary, and subsequently converted to wet weight using available data (Ohio Division of Wildlife 2016). Prey items were grouped into the following eight diet categories: shiners (primarily emerald shiner (*Notropis atherinoides*)), clupeids (primarily gizzard shad (*Dorosoma cepedianum*)), *Morone* spp. (primarily white perch (*Morone americana*)), rainbow smelt (*Osmerus mordax*), yellow perch, round goby (*Neogobius melanostomus*), miscellaneous fish (primarily rainbow trout (*Oncorhynchus mykiss*)), and invertebrates (burrowing mayflies, other ephemeropterans, amphipods, *Bythotrephes longimanus*, chironomids, other dipterans, isopods, and trichopterans). Wet weights of individual prey were used to calculate percent diet composition of each prey category for each walleye, and mean percent diet composition was calculated across individuals for each month (May through October) (Table 1). Percent composition of unidentified fish was distributed proportionally across the identified finfish species each month. Diet composition for April and November was assumed to be equal to that for May and October, respectively. Diet composition for December through March was assumed to be equal to the average diet composition for months May and October. A total of 7860 walleye stomachs were examined. Of these, 3449 contained prey items and 4411 were empty.

Energy densities for gizzard shad, *Morone* spp., yellow perch, and round goby from Lake Erie were taken from Kershner (1996) and Bunnell et al. (2005) (Table 2). Energy densities for

miscellaneous fish and invertebrates from Lake Erie were based on previous studies (Cummins and Wuycheck 1971; Pothoven et al. 2017). To determine energy densities of Lake Erie emerald shiner and rainbow smelt, we used bottom trawling to sample emerald shiner and rainbow smelt from Lake Erie during May-October 2005-2007. In total, three composite samples of emerald shiners and five composite samples of rainbow smelt were homogenized in a blender. Number of emerald shiners and rainbow smelt per composite ranged from 12 to 25 fish and from 10 to 23 fish, respectively. Energy density, in J/g wet weight, was determined for each composite using a Parr 1261 isoperibol bomb calorimeter (Parr Instrument Company, Moline, IL), following the procedure outlined by Pothoven et al. (2012). For both emerald shiners and rainbow smelt, energy densities were averaged across the composite samples to determine mean energy density (Table 2). Energy densities for Lake Erie walleye were determined by capturing 41 walleye in Lake Erie using gill nets during May 2005 (Pothoven et al. 2008). Upon capture, each walleye was individually wrapped in foil, bagged, and frozen until processing. In the laboratory, frozen fish were homogenized using a commercial blender. To determine dry-to-wet mass, we dried a 20-30 g subsample of ground fish tissue at 70°C to a constant mass (about 3 days). For energy density (J/g wet weight), the dried material was ground with a mortar and pestle, and then a 1-g subsample was combusted in a Parr 1261 isoperibol bomb calorimeter that was standardized with benzoic acid. To estimate within-fish variability for bomb calorimetry, we initially quantified energy density in triplicate. The mean coefficient of variation for individual fish was 1.02% (n = 10), thus, we subsequently quantified energy density using only one subsample per fish following the procedure outlined by Pothoven et al. (2017). Walleye energy density increased with increasing walleye weight, and therefore we fitted a straight line, using simple linear regression analysis, to walleye energy density as a function of walleye weight. Walleye weights

ranged from 487 to 6426 g. We used the fitted regression line $ED = 7776 + 0.231W$, where ED = walleye energy density (J/g wet weight) and W = walleye wet weight (g), for the bioenergetics modeling of Lake Erie walleye. Because energy density did not significantly differ between the sexes of Lake Erie walleye, this fitted regression line was used for both sexes. Similarly, energy density did not significantly differ between the sexes of Lake Huron walleye, and therefore the walleye energy density function used by Pothoven et al. (2017) was applied to both sexes in our Lake Huron simulations.

Because 15 October represented the typical midpoint date for the duration of the Ohio DNR fall gillnet survey in Lake Erie, we assigned 15 October as the starting day for the Lake Erie bioenergetics model simulations. Spawning day was assigned 10 April for Lake Erie walleye (Ohio Division of Wildlife 2011, 2012, 2013). Again, spawning was simulated by females and males losing 22.1% and 2.9%, respectively, of their body weight on the spawning day. Females and males were assumed to mature by ages 4 and 3 years, respectively (Ohio Division of Wildlife 2011, 2012, 2013).

Bioenergetics modeling: exchanging temperature regimes between areas and sexes

Walleye growth is affected by both food availability and the temperature regime experienced by the walleye. One of the steps in identifying which of these two factors is the more influential determinant of walleye growth in the Lake Huron – Lake Erie system is to assess the scope for changes in walleye growth brought about by geographic variation in temperature regimes. Results from this scoping exercise can then be compared with observed geographic differences in walleye growth within the Lake Huron – Lake Erie system solely attributable to food availability, and then relative importance of the two factors can be determined. To assess the

scope for changes in walleye growth brought about by geographic variation in temperature regimes, we performed the following set of bioenergetics model simulations. For a specific combination of geographic area and sex, walleye growth was predicted using the food consumption schedule for that specific combination of geographic area and sex (as determined by the fit-to-growth simulations) but replacing the temperature regime for that specific combination of geographic area and sex with the temperature regimes for the other three geographic areas and the same sex. The food consumption schedule was the total amount of food consumed by an average walleye over the course of the year for each of the 10 simulation years from age 3 through age 13, as determined by the 10 fit-to-growth simulations. Observed weight at age 3 was the starting walleye weight, and walleye growth was predicted over ages 4-13. For each age between 4 and 13, predicted walleye weight was compared with observed walleye weight. For example, consider the case of northern Lake Huron female walleye. We predicted female walleye growth from age 3 through age 13, using the feeding schedule for northern Lake Huron females and maintaining bioenergetics model inputs specific for northern Lake Huron females other than temperature regime, under the following three temperature regimes: (i) southern Lake Huron female walleye, (ii) Saginaw Bay female walleye, and (iii) Lake Erie female walleye. The starting weight for each of these three scenarios was the observed weight at age 3 for Lake Huron female walleye. For each of the three scenarios, predicted weight at age was compared with observed weight at age for Lake Huron female walleye. Percent changes from observed weight at age were recorded.

In our bioenergetics model simulations, we kept food consumption constant while subjecting the fish to temperature regimes from other geographic areas in order to assess the effect of geographic variation in temperature regimes on fish growth. Our approach was similar to that of

Kao et al. (2015a, 2015b), who inputted the present-day food consumption schedule into fish bioenergetics models while subjecting the fish to a temperature regime predicted for the future using climate change models. The goal of the Kao et al. (2015a, 2015b) studies was to forecast the effects of climate change on fish growth in the Laurentian Great Lakes.

To assess the effect of variability in temperature regimes between the sexes on walleye growth, we performed the following set of bioenergetics model simulations. For a specific combination of geographic area and sex, walleye growth was predicted using the food consumption schedule for that specific combination of geographic area and sex (as determined by the fit-to-growth simulations) but replacing the temperature regime for that specific combination of geographic area and sex with the temperature regime for that specific geographic area but for the opposite sex. Again, observed weight at age 3 was the starting walleye weight, and walleye growth was predicted over ages 4-13. Predicted walleye weight was compared with observed walleye weight for ages 4-13. Consider the case of northern Lake Huron female walleye as an example. We predicted female walleye growth from age 3 through age 13, using the feeding schedule for northern Lake Huron females and maintaining bioenergetics model inputs specific for northern Lake Huron females other than temperature regime, but we replaced the temperature regime for northern Lake Huron female walleye with the temperature regime for northern Lake Huron male walleye. Initial weight was the observed weight at age 3 for Lake Huron female walleye. Predicted weight at age was compared with observed weight at age for Lake Huron female walleye over ages 4-13, and percent changes from observed weight at age were recorded. We did not assess the effect of variation in food availability between the sexes on the sex difference in walleye growth, because factors other than a sex difference in food availability likely were responsible for faster growth by females compared with males (Bozek et

al. 2011; Rennie and Venturelli 2015). In contrast, assessment of the effect of a sex difference in temperature regimes on the sex difference in growth was a secondary goal of our study.

Results

Temperature regimes

Temperature data downloaded from temperature loggers implanted in 81 individuals (16% of total number of implanted individuals) were used to derive temperature regimes experienced by walleye. Temperature data from 5 females and 3 males were used to characterize the temperature regimes for walleye inhabiting northern Lake Huron, whereas the temperature regimes for walleye inhabiting southern Lake Huron were based on data from 7 females and 2 males. In addition, temperature data from 18 females and 17 males and from 13 females and 16 males were used to characterize the temperature regimes for walleye residing in Saginaw Bay and Lake Erie, respectively. Ages of the walleye from Lake Huron ranged from 4 to 14 years, and only one of these walleye was an age-14 fish. Similarly, ages of walleye from Lake Erie ranged from 4 to 13 years.

On average, temperatures experienced by the walleye were lowest for northern Lake Huron walleye and highest for Lake Erie walleye (Fig. 2). Over the course of the calendar year, mean temperatures experienced by female and male walleye from northern Lake Huron were 8.7 and 8.6 °C, respectively. Temperature regimes were slightly higher for southern Lake Huron walleye compared with northern Lake Huron walleye, with female and male walleye from southern Lake Huron experiencing average temperatures of 9.3 and 8.9 °C, respectively, over the course of the calendar year. Saginaw Bay walleye experienced higher temperatures than walleye from the main basin of Lake Huron, as female and male walleye from Saginaw Bay experienced mean

temperatures of 10.8 and 10.5 °C, respectively, during the year (Fig. 2). Temperatures experienced by female and male walleye from Lake Erie averaged 11.1 and 11.5 °C, respectively. Thus, on average, temperature regimes for female walleye were higher than those for male walleye in the main basin of Lake Huron and Saginaw Bay, whereas the temperature regime for male walleye from Lake Erie averaged a higher value than the temperature regime for female walleye from Lake Erie. Peak temperatures experienced by female walleye were 20.6, 21.3, 23.8, and 23.4 °C in northern Lake Huron, southern Lake Huron, Saginaw Bay, and Lake Erie, respectively (Fig. 2). Peak temperatures experienced by male walleye were 20.6, 21.8, 22.5, and 25.1 °C in northern Lake Huron, southern Lake Huron, Saginaw Bay, and Lake Erie, respectively.

Growth trajectories

Walleye attained much larger sizes in Lake Erie than in Lake Huron (Fig. 3; Table 3). Age-3 female walleye in Lake Erie were 2.5 times greater in weight than age-3 female walleye in Lake Huron, and age-3 male walleye in Lake Erie were 2.2 times greater in weight than age-3 male walleye in Lake Huron. At age 8, female and male walleye in Lake Erie were 84% and 72%, respectively, greater in weight than female and male walleye in Lake Huron (Fig. 3). At age 13, mean weight of female walleye, as estimated from our analysis, was 3.2 kg in Lake Erie compared with 2.1 kg in Lake Huron. For males, we estimated mean weight in Lake Erie to be 1.9 kg compared with 1.2 kg in Lake Huron. Averaging across ages 3-13, female walleye from Lake Erie exceeded female walleye from Lake Huron in weight by 92%. Similarly, averaging across ages 3-13, male walleye from Lake Erie were 78% greater in weight than male walleye

from Lake Huron. Averaging across ages and sexes, walleye growth in Lake Erie was 85% greater than walleye growth in Lake Huron.

Females grew substantially faster than males in both Lake Huron and Lake Erie (Fig. 3).

Averaging across ages 3-13, female walleye from Lake Huron were 47% greater in weight than Lake Huron male walleye. Similarly, averaging across ages 3-13, female walleye from Lake Erie were 56% greater in weight than Lake Erie male walleye.

Based on results from t tests for paired comparisons, Lake Erie female walleye at a given age were significantly greater in weight than Lake Huron female walleye ($t = 28.74$; $df = 10$; $P < 0.0001$). Likewise, Lake Erie male walleye at a given age were significantly greater in weight than Lake Huron male walleye ($t = 56.74$; $df = 10$; $P < 0.0001$). At a given age, female walleye were significantly greater in weight than male walleye in both Lake Huron ($t = 5.56$; $df = 10$; $P = 0.0002$) and Lake Erie ($t = 9.45$; $df = 10$; $P < 0.0001$).

Bioenergetics modeling: fit to observed growth

On average, the rate of food consumption by Lake Erie walleye was nearly double the rate of food consumption by Lake Huron walleye (Fig. 4). At age 13, cumulative consumption by female and male walleye from Lake Erie was 59.9 and 32.4 kg, respectively. In contrast, cumulative consumption at age 13 by female and male walleye from Lake Huron ranged from 32.0 kg and 17.2 kg, respectively, in northern Lake Huron to 42.5 kg and 23.5 kg, respectively, in Saginaw Bay (Fig. 4). Thus, cumulative consumption at age 13 in Saginaw Bay was roughly 30% greater than cumulative consumption in the main basin of Lake Huron. Averaging cumulative consumption across ages 4-13, mean cumulative consumption for Lake Erie walleye

was greater than that for Lake Huron (averaging across the three geographic areas) by factors of 2.0 and 1.8 for females and males, respectively.

On average, the rate of food consumption by female walleye exceeded the rate of food consumption by male walleye by a factor of 1.6 (Fig. 4). Averaging cumulative consumption across ages 4-13, mean cumulative consumption by females was greater than that by males by factors of 1.6, 1.6, 1.6, and 1.7 for northern Lake Huron, southern Lake Huron, Saginaw Bay, and Lake Erie, respectively.

Bioenergetics modeling: exchanging temperature regimes between areas and sexes

Averaging across ages 4-13, changes in weight at age resulting from exchanging temperature regimes between geographical areas did not exceed 16% for female walleye and did not exceed 29% for male walleye (Figs. 5-8). For northern Lake Huron, weight at age for female and male walleye decreased, on average, by 15% and 26%, respectively, when the Lake Erie temperature regimes were imposed (Fig. 5). In contrast, weight at age decreased, on average, by only 5% for both female and male walleye when the southern Lake Huron temperature regimes were imposed in northern Lake Huron. For southern Lake Huron, weight at age for female and male walleye decreased, on average, by 11% and 23%, respectively, when the Lake Erie temperature regimes were imposed (Fig. 6). However, weight at age increased by 5%, on average, for both female and male walleye when the northern Lake Huron temperatures were imposed in southern Lake Huron. For Saginaw Bay, weight at age of female and male walleye increased, on average, by 16% and 21% when northern Lake Huron temperatures were imposed (Fig. 7). When Lake Erie temperatures were imposed in Saginaw Bay, weight at age of female and male walleye decreased, on average, by < 1% and 10%, respectively. For Lake Erie, weight at age of female

and male walleye increased, on average, by 14% and 29%, respectively, when northern Lake Huron temperatures were used in the bioenergetics model simulations (Fig. 8). In contrast, weight at age of female and male walleye increased, on average, by only 1% and 9%, respectively, when the Saginaw Bay temperature regimes were imposed.

Changes in walleye weight at age were less than 7% when temperature regimes were exchanged between sexes in the bioenergetics model simulations (Fig. 9). For northern Lake Huron, weight at age of females increased by 1%, on average, while weight at age of males decreased by 2%, on average, when temperature regimes were exchanged between the sexes. For southern Lake Huron, weight at age of females increased by 3%, on average, while weight at age of males decreased by 3%, on average, when temperatures were swapped between the sexes. For Saginaw Bay, weight at age of females increased by 3%, on average, whereas weight at age of males decreased by 3%, on average, when temperatures were exchanged between the sexes (Fig. 9). Finally, for Lake Erie, weight at age of females decreased by 3%, on average, while weight at age of males increased by 5%, on average, when temperature regimes were exchanged between the sexes.

Discussion

The observed difference in temperature regimes experienced by walleye between the two lakes made no contribution whatsoever toward the faster walleye growth in Lake Erie compared with Lake Huron. Data from our iButton temperature loggers implanted in fish indicated that Lake Erie walleye experienced higher temperatures than Lake Huron walleye. Thus, if food availability, as indexed by food consumption, did not differ between the two lakes and all other factors (factors other than temperature regime) affecting growth were identical between the two

lakes, then walleye growth would have been expected to be faster in Lake Huron than in Lake Erie. Moreover, results from our bioenergetics model simulations showed that walleye growth in Lake Huron would decrease if Lake Erie temperature regimes were imposed in Lake Huron, and that walleye growth in Lake Erie would increase if Lake Huron temperature regimes were imposed in Lake Erie.

We conclude that faster walleye growth in Lake Erie compared with walleye growth in Lake Huron was most likely due to much greater food availability in Lake Erie compared with that in Lake Huron. This conclusion was supported by the available data from prey fish hydroacoustics surveys from both lakes. Lakewide mean pelagic prey fish biomass density in Lake Huron was about 8 kg/ha during 2011-2012, based on results from the lakewide hydroacoustics survey (Warner et al. 2013). The dominant pelagic prey fish species in Lake Huron included rainbow smelt and bloater (*Coregonus hoyi*), and each species contributed approximately 50% to the overall biomass density. In contrast to Lake Huron, acoustic estimates of mean biomass density of yearling and older rainbow smelt in central and eastern Lake Erie during 2011-2012 ranged from 69 to 85 kg/ha (Forage Task Group 2012, 2013; Patrick Kocovsky, US Geological Survey, Great Lakes Science Center, personal communication). Our conclusion was also supported by the available data on lake primary production. Estimates of annual primary production in Lake Erie and Lake Huron during 2008, the only recent year for which these estimates were available, were 5.4 kt C/ha and 0.8 kt C/ha, respectively, where 1 kt C = 1 kilotonne C = 1000 metric tons of carbon fixed by photosynthesis during the year (Great Lakes Primary Production Model 2017). Thus, from a primary production standpoint, Lake Erie had potential to support a substantially higher prey fish biomass density than Lake Huron.

The higher food availability in Lake Erie compared with Lake Huron was not due to the biomass density of walleye in Lake Huron being substantially greater than that in Lake Erie. As density of a fish population increases, intraspecific competition for food within the fish population is expected to increase, and consequently growth of individual fish within the population would be expected to decrease (Lorenzen and Enberg 2002). Thus, as biomass density of the fish population increases, availability of food for an individual fish in the population tends to decrease, and growth of individual fish tends to decrease as well. Adult walleye population biomass in Saginaw Bay and the main basin of Lake Huron averaged 1.6 kt during 2011-2012 (Fielder and Bence 2014). Assuming that the bulk of the walleye population in the main basin of Lake Huron inhabits waters corresponding to bottom depths less than 23 m deep, and including all of Saginaw Bay as habitable by walleye, the surface area available to walleye in the main basin of Lake Huron and Saginaw Bay would be 7057 km² (Wang et al. 2015). This assumption appeared to be a reasonable one (Fielder and Bence 2014). Thus, adult walleye biomass density in the habitable region of Lake Huron's main basin and Saginaw Bay was estimated to be 2 kg/ha. Adult walleye population biomass in Lake Erie averaged 65.4 kt during 2011-2012 (Walleye Task Group 2016), and all of Lake Erie's surface area of 25690 km² was considered habitable by walleye (Ryan et al. 2003; Pandit et al. 2013). Thus, adult walleye biomass density in Lake Erie was estimated to be 25 kg/ha, a value more than an order of magnitude higher than that for Lake Huron. Clearly, the higher food availability in Lake Erie was not attributable to greater adult walleye biomass density in Lake Huron compared with that for Lake Huron.

We also conclude that food availability was of considerably more importance in determining walleye growth in the Lake Huron – Lake Erie system than temperature regime. The faster

walleye growth in Lake Erie than in Lake Huron was solely attributable to greater food availability in Lake Erie than in Lake Huron. On average, Lake Erie walleye grew 85% faster than Lake Huron walleye. Our bioenergetics modeling results indicated that variability in temperature regimes among geographic areas of the Lake Huron – Lake Erie system could account for up to a 16% change in the growth of female walleye and a 29% change in the growth of male walleye. More specifically, we showed that variation in temperature regimes between the three basins of Lake Huron could account for as high as a 16% change in growth of female walleye and as high as a 21% change in growth of male walleye. In addition, variation in temperature regimes between the two lakes could account for as high as a 15% change in growth of female walleye and as high as a 29% change in growth of male walleye. Clearly, of these two factors affecting walleye growth, food availability was the dominant factor in the Lake Huron – Lake Erie system.

The faster walleye growth in Lake Erie compared with that in Lake Huron was not explained by differences in prey and/or walleye energy densities between the two lakes. If the ratio of walleye energy density to prey energy density was greater for Lake Huron than for Lake Erie, then walleye would have to consume more food in Lake Huron than in Lake Erie to achieve the same amount of growth, all other factors being equal. Thus, this scenario would contribute toward faster walleye growth in Lake Erie than in Lake Huron. However, the ratio of walleye energy density to prey energy density for Lake Erie was greater than that for Lake Huron. Averaging across ages 3-13, walleye energy densities were 6311 and 8288 J/g in Lake Huron and Lake Erie, respectively. Weighted (weighting by the proportion of each prey type eaten) averages of prey energy densities were 4888, 4251, and 5350 J/g in the main basin of Lake Huron, Saginaw Bay, and Lake Erie, respectively. Consequently, values of the ratio of walleye

597 energy density to prey energy density were 1.29, 1.48, and 1.55 for the main basin of Lake
598 Huron, Saginaw Bay, and Lake Erie, respectively. Thus, Lake Erie walleye had to consume
599 more food than Lake Huron walleye to achieve the same amount of growth, all other factors
600 being equal. And Lake Erie walleye grew nearly twice as fast as Lake Huron walleye. In sum,
601 differences in walleye and prey energy densities between the two lakes did not contribute toward
602 the faster walleye growth in Lake Erie.

603 Our research has provided new insights into the effect of variation in temperature regimes
604 between the sexes on the difference in walleye growth between the sexes. For all three basins of
605 Lake Huron, females experienced slightly higher temperatures, on average, than males.
606 Exchanging temperature regimes between the sexes of Lake Huron walleye resulted in slight (<
607 5%) increases in growth of females and slight decreases in growth of males. Thus, sex
608 differences in temperature regimes could not account for any portion of the observed difference
609 in growth between the sexes of Lake Huron walleye, as observed weight at age of females
610 averaged 47% higher than that of males over ages 3-13. For Lake Erie, males experienced
611 slightly higher temperatures than females. Exchanging temperature regimes between the sexes
612 of Lake Erie walleye led to a 3% decrease, on average, in growth of females and a 5% increase,
613 on average, in growth of males. Thus, the sex difference in temperature regimes could account
614 for females being about 8% greater in weight at age than males from Lake Erie. Females were
615 observed to be 56% greater, on average, in weight at age than males from Lake Erie. We
616 conclude that the sex difference in temperature regimes explained 15% of the observed
617 difference in growth between the sexes of Lake Erie walleye, but none of the observed sex
618 difference in growth of Lake Huron walleye. More research is needed to identify the
619 predominant causes underlying sexually dimorphic growth in fish. We also note that the slightly

higher temperature regime experienced by male walleye compared with female walleye in Lake Erie was expected based on results reported by Wang et al. (2007). Using data from a long-term mark and recapture study, these researchers showed that male walleye were more likely to reside in the western basin of Lake Erie throughout the year than female walleye. On average, surface water temperatures in the western basin are higher than those in the central or eastern basin of Lake Erie (Moukomla and Blanken 2016), and therefore higher temperatures experienced by males would be presumed. Reasons for the slightly higher temperatures experienced by females compared with males in Lake Huron were unclear.

One caveat for our findings was that the growth trajectories that we developed for walleye from Lake Huron were specifically for the Saginaw Bay walleye stock, which inhabits Saginaw Bay and the main basin of Lake Huron (Fielder and Bence 2014; Hayden et al. 2014). Walleye stocks also inhabit both Georgian Bay and the North Channel of Lake Huron, but these walleye stocks do not mix with the Saginaw Bay stock (Fielder et al. 2010; Stepien et al. 2010). Walleye grew faster in Georgian Bay and the North Channel than in the main basin of Lake Huron during 1982-2003 (Chu and Koops 2007), and this difference in growth appears to hold true today as well (Arunas Liskauskas and Adam Cottrill, Ontario Ministry of Natural Resources and Forestry, personal communication).

Another caveat for our findings was that the temperature regimes that we developed for walleye residing in the main basin of Lake Huron were based on relatively low numbers of recaptured walleye. Temperature regimes for the northern main basin of Lake Huron were based on data from 5 females and 3 males, and temperature regimes for the southern main basin of Lake Huron were based on data from 7 females and 2 males. However, despite the low sample sizes with regard to number of walleye recaptured, the temperature regimes did indeed show that

walleye experienced slightly higher temperatures in the southern main basin than in the northern main basin and that walleye experienced higher temperatures in Saginaw Bay than in the main basin of Lake Huron, as would be expected based on the long-term observations of surface water temperatures by Moukomla and Blanken (2016). To perform a bioenergetics analysis for walleye growth in Georgian Bay and the North Channel of Lake Huron analogous to that from our study, temperature regimes for Georgian Bay walleye and North Channel walleye would need to be developed. This would involve implanting Georgian Bay walleye and North Channel walleye with temperature loggers.

Our results should be useful in refining and updating models for quantifying the role of walleye as a piscivore in the Lake Huron and Lake Erie ecosystems. He et al. (2015) coupled age-structured population models with fish bioenergetics models to assess the effect of piscivory by Chinook salmon (*Oncorhynchus tshawytscha*), lake trout (*Salvelinus namaycush*), walleye, and lake whitefish (*Coregonus clupeaformis*) on Lake Huron prey fish populations. For modeling purposes, these researchers assumed that the temperature regime experienced by walleye was equivalent to the long-term mean temperature profile from daily water temperature records kept at the Bay Metro Water Treatment Plant, which drew water directly from Saginaw Bay. Their modeling effort could be refined and updated by applying the temperature regimes developed in our study based on data from iButton temperature loggers implanted in Lake Huron walleye. With regard to assessing the role of piscivory by the Lake Erie walleye population on prey fish populations, previous efforts have relied on temperature regimes derived from measurements of water temperatures in Lake Erie (Hartman and Margraf 1992; Kershner et al. 1999), rather than derived from in situ measurements from temperature loggers implanted in Lake Erie walleye. Application of our temperature regimes based on data from the iButton

temperature loggers would lead to more accurate estimates of prey fish consumption by the Lake Erie walleye population. Moreover, the two above-mentioned bioenergetics model applications to the Lake Erie walleye population were not based on energy density determinations specifically for Lake Erie walleye. Rather, energy density of Lake Erie walleye was assumed to equal that of a typical value for fishes (Winberg 1956; Hewett and Johnson 1992). Use of our energy density determinations specifically for Lake Erie walleye would improve the accuracy of the bioenergetics model applications to the Lake Erie walleye population. Finally, detailed studies on walleye diet in Lake Erie, like the studies conducted during the late 1970s and 1980s (Knight et al. 1984; Hartman and Margraf 1992), have rarely been published since 1995. Our study provided a detailed diet schedule for Lake Erie walleye, from spring through fall based on more than 3000 non-empty stomachs, by summarizing stomach contents data over years 2000-2015. This diet schedule will be invaluable in updating and refining models used to quantify the effects of piscivory by the Lake Erie walleye population on Lake Erie prey fish populations. Once the models for quantifying piscivory by the walleye population have been refined, these new models would contribute toward ecosystem-based management, as advocated by Link (2002), Pikitch et al. (2004), Tsehaye et al. (2014), and Grüss et al. (2017), of the important fisheries operating in both lakes.

Our study was innovative, from two perspectives. First, our study represented the first assessment of the effect of a sex difference in temperature regimes on the sex difference in growth within a fish population. We found that the sex difference in temperature regimes for Lake Huron walleye made no contribution at all to the observed faster growth of females compared with males. In the case of Lake Erie walleye, the sex difference in temperature regimes accounted for 15% of the observed difference in growth between the sexes. Thus,

although the sex difference in temperature regimes contributed to the observed sex difference in growth of Lake Erie walleye, it was not the primary driver of the sexually dimorphic walleye growth. Second, we were the first to evaluate the effects of both interbasin and interlake variability in temperature regimes on fish growth in a two-lake system with a waterway connection between the two lakes. Results showed that interbasin and interlake variability in temperature regimes had just a moderate effect on walleye growth in the Lake Huron – Lake Erie system. Further, food availability was the dominant factor affecting walleye growth in this two-lake system. Coupling of archival temperature tagging and acoustic telemetry with fish bioenergetics modeling should continue to provide new insights into the relative importance of temperature regime as a factor influencing fish growth in fish populations from multi-lake systems around the world.

Acknowledgements

The authors would like to convey our appreciation to the staffs at the Sandusky and Fairport Harbor Fisheries Research Stations (Ohio DNR), Lake Erie Management Unit (OMNRF), Waterford Fisheries Station - West (Michigan DNR), Lake Erie and Hammond Bay Biological Stations (US Geological Survey Great Lakes Science Center), and Lower Great Lakes Fish and Wildlife Conservation Office (US Fish and Wildlife Service) for maintaining the acoustic receiver network throughout the duration of the study. We are specifically grateful to K. Murchie (Carleton University), J. Ives (Great Lakes Fishery Commission), and S. Zahn Seegert (Great Lakes Fishery Commission) for their assistance in the field implanting of acoustic transmitters, to C. Holbrook (Hammond Bay Biological Station) for providing study design recommendations, and to the staff at the Fairport Harbor Fisheries Research Station for

maintaining the Lake Erie walleye diet database. We are also grateful to the Great Lakes Fishery Commission for distributing rewards for captured walleye implanted with temperature loggers. We thank Danielle Forsythe for her assistance with calculation of lake surface areas. Use of trade, product, or firm names does not imply endorsement by the United States Government. This work was funded by the Great Lakes Fishery Commission by way of Great Lakes Research Initiative appropriations (GL-00E23010) to CCK. This article is Contribution 33 of the Great Lakes Acoustic Telemetry Observation System (GLATOS).

References

- Bergstedt, R.A., Argyle, R.L., Seelye, J.G., Scribner, K.T., and Curtis, G.L. 2003. In situ determination of the annual thermal habitat use by lake trout (*Salvelinus namaycush*) in Lake Huron. J. Gt. Lakes Res. **29**(Suppl. 1): 347-361.
- Bozek, M.A., Baccante, D.A., and Lester, N.P. 2011. Walleye and sauger life history. In Biology, management, and culture of walleye and sauger. Edited by B.A. Barton. American Fisheries Society, Bethesda. pp. 233-301.
- Bunnell, D.B., Johnson, T.B., and Knight, C.T. 2005. The impact of introduced round gobies (*Neogobius melanostomus*) on phosphorus cycling in central Lake Erie. Can. J. Fish. Aquat. Sci. **62**: 15-29.
- Chipps, S.R., and Wahl, D.H. 2008. Bioenergetics modeling in the 21st century: reviewing new insights and revisiting old constraints. Trans. Am. Fish. Soc. **137**: 298-313.
- Chu, C., and Koops, M.A. 2007. Life history parameters of Great Lakes populations of lake trout, lake whitefish, bloater, walleye, and yellow perch. Can. Manuscr. Rep. Fish. Aquat. Sci. 2811.

- 735 Cooke, S.J., Midwood, J.D., Thiem, J.D., Klimley, P., Lucas, M.C., Thorstad, E.B., Eiler, J.,
736 Holbrook, C., and Ebner, B.C. 2013. Tracking animals in freshwater with electronic tags:
737 past, present and future. *Anim. Biotelem.* **1**: 5.
- 738 Cummins, K.W., and Wuycheck, J.C. 1971. Caloric equivalents for investigations in ecological
739 energetics. *Mitt. Int. Ver. Theor. Angew. Limnol.* **18**: 1-151.
- 740 Deslauriers, D., Chipps, S.R., Breck, J.E., Rice, J.A., and Madenjian, C.P. 2017. Fish
741 Bioenergetics 4.0: An R-based modeling application. *Fisheries* **42**: 586-596.
- 742 Donaldson, M.R., Cooke, S.J., Patterson, D.A., Hinch, S.C., Robichaud, D., Hanson, K.C.,
743 Olsson, I., Crossin, G.T., English, K.K., and Farrell, A.P. 2009. Limited behavioural
744 thermoregulation by adult upriver-migrating sockeye salmon (*Oncorhynchus nerka*) in the
745 Lower Fraser River, British Columbia. *Can. J. Zool.* **87**: 480-490.
- 746 Enberg, K., Dunlop, E.S., and Jørgensen, C. 2008. Fish growth. *In Encyclopedia of ecology.*
747 *Edited by S.E. Jørgensen and B.D. Fath.* Elsevier, Oxford. pp. 1564-1572.
- 748 Feiner, Z.S., and Höök, T.O. 2015. Environmental biology of percid fishes. *In Biology and*
749 *culture of percid fishes: principles and practices.* *Edited by P. Kestemont, K. Dabrowski,*
750 *and R.C. Summerfelt.* Springer, Dordrecht. pp. 61-100.
- 751 Fielder, D.G. 2014. Mortality, exploitation, movement, and stock size of Saginaw Bay
752 Walleyes, 1981-2011: 31 years of tag return analysis. Michigan Department of Natural
753 Resources, Fisheries Report 04, Lansing, Mich.
- 754 Fielder, D.G., and Bence, J.R. 2014. Integration of auxiliary information in statistical catch-at-
755 age (SCA) analysis of the Saginaw Bay stock of Walleye in Lake Huron. *N. Am. J. Fish.*
756 *Manage.* **34**: 970-987.

- Fielder, D.G., and Thomas, M.V. 2014. Status and trends of the fish community of Saginaw Bay, Lake Huron 2005-2011. Michigan Department of Natural Resources, Fisheries Report 03, Lansing, Mich.
- Fielder, D.G., Liskauskas, A.P., Gonder, D.J.A., Mohr, L.C., and Thomas, M.V. 2010. Status of walleye in Lake Huron. *In* Status of walleye in the Great Lakes: proceedings of the 2006 Symposium. Great Lakes Fish. Comm. Tech. Rep. 69. pp. 71-90.
- Fielder, D.G., Kolb, T.L., Goniew, T.M., Wesander, D.L., and Schrouder, K.S. 2014. Fisheries of Saginaw Bay, Lake Huron 1986-2010. Michigan Department of Natural Resources, Fisheries Report 02, Lansing, Mich.
- Forage Task Group. 2012. Report of the Lake Erie Forage Task Group to the Standing Technical Committee of the Lake Erie Committee, Great Lakes Fishery Commission, Ann Arbor, Mich. Available from: http://www.glfc.org/lakecom/lec/FTG_docs/annual_reports/FTG_report_2012.pdf. [accessed 1 June 2017].
- Forage Task Group. 2013. Report of the Lake Erie Forage Task Group to the Standing Technical Committee of the Lake Erie Committee, Great Lakes Fishery Commission, Ann Arbor, Mich. Available from: http://www.glfc.org/lakecom/lec/FTG_docs/annual_reports/FTG_report_2013.pdf. [accessed 1 June 2017].
- Great Lakes Information Network. 2017. Great Lakes Commission, Ann Arbor, Mich. Available from: <http://maps.glin.net>. [accessed 1 December 2017].

- 778 Great Lakes Primary Production Model. 2017. Michigan Tech Research Institute, Michigan
779 Technological University, Ann Arbor, Mich. Available from:
780 <http://www.mtri.org/glppm.html>. [accessed 1 June 2017].
- 781 Grüss, A., Rose, K.A., Simons, J., Ainsworth, C.H., Babcock, E.A., Chagaris, D.D., De Mutsert,
782 K., Froeschke, J., Himchak, P., Kaplan, I.C., O'Farrell, H., and Zetina Rejon, M.J. 2017.
783 Recommendations on the use of ecosystem modeling for informing ecosystem-based
784 fisheries management and restoration outcomes in the Gulf of Mexico. *Mar. Coast. Fish.* **9**:
785 281-295.
- 786 Hartman, K.J., and Margraf, F.J. 1992. Effects of prey and predator abundances on prey
787 consumption and growth of walleyes in western Lake Erie. *Trans. Am. Fish. Soc.* **121**: 245-
788 260.
- 789 Hayden, T.A., Holbrook, C.M., Fielder, D.G., Vandergoot, C.S., Bergstedt, R.A., Dettmers, J.M.,
790 Krueger, C.C., and Cooke, S.J. 2014. Acoustic telemetry reveals large-scale migration
791 patterns of walleye in Lake Huron. *PLoS ONE* **9**: e11833.
- 792 Hayward, R.S., and Margraf, F.J. 1987. Eutrophication effects on prey size and food available
793 to yellow perch in Lake Erie. *Trans. Am. Fish. Soc.* **116**: 210-223.
- 794 He, J.X., Bence, J.R., Madenjian, C.P., Pothoven, S.A., Dobiesz, N.E., Fielder, D.G., Johnson,
795 J.E., Ebener, M.P., Cottrill, R.A., Mohr, L.C., and Koproski, S.R. 2015. Coupling age-
796 structured stock assessment and fish bioenergetics models: a system of time-varying models
797 for quantifying piscivory patterns during the rapid trophic shift in the main basin of Lake
798 Huron. *Can. J. Fish. Aquat. Sci.* **72**: 7-23.

- 799 Hewett, S.W., and Johnson, B.L. 1992. Fish bioenergetics model 2: an upgrade of a generalized
800 bioenergetics model of fish growth for microcomputers. University of Wisconsin Sea Grant
801 Institute, Technical Report WIS-SG-92-250, Madison, Wisc.
- 802 Jobling, M. 1994. Fish bioenergetics. Chapman and Hall, London.
- 803 Kao, Y.-C., Madenjian, C.P., Bunnell, D.B., Lofgren, B.M., and Perroud, M. 2015a.
804 Temperature effects induced by climate change on the growth and consumption by
805 salmonines in Lakes Michigan and Huron. *Environ. Biol. Fishes* **98**: 1089-1104.
- 806 Kao, Y.-C., Madenjian, C.P., Bunnell, D.B., Lofgren, B.M., and Perroud, M. 2015b. Potential
807 effects of climate change on the growth of fishes from different thermal guilds in Lakes
808 Michigan and Huron. *J. Gt. Lakes Res.* **41**: 423-435.
- 809 Kershner, M.W. 1996. Walleye predatory demand and prey supply in Lake Erie: exploring
810 environmental variation and its historical effects. Ph.D. Dissertation, Ohio State University,
811 Columbus, Ohio.
- 812 Kershner, M.W., Schael, D.M., Knight, R.L., Stein, R.A., and Marschall, E.A. 1999. Modeling
813 sources of variation for growth and predatory demand of Lake Erie walleye (*Stizostedion*
814 *vitreum*), 1986-1995. *Can. J. Fish. Aquat. Sci.* **56**: 527-538.
- 815 Kitchell, J.F., Stewart, D.J., and Weininger, D. 1977. Applications of a bioenergetics model to
816 yellow perch (*Perca flavescens*) and walleye (*Stizostedion vitreum vitreum*). *J. Fish. Res.*
817 *Board Can.* **34**: 1922-1935.
- 818 Knight, R.L., Margraf, F.J., and Carline, R.F. 1984. Piscivory by walleyes and yellow perch in
819 western Lake Erie. *Trans. Am. Fish. Soc.* **113**: 677-693.
- 820 Landsman, S.J., Nguyen, V.M., Gutowsky, L.F.G., Gobin, J., Cook, K.V., Binder, T.R., Lower,
821 N., McLaughlin, R.L., and Cooke, S.J. 2011. Fish movement and migration studies in the

- 822 Laurentian Great Lakes: Research trends and knowledge gaps. *J. Gt. Lakes Res.* **37**: 365-
823 379.
- 824 Link, J.S. 2002. What does ecosystem-based fisheries management mean? *Fisheries* **27(4)**: 18-
825 21.
- 826 Lorenzen, K., and Enberg, K. 2002. Density-dependent growth as a key mechanism in the
827 regulation of fish populations: evidence from among-population comparisons. *Proc. R. Soc.*
828 *Lond. B Biol. Sci.* **269**: 49-54.
- 829 Lucas, M.C., and Baras, E. 2000. Methods for studying spatial behavior of freshwater fishes in
830 the natural environment. *Fish Fish.* **1**: 283-316.
- 831 Madenjian, C.P. 2011. Bioenergetics in ecosystems. *In* Encyclopedia of fish physiology: from
832 genome to environment. *Edited by* A.P. Farrell. Academic Press, San Diego. pp. 1675-
833 1680.
- 834 Madenjian, C.P., Noguchi, G.E., Haas, R.C., and Schrouder, K.S. 1998. Sexual difference in
835 polychlorinated biphenyl accumulation rates of walleye (*Stizostedion vitreum*). *Can. J. Fish.*
836 *Aquat. Sci.* **55**: 1085-1092.
- 837 Madenjian, C.P., Cochran, P.A., and Bergstedt, R.A. 2003. Seasonal patterns in growth, blood
838 consumption, and effects on hosts by parasitic-phase sea lampreys in the Great Lakes: an
839 individual-based model approach. *J. Gt. Lakes Res.* **29(Suppl. 1)**: 332-346.
- 840 Madenjian, C.P., O'Connor, D.V., Pothoven, S.A., Schneeberger, P.J., Rediske, R.R., O'Keefe,
841 J.P., Bergstedt, R.A., Argyle, R.L., and Brandt, S.B. 2006. Evaluation of a lake whitefish
842 bioenergetics model. *Trans. Am. Fish. Soc.* **135**: 61-75.
- 843 Melstrom, R.T., and Lupi, F. 2013. Valuing recreational fishing in the Great Lakes. *N. Am. J.*
844 *Fish. Manage.* **33**: 1184-1193.

- 845 Moukomla, S., and Blanken, P.D. 2016. Remote sensing of the North American Laurentian
846 Great Lakes' surface temperature. *Remote Sens.* **8**: 286.
- 847 Ogle, D.H. 2016. FSA: Fisheries Stock Analysis. R package version 0.8.11.
- 848 Ohio Division of Wildlife. 2011. Ohio's Lake Erie fisheries, 2010. Annual status report.
849 Federal Aid in Fish Restoration Project F-69-P. Ohio Department of Natural Resources,
850 Division of Wildlife, Lake Erie Fisheries Units, Fairport and Sandusky, Ohio.
- 851 Ohio Division of Wildlife. 2012. Ohio's Lake Erie fisheries, 2011. Annual status report.
852 Federal Aid in Fish Restoration Project F-69-P. Ohio Department of Natural Resources,
853 Division of Wildlife, Lake Erie Fisheries Units, Fairport and Sandusky, Ohio.
- 854 Ohio Division of Wildlife. 2013. Ohio's Lake Erie fisheries, 2012. Annual status report.
855 Federal Aid in Fish Restoration Project F-69-P. Ohio Department of Natural Resources,
856 Division of Wildlife, Lake Erie Fisheries Units, Fairport and Sandusky, Ohio.
- 857 Ohio Division of Wildlife. 2016. Ohio's Lake Erie fisheries, 2015. Annual status report.
858 Federal Aid in Fish Restoration Project F-69-P. Ohio Department of Natural Resources,
859 Division of Wildlife, Lake Erie Fisheries Units, Fairport and Sandusky, Ohio.
- 860 Pandit, S.N., Zhao, Y., Ciborowski, J.J.H., Gorman, A., and Knight, C.T. 2013. Suitable habitat
861 model for walleye (*Sander vitreus*) in Lake Erie: Implications for inter-jurisdictional harvest
862 quota allocations. *J. Gt. Lakes Res.* **39**: 591-601.
- 863 Peat, T.B., Hayden, T.A., Gutowsky, L.F.G., Vandergoot, C.S., Fielder, D.G., Madenjian, C.P.,
864 Murchie, K.J., Dettmers, J.M., Krueger, C.C., and Cooke, S.J. 2015. Seasonal thermal
865 ecology of adult walleye (*Sander vitreus*) in Lake Huron and Lake Erie. *J. Therm. Biol.* **53**:
866 98-106.

- 867 Persson, L., and Greenberg, L.A. 1990. Juvenile competitive bottlenecks: the perch (*Perca*
868 *fluviatilis*)–roach (*Rutilus rutilus*) interaction. *Ecology* **71**: 44-56.
- 869 Pikitch, E.K., Santora, C., Babcock, E.A., Bakun, A., Bonfil, R., Conover, D.O., Dayton, P.,
870 Doukakis, P., Fluharty, D., Heneman, B., Houde, E.D., Link, J., Livingston, P.A., Mangel,
871 M., McAllister, M.K., Pope, J., and Sainsbury, K.J. 2004. Ecosystem-based fishery
872 management. *Science* **305**: 346-347.
- 873 Pincock, D.G. 2009. Detection performance of lines of VR2W/VR3 receivers. Available from:
874 http://vemco.com/vp-content/uploads/2012/11/line_performance.pdf. [accessed 1 June
875 2017].
- 876 Pothoven, S.A., Ludsin, S.A., Höök, T.O., Fanslow, D.L., Mason, D.M., Collingsworth, P.D.,
877 and Van Tassell, J.J. 2008. Reliability of bioelectrical impedance analysis for estimating
878 whole-fish energy density and percent lipids. *Trans. Am. Fish. Soc.* **137**: 1519-1529.
- 879 Pothoven, S.A., Bunnell, D.B., Madenjian, C.P., Gorman, O.T., and Roseman, E.F. 2012.
880 Energy density of bloaters in the Upper Great Lakes. *Trans. Am. Fish. Soc.* **141**: 772-780.
- 881 Pothoven, S.A., Madenjian, C.P., and Höök, T.O. 2017. Feeding ecology of the walleye
882 (*Percidae*, *Sander vitreus*), a resurgent piscivore in Lake Huron (Laurentian Great Lakes)
883 after shifts in the prey community. *Ecol. Freshw. Fish* **26**: 676-685.
- 884 Quinn, T.J., II, Deriso, R.B. 1999. Quantitative fish dynamics. Oxford University Press, New
885 York.
- 886 Rennie, M.D., and Venturelli, P.A. 2015. The ecology of lifetime growth in percid fishes. *In*
887 *Biology and culture of percid fishes: principles and practices*. Edited by P. Kestemont, K.
888 Dabrowski, and R.C. Summerfelt. Springer, Dordrecht. pp. 499-536.

- 889 Roseman, E.F., Knight, R.L., Wright, E., Einhouse, D., Kayle, K., Newman, K., and Hoopes,
890 R.L. 2008. Ecology and international governance of Lake Erie's percid fisheries. *In*
891 International governance of fisheries ecosystems: learning from the past, finding solutions
892 for the future. *Edited by* M.G. Schechter, N.J. Leonard, and W.W. Taylor. American
893 Fisheries Society, Bethesda. pp. 145-169.
- 894 Ryan, P.A., Knight, R., MacGregor, R., Towns, G., Hoopes, R., and Culligan, W. 2003. Fish-
895 community goals and objectives for Lake Erie. Great Lakes Fishery Commission, Special
896 Publication 03-02, Ann Arbor, Mich.
- 897 Stepien, C.A., Murphy, D.J., Lohner, R.N., Haponski, A.E., and Sepulveda-Villet, O.J. 2010.
898 Status and delineation of walleye (*Sander vitreus*) genetic stock structure across the Great
899 Lakes. *In* Status of walleye in the Great Lakes: proceedings of the 2006 Symposium. Great
900 Lakes Fish. Comm. Tech. Rep. 69. pp. 189-223.
- 901 Tsehaye, I., Jones, M.L., Bence, J.R., Brenden, T.O., Madenjian, C.P., and Warner, D.M. 2014.
902 A multispecies statistical age-structured model to assess predator-prey balance: application
903 to an intensively managed Lake Michigan pelagic fish community. *Can. J. Fish. Aquat. Sci.*
904 **71**: 627-644.
- 905 Vandergoot, C.S., Murchie, K.J., Cooke, S.J., Dettmers, J.M., Bergstedt, R.A., and Fielder, D.G.
906 2011. Evaluation of two forms of electroanesthesia and carbon dioxide for short-term
907 anesthesia in walleye. *N. Am. J. Fish. Manage.* **31**: 914-922.
- 908 Walleye Task Group. 2016. Report of the Lake Erie Walleye Task Group to the Standing
909 Technical Committee of the Lake Erie Committee, Great Lakes Fishery Commission, Ann
910 Arbor, Mich. Available:

- 911 http://www.glfc.org/lakecom/lec/WTG_docs/annual_reports/WTG_report_2016.pdf.
912 [accessed 1 June 2017].
- 913 Wang, H.-Y., Rutherford, E.S., Cook, H.A., Einhouse, D.W., Haas, R.C., Johnson, T.B.,
914 Kenyon, R., Locke, B., and Turner, M.W. 2007. Movement of walleyes in Lakes Erie and
915 St. Clair inferred from tag return and fisheries data. *Trans. Am. Fish. Soc.* **136**: 539-551.
- 916 Wang, H.-Y., Cook, H.A., Einhouse, D.W., Fielder, D.G., Kayle, K.A., Rudstam, L.G., and
917 Höök, T.O. 2009. Maturation schedules of walleye populations in the Great Lakes region:
918 comparison of maturation indices and evaluation of sampling-induced biases. *N. Am. J.*
919 *Fish. Manage.* **29**: 1540-1554.
- 920 Wang, L., Riseng, C.M., Mason, L.A., Wehrly, K.E., Rutherford, E.S., McKenna, J.E., Jr,
921 Castiglione, C., Johnson, L.B., Infante, D.M., Sowa, S., Robertson, M., Schaeffer, J.,
922 Khoury, M., Gaiot, J., Hollenhorst, T., Brooks, C., and Coscarelli, M. 2015. A spatial
923 classification and database for management, research, and policy making: The Great Lakes
924 aquatic habitat framework. *J. Gt. Lakes Res.* **41**: 584-596.
- 925 Warner, D.M., O'Brien, T.P., Farha, S., Schaeffer, J.S., and Lenart, S. 2013. Status and trends
926 of pelagic prey fishes in Lake Huron, 2012. U.S. Geological Survey, Great Lakes Science
927 Center, Ann Arbor, Mich. Available from:
928 http://www.glsc.usgs.gov/sites/default/files/product_files/Pelagic%20Preyfish%20Huron%202012.pdf.
929 [accessed 1 June 2017].
- 930 Werner, E.E. 1974. The fish size, prey size, handling time relation in several sunfishes and
931 some implications. *J. Fish. Res. Board Can.* **31**: 1531-1536.
- 932 Winberg, G.G. 1956. Rate of metabolism and food requirements of fishes. *Fish. Res. Board*
933 *Can. Transl. Ser. No.* 194.

Table 1. Diet schedule for adult walleye from Lake Erie.

Month	Shiners	Clupeids	<i>Morone</i> spp.	Rainbow smelt	Yellow perch	Round goby	Other fish	Invertebrates
April - May	0.39	0.19	0.02	0.37	0.00	0.02	0.01	0.00
June	0.65	0.04	0.04	0.21	0.01	0.03	0.00	0.02
July	0.47	0.00	0.00	0.34	0.08	0.08	0.00	0.03
August	0.25	0.23	0.10	0.14	0.09	0.19	0.00	0.00
September	0.21	0.60	0.12	0.01	0.04	0.01	0.01	0.00
October - November	0.19	0.74	0.01	0.06	0.00	0.00	0.00	0.00
December - March	0.29	0.47	0.01	0.22	0.00	0.01	0.00	0.00

Note: Numeric table entries represent estimated proportions in diet, on a wet weight basis.

Table 2. Energy densities for prey of adult walleye from Lake Erie.

Prey category	Energy density (J/g)	Source
Shiners	6372	This study
Clupeids	5014	Kershner (1996)
<i>Morone</i> spp.	3760	Kershner (1996)
Rainbow smelt	5241	This study
Yellow perch	3823	Kershner (1996)
Round goby	3621	Bunnell et al. (2005)
Other fish	4512	Pothoven et al. (2017)
Invertebrates	3134	Cummins and Wuycheck (1971)

Note: Numeric table entries represent energy densities, on a wet weight basis.

Table 3. Parameter estimates from fitting von Bertalanffy growth curves to total length and age data for walleye caught in Saginaw Bay of Lake Huron during September and walleye caught in western and central Lake Erie during October, years 2010-2012.

Lake and sex	L_{∞} (mm)	K (year ⁻¹)	t_0 (years)
Lake Huron females	629 (14)	0.226 (0.015)	-0.596 (0.125)
Lake Huron males	509 (5)	0.361 (0.017)	-0.079 (0.090)
Lake Erie females	661 (1)	0.374 (0.004)	-0.113 (0.020)
Lake Erie males	558 (1)	0.468 (0.006)	-0.193 (0.025)

Note: L_{∞} = asymptotic average total length, K = Brody growth coefficient, and t_0 = age at which average total length is equal to zero. Standard error enclosed within parentheses.

Figure captions

Fig. 1. Map of study area. Solid circles denote locations for acoustic receivers and solid triangles denote tagging locations. Base map data taken from the Great Lakes Information Network (GLIN) (2017).

Fig. 2. Temperature regimes experienced by walleye in the northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie, by sex. Each bar represents one standard error. Error bar width may be less than the diameter of the symbol marker used to denote mean temperature, and therefore not visible. See Temperature regimes subsection of Materials and methods section for more details.

Fig. 3. Growth trajectories for walleye in Lake Huron and Lake Erie, by sex. Estimated weight at age based on gillnet surveys in Saginaw Bay (Lake Huron) during September 2010-2012 and gillnet surveys in Lake Erie during October 2010-2012. Each bar represents one standard error. Error bar width may be less than the diameter of the symbol marker used to denote mean weight at age, and therefore not visible. See Growth trajectories subsection of Materials and methods section for more details.

Fig. 4. Cumulative food consumption by walleye, beginning at age 3, in the northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie, by sex. Food consumption estimated by application of a walleye bioenergetics model (Kitchell et al. 1977;

Hewett and Johnson 1992) to observed growth trajectories. See Bioenergetics modeling: fit to observed growth subsection of Materials and methods section for more details.

Fig. 5. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in the northern main basin of Lake Huron with the temperature regime for southern main basin of Lake Huron walleye (upper panel), Saginaw Bay walleye (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

Fig. 6. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in the southern main basin of Lake Huron with the temperature regime for northern main basin of Lake Huron walleye (upper panel), Saginaw Bay walleye (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

Fig. 7. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in Saginaw Bay with the temperature regime for walleye from the northern main basin of Lake Huron (upper panel), walleye from the southern main basin of Lake Huron (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

Fig. 8. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in Lake Erie with the temperature regime for walleye from the northern main basin of Lake Huron (upper panel), walleye from the southern main basin of Lake Huron (middle panel), and Saginaw Bay walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

Fig. 9. Predicted changes in walleye weight at age by exchanging temperature regimes experienced by the walleye between the sexes for each of the four geographic areas: northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

Fig. 1. Map of study area. Solid circles denote locations for acoustic receivers and solid triangles denote tagging locations.

196x260mm (300 x 300 DPI)

Fig. 2. Temperature regimes experienced by walleye in the northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie, by sex. Each bar represents one standard error. Error bar width may be less than the diameter of the symbol marker used to denote mean temperature, and therefore not visible. See Temperature regimes subsection of Materials and methods section for more details.

390x300mm (300 x 300 DPI)

Fig. 3. Growth trajectories for walleye in Lake Huron and Lake Erie, by sex. Estimated weight at age based on gillnet surveys in Saginaw Bay (Lake Huron) during September 2010-2012 and gillnet surveys in Lake Erie during October 2010-2012. Each bar represents one standard error. Error bar width may be less than the diameter of the symbol marker used to denote mean weight at age, and therefore not visible. See Growth trajectories subsection of Materials and methods section for more details.

390x300mm (300 x 300 DPI)

Fig. 4. Cumulative food consumption by walleye, beginning at age 3, in the northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie, by sex. Food consumption estimated by application of a walleye bioenergetics model (Kitchell et al. 1977; Hewett and Johnson 1992) to observed growth trajectories. See Bioenergetics modeling: fit to observed growth subsection of Materials and methods section for more details.

390x300mm (300 x 300 DPI)

Fig. 5. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in the northern main basin of Lake Huron with the temperature regime for southern main basin of Lake Huron walleye (upper panel), Saginaw Bay walleye (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

285x369mm (300 x 300 DPI)

Fig. 6. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in the southern main basin of Lake Huron with the temperature regime for northern main basin of Lake Huron walleye (upper panel), Saginaw Bay walleye (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

285x369mm (300 x 300 DPI)

Fig. 7. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in Saginaw Bay with the temperature regime for walleye from the northern main basin of Lake Huron (upper panel), walleye from the southern main basin of Lake Huron (middle panel), and Lake Erie walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

285x369mm (300 x 300 DPI)

Fig. 8. Predicted changes in walleye weight at age by replacing the temperature regime experienced by walleye in Lake Erie with the temperature regime for walleye from the northern main basin of Lake Huron (upper panel), walleye from the southern main basin of Lake Huron (middle panel), and Saginaw Bay walleye (lower panel), by sex. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

285x369mm (300 x 300 DPI)

Fig. 9. Predicted changes in walleye weight at age by exchanging temperature regimes experienced by the walleye between the sexes for each of the four geographic areas: northern main basin of Lake Huron, southern main basin of Lake Huron, Saginaw Bay, and Lake Erie. See Bioenergetics modeling: exchanging temperature regimes between areas and sexes subsection of the Materials and methods section for more details.

390x300mm (300 x 300 DPI)