Canadian Journal of Forest Research
 Revue canadienne de recherche forestière

Estimating stand-level economic impacts of black bear damage to intensively managed forests

Journal:	<i>Canadian Journal of Forest Research</i>
Manuscript ID	cjfr-2017-0314.R1
Manuscript Type:	Article
Date Submitted by the Author:	10-Feb-2018
Complete List of Authors:	Kline, Kristina; Oregon State University, Forest Ecosystems and Society Taylor, Jimmy; USDA, National Wildlife Research Center, Oregon Field Station Morzillo, Anita; University of Connecticut, Natural Resources and the Environment
Keyword:	black bear, Douglas-fir, economic impacts, peeling, timber damage
Is the invited manuscript for consideration in a Special Issue? :	N/A

SCHOLARONE™
 Manuscripts

Estimating stand-level economic impacts of black bear damage to intensively managed forests

Kristina N. Kline^{a,1,*}, Jimmy D. Taylor^b, Anita T. Morzillo^c

^aDepartment of Forest Ecosystems and Society, Oregon State University, 321 Richardson Hall
Corvallis, Oregon 97331, U.S.A. knkline11@gmail.com. ph. 720-308-1082

^bUSDA APHIS, Wildlife Services, National Wildlife Research Center, 321 Richardson Hall
Corvallis, Oregon 97331, U.S.A. jimmy.d.taylor@aphis.usda.gov

^cDepartment of Natural Resources and the Environment, University of Connecticut, 1376 Storrs
Rd. Storrs, Connecticut 06269, U.S.A. anita.morzillo@uconn.edu

¹Present address: 361 S. Camino del Rio #198, Durango, CO 81303
*Corresponding author

Abstract

Black bears (*Ursus americanus*) peel conifers in early spring to forage on energy-rich vascular tissues resulting in damage to timber stands. The objective of our study was to develop and demonstrate a conceptual framework and methods for estimating stand-level volume and economic losses from black bear damage. We created tree lists from surveys of healthy and bear-damaged trees in timber stands of western Washington and Oregon. The forest growth model, Forest Vegetation Simulator (FVS) was used to project stand volume under two damage scenarios and an undamaged scenario. One damage scenario (Salvage) accounted for mortality and volume losses of fully and partially girdled trees; a second scenario (Total Loss) assumed complete loss of all trees peeled by black bears, regardless of peeling severity. The Fuel Reduction Cost Simulator (FRCS) was applied to estimate the value of logs delivered to the mill after accounting for logging and hauling costs associated with harvest. Present value of stands was calculated to translate volume losses into economic losses associated with bear damage. Economic losses ranged from 4-16% (Salvage) and 17-46% (Total Loss) of net present value. Our approach can be adapted for other forest settings and for forest management plans that assess wildlife damage.

Keywords: black bear; Douglas-fir; economic impacts; peeling; timber damage

64 **Introduction**

65 American black bears (*Ursus americanus*) damage trees after canopy closure in
66 intensively managed forests of the Pacific Northwest, peeling bark from conifers in early spring
67 to forage on phloem and cambial tissues (hereafter vascular tissues). These tissues provide
68 energy-rich soluble sugars for black bears at a time of year when similarly attractive energy
69 sources such as salmonberry (*Rubus spectabilis*), red huckleberry (*Vaccinium parvifolium*), and
70 blackberry (*Rubus ursinus*) are scarce. The presence of up to 3.5% soluble sugars in the phloem
71 of trees at this time of year provides fructose, sucrose and glucose, which help black bears meet
72 their energy needs following winter dormancy (Kimball et al. 1998a, Kimball et al. 1998b,
73 Radwan 1969, Ziegler 2004). Although first reported in the early 1900's (Pierson 1966), it
74 was not until the increase in intensive forest management in the 1940's that bark peeling by
75 black bears was identified as a problem for timber production in the Pacific Northwest (Pierson
76 1966).

77 In western Oregon and Washington, black bears typically damage Douglas-fir
78 (*Pseudotsuga menziesii*) trees in stands that are intensively managed for timber production
79 (Schmidt and Gourley 1992). Management activities such as thinning and fertilization increase
80 tree growth and volume, which in turn increases sugar concentrations in vascular tissues
81 (Kimball et al. 1998b). This makes trees more attractive as forage to black bears. Peeling occurs
82 at varying severities, which result in a variety of damage impacts. A tree will eventually die if
83 fully girdled (i.e., peeled around the entire circumference of the trunk). A tree that is partially
84 girdled becomes more susceptible to insect infestations, fungal decay (Kanaskie et al. 1990), and
85 windfall (Witmer et al. 2000), thereby reducing the likelihood of surviving to harvest age. Miller

et al. (2007) evaluated survival of trees damaged by black bears in Washington, and reported mortality rates for partially girdled trees at 17% over a 16-year period. An ongoing, long-term study of simulated bole damage in Capitol Forest, Washington, reported mortality rates of between 5-28% for partially girdled trees, with mortality rates increasing with the percentage of the bole circumference damaged (Connie Harrington, United States Forest Service USFS, Unpublished data). Partially girdled trees that survive to harvest age also may experience a reduction in the volume of merchantable timber produced (Lowell et al. 2010, Pierson 1966).

To our knowledge, only two studies have quantified timber volume losses due to black bear damage. In a survey of black bear-damaged trees sent to mill in Washington, Lowell et al. (2010) found a 6.4% loss in volume, on average, for trees partially girdled as a result of black bear damage. Additionally, Lowell et al. (2010) found the value of such partially girdled trees was 5% less than undamaged trees. Elsewhere, Pierson (1966) reported average volume losses from a survey of 100 Douglas-fir trees with varying damage severities. Trees with bark removed from less than 50% of the trunk circumference lost an average of 7% merchantable volume, while trees with bark removed from more than 50% of the trunk circumference lost an average of 10% merchantable volume (Pierson 1966). While informative, these studies are limited to a focus at the scale of individual trees.

Our objective was to further our understanding of black bear damage to timber stands by developing and demonstrating a conceptual framework and methods for estimating stand-level volume losses and economic losses from black bear damage. Previous studies on economic impacts of black bear damage to trees focused at the regional level and used aerial counts of conifers with red crowns as an index of bear damage (Nolte and Dykzeul 2002, Taylor et al.

2014). Aerial estimates may overestimate bear damage because other sources, such as root rot, cause red crowns (Kanaskie et al, 1990). They also may underestimate black bear damage because partial peeling affects wood quality but does not cause crowns to turn red (Kanaskie et al, 1990). Therefore, our study builds upon previous research by addressing volume losses to black bear damage at the stand level in three ways: 1) including the impacts of volume lost from partially girdled trees, 2) accounting for volume lost in relation to the severity of individual tree damage, and 3) accounting for stand growth and yield projected to harvest.

Methods

Study sites

Study sites consisted of four intensively managed Douglas-fir stands on private land within the Coast Range and the western Cascades of Oregon and Washington (Figure 1). The western Cascades ecoregion consists of a mild maritime climate with cool, wet winters, and hot, dry summers (Immell et al. 2013). Average annual rainfall ranges from 107-226 cm and average annual snowfall ranges from 18-592 cm occurring above 1,220 meters (Oregon Dept. of Fish and Wildlife 2006). Stands were dominated by Douglas-fir, with co-dominants including western hemlock (*Tsuga heterophylla*), western redcedar (*Thuja plicata*), and red alder (*Alnus rubra*). Understory vegetation was comprised of salmonberry (*Rubus spectabilis*), vine maple (*Acer circinatum*), swordfern (*Polystichum munitum*), foxglove (*Digitalis purpurea*), Oxalis (*Oxalis oregana*), Oregon grape (*Mahonia nervosa*), snowberry (*Symphoricarpos albus*), devil's club (*Oplopanax horridus*), stinging nettle (*Urtica dioica*), Trillium (*Trillium ovatum*), lady fern (*Athyrium filix-femina*), pacific bleeding heart (*Dicentra formosa*), and miner's lettuce (*Claytonia perfoliata*).

The Coast Range ecoregion consists of a maritime climate, with mild, wet winters and warm, dry summers (Cushman and McGarigal 2003). Average annual precipitation ranges from 152-249 cm (Oregon Dept. of Fish and Wildlife 2006). Stands are dominated by Douglas-fir, with co-dominants including western hemlock, western redcedar, and red alder. Understory vegetation was comprised of salmonberry, salal (*Gaultheria shallon*), vine maple, swordfern, foxglove, Oxalis, Oregon grape and snowberry.

Stands were even-aged plantations and managed for timber production using intensive silvicultural practices. Characteristics varied by stand and each was considered by managing foresters to contain high levels of black bear damage. The WA Coast Range site (CR-WA) was located in Pacific County, approximately one kilometer south of the Lower Chehalis State Forest. The western Cascades site in WA (WC-WA) was located in Lewis County, approximately one kilometer north of the Gifford Pinchot National Forest. These stands had not been pre-commercially thinned.

The OR Coast Range site (CR-OR) was located in Lincoln County, approximately one kilometer east of the Siuslaw National Forest. This stand was pre-commercially thinned at age 13 to a density of 733 trees per hectare. The western Cascades site in OR (WC-OR) was located in Lane County, approximately one kilometer north of Oregon Department of Forestry (ODF) land. This stand was pre-commercially thinned at age 14 to a density of 650 trees per hectare.

Sampling design

We collected data in June 2015, and sampled 10% of each stand. We used fixed 0.04 ha circular plots to survey each stand. With fixed circular plots, we were able to observe the full

circumference of each tree, ensuring all damage was captured regardless of where damage occurred on the tree.

We evenly spaced plots at 50 meters throughout the stand in a grid-like arrangement (Figure 2). Within each plot, we measured diameter at breast height (dbh) of every tree greater than 10 cm dbh. We measured height and height-to-live-crown-base (HCB) of every tenth tree as well as every black bear-damaged tree. For trees damaged by black bear, we noted the condition of the crown (red, yellow, green, or no needles), and measured the percentage of the circumference peeled.

Imputation of tree-level attributes

Imputation of values for certain tree-level attributes, such as height, is a necessary component of forest inventory (Garber et al. 2009). Measuring heights of all individual trees is a time consuming task (Wang and Hann 1988) and imputation greatly increases the efficiency at which a stand can be sampled. Additionally, forest modeling requires height estimates for every tree in order to calculate volume. Height imputations involve the use of both dbh and height variables, as a strong relationship exists between the two (Curtis 1967). For each stand, we used the following height-diameter equation, which is commonly used in the Pacific Northwest, to relate tree height to dbh and impute missing heights (see Curtis 1967 for details):

$$[1] \quad H = 1.37 + \beta_1 * \exp(\beta_2/dbh) + \varepsilon_1$$

where H is the height of the tree in meters, 1.37 is the height in meters above ground at which dbh is measured, $\exp$ is the base of the natural logarithm, dbh is the diameter at breast height of the tree in centimeters, β_1 and β_2 are parameters to be estimated from the data, and ε_1 is the

residual error with $\varepsilon_1 \sim N(0, \sigma_1^2)$. Derived from equations fit with our field data, β_1 was the upper asymptote for predicted heights, and β_2 determined the shape of the approach to this upper asymptote. We then log transformed this equation in order to obtain initial parameter estimates through simple linear regression. Due to differences in tree density and stand structure, we fit individual equations for each stand. We completed all statistical analyses using SAS software, Version 9.2 (2011 SAS Institute, Cary NC).

To impute missing height-to-live-crown-base (HCB) measurements we used the following equation:

$$[2] \quad \text{HCB} = \beta_1 * e^{(\beta_2 / \text{dbh})}$$

where HCB is the height from the base of the trunk to the base of live crown measured in meters, e is the base of the natural logarithm and dbh is the diameter at breast height of the tree in centimeters. Data used to fit the models were limited to living trees only, as dead trees had no HCB values. For all stands, the β_1 and β_2 parameters for damaged and undamaged trees were the same.

Estimating stand volume and economic value

For analysis of bear damage impacts we used a combination of existing models to estimate forest growth and yield, harvest costs, and present value (Figure 3). We first transformed tree-level data (Figure 3a) into tree lists to construct stand-level projections (Figure 3b). Tree lists consisted of the following tree-level variables: tree number, species, dbh, height, crown ratio, and expansion factor. We calculated crown ratio by subtracting the HCB from the full height of the tree to obtain the crown height, and then dividing the crown height by the full height of the tree. We calculated expansion factor (the number of trees per hectare a given plot

tree represents) by taking the denominator of the plot size (same for all stands), which was 0.4 ha or 1/10 acre, and dividing it by the number of plots sampled in that stand. For example in our stand with 37 plots, the expansion factor was 2.07 (10/37). Following that, we input tree lists for each stand into the Forest Vegetation Simulator (FVS) growth and yield model (Figure 3c; Dixon 2002) within the Landscape Management System (LMS) interface, Version 2.1 (Nelson et al. 1999). FVS is a distance-independent, individual tree growth and yield model (Dixon 2002) capable of simulating a wide range of silvicultural treatments for most tree species, forest types, and stand conditions (Teck et al. 1997, Crookston and Havis 2002, Crookston and Dixon 2005, U.S. Dept. of Agriculture 2016). While Ceder and Marzluff (2002) used FVS with the LMS interface to evaluate wildlife habitat, our use of FVS within LMS was consistent with traditional harvest planning (Crookston and Dixon 2005). We used the Pacific Northwest Coast (PN) variant of FVS for Coast Range sites, and the West Cascades (WC) variant for western Cascades sites. Both are accepted methods in the Pacific Northwest region, and applicable to the Douglas-fir stands of focus in this study (Dixon 2002).

Use of alternative scenarios in ecological and economic modeling is useful for exploring the range in plausible outcomes in the face of uncertainty (Francis and Hamm 2011). In our study, we developed two scenarios to explore estimated loss in timber volume due to black bear damage in each stand. In this first scenario, we assumed a percentage of partially girdled trees would die, conditional on the severity of damage. For individual trees that survived, we assumed a percentage of volume would be lost (7-10%; Pierson 1966), also conditional on the severity of damage. Under scenario one, damaged trees with standing live volume were harvested and processed, thus we refer to this as the “Salvage” scenario. We acknowledge that the term

salvage in forestry commonly refers to a logging operation following a larger natural disturbance (e.g., fire, ice damage, windstorm). In this case, we apply the term in a similar fashion to account for removal and processing of all merchantable value from black bear-damaged trees. We based the second scenario on personal communications with some of our cooperating landowners who felt that bear-damaged trees provided no economic return. In this second scenario, any black bear damage resulted in a complete loss of the damaged tree's volume, regardless of whether the tree was partially girdled or killed. We assumed that the costs required to salvage a black bear-damaged tree at harvest offset any profit that would be made from the tree, resulting in zero monetary value gained from harvest. Thus, we referred to this as the "Total Loss" scenario.

Salvage scenario

To account for tree mortality and individual tree volume loss due to bear peeling in the growth and yield model under this scenario, we applied Connie Harrington's estimates (USFS, Unpublished data) of percent mortality and Pierson's (1966) estimates of volume lost at the tree level to our field data. Harrington's estimates (USFS, Unpublished data) were derived from five different categories of bole circumference damage: 20, 40, 60, 80 and 90% of circumference. Our categories of damage severity from field data ranged from 10-100% in 5% increments, thus we re-categorized them to match those used by Harrington. For trees in the 20, 40, and 60% damage categories, Harrington's cumulative mortality rates leveled off around 6% after six years. For our data, we assumed that mortality rates for these three categories (20, 40, and 60) would remain at 6% in the future. We designated this as a single damage category, further referred to as the low-damage category. This category contained all trees with observed damage $\leq 60\%$ of the circumference. For trees in the 80 and 90% categories, Harrington (USFS,

Unpublished data) showed that mortality rates increased linearly over time. For our data, we assumed that mortality rates in these categories would continue on these linear trajectories into the future. With this assumption, we fit a linear model for each category in order to estimate mortality rates past eight years. This was necessary as stands were to be projected in the growth model 15-45 years into the future. The equation fit for the 90% category was:

$$[7] \quad m = -17.9750 + 3.6083 * yr$$

The equation fit for the 80% category was:

$$[8] \quad m = -8.4028 + 1.4583 * yr$$

where m = % mortality at harvest age and yr = the number of years since damage occurred. The yr value was obtained by adding the number of years since damage occurred on average in each stand at present to the number of years each stand was to be projected in the growth model. We assigned all trees with observed damage between 60-80% circumference peeled to the 80% category. We assigned all trees with observed damage between 80-99% circumference peeled to the 90% category.

To estimate each stand's volume, we implemented two thinning treatments in the growth model. We used thinning treatments as a surrogate to simulate black bear peeling, as the model does not contain direct inputs for wildlife damage. In the first thinning, we removed all observed black bear-killed trees before growing the stand to harvest age. After we grew each stand to harvest age in the model, we implemented an additional thinning. To implement the second thinning, we calculated the proportion of partially girdled trees in each stand that fell into each of the three damage categories (low-damage, 80%, and 90%). We then removed the percentage of

trees in each damage category that would have died over time from black bear damage. We based these removal percentages on mortality rates derived from the linear models we fit for each stand (Table 1). We identified removal trees by species (Douglas-fir) and dbh (trees with the mean dbh of black bear-damaged trees).

To compute losses in volume of surviving partially girdled trees, we calculated the standing volume of all trees remaining at harvest with $\leq 50\%$ of their circumference damaged, and then multiplied it by 7% to obtain the first volume reduction value. Then, we calculated the standing volume of all trees remaining at harvest with $> 50\%$ of their circumference damaged, and multiplied the volume of damaged trees by 10% to obtain the second volume reduction value. We added these two volume reductions together and then subtracted from the total stand volume at harvest to obtain a recoverable stand volume after accounting for black bear damage.

Total Loss scenario

To simulate this scenario in the growth model, we implemented two thinning treatments in each stand. In the first thinning, we removed all observed black bear-killed trees. We then projected each stand to harvest age, and implemented a second thinning. In this second thinning, we removed all remaining partially girdled trees from each stand. We selected removal trees by species (Douglas-fir) and by dbh (trees with the mean dbh of black bear damaged trees). We removed partially girdled trees after each stand was projected to harvest because they are usually left to grow until harvest. In this scenario, they become a complete loss at harvest because the value of recoverable volume is assumed equal to harvesting costs.

Undamaged scenario

We developed an “Undamaged” scenario for each stand to serve as a control for comparison of the two damaged scenarios. To simulate undamaged stands, we treated black bear-killed trees as undamaged living trees. We originally assigned black bear-killed trees crown ratio values of zero. To include them as living trees, we imputed their crown ratios using the HCB equations described above specific to each stand. We then projected stands to harvest age in the growth model and calculated volume of surviving trees at harvest.

Present stand value

We used present value of each stand to translate volume losses into economic losses. Present value estimations require knowledge of volume at harvest (Figure 3d; as obtained from FVS outputs, i.e., Figure 3c) and the value of logs delivered to the mill (pond value). These estimations also require knowledge of the logging and hauling costs that are subtracted from the value of logs delivered to the mill. To estimate logging and hauling costs associated with each stand at harvest, we input volume at harvest values for each stand under all scenarios into the Fuel Reduction Cost Simulator (FRCS; Figure 3e) (Fight et al. 2006), specifically the FRCS-West variant. Data inputs included stand slope, average yarding distance from the stand to a roadside landing, stand area, elevation, harvesting system used, number of large trees/ha, and mean volume/large tree. We derived large trees/ha and mean volume/large tree values from FVS output (Figure 3f) by dividing total volume/ha by trees/ha. We derived average yarding distance by measuring the distance from the GIS calculated centroid of each stand to the nearest road in GIS. Spatial road data were obtained from cooperating landowners. We derived slope values from digital elevation model layers in GIS using the Spatial Analyst Slope Tool.

We performed the FRCS simulation using the Special “Billion-Ton” Processing Rules. These rules designate a harvesting method based on each stand’s slope and volume/ha. If the slope was $\leq 40\%$ then two alternatives of a ground-based logging system were considered by the model: mechanical whole-tree harvesting with feller-bunchers and skidders used to transport bunches, or manual whole-tree harvesting with chainsaws and skidders used to transport whole trees (Dykstra 2010, Fight et al. 2006). FRCS completes calculations for both possible alternatives and selects the lower-cost alternative (Dykstra 2010). If the slope was $>40\%$ the simulator used manual felling and cable yarding as the harvesting system (Dykstra 2010). Based on stand slopes, we harvested CR-WA and WC-WA in the simulation using a ground-based mechanical system. We harvested CR-OR and WC-OR in the simulation using a system of manual felling with chainsaws and cable yarding. We simulated all stands as clear-cuts and included loading costs.

The following Land Expectation Value (LEV) Equation (Figure 3g) was applied to estimate the present value of each stand (Figure 3h) under all scenarios:

$$[9] \quad PV = (Vh * SP) / (1 + i)^y$$

where PV = the present value of the stand in dollars, Vh = the total volume of the stand at harvest age, SP = the stumpage price which is the pond value (i.e., log value) minus logging and hauling costs, i = the discount interest rate, and y = the number of years from present to harvest age (years projected). To determine an average log value per thousand board feet (MBF) we used output from the growth model to calculate a distribution of volumes by log grade at harvest in each stand. We then calculated a weighted mean log value per MBF based on this distribution for

each stand (Table 2). We distributed stand volume among the following six log grades: special mill, #2 sawmill, #3 sawmill, #4 sawmill, chip-and-saw, and pulp logs. We used the most current market value for each grade in the calculation of weighted mean price per Douglas-fir MBF. We used a discount interest rate of 5%, because the most common interest rates used in these calculations are 4-6% (Darius Adams, Oregon State University, Personal communication).

Results

Stand-level inputs differed by site (Table 3) and resulted in different levels of volume at harvest (Table 4). The WC-OR stand contained the highest levels of black bear damage with 42.4% of the stand damaged. The CR-WA stand contained 16.2% damage, WC-WA contained 13.5% damage, and CR-OR contained 8.5% damage. Volume losses in the Salvage and Total Loss scenarios ranged from 4-15% and 16-43%, respectively (Table 4). Volume losses in the Total Loss scenario were on average four times greater than volume losses in the Salvage scenario. Economic losses in the Salvage scenario ranged from \$472/ha to \$1,635/ha while economic losses in the Total Loss scenario ranged from \$2,416/ha to \$4,978/ha (Table 5).

Discussion

Economic impacts of black bear damage

Wildlife damage is an ongoing concern for intensively managed timber resources, and tools are needed to improve assessment to damage impacts. Our approach and estimates of economic loss to black bear damage advanced existing methods and estimates by including the additional impacts of partially girdled trees and accounting for loss based on the severity of individual tree damage. Our two damage scenarios reflected how different landowners might interpret the losses they incur from severe black bear damage on their lands. In our Salvage

scenario, damaged stands retained 84-96% of the value of undamaged stands. In our Total Loss scenario, losses were on average four times greater, and black bear damaged stands retained 54-83% of the value of undamaged stands (Table 5). These economic losses can be transformed into the perspective of average timber management costs (Table 6). In the Salvage scenario, losses from black bear damage equaled the costs for landowners to prepare and plant 5-22 ha of industrial timberland. In the Total Loss scenario, losses from black bear damage equaled the costs for landowners to prepare and plant 17-66 ha of industrial timberland. Losses under the Total Loss scenario also were equivalent to the costs required to pre-commercially thin 51-195 ha of industrial timberland. For small landowners, the profits from harvesting timber might be applied toward the management of other timber stands on their lands. If landowners are relying on harvest profits to fund ongoing intensive management of other stands among their lands, black bear damage may hinder the ability to complete concurrent intensive management of other stands. Moreover, if landowners have already expended funds on preventive management of black bears, then future losses may be even greater if timber stands continue to incur severe damage.

Implications for management

Black bears are valued highly in Oregon and Washington as game animals and as an important wildlife species by the general public (Oregon Dept. of Fish and Wildlife 2012). For those areas that continue to experience severe damage, additional management considerations are suggested. Damage may be mitigated by continued trapping efforts in these stands with emphasis on selective removal of damaging black bears. There is opportunity for private landowners to team up with local hunters by providing access and location information for black

bear damage on their lands. Supplemental feeding also has been used in select areas to reduce bear damage to conifers (Ziegltrum 2004). For areas that seem to experience higher levels of damage post-thinning or post-fertilization, it may be beneficial to delay thinning or fertilization until stands are past the most susceptible age for peeling (Barnes and Engeman 1995, Schmidt and Gourley 1992). Management costs could be quantified and contrasted with projected losses to bear damage. Our methodology can be incorporated into forest planning to help landowners assess whether the volume losses incurred from black bear damage can be offset by management techniques such as delayed thinning, depredation hunting, and supplemental feeding.

The four stands we sampled contained what is considered severe black bear damage (> 25 black bear-damaged trees/ha), yet covered a range of severities, from 8.5% to 42% damage. This understanding of economic losses associated with a range of severities will allow landowners to make economically favorable planning decisions for managing both black bears and timberlands to prevent severe black bear damage in the future and meet forestland management goals. On-the-ground monitoring of black bear damage frequency and severity across western Oregon and Washington at the stand level will provide an understanding of these changes over time as a result of black bear and forest management decisions. Although demonstrated using data from western Oregon and Washington, our approach and conceptual framework incorporates variables and methods that are applicable and transferrable to forested landscapes where black bear damage occurs in other locations, or for other wildlife species known to damage standing timber.

Opportunities for future research

Our growth models and economic models provide opportunities for future research on black bear damage impacts. First, our estimates represented a snapshot in time of black bear

damage observed in a single year of the timber rotation. We do not currently have the ability to accurately predict what levels of new damage, if any, will occur in these stands over the next 15-45 years. Additionally, in this case, we were primarily focused on understanding losses in these stands in their current state. Therefore, our models account only for black bear damage that has occurred in these stands between stand initiation (i.e., time of planting) and 2015. Repeated observations of the same plots over multiple years would improve future estimates of economic loss by black bears.

Another opportunity for future research is developing a tool within forest growth and yield models that simulates and captures complex tree growth nuances that are associated with black bear damage. Black bear damage in our models was treated as analogous to a thinning treatment. When removing black bear-killed trees, the model treated these trees as if killed that year. That treatment in turn affects the predicted growth of surviving trees in the model. In reality, however, our field data revealed that the majority of the black bear-killed trees had been dead or dying for multiple years. As a result, the remaining surviving trees likely had already responded to relinquished resources and growing space around them. Additionally, the model simulates thinning uniformly across the stand, whereas black bear damage imposes a relatively clustered pattern of thinning. The clustered pockets of dead trees resulting from black bear damage initiate a different response in the future growth of the stand than uniformly spaced mortality. With the development of a black bear-damage tool, modeling damage impacts over time would better reflect forest response to black bear damage, and perhaps could be applied to timber damage patterns resulting from other wildlife species.

A final opportunity for future research lies in understanding mortality rates and growth rates of damaged trees over a larger temporal scale. Miller et al.'s (2007) study of the growth of black bear-damaged trees found that partially girdled trees averaged 29-33% faster diameter growth than nearby undamaged trees. Additionally, Harrington's ongoing study (USFS, Unpublished data) of tree growth eight years following simulated bole damage showed an increase in diameter growth of partially girdled trees as well as initial decreases in height growth. We chose not to account for changes in height or diameter growth of damaged trees in this initial model because the eight years of reported values were not convincingly sufficient to project 15-45 years into the future. Instead, to focus the scope of our study, we decided that it was more imperative to account for tree mortality over time from black bear damage wounds. Understanding damaged tree mortality and growth rates for a full timber rotation will provide more accurate estimates of losses to black bear damage at a time scale that more precisely reflects forestland management.

Acknowledgements

We thank Doug Maguire for providing comments on earlier versions of this manuscript as well as guidance with forest growth and yield modeling and analysis. We thank Darius Adams for guidance with economic modeling and analysis; Ariel Muldoon for statistical support; Mike Tucker, Don Irons, and Jeff Light for site access; and Luke Kline, Blake Murden, and Rob Nagel for field assistance. We thank Georg Ziegltrum, Alan Kanaskie, Rob Flowers and Mark Gourley for their continued support and interest in this study. Funds were provided by Oregon Forest Industries Council and Washington Forest Protection Association.

References

- Askham, L.R. 1992. Voles. *In* Silvicultural approaches to animal damage management in Pacific Northwest forests. *Edited by* C. Black. USDA Forest Service Pacific Northwest Research Station, Portland, Oreg. pp 187-204.
- Barnes, V.G., and Engeman, R.M. 1995. Black bear damage to lodgepole pine in central Oregon. *Northwest Nat.* 76:127-129.
- Black, H.C., and Lawrence, W.H. 1992. Animal Damage Management in Pacific Northwest Forests 1901-1990. *In* Silvicultural approaches to animal damage management in Pacific Northwest forests. *Edited by* C. Black. USDA Forest Service Pacific Northwest Research Station, Portland, Oreg. pp 23-56.
- Black, H. C., Dimock II, E.J., Evans, J., and Rochelle, J.A. 1979. Animal damage to coniferous plantations in Oregon and Washington. Part I: a survey, 1963-1975. *Oreg. St. Univ. Sch. of Forestry Res. Bull.* 25.
- Ceder, K.R., and Marzluff, J.M., 2002. Linking tools of forest and wildlife managers: wildlife habitat evaluation using the landscape management system. *In* Proceedings of the 2nd Forest Vegetation Simulator Conference, Fort Collins, Colo., 12–14 February 2002. *Edited by* N.L. Crookston and R.N. Havis. USDA Forest Service, Rocky Mountain Research Station, Ogden, Uta. pp. 200–208.
- Crookston, N.L., and Dixon, G.E. 2005. The forest vegetation simulator: A review of its structure, content, and applications. *Comp. and Elect. in Agric.* 49:60-80.
- Crookston, N.L., and Havis, R.N. 2002. Second Forest Vegetation Simulator Conference. *In* Proceedings of the 2nd Forest Vegetation Simulator Conference, Fort Collins, Colo., 12–

- 14 February 2002. *Edited by* N.L. Crookston and R.N. Havis. USDA Forest Service, Rocky Mountain Research Station, Ogden, Uta. pp 1-208.
- Curtis, R.O. 1967. Height-diameter and height-diameter-age equations for second-growth Douglas-fir. *For. Sci.* 13:365-375.
- Cushman, S.A., and McGarigal, K. 2003. Landscape-level patterns of avian diversity in the Oregon Coast Range. *Ecol. Monogr.* 73:259-281.
- Dixon, G.E. 2002. Essential FVS: A users guide to the forest vegetation simulator. USDA Forest Service Intern. Rep.
- Dykstra, D. 2010. Batch processing in FRCS. USDA Forest Service Pacific Northwest Research Station User Guide for FRCS Fact sheet.
- Fight, R.D., Hartsough, B.R., and Noordijk, P. 2006. Users guide for FRCS: fuel reduction cost simulator software. USDA Forest Service Gen. Tech. Rep. PNW-GTR-668.
- Flowers, R., Norlander, D., and Kanaskie, A. 2014. Memorandum: 2014 aerial survey of tree mortality in northwest Oregon. *Oreg. Dep. of Forestry Rep.*
- Francis, S. R., and Hamm, J. 2011. Looking forward: using scenario modeling to support regional land use planning in Northern Yukon, Canada. *Ecol. Soc.* 16:18.
- Garber, S.M., Temesgen, H., Monleon, V.J., and Hann, D.W. 2009. Effects of height imputation strategies on stand volume estimation. *Can. J. For. Res.* 39:681-690.
- Immell, D., Jackson, D.H., and Boulay, M.C. 2013. Denning ecology of American black bears in the Cascade Mountains of Oregon. *Ursus* 24:1-12.
- Kanaskie, A., Irwin, G., Chetock, J., and Overhulser D. 1990. Black bear damage to forest trees in northwest Oregon 1988-1989. *Oreg. Dep. Forest. Pest Manag. Rep.*

- Kimball, B.A., Nolte, D.L., Engeman, R.M., Johnston, J.J., and Stermitz, F.R. 1998a. Chemically mediated foraging preference of black bear (*Ursus americanus*). *J. Mammal.* 79:448-456.
- Kimball, B.A., Turnblom, E.C., Nolte, D.L., Griffin, D.L., and Engeman, R.M. 1998b. Effects of thinning and nitrogen fertilization on sugars and terpenes in Douglas-fir vascular tissues: implications for black bear foraging. *For. Sci.* 44:599-602.
- Lowell, E.C., Dykstra, D., and McFadden, G. 2010. Effects of bear damage on Douglas-fir lumber recovery. *West. J. Appl. Forest.* 25:73-80.
- Miller, R.E., Anderson, H.W., Reukema, D.L., and Max, T.A. 2007. Growth of bear-damaged trees in a mixed plantation of Douglas-fir and red alder. *USDA For. Serv. Pacific Northwest Res. Stn. Res. Paper PNW-RP-571.*
- Nelson, C.E., McCarter, J.B., Wilson, J., Baker, P.J., Hall, J.S., and Oliver, C.D. 1999. *Landscape Management System User's Manual.* Univ. of Wash. Coll. of For. Resour. Intern. Rep.
- Nolte, D.L., and Dykzeul, M. 2002. Wildlife impacts on forest resources. *In* Proceedings of the 3rd National Wildlife Research Center Special Symposium: Human Conflicts with Wildlife: Economic Considerations, Fort Collins, Colo. 1-3 August 2000. *Edited by* L. Clark, J. Hone, J.A. Shivik, R.A. Watkins, K.C. Vercauteren, and J.K. Yoder. U.S. Department of Agriculture, Wildlife Services, National Wildlife Research Center, Fort Collins, Colo. pp.163-168.
- Oregon Department of Fish and Wildlife. 2012. Oregon black bear management plan. *Oreg. Dep. Fish and Wildl. Rep.*

- Oregon Department of Fish and Wildlife. 2006. Oregon conservation strategy. Oreg. Dep. Fish and Wildl. Rep.
- Pierson, D.J. 1966. Washington's black bear problem. West. Assoc. Game and Fish Commiss. 46:96-100.
- R Development Core Team. 2010. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Radwan, M.A. 1969. Chemical composition of vascular tissue of four tree species in relation to feeding by black bear. For. Sci. 15:11-16.
- SAS Institute Inc. 2011. Base SAS® 9.2 Procedures Guide. Cary, NC: SAS Institute Inc.
- Schmidt, W., and Gourley, M. 1992. Black bear. *In* Silvicultural approaches to animal damage management in Pacific Northwest forests. *Edited by* H. Black. USDA Forest Service Pacific Northwest Research Station, Portland, Oreg. pp 309-311.
- Taylor, J., Morzillo, A., and Anderson, A. 2014. Estimating the total economic impact of black bear peeling in western Oregon using GIS and REMI. *In* Proceedings of the 26th Vertebrate Pest Conference, Waikoloa Village, Hawaii. 3-6 March 2014. *Edited by* R. M. Timm and J. M. O'Brien. University of Calif., Davis. pp 419-422.
- Teck, R., Moeur, M., and Adams, J. 1997. Gen. Tech. Re INT-GTR-373. *In* Proceedings of the Forest Vegetation Simulator Conference, Fort Collins, Colo. 3-7 February 1997. *Edited by* R. Teck and M. Moeur. USDA Forest Service, Intermountain Research Station, Ogden, Uta. pp 1-222.

- U.S. Department of Agriculture, Forest Service. 2016. "What is FVS?". US Forest Service Forest Management Service Center. Available from <https://www.fs.fed.us/fmfc/fvs/whatis/index.shtml> [last accessed 5 July 2017].
- Wang, C., and Hann, D.W. 1988. Height-diameter equations for sixteen tree species in the central western Willamette Valley of Oregon. Oreg. St. Univ. Coll. of Forest. For. Res. Lab. Paper 51.
- Witmer, G., Nolte, D., and Stewart, W. 2000. Integrated pest management of black bear reforestation damage. USDA National Wildlife Research Center, Staff Publications Paper 824.
- Ziegler, G.J. 2004. Efficacy of black bear supplemental feeding to reduce conifer damage in western Washington. J. Wildl. Manage. 68:470-4.

Tables

Table 1. Percent mortality values used for each stand at harvest age, as derived from the percent mortality linear models fit from Connie Harrington’s data (USFS, Unpublished data).

Stand	Yrs Projected	Perc. Mort. for 80%*	Perc. Mort. for 90%†	Perc. Mort. for Low Damage‡
CR-WA	45	65	100	6
CR-OR	15	21	54	6
WC-OR	40	54	100	6
WC-WA	30	40	100	6

* Percent of partially girdled trees with between 60 and 80% of their circumference peeled that would die in each stand after being projected to harvest age. Value derived from the following linear regression equation: $m = -8.4028 + 1.4583 \cdot yr$ where $m = \% \text{ mortality}$ at harvest age and $yr =$ the number of years since damage occurred. $R^2 = 0.9483$.

† Percent of partially girdled trees with between 80 and 99% of their circumference peeled that would die in each stand after being projected to harvest age. Value derived from the following linear regression equation: $m = -17.9750 + 3.6083 \cdot yr$. $R^2 = 0.9771$.

‡ Percent of partially girdled trees with less than 60% of their circumference peeled that would die in each stand after being projected to harvest age.

Table 2. Distribution of volume by different log grades at harvest, and weighted mean log values per Douglas-fir MBF used in present value calculations.

Stand	Log Grade	Log Value/MBF	% Total Volume	Weighted Mean Value/MBF
CR-WA	chip-and-saw	\$187	2	\$599
	special mill	\$700	30	
	pulp logs	\$107	1	
	#2 sawmill	\$605	35	
	#3 sawmill	\$550	24	
	#4 sawmill	\$525	8	
CR-OR	chip-and-saw	\$187	4	\$533
	pulp logs	\$107	2	
	#2 sawmill	\$605	26	
	#3 sawmill	\$550	48	
	#4 sawmill	\$525	20	
WC-OR	chip-and-saw	\$187	5	\$517
	pulp logs	\$107	2	
	#2 sawmill	\$605	1	
	#3 sawmill	\$550	69	
	#4 sawmill	\$525	23	
WC-WA	chip-and-saw	\$187	6	\$517
	pulp logs	\$107	2	
	#2 sawmill	\$605	6	
	#3 sawmill	\$550	63	
	#4 sawmill	\$525	24	

Table 3. Stand level-details for each study site in western Oregon and Washington, 2015.

Stand	Age	Size (ha)	Site Index	DBH (SE) [#]	Ht (SE) [^]	Elev (m)	TPH [*]	Plots [†]
CR-WA	17	15	132	18.3 (0.041)	11.5 (0.155)	170-280	936	37
CR-OR	29	8	131	27.4 (0.124)	22.1 (0.475)	380-520	642	20
WC-OR	15	6	115	17.3 (0.052)	10.7 (0.269)	380-1750	724	15
WC-WA	33	13	111	24.1 (0.099)	15.7 (0.452)	790-920	684	32

[#]Mean DBH in cm

[^]Mean height in meters

^{*}Stand density in trees/ha

[†]Number of 0.04 ha sample plots surveyed in each stand

Table 4. Volume at harvest and present stand values for each of four sample stands under two bear damage scenarios and an undamaged scenario.

Stand	Harvest Age	<i>Undamaged</i>		<i>Salvage Scenario</i>		<i>Total Loss Scenario</i>	
		Total Vol at Harvest (m ³)	Present Value of Stand	Total Vol at Harvest (m ³)	Present Value of Stand	Total Vol at Harvest (m ³)	Present Value of Stand
CR-WA	62	23536	\$250,072	21309	\$225,542	16720	\$175,407
CR-OR	44	5307	\$181,854	5091	\$174,533	4440	\$150,049
WC-OR	55	4335	\$42,659	3686	\$35,812	2486	\$23,131
WC-WA	63	7117	\$124,280	6794	\$118,142	5375	\$92,867

Table 5. Economic losses to bear damage for each of four sample stands under two bear damage scenarios.

Stand	% Damaged	TPH Damaged	<i>Salvage Scenario</i>			<i>Total Loss Scenario</i>		
			Economic Loss	Loss/ha	% Value of Undamaged Stand	Economic Loss	Loss/ha	% Value of Undamaged Stand
CR-WA	16.2	148	\$24,530	\$1,635	90	\$74,666	\$4,978	70
CR-OR	8.5	55	\$7,321	\$915	96	\$31,805	\$3,976	83
WC-OR	42.4	310	\$6,846	\$1,141	84	\$19,528	\$3,255	54
WC-WA	13.5	93	\$6,138	\$472	95	\$31,413	\$2,416	75

Table 6. Average cost/ha for common timber management activities in western Oregon and Washington, 2015.

Management Activity	Cost per Hectare
Site preparation	\$644.94
Site planting	\$480.13
Decadal management	\$156.43
Precommercial-thin	\$383.26
Fertilization	\$179.18

Figures

Figure 1. Locations of the four study sites in the western Cascades and Coast Range of Oregon (CR-OR and WC-OR) and Washington (WC-WA and CR-WA).

Figure 2. Illustration of arrangement of 0.04 ha circular plots. Plots were evenly spaced across each stand in a grid-like system.

Figure 3. Flowchart illustrating various growth model and economic model inputs and outputs for simulating impacts of bear damage to private timberlands in western Oregon.