Canadian Journal of Fisheries and Aquatic Sciences
Journal canadien des sciences halieutiques et aquatiques

**Evaluating benefits of stocking on sockeye recovery
projections in a nutrient-enhanced mixed life history
population**

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0438.R1
Manuscript Type:	Article
Date Submitted by the Author:	02-Mar-2018
Complete List of Authors:	van Poorten, Brett; BC Ministry of Environment; University of British Columbia Sauder School of Business, Institute for the Oceans and Fisheries Harris, Shannon; British Columbia Ministry of Environment Hebert, Allison; British Columbia Ministry of Environment
Is the invited manuscript for consideration in a Special Issue? :	N/A
Keyword:	Sockeye, Kokanee, Density-dependent, Hatchery supplementation, heritability

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

**Evaluating benefits of stocking on sockeye recovery projections
in a nutrient-enhanced mixed life history population**

Brett T. van Poorten, Shannon Harris and Allison Hebert

British Columbia Ministry of Environment and Climate Change Strategy, 2202 Main
Mall, Vancouver, B.C., Canada, V6T 1Z4

Direct Correspondence to the senior author
Email: Brett.vanPoorten@gov.bc.ca
Phone: 604-222-6761

Abstract

Dam construction often blocks migration of anadromous sockeye salmon (*Oncorhynchus nerka*), resulting in a residualized population that is often managed as landlocked kokanee. Anadromy resumes when a reconnection to the ocean is established, however there may be selective pressure acting on threshold trait(s) affecting smolt timing and probability. While there may be interest in predicting persistence of anadromous and residual sockeye forms, this is difficult because the heritability of smolting in these populations is poorly known. We develop a fully density-dependent age-structured model to project abundances for both anadromous and resident sockeye. The model considers trophic interactions due to nutrient variation and the density dependent consequences for smolting and adult returns. Moreover, it asks how each life history type will persist if a hatchery were used to promote anadromous sockeye through artificial selection. We show hatchery supplementation is unlikely to significantly impact anadromous or resident sockeye, although there is substantial prediction uncertainty suggesting in projections. Our study suggests that providing passage for previous land-locked anadromous populations will lead to the return of anadromous fish in the short-term, but long-term prospects are far from certain.

52 **Introduction**

53 Dams and reservoirs are an important component of the landscape and while they
54 serve an important socio-economic function, they also pose a threat to ecological
55 integrity of the watershed (Freeman et al. 2003; Liermann et al. 2012). Dams and
56 reservoirs disrupt littoral habitat and productivity, alter hydrodynamic processes,
57 alter reservoir nutrient dynamics and complicate or remove connectivity within the
58 watershed (Friedl and Wüest 2002; Hall et al. 2011). The loss of connectivity is
59 especially important in systems that have traditionally supported anadromous fish
60 species, which annually transport large quantities of marine-derived nutrients to
61 support freshwater and surrounding terrestrial ecosystems (Schindler et al. 2005;
62 Scheuerell et al. 2007; Hall et al. 2012). Additionally, the impact of dams to the
63 cultural identity of indigenous communities is substantial as members of First
64 Nations who were traditionally connected to the fish and flowing water must look
65 elsewhere to satisfy this cultural and spiritual loss (Baxter 1977; Gregory and
66 Trousdale 2009). Restoration of fish passage is often technically feasible but can
67 often be expensive and decisions on whether and where to invest in passage must
68 appropriately weigh environmental, economic, social and cultural values.

69 Managing fish populations usually involves taking actions that directly or indirectly
70 affect the numbers or size of animals in the system. This can occur either due to
71 stocking, changes in fisheries management, which results in selective harvest of fish,
72 or due to enhancement of stocks or habitat (Walters and Martell 2004). However,
73 the implications for size structure and abundance are not independent (He et al.

2015). Density dependent processes mean that changes in the abundance or size-structure of the population result in changes in the food resources available per individual, which in turn result in growth or survival implications (Lorenzen 2008). On longer time scales, changes in size at age of mature individuals could result in more or larger eggs produced per female, which affects density dependent survival of resulting larvae in the next generation (Beverton and Holt 1957). Moreover, these simple changes can be propagated through the food web through competitors, predators and prey. The outcomes of any management decision can be complex and counterintuitive (Hilborn and Walters 1992).

Augmenting fish populations with hatchery-reared, wild-stock fish is a frequently used tactic in recovering low abundance populations. The basic idea is to improve survival of young fish and augment abundance, but the detrimental effects of hatcheries are often overlooked (Levin et al. 2001; Araki and Schmid 2010). Negative consequences of stocking include increased competition with wild-spawned fish, inbreeding depression, introgression and disease transfer, among others (Fleming and Gross 1993; Reisenbichler and Rubin 1999; Weber and Fausch 2003; Araki et al. 2007; Araki and Schmid 2010). Many of these impacts are often not observed in the hatchery; persistent differences between wild and hatchery fish are often only evident by their performance following release in the wild (Araki et al. 2008; Thrower and Hard 2009). Among anadromous species like Pacific salmon (*Oncorhynchus spp.*), the goal of stocking is often to improve abundance of adults once stocked fish smolt to the ocean, grow and return, but impacts of increased densities of pre-smolting juveniles (i.e. rearing juveniles in the donor stream or

lake) on the remainder of the freshwater ecosystem are often overlooked. This is particularly poignant given uncertain density dependent responses in growth and recruitment.

Sockeye salmon (*O. nerka*) have a broad suite of life history types due to a combination of phenotypic and genotypic differences among populations and individuals. Anadromous sockeye salmon rear in freshwater for one to three years prior to smolting, migrate to the ocean and remain there for one to three years before returning to freshwater to spawn and die, typically between ages 3-5. Some sockeye, termed residents, may delay smolting or mature early without migrating to the ocean (this predominantly occurs in males). Early maturing resident sockeye males interbreed with anadromous sockeye by sneaking fertilization, but are morphologically distinct in that they are much smaller and do not attain the typical red colouration at spawning because they are inefficient at utilizing the lower concentration of carotenoids in a freshwater diet (Craig and Foote 2001). Finally, there are kokanee salmon, the freshwater form of *O. nerka*, which do not smolt and may live sympatrically with sockeye (Taylor et al. 1996). Kokanee live for 3-5 years in freshwater before spawning. Kokanee are thought to maintain genetic isolation from sockeye due to differences in spawning habitat and size: while they have re-evolved the ability to turn red at maturity, they are much smaller (Wood and Foote 1996; Craig and Foote 2001; Dodson et al. 2013).

Among resident and anadromous sockeye, the choice to smolt is thought to be a function of a threshold trait(s) which initiates physiological changes to the body

once certain conditions are met (Dodson et al. 2013), however, when anadromy is blocked, sockeye may all become residents, as anadromy is not obligatory for their life history (Ricker 1959). Interestingly, even kokanee populations that have been isolated for thousands of years are capable of smolting and surviving in the marine environment before returning to their native rivers, albeit with lower marine survival than their anadromous sockeye counterparts (reviewed in Godbout et al. 2011). Godbout et al. (2011) reported the return of anadromy in a previously isolated *O. nerka* population after passage was allowed after nearly 90 years. This population is a deep-water spawner, so neither resident nor anadromous individuals develop red colouration, calling into question whether these are true sockeye or kokanee. Although elders of the Katzie First Nation tell stories of red sockeye being caught in the upper Alouette River (above what is now the Alouette Reservoir; Diamond 1955), this phenotype no longer exists. Interestingly, only a low proportion of this population smolts each year, calling into question the heritability and selective pressure of smolting. While several First Nations and stakeholders wish to see strong annual sockeye returns to the reservoir each year, the decision to invest in permanent fish passage (e.g. a fish ladder) will partially depend on heritability of smolting and in turn, projected sockeye returns.

This study examines the density dependent consequences of stocking in a reservoir with a mixed anadromous-resident sockeye population. We use existing data to estimate various key population parameters, including density dependent growth and survival rates. We project anadromous sockeye returns and abundance of resident sockeye given the full range of uncertainty in all parameters as well as

heritability of smolting. Here, heritability of smolting is used as a surrogate for heritability on liability traits, which are a component of complex and relatively uncertain threshold traits leading to alternative migratory tactics (Dodson et al. 2013). The model will be used to provide an evaluation of the value of stocking to promote smolting and provide medium-term (25-year) projections for recovery of the anadromous life history phenotype.

Methods

Study System

This study centers on the sockeye population rearing in Alouette Reservoir (49° 17'N, 122° 29'W), in southwestern British Columbia (BC). This population exhibits both freshwater resident and anadromous phenotypes. To underline ambiguity in true origins of the population, we refer to all *O. nerka* in the population as 'nerkids'. Alouette Lake Reservoir is a medium-sized reservoir (1666 ha at full-pool) approximately 50 km west of Vancouver, Canada. The reservoir is 17 km long, consisting of two basins separated by a shallow isthmus. The lake is home to numerous cool- and cold-water species, and supports a recreational fishery on rainbow trout (*O. mykiss*), cutthroat trout (*O. clarkii*), bull trout (*Salvelinus confluentus*) and resident nerkids, although the majority of fishing effort targets nerkids.

Alouette Reservoir was created in 1928 by completion of the Alouette Dam at the south end of the lake, effectively blocking access to the reservoir from the lower

Alouette River. Most flow is now redirected north to a 1 km long tunnel linking to Stave Reservoir, where stored water is used for hydroelectric power generation. The remainder of water is released through a low-level outlet in the dam to the Alouette River, although water can also be released over a crest gate and a free crest weir when water levels exceed 125.5 masl. Completion of the dam meant a cessation of anadromy for several Pacific salmon species, including sockeye salmon, as there was no provision for passage at the dam. Most salmon species were extirpated from the reservoir following completion of the dam with the exception of sockeye, where only obligate residents remained. This also resulted in a loss of annual salmon returns to the local Katzie First Nation.

Productivity in Alouette Reservoir is thought to have followed the typical boom-and-bust nutrient cycle of reservoir formation ('cultural oligotrophication'; Stockner et al. 2000) following dam construction primarily due to a disruption of natural hydrology, sedimentation in the new reservoir and loss of carbon production from the littoral community. Additionally, the loss of marine-derived nutrients transported by anadromous salmon have also contributed to declines in productivity (Larkin and Slaney 1996; Cederholm et al. 1999). Nutrient restoration in Alouette Reservoir began in 1999 with the primary objective of improving the recreational fishery for resident nerkids. Since that time, inorganic agricultural grade fertilizer has been added weekly from approximately May to September each year.

184 The Alouette system has been carefully monitored to promote production of a food
185 web that leads to production of nerkids. Phosphorus and nitrogen levels, as well as
186 phyto- and zooplankton densities and composition were monitored bi-monthly in
187 the first four years and monthly thereafter, in an attempt to maximize edible
188 phytoplankton while minimizing the risk of plankton bloom. The amount and ratio
189 of phosphorus to nitrogen has varied over the course of the program to adjust to
190 varying annual growing conditions and adaptively determine the optimal strategy
191 aimed at promoting the growth of edible plankton and therefore fish. Gill nets are
192 used in the fall to monitor resident nerkid size structure and hydroacoustic surveys
193 have been conducted concurrent with gill net sampling to provide an index of
194 abundance. A single beam hydroacoustic survey was used from 1998 to 2001, but
195 this was switched to a split beam survey thereafter. Finally, access point creel
196 surveys have been conducted in 1996 and 2014, providing estimates of total harvest
197 from recreational anglers; there is no other fishery on the reservoir.

198 Annual spills of water over the free crest weir began in 2005 and continued annually
199 from approximately April 15 to June 15 in an effort to re-establish smolting in the
200 nerkid population. Abundance of smolting nerkids was monitored with mark
201 recapture using two rotary screw traps in the lower Alouette River. Numbers of
202 nerkid smolts emigrating from the reservoir has varied annually from 720 to 62,000
203 (Borick-Cunningham and Smith 2016) and returns have varied annually from 0 to
204 115 (Mathews et al. 2016). Returning adult nerkids have been transported to the
205 reservoir through a trap and truck system. Genetic comparisons of smolting and
206 resident nerkids has shown these two phenotypes to be considered the same

population and not genetically distinct. Examination of the next generation do not conclusively demonstrate intermixing of the two phenotypes (Godbout et al. 2010). Re-anadromization of sockeye was originally seen as a great success for members of the Katzie First Nation and local stakeholders, but the lack of improvement in the annual number of smolts and the low mean marine survival ($<1\%$) has been disappointing and led to calls by the Katzie people and local stakeholders for further action to be taken.

To further promote smolting in Alouette nerkids, stakeholders are proposing hatchery intervention. The annual supplemental stocking of nerkid fry is proposed to occur over the short-term (two generations) using returning adult Alouette ocean-run nerkids as parental stock. The expectation is that annual stocking of fry derived from anadromous nerkids may artificially select for smolting, with the objective of increasing the proportion of nerkids that smolt. Before investing significant time and money on a stocking program it is important to evaluate the potential success and effects of the program to resident and anadromous nerkids. At present, the density dependent impacts of additional age-0 fry on the growth and abundance of resident nerkids and the resultant fishery are unknown. Likewise, with no prior information on the heritability of smolting and the potential outcome of artificially selecting for smolting, it is difficult to predict how anadromous nerkid returns will continue over time.

227   **Model Development**

228   A two-phenotype nerkid model was developed to predict the strength of density
229   dependent processes and evaluate the potential outcomes that stocking and changes
230   in the probability of smolting may have on both the anadromous phenotype and the
231   freshwater recreational fishery. Derivation of Beverton-Holt recruitment
232   parameters and state dynamics model are described in Table 1 and parameters of
233   the models are listed in Table 2. The state-dynamics model is conditioned on smolt
234   estimates from the downstream smolt mark-recapture program (Mathews et al.
235   2016), returning smolted nerkid count data from the trap and truck program below
236   the Alouette dam (Borick-Cunningham and Smith 2016) and zooplankton densities
237   (Hebert et al. 2015) from the nutrient restoration program. The model is fit to
238   hydroacoustic, gill net and fishery data collected as part of the nutrient restoration
239   program to parameterize the demography of the nerkid population. The fitted
240   model was then used to make projections of both phenotypes for 25 years
241   considering uncertainty in the heritability of smolting.

242   A key component of this model is the stock-recruitment function, which is driven by
243   density dependent compensation and carrying capacity in juvenile mortality as
244   conditions change in the reservoir (Eqs. T1.3-T1.6). This model is derived from
245   arguments in Walters and Korman (1999) and fully described in van Poorten et al.
246   (*in press*). In it, the maximum survival rate, α , is a function of a base mortality and a
247   food-dependent mortality rate, causing maximum juvenile survival to increase in
248   years of abundant food supply. The carrying capacity parameter, β , is also

influenced by food density, with maximum recruitment increasing as a quadratic function of food density.

The model includes a combination of parameters that are estimated (T1.1) and assumed constant (T1.2). All indices, parameters and variables are described in Table 2. Recruitment from eggs to age-1 is divided into two stanzas (Moussalli and Hilborn 1986): one predicts the number of fry (age-0.5; T1.3-T1.4) and the second predicts the number of yearlings (age-1; T1.5-T1.6). In both functions, recruitment is limited by zooplankton availability and competition among the recruiting cohort using functions suggested in Walters and Korman (1999) and van Poorten et al. (*in press*). Recruitment to age-0.5 fry is multiplied by a lognormally distributed process error term. Initial growth is assumed to have been constant resulting in length-at-age following a standard von Bertalanffy function (T1.7). Numbers at age are predicted for two annual time periods: one in the autumn to correspond with timing of smolting, which is initialized using estimated abundances ($N_{t,a}$; T1.8) and one the following spring ($N_{(sp)t,a}$; T1.9) to correspond to sampling dates. Consumption each year is assumed to vary in response to available zooplankton and competition among nerkids, assuming all size-classes feed on the same zooplankton resource. This results in asymptotic size varying over time, while metabolic rate remaining constant (Walters and Post 1993; van Poorten and Walters 2016; T1.10), resulting in size-at-age varying each year (T1.11). Vulnerability to harvest is assumed to be a logistic function of length (T1.12). The fishery is assumed to occur primarily in summer, so total mortality during summer is half of annual natural mortality and all of size-adjusted fishing mortality (T1.13). Nerkids spawn in autumn and hatch in

272 spring, therefore, number of fry in fall is calculated based on egg abundance the
273 previous year using the first recruitment stanza calculation adjusted with lognormal
274 process error (T1.14). Yearlings the following spring are calculated based on
275 predicted fall fry abundance and the second recruitment stanza calculation (T1.15).
276 In subsequent years, fall abundance of nerkids in the reservoir is reduced as some
277 fish smolt out of the system in early spring. Egg production the following year is
278 based on size-based fecundity of age-3 and 4 resident nerkids (T1.16; McGurk,
279 2000) plus eggs contributed by returning anadromous nerkids (T1.17-T1.18).
280 Anadromous nerkid fecundity is based on a power function with length using
281 parameters from Pitt Lake sockeye (McGurk 2000). Because resident nerkids are
282 semelparous, the model assumes a fixed proportion of fish mature at age-3 (m_3 , set
283 at 0.9) and the remaining fish spawn the following year (T1.19).

284 Annual numbers of smolting fish are modelled as normally distributed random
285 variables with means and standard deviations set based on annual Peterson mark-
286 recapture estimates of abundance based on rotary screw traps set downstream of
287 the dam (Mathews et al. 2016). Trapping downstream of the dam was not conducted
288 in 2015; therefore no estimate of smolt outmigration was available for that year.
289 Smolt abundance in 2015 was set as a normally distributed random variable with
290 prior distributions of mean and standard deviation based on the empirical
291 distributions of each over the entire time series of smolt estimates for the system.

292 Model parameters (T1.1) were estimated using a state-space Bayesian approach
293 that jointly fit to observations of length-at-age, hydroacoustic estimates of

abundance, and annual estimates of fishery harvest. Annual length-at-age observations for individual fish captured in gill net sampling ($l_{t,a,i}$) were assumed to be normally distributed with observation error $\tau_{(l)t,a}$ (T1.21-T1.22)

$$1) \quad l_{t,a,i} \sim N(\hat{l}_{t,a}, \tau_{(l)t,a})$$

Log-transformed hydroacoustic abundance estimates were assumed to follow a normal distribution. Annual autumn hydroacoustic estimates of abundance were divided into age-0 and age-1 and older nerkids based on modal separation of annual target strength distributions. Therefore model predictions of log-transformed autumn abundance were fit separately to each age group based on abundance in autumn

$$2) \quad \log(N_{(H)t,0}) = N(\log(N_{t,a=0}), \tau_{(H)0})$$

$$3) \quad \log(N_{(H)t,1+}) = N(\log(\sum_{a=1}^A N_{t,a}), \tau_{(H)1+})$$

where precisions for age-0 and 1+ age groups observation errors are described in T1.23-T1.26.

Total fishery harvest has only been evaluated twice since 1998 in Alouette Reservoir using creel surveys: once in 2002 (McCusker and Wilson 2002) and once in 2014 (Hebert et al. 2016). Log-transformed total annual harvest estimates were assumed to be normally distributed

$$4) \quad \log(C_t) = N(\log(\sum_{a=0}^A \hat{C}_{t,a}), \tau_{(C)})$$

where harvest precision in observation is directly estimated.

Prior probability distributions for all estimated parameters are shown in Table 3. We used JAGS 3.4.0 (Plummer 2003) to numerically approximate the posterior probability distribution using Markov Chain Monte Carlo (MCMC) simulation. Posterior distributions were calculated from 2,500,000 iterations after a burn-in of 1,000,000 and thinned to provide a final sample of 1,000 from three separate chains. Convergence could not be rejected using Gelman-Rubin convergence diagnostics and visual inspection of MCMC chains.

Future projections

The two-phenotype nerkid model was used to project the system 25 years into the future using estimated parameters. Zooplankton densities were assumed to remain constant at a level equal to the mean densities observed from 2011-2015. This is consistent with the goals of the nutrient restoration program. Anadromous nerkid-derived offspring are assumed to be stocked as fry directly into the reservoir for the entire 25 years. All smolts leaving the system are assumed to be age-1 for simplicity.

Nerkid fry descended from returning adults stocked into the reservoir are assumed to smolt at a rate dictated by the narrow-sense heritability (h^2) of smolting (Morrisey et al. 2014; Table 4). The proportion of stocked fry that will smolt is given in T4.1. Note that returning anadromous nerkids are selected as broodstock in the hatchery, therefore the proportion of hatchery fry that smolt is related to the difference between the proportion of the previous generation that smolted from selected parents ($=1.0$) and the general population ($=p_{(smolt)t-4}$). Resident nerkids in future generations will also be influenced by the selection differential for smolting

based on the difference between the proportions of resident nerkids that smolted the previous generation and the relative genetic success of smolting (the proportion of eggs laid that were derived from anadromous nerkids; T4.2). The number of smolts each year is then a combination of hatchery and resident nerkid smolts (T4.3). The number of anadromous nerkids returning two years later are given by the smolts leaving the reservoir multiplied by a total marine survival rate (T4.4).

Projected dynamics of the age-structured model proceed as in Table 1, except additional fish may be stocked into the reservoir as fry to the 0+ population. These fish then compete for resources with natural-spawned offspring, thereby reducing survival to age-1 through increased density dependent competition. For ease of interpretation, all smolts are assumed to be age-1.

Future projections of anadromous nerkid returns and the resident nerkid population are dependent on two key parameters: marine survival rate and heritability of smolting. Marine survival was fixed at 0.25% based on the geometric mean observed marine survival rate for Alouette anadromous nerkids (Alouette River Management Society 2015). We are not aware of any estimates of heritability of smolting in sockeye; heritability was therefore assumed to be a uniform beta distributed random variable with both shape parameters set to 1.0 (i.e. uniform from 0-1).

355 Results

356 Inorganic fertilizer added into Alouette Reservoir starting in 1999 accounted for a
357 substantive proportion of the total nitrogen and phosphorus in the system (Fig. 1).
358 Total nitrogen and phosphorus loading in the reservoir varied as researchers
359 refined the relative amounts of nitrogen and phosphorus to best produce edible
360 phytoplankton. This had an impact on zooplankton densities, although seasonal
361 averages varied widely across years. Increased nutrient load in the reservoir
362 resulted in marked increases in mean annual zooplankton densities, initially in all
363 cladocerans and later focused on *Daphnia* sp. (Fig. 1). Nerkid growth and size-at-age
364 immediately increased as a result of increased zooplankton densities (Fig. 2). Larger
365 nerkids released more eggs, resulting in a lagged increase in abundance of age-0
366 nerkids due to both increased egg abundance and increased survival of age-0 fish.
367 Older age-classes also increased over time due to improved abundance and survival
368 of age-0 nerkids (Fig. 2). As abundance of both zooplankton and nerkids began to
369 stabilize, nerkid body size also stabilized with size-at-age slightly larger than before
370 nutrient restoration.

371 The model was able to estimate density dependent processes relating to survival
372 from egg to fry, and survival from fry to smolt. Carrying capacity for both stanzas
373 increased three-fold in the first two years following the onset of nutrient
374 restoration, declined as zooplankton densities declined and fry density increased
375 and stabilized at an intermediate level after 2008 (Fig. 3). Carrying capacities closely
376 track zooplankton density, with the strength of the zooplankton effect influenced by

density of age-0+ or age-1, respectively. Carrying capacity for both stanzas was similar, which reflects the difficulty in updating the prior prediction of p_1 - p_3 , which allocate Beverton-Holt parameters across the two stanzas.

If the nutrient restoration program maintains zooplankton density at densities seen in recent years, the model predicts resident nerkid vulnerable abundance to remain approximately similar to what has been seen to date (Fig. 4). Unfortunately, median anadromous nerkid returns are projected to decline over time due to low marine survival. However, these projections include uncertainty in heritability, which contribute variation in the annual distribution of predicted vulnerable resident and anadromous nerkid returns. If heritability of smolting is actually low, anadromous nerkid returns will persist over time despite lower survival in the marine environment, albeit at low abundance (Fig 5; left panels). In this scenario, returning fish are primarily smolts from the abundant resident nerkid phenotype, rather than a persistent anadromous phenotype. If heritability is actually high, anadromous nerkid returns will rapidly decline and that life history form will be lost because marine survival is too low to maintain a sustainable anadromous run for this population. Both scenarios show minimal influence on resident nerkid persistence because of low inputs from anadromous nerks each year (Fig 5; top panels).

If fry derived from anadromous nerks are stocked into the reservoir, abundance of resident nerks vulnerable to the recreational fishery is projected to decline over 25 years and the extent of the decline is directly related to stocking rate (Fig. 6a). Note that vulnerable abundance is a function both of numbers, which decline with

increased competition among fry, and size, which increases with reduced competition following smolting. Once stocking rate exceeds 35,000 anadromous nerkid fry, there is a 5% probability of extirpation of the resident phenotype and this probability increases with stocking. The range of projected resident nerkid abundance is substantial, with 95% credible intervals ranging from a high approximately 50,000 vulnerable resident nerkids at a stocking rate of 80,000 nerkid fry, down to complete extirpation. Mean anadromous nerkid returns 25 years in the future are projected to moderately increase with stocking rate, though again, uncertainty around projections is substantial (Fig. 6b). For example, stocking 50,000 anadromous nerkid fry results a 95% probability that anadromous nerkid returns in 25 years may range between 10 and over 3,300. Even at this high stocking rate, there is a 50% chance that projected returns 25 years in the future will be 161 anadromous nerkids or less.

Discussion

Marine-derived nutrients are an important subsidy to oligotrophic coastal systems. This is especially so for reservoirs such as Alouette Reservoir, which are nutrient deprived due to the effect of dams on within-reservoir nutrient dynamics (Scott et al. 2017). Ordinarily, additional sockeye returning to the system would increase natural nutrient levels and consequent prey density resulting in improvements to production and carrying capacity (Cederholm et al. 1999; Naiman et al. 2002; Schindler et al. 2003). However the nutrient restoration program offsets this benefit because nutrient levels are managed to maximize edible phytoplankton

concentrations. Additional marine derived nutrients would be largely matched by a decline in added inorganic fertilizer. In fact, it would take substantial anadromous nerkid returns to exceed added nutrient concentration in the system (Scott et al. 2017). The overall impact is that improved anadromous returns only add competitors to the system without improving growing conditions. An important lesson from the Alouette case study is that sockeye restoration targets should carefully consider trophic interactions, especially in systems already undergoing nutrient restoration.

Despite nearly ten years of anadromous nerkid returns, smolting rates from Alouette Reservoir are often still low, either suggesting this is a kokanee population with low smolting heritability or a sockeye population where conditions favour residency over anadromy. This work highlights the substantial value of information with respect to understanding the mechanisms and thresholds related to smolting when considering restoring or enhancing anadromous sockeye stocks (Hansen et al. 2016). Barriers to migration are being re-examined and solutions such as dam removal, trap-and-truck and fish ladders are being considered at many dams throughout North America (Anderson et al. 2014). Predicting outcomes of reintroductions is difficult in light of uncertainty in heritability and often only includes extreme possibilities (e.g. 100% anadromy or 100% residency; Hansen et al. 2016). We expect this work will guide future attempts to restore efforts to restore anadromy using stranded nerkid phenotypes.

442 Kokanee and sockeye populations occur sympatrically throughout their range, yet
443 these pairs are consistently genetically distinct, partially because of reproductive
444 isolation due to their sexual dimorphism (Foote et al. 1989; Taylor et al. 1996).
445 Taylor et al. (1996) suggested that although male kokanee may sneak fertilizations
446 with female sockeye (Foote and Larkin 1988), there must be substantial selective
447 pressure against kokanee-sockeye 'hybrids'. Although there is a substantial
448 polymorphism in body size between resident and anadromous nerkids in Alouette
449 Reservoir, there is no difference in colour (both are drab olive or silver),
450 presumably because of deep-water spawning (Moreira and Taylor 2015).
451 Performance of sockeye-kokanee hybrids has been shown to be inferior in salinity
452 tolerance, swimming performance, growth, and development rate (Wood and Foote
453 1990; Taylor and Foote 1991; Foote et al. 1992). Although Alouette nerkids have
454 smolted, returned and been released back into the reservoir for almost 10 years,
455 there have been no observations from genetic analysis of offspring from these
456 anadromous fish and no conclusive evidence of hybrids, although sample sizes were
457 low (Godbout et al. 2011; Alouette River Management Society 2015). The model
458 presented here assumes no intermixing, but neither does it assume Allee effects.
459 Given the low anadromous nerkid return rates, it may be that re-establishment in
460 the Alouette Reservoir may only be possible following one or more anomalously
461 good smolting or marine survival events, or through hatchery supplementation.

462 Our results show that sustained stocking is unlikely to dramatically improve
463 anadromous nerkid returns over the first 25 years, even at substantial stocking
464 rates. Sockeye salmon are particularly difficult to culture largely due to the potential

for disease outbreak (Meyers et al. 1990). While stocking may be used as a tool to select for anadromy if there is some heritability in smolting, selective pressure for smolting will return to normal once hatchery supplementation is ceased. If heritability were known to be high and marine survival even marginally improved, the chance of successful restoration of anadromy may be higher, though this would also likely spell the demise of the resident nerkid life history. Similar instances where previously landlocked steelhead (*O. mykiss*) allowed to smolt revealed a low prevalence of smolting due to low heritability (Thrower et al. 2004; Pearse et al. 2009). This highlights the need to determine heritability of smolting to evaluate the benefits (in terms of sockeye returns) against costs of restoration or installation of permanent fish passage.

Arguably the most uncertain process in any Pacific salmon life cycle model is marine survival, which is essentially a catch-all for multiple, but unmeasured impacts on anadromous fish. Alouette anadromous nerkids migrate outside times of major commercial fishery openings (Godbout et al. 2011), so the causes of marine survival variation are largely unknown. Marine survival for Alouette anadromous nerkids has ranged from 0 to 1.3% since outmigration began, which is low among nearby stocks (DFO 2010, 2017). If marine survival for Alouette anadromous nerkids were to improve, the outcome for this phenotype would be better than the median prediction from our models. This again serves to underscore the tremendous uncertainty in restoration potential for this stock and the need to invest in determining heritability to reduce this uncertainty.

Our work clearly demonstrates high uncertainty in expected anadromous nerkid returns if smolting continues. This suggests that from a purely ecological and economic standpoint, investing in permanent upstream fish passage infrastructure may not be prudent. However, this is one of many examples where the historic construction of dams has directly affected one or more First Nations (Notzke 1994). The decision to invest in anadromy should and does weigh ecological, economic, social and cultural values; the cultural benefit to restoring anadromy would be substantial. This work highlights not only the uncertain ecological benefit of stocking anadromous nerkids into this system, but also the uncertain cultural gain. Results here should reinforce how important it is to understand the factors that contribute to smolting in order to contribute to a more certain and predictable outcome, both for the hydro-electric utility who must invest in fish passage, but also for the Katzie First Nation that has waited so long for salmon to return.

Acknowledgements

This work builds on substantial and sustained investments of time and money by several groups. The Alouette River Sockeye Reanadromization Project is a collaboration of the Katzie First Nation, Alouette River Management Society, BC Ministry of Environment and Climate Change Strategy, Fisheries and Oceans Canada and BC Hydro. Funding for the various programs is provided by BC Hydro, BC Ministry of Environment and Climate Change Strategy and the Fish and Wildlife Compensation Program. The Alouette Nutrient Restoration Program is conducted by the BC Ministry of Environment, with assistance from Ministry of Justice and

509 Attorney General – Fraser Regional Corrections Centre and BC Ministry of Forests,
510 Lands, Natural Resource Operations and Rural Development. Smolt enumeration is
511 managed by Katzie First Nation and LGL Environmental Consultants; the sockeye
512 transplant program is managed by the Alouette River Management Society and the
513 Fraser Regional Correction Center. We are indebted to all these organizations and
514 individuals for the substantial data and support that led to this work.

Draft

References

- Alouette River Management Society. 2015. Alouette sockeye adult enumeration: Implementation year 7. Fish and Wildlife Compensation Program - Coastal. Burnaby, BC.
- Anderson, J.H., Pess, G.R., Carmichael, R.W., Ford, M.J., Cooney, T.D., Baldwin, C.M., and McClure, M.M. 2014. Planning Pacific salmon and steelhead reintroductions aimed at long-term viability and recovery. *North Am. J. Fish. Manag.* **34**: 72–93. doi: 10.1080/02755947.2013.847875.
- Araki, H., Berejikian, B.A., Ford, M.J., and Blouin, M.S. 2008. Fitness of hatchery-reared salmonids in the wild. *Evol. Appl.* **1**: 342–355.
- Araki, H., Cooper, B., and Blouin, M.S. 2007. Genetic effects of captive breeding cause a rapid, cumulative fitness decline in the wild. *Science* **318**: 100–103.
- Araki, H., and Schmid, C. 2010. Is hatchery stocking a help or harm? *Aquaculture* **308**: S2–S11. Elsevier B.V. doi: 10.1016/j.aquaculture.2010.05.036.
- Baxter, R.M. 1977. Environmental effects of dams and impoundments. *Annu. Rev. Ecol. Syst.* **8**: 255–283.
- Beverton, R.J.H., and Holt, S.J. 1957. On the dynamics of exploited fish populations. Chapman & Hall, London.
- Borick-Cunningham, G., and Smith, S. 2016. Alouette sockeye adult enumeration project - (Bridging Year 2015). Fish and Wildlife Compensation Program -

- 535 Coastal. Burnaby, BC.
- 536 Cederholm, C.J., Kunze, M.D., Murota, T., and Sibatani, A. 1999. Pacific salmon
537 carcasses: essential contributions of nutrients and energy for aquatic and
538 terrestrial ecosystems. *Fisheries* **24**: 6–15.
- 539 Craig, J.K., and Foote, C.J. 2001. Countergradient variation and secondary sexual
540 color: phenotypic convergence promotes genetic divergence in carotenoid use
541 between sympatric anadromous and nonanadromous morphs of sockeye
542 salmon (*Oncorhynchus nerka*). *Evolution* **55**: 380–391.
- 543 DFO. 2010. Assessment of Cultus Lake sockeye salmon in British Columbia in 2009
544 and evaluation of recent recovery activities. Canadian Science Advisory
545 Secretariat Science Advisory Report. Ottawa, ON.
- 546 DFO. 2017. Pre-season run size forecasts for Fraser River sockeye (*Oncorhynchus*
547 *nerka*) and pink (*O. gorbuscha*) salmon in 2017. Canadian Science Advisory
548 Secretariat Science Advisory Report. Ottawa, ON.
- 549 Diamond, J. 1955. The faith of a coast salish indian. British Columbia Provincial
550 Museum, Victoria, B.C.
- 551 Dodson, J.J., Aubin-Horth, N., Theriault, V., and Paez, D.J. 2013. The evolutionary
552 ecology of alternative migratory tactics in salmonid fishes. *Biol. Rev.* **88**: 602–
553 625.
- 554 Fleming, I.A., and Gross, M.R. 1993. Breeding success of hatchery and wild coho
555 salmon (*Oncorhynchus kisutch*) in competition. *Ecol. Appl.* **3**: 230–245.
- 556 Foote, C.J., and Larkin, P.A. 1988. The role of male choice in the assortative mating of

- 557   anadromous and non-anadromous sockeye salmon (*Oncorhynchus nerka*).
558   Behaviour **106**: 43–62.
- 559   Foote, C.J., Wood, C.C., Clarke, W.C., and Blackburn, J. 1992. Circannual cycle of
560   seawater adaptability in *Oncorhynchus nerka*: genetic differences between
561   sympatric sockeye salmon and kokanee. Can. J. Fish. Aquat. Sci. **49**: 99–109.
- 562   Foote, C.J., Wood, C.C., and Withler, R.E. 1989. Biochemical genetic comparison of
563   sockeye salmon and kokanee, the anadromous and nonanadromous forms of
564   *Oncorhynchus nerka*. Can. J. Fish. Aquat. Sci. **46**: 149–158.
- 565   Freeman, M.C., Pringle, C.M., Greathouse, E.A., and Freeman, B.J. 2003. Ecosystem-
566   level consequences of migratory faunal depletion caused by dams. In American
567   Fisheries Society Symposium 35. pp. 255–266.
- 568   Friedl, G., and Wüest, A. 2002. Disrupting biogeochemical cycles – Consequences of
569   damming. Aquat. Sci. **64**: 55–65.
- 570   Godbout, L., Wood, C.C., Withler, R., and Menard, D. 2010. Smolt parental lineage
571   assessment to evaluate successful spawn of returned 2009 Alouette sockeye
572   spawners. Fish and Wildlife Compensation Program - Coastal. Burnaby, BC.
- 573   Godbout, L., Wood, C.C., Withler, R.E., Latham, S., Nelson, R.J., Wetzel, L., Barnett-
574   Johnson, R., Grove, M.J., Schmitt, A.K., McKeegan, K.D., and Bradford, M. 2011.
575   Sockeye salmon (*Oncorhynchus nerka*) return after an absence of nearly 90
576   years: a case of reversion to anadromy. Can. J. Fish. Aquat. Sci. **68**: 1590–1602.
577   doi: 10.1139/f2011-089.
- 578   Gregory, R., and Trousdale, W. 2009. Compensating aboriginal cultural losses: An

- 579 alternative approach to assessing environmental damages. *J. Environ. Manage.*
580 **90**: 2469–2479.
- 581 Hall, C.J., Jordaan, A., and Frisk, M.G. 2011. The historic influence of dams on
582 diadromous fish habitat with a focus on river herring and hydrologic
583 longitudinal connectivity. *Landsc. Ecol.* **26**: 95–107.
- 584 Hall, C.J., Jordaan, A., and Frisk, M.G. 2012. Centuries of anadromous forage fish loss:
585 consequences for ecosystem connectivity and productivity. *Bioscience* **62**: 723–
586 731.
- 587 Hansen, A.G., Gardner, J.R., Beauchamp, D.A., Paradis, R., and Quinn, T.P. 2016.
588 Recovery of sockeye salmon in the Elwha River, Washington, after dam
589 removal: dependence of smolt production on the resumption of anadromy by
590 landlocked kokanee. *Trans. Am. Fish. Soc.* **145**: 1303–1317.
- 591 He, J.X., Bence, J.R., Madenjian, C.P., Pothoven, S.A., Dobiesz, N.E., Fielder, D.G.,
592 Johnson, J.E., Ebener, M.P., Cottrill, R.A., Mohr, L.C., and Koproski, S.R. 2015.
593 Coupling age-structured stock assessment and fish bioenergetics models: a
594 system of time-varying models for quantifying piscivory patterns during the
595 rapid trophic shift in the main basin of Lake Huron. *Can. J. Fish. Aquat. Sci.* **17**:
596 1–17.
- 597 Hebert, A.S., Andrusak, G.F., Harris, S.L., and Weir, T. 2016. Alouette Reservoir
598 nutrient restoration program 2014-15. Ecosystems Branch, BC Ministry of
599 Environment. Victoria, BC
- 600 Hebert, A.S., Andrusak, G.F., Harris, S.L., Weir, T., and Vidmanic, L. 2015. Alouette

- 601 reservoir nutrient restoration project: 5-year review, 2009-2013. Ecosystems
602 Branch, BC Ministry of Environment. Victoria, BC.
- 603 Hilborn, R., and Walters, C.J. 1992. Quantitative fisheries stock assessment: Choice,
604 dynamics and uncertainty. Chapman and Hall, New York.
- 605 Larkin, G.A., and Slaney, P.A. 1996. Trends in marine-derived nutrient sources to
606 south coastal British Columbia streams: impending implications to salmonid
607 production. BC Ministry of Environment, Lands and Parks and BC Ministry of
608 Forests. Victoria, BC.
- 609 Levin, P.S., Zabel, R.W., and Williams, J.G. 2001. The road to extinction is paved with
610 good intentions: negative association of fish hatcheries with threatened salmon.
611 Proc. R. Soc. B Biol. Sci. **268**: 1153–1158.
- 612 Liermann, C.R., Nilsson, C., Robertson, J., and Ng, R.Y. 2012. Implications of dam
613 obstruction for global freshwater fish diversity. Bioscience **62**: 539–548.
- 614 Lorenzen, K. 2008. Fish population regulation beyond “stock and recruitment”: the
615 role of density-dependent growth in the recruited stock. Bull. Mar. Sci. **83**: 181–
616 196.
- 617 Mathews, M.A., Smith, J.J., and Bocking, R.C. 2016. Evaluation of the migration
618 success of *O. nerka* (Kokanee / Sockeye) from the Alouette Reservoir, 2015.
619 Fish and Wildlife Compensation Program - Coastal. Burnaby, BC.
- 620 McCusker, M., and Wilson, G. 2002. Alouette Reservoir creel survey, April to October
621 2002. Vancouver, B.C. BC Hydro Power Supply Environmental Services.
622 Burnaby, BC.

- 623 McGurk, M.D. 2000. Comparison of fecundity–length–latitude relationships between
624 nonanadromous (kokanee) and anadromous sockeye salmon (*Oncorhynchus*
625 *nerka*). Can. J. Zool. **78**: 1791–1805.
- 626 Meyers, T.R., Thomas, J.B., Follett, J.E., and Saft, R.R. 1990. Infectious hematopoietic
627 necrosis virus: trends in prevalence and the risk management approach in
628 Alaskan sockeye salmon culture. J. Aquat. Anim. Health **2**: 85–98.
- 629 Moreira, A.L., and Taylor, E.B. 2015. The origin and genetic divergence of “ black ”
630 kokanee , a novel reproductive ecotype of *Oncorhynchus nerka*. **1595**: 1584–
631 1595.
- 632 Morrissey, M.B., de Villemereuil, P., Doligez, B., and Gimenez, O. 2014. Bayesian
633 approaches to the quantitative genetic analysis of natural populations. In
634 Quantitative Genetics in the Wild. Edited by A. Charmantier, D. Garant, and
635 L.E.B. Kruuk. Oxford University Press, Oxford. pp. 228–253.
- 636 Moussalli, E., and Hilborn, R. 1986. Optimal Stock Size and Harvest Rate in
637 Multistage Life History Models. Can. J. Fish. Aquat. Sci. **43**: 135–141. doi:
638 10.1139/f86-014.
- 639 Naiman, R.J., Bilby, R.E., Schindler, D.E., and Helfield, J.M. 2002. Pacific salmon,
640 nutrients, and the dynamics of freshwater and riparian ecosystems. Ecosystems
641 **5**: 399–417.
- 642 Notzke, C. 1994. Aboriginal peoples and natural resources in Canada. Captus Press.
- 643 Pearse, D.E., Hayes, S.A., Bond, M.H., Hanson, C. V, Anderson, E.C., MacFarlane, R.B.,
644 and Garza, J.C. 2009. Over the falls? Rapid evolution of ecotypic differentiation

- 645 in steelhead/ rainbow trout (*Oncorhynchus mykiss*). J. Hered. **100**: 515–525.
- 646 Plummer, M. 2003. JAGS: A program for analysis of Bayesian graphical models using
647 Gibbs sampling. In Proceedings of the 3rd International Workshop on
648 Distributed Statistical Computing.
- 649 van Poorten, B.T., Arlinghaus, R., Daedlow, K., and Haertel-Borer, S.S. 2011. Social-
650 ecological interactions, management panaceas, and the future of wild fish
651 populations. Proc. Natl. Acad. Sci. **108**: 12554–12559. doi:
652 10.1073/pnas.1013919108/-
653 /DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1013919108.
- 654 van Poorten, B.T., and Walters, C.J. 2016. How can bioenergetics help us predict
655 changes in fish growth patterns? Fish. Res. **180**: 23–30. Elsevier B.V. doi:
656 10.1016/j.fishres.2015.07.031.
- 657 Reisenbichler, R.R., and Rubin, S.P. 1999. Genetic changes from artificial propagation
658 of Pacific salmon affect the productivity and viability of supplemented
659 populations. ICES J. Mar. Sci. **56**: 459–466.
- 660 Ricker, W.E. 1959. Additional observations concerning residual sockeye and
661 kokanee (*Oncorhynchus nerka*). J. Fish. Res. Board Canada **16**: 897–902.
- 662 Scheuerell, M.D., Moore, J.W., Schindler, D.E., and Harvey, C.J. 2007. Varying effects of
663 anadromous sockeye salmon on the trophic ecology of two species of resident
664 salmonids in southwest Alaska. Freshw. Biol. **52**: 1944–1956.
- 665 Schindler, D.E., Leavitt, P.R., Brock, C.S., Johnson, S.P., and Quay, P.D. 2005. Marine-
666 derived nutrients, commercial fisheries, and production of salmon and lake

- algae in Alaska. *Ecology* **86**: 3225–3231.
- Schindler, D.E., Scheuerell, M.D., Moore, J.W., Gende, S.M., Francis, T.B., and Palen, W.J. 2003. Pacific salmon and the ecology of coastal ecosystems. *Front. Ecol. Environ.* **1**: 31–37.
- Scott, D.C., Harris, S.L., Hebert, A.S., and van Poorten, B.T. 2017. Nutrient dynamics in a highly managed reservoir system: considering anadromous sockeye salmon (*Oncorhynchus nerka*) and nutrient restoration. *Lake Reserv. Manag.* **33**: 14–22.
- Stockner, J.G., Rydin, E., and Hyenstrand, P. 2000. Cultural oligotrophication: causes and consequences for fisheries resources. *Fisheries* **25**: 7–14.
- Taylor, E.B., and Foote, C.J. 1991. Critical swimming velocities of juvenile sockeye salmon and kokanee, the anadromous and non-anadromous forms of *Oncorhynchus nerka* (Walbaum). *J. Fish Biol.* **38**: 407–419.
- Taylor, E.B., Foote, C.J., and Wood, C.C. 1996. Molecular genetic evidence for parallel life-history evolution within a Pacific salmon (sockeye salmon and kokanee, *Oncorhynchus nerka*). *Evolution* **50**: 401–416.
- Thrower, F., Guthrie, C.I., Nielsen, J., and Joyce, J. 2004. A comparison of genetic variation between an anadromous steelhead, *Oncorhynchus mykiss*, population and seven derived populations sequestered in freshwater for 70 years. *Environ. Biol. Fishes* **69**: 111–125.
- Thrower, F.P., and Hard, J.J. 2009. Effects of a single event of close inbreeding on growth and survival in steelhead. *Conserv. Genet.* **10**: 1299–1307.
- Walters, C., and Korman, J. 1999. Linking recruitment to trophic factors: revisiting

- 689 the Beverton-Holt recruitment model from a life history and multispecies
690 perspective. *Rev. Fish Biol. Fish.* **9**: 187–202.
- 691 Walters, C., and Martell, S. 2004. Fisheries ecology and management. Princeton
692 University Press, Princeton.
- 693 Walters, C.J., and Post, J.R. 1993. Density-dependent growth and competitive
694 asymmetries in size-structured fish populations: A theoretical model and
695 recommendations for field experiments. *Trans. Am. Fish. Soc.* **122**: 34–45.
- 696 Weber, E.D., and Fausch, K.D. 2003. Interactions between hatchery and wild
697 salmonids in streams: differences in biology and evidence for competition. *Can.*
698 *J. Fish. Aquat. Sci.* **60**: 1018–1036.
- 699 Wood, C.C., and Foote, C.J. 1990. Genetic differences in the early development and
700 growth of sympatric sockeye salmon and kokanee (*Oncorhynchus nerka*), and
701 their hybrids. *Can. J. Fish. Aquat. Sci.* **47**: 2250–2260.
- 702 Wood, C.C., and Foote, C.J. 1996. Evidence for sympatric genetic divergence of
703 anadromous and nonanadromous morphs of sockeye salmon (*Oncorhynchus*
704 *nerka*). *Evolution* **50**: 1265–1279.

705

Figures

Figure 1: Total nitrogen (top panel) and phosphorus (middle panel) loads in Alouette Reservoir from 1998 to 2015, separated into natural loading and mass added through the nutrient restoration program. Bottom panel shows the associated zooplankton biomass over the same years, broken up into copepods, daphnia and other cladocerans.

Figure 2: Median (lines) and 95% credible interval (shaded area) of predicted length-at-age (top panel) for nerkids in Alouette Reservoir from 1998-2015. Observed (circles) and predicted (lines and shaded area) abundance of age-0 (middle panel) and age-1 to -4 nerkids (bottom panel) in the hydroacoustic surveys from 1998-2015. Hydroacoustic surveys are assumed to be non-selective for age.

Figure 3: Log-transformed carrying capacity for nerkids in Alouette Reservoir from 1998-2015 for two stanzas: (top panel) egg to fry; and (bottom panel) fry to age-1.

Figure 4: Time series of vulnerable resident nerkid abundance (top panel) and anadromous nerkid returns (bottom panel) from 1998-2040. Posterior distributions of each are characterized by median estimates (solid line) and projections (dotted line) and 95% credible intervals (shaded area). Resident nerkid returns from 1998-2015 (dots) are observed data. Projections assume uniform distribution of heritability from 0 to 1.

Figure 5: Time series of vulnerable resident nerkid abundance (top panels) and anadromous nerkid returns (bottom panels) from 1998-2040. Posterior distributions are characterized as above. Projections assume heritability is known at either 0.2 (left panels) or 0.8 (right panels).

Figure 6: Projected abundance at year-25 of vulnerable of resident nerkids (A) and anadromous nerkid returns (B) after constant stocking at a particular rate. Solid line represents median predictions, shaded areas represent different credible intervals.

Draft

Table 1: Statistical catch-age model of the mixed resident-anadromous nerkid system including two-stage Beverton-Holt recruitment and density-dependent growth.

Parameters	
T1.1	$\theta = \left(L_{\infty,1}, K, L_0, c^{(1)}, c^{(2)}, c^{(3)}, p_1, p_2, p_3, \{N_{0,a}\}_{a=1}^{a=A}, M, F, \lambda, \tau, \xi, \psi, \chi, \{\varepsilon_t\}_{t=1}^{t=T} \right)$
T1.2	$\varphi = (a_e, b_e, L_{50(f)}, \sigma_{(f)}, m_3, \rho)$
Multi-stanza stock recruit variables	
T1.3	$\alpha_{(1)t} = \exp \left(-c^{(1)}p_1 - \frac{c^{(2)}p_2}{z_t} \right)$
T1.4	$\beta_{(1)t} = \frac{c^{(3)}p_3}{c^{(1)}p_1z_t^2 + c^{(2)}p_2z_t} (1 - a_{(1)t})$
T1.5	$\alpha_{(1+)t} = \exp \left(-c^{(1)}(1 - p_1) - \frac{c^{(2)}(1 - p_2)}{z_t} \right)$
T1.6	$\beta_{(1+)t} = \frac{c^{(3)}(1 - p_3)}{c^{(1)}(1 - p_1)z_t^2 + c^{(2)}(1 - p_2)z_t} (1 - a_{(1+)t})$
Initial population	
T1.7	$\hat{l}_{t=1,a} = \begin{cases} L_0 & a = 0 \\ L_0 e^{-Ka} + L_{\infty,1}(1 - e^{-Ka}) & a \geq 1 \end{cases}$
T1.8	$\hat{N}_{t=1,a} = N_{0,a}$
T1.9	$\hat{N}_{(sp)t=2,a} = \begin{cases} \frac{N_{0,a}\alpha_{(1+)1}}{1 + N_{0,a}\beta_{(1+)1}} & a = 1 \\ N_{0,a-1}e^{-M/2} & a \geq 2 \end{cases}$
State dynamics	
T1.10	$L_{\infty,t} = \frac{\tau + \lambda z_t}{1 + \xi \sum_{a=0}^A (N_{t,a} L_{t-1,a}^2)}$
T1.11	$\hat{l}_{t,a} = \begin{cases} L_0 & a = 0 \\ \hat{l}_{t-1,a-1} + L_{\infty,t}(1 - e^{-K}) & a \geq 1 \end{cases}$
T1.12	$v_{t,a} = \left[1 + \exp \left(-\frac{(\hat{l}_{t,a} - L_{50(f)})}{\sigma_{(f)}} \right) \right]^{-1}$
T1.13	$Z_{a,t} = M/2 + v_{t,a}F$

$$T1.14 \quad \hat{N}_{t,a} = \begin{cases} \frac{\hat{E}_{t-1}\alpha_{(1)t}}{1 + \hat{E}_{t-1}\beta_{(1)t}} e^{\varepsilon_t - 0.5\sigma_{(R)}^2} & a = 0; \varepsilon_t = N(0, \sigma_{(R)}) \\ (\hat{N}_{(sp)t,a-1} - \hat{s}_{t,a-1}) \frac{v_{t,a}F}{Z_{a,t}} e^{-Z_{a,t}} & 1 < a < A \\ \hat{N}_{(sp)t,a-1} \frac{v_{t,a}F}{Z_{a,t}} e^{-Z_{a,t}} & a = A \end{cases}$$

$$T1.15 \quad \hat{N}_{(sp)t+1,a} = \begin{cases} \frac{\hat{N}_{t,a}\alpha_{(1+)t}}{1 + \hat{N}_{t,a}\beta_{(1+)t}} & a = 1 \\ \hat{N}_{t,a}e^{-(M/2)} & 1 < a < A - 2 \\ \hat{N}_{t,a}(1 - m_3)e^{-(M/2)} & A - 1 \end{cases}$$

$$T1.16 \quad \hat{f}_{t,a} = a_e + \hat{l}_{t,a} \exp(b_e) \quad \{a = A - 1, A\}$$

$$T1.17 \quad \hat{E}_{(S)t} = \hat{f}_s \hat{N}_{(S)t} / 2$$

$$T1.18 \quad \hat{E}_t = (\hat{f}_3 \hat{N}_{t,3} m_3 + \hat{f}_A \hat{N}_{t,A}) / 2 + \hat{E}_{(S)t}$$

Fishery catch-at-age observations

$$T1.19 \quad \hat{C}_{t,a} = \hat{N}_{t,a} (1 - e^{-v_{t,a}F_t})$$

Process and observation error

$$T1.20 \quad \sigma_{(R)} = \rho\psi$$

$$T1.21 \quad \sigma_{(l)t,a} = \hat{l}_{t,a} c v_l$$

$$T1.22 \quad \tau_{(l)t,a} = \sigma_{(l)t,a}^{-2}$$

$$T1.23 \quad \sigma_{(H)a=0} = (1 - \rho)\psi\chi$$

$$T1.24 \quad \tau_{(H)a=0} = \sigma_{(H)a=0}^{-2}$$

$$T1.25 \quad \sigma_{(H)a=1+} = (1 - \rho)\psi(1 - \chi)$$

$$T1.26 \quad \tau_{(H)a=1+} = \sigma_{(H)a=1+}^{-2}$$

Table 2: Notation for the statistical catch-age models. Values of ++ indicate estimated parameters. Parameter descriptions include units in parenthesis.

Symbol	Value	Description
Indices		
t	$\{1,2,\dots,T\}$	Time step ($T=15$)
a	$\{1,2,\dots,A\}$	Age-class ($A=4$)
Model parameters		
$L_{\infty,1}$	++	Asymptotic length in first year (mm)
K	++	von Bertalanffy metabolic parameter (yr^{-1})
L_0	++	Age at age-0 (mm)
$c^{(1)}$	++	Base mortality parameter
$c^{(2)}$	++	Mortality per time spent foraging (N_{prey}^{-1})
$c^{(3)}$	++	Carrying capacity parameter (fish^{-1})
p_1, p_2, p_3	++	Proportion of $c^{(1)}, c^{(2)}, c^{(3)}$ attributed to first or second stanza
$N_{0,a}$	++	Initial abundance at age- a
M	++	Instantaneous natural mortality rate (yr^{-1})
F	++	Instantaneous fishing mortality rate (yr^{-1})
λ	++	Food-dependence parameter of density-dependent growth ($\text{mm } N_{\text{prey}}^{-1}$)
τ	++	Alternate food parameter of density-dependent growth (N_{prey})
ξ	++	Competition-dependence parameter of density-dependent growth (mm^{-2})
ψ	++	Total (observation + process) error
χ	++	Proportion of acoustic observation error attributed to age-0
a_e	-7.151	Egg-length multiplier (egg fl^{-1})
b_e	2.375	Egg-length exponent
$L_{50(f)}$	225	Length-at-50% vulnerability to angling
$\sigma_{(f)}$	20	Angling vulnerability variance
m_3	0.9	Proportion of kokanee mature at age-3
ρ	0.75	Proportion of total error attributed to process error
Sv_m	0.0025	Marine survival for sockeye from smolt to return
Derived variables		
$L_{\infty,t}$		Asymptotic length in year- t (mm)
$\hat{l}_{t,a}$		Length-at-age in year- t (mm)
$\hat{f}_{t,a}$		Fecundity at age in year- t (eggs/female)
$v_{t,a}$		Length-based vulnerability to angling in year- t

$\alpha_{(1)t}, \alpha_{(1+)t}$	Maximum survival to age-1 or -1+, respectively
$\beta_{(1)t}, \beta_{(1+)t}$	Carrying capacity parameter to age-1 and 1+, respectively (fish ⁻¹)
$Z_{a,t}$	Total fishing mortality at age in year- t
$\sigma_{(R)}$	Process error in age-0+ recruitment
$\sigma_{(H)a=0}, \sigma_{(H)a=1+}$	Observation error in age-0 and -1+ abundance
$\sigma_{(l)t,a}$	Observation error in length-at-age
$\tau_{(C)}$	Precision in observation error for annual catch
State variables	
$\hat{N}_{t,a}$	Number of age- a fish in fall of year- t
$\hat{N}_{(sp)t,a}$	Number of age- a fish in spring of year- t
V_t	Vulnerable abundance in year- t
$\hat{C}_{t,a}$	Catch at age in year- t
$\hat{E}_{(S)t}$	Sockeye eggs in year- t
$\hat{E}_t$	Total (sockeye + kokanee) eggs in year- t
Observations	
$I_{t,a}$	Hydroacoustic survey index in year- t for ages $a=0$ or $a=1-4$ combined
$C_{t,a}$	Catch of age- a fish in year- t in recreational fishery
$l_{t,a,i}$	Length-at-age in year- t for individual i
Management Controls	
x_t	Spring age-0 stocked fry descended from returning sockeye (fish)
Uncertain states	
h^2	Heritability of smolting

Table 3: Prior probability density functions (PDF) and posterior predictive distributions for each estimated parameter. Prior PDFs are described as either normal $N(\mu, \tau)$, uniform $U(\text{upper}, \text{lower})$, beta $B(\alpha, \beta)$ or gamma $G(\text{shape}, \text{rate})$ distributions. Posterior distributions for parameters are reported with mean and standard deviation with upper and lower 95% credible intervals.

Parameter	Prior PDF	Posterior PDF	95% Credible interval
$L_{\infty,1}$	$U(100, 1000)$	287.6 (7.0)	274.6-300.7
K	$U(0, 2)$	0.5 (0.0)	0.5-0.5
L_0	$N(60, 6)$	13.1 (3.4)	6.4-20.1
c_1	$U(0, 10)$	1.8 (6.4)	0.7-3.3
c_2	$U(0, 10)$	0.1 (0.1)	0.0-0.4
c_3	$U(0, 1)$	0.0 (0.0)	0.0-0.0
p_1	$B(2, 2)$	0.4 (0.2)	0.1-0.7
p_2	$B(2, 2)$	0.6 (0.2)	0.1-0.9
p_3	$B(2, 2)$	0.5 (0.2)	0.1-0.9
$N_{0,0}$	$U(0, 500,000)$	33,164.4 (12,258.4)	15,297.1-63,786.8
$N_{0,1}$	$U(0, 500,000)$	14,062.9 (9,033.4)	999.7-33,249.2
$N_{0,2}$	$U(0, 500,000)$	5,701.3(4,682.9)	161.2-17,468.1
$N_{0,3}$	$U(0, 500,000)$	6,399.4 (4,677.5)	313.0-17,543.3
$N_{0,4}$	$U(0, 500,000)$	20,385.6 (4,370.1)	12,060.2-29,147.2
M	$N(0.6, 0.6)$	0.9 (0.1)	0.6-1.1
F	$U(1, 1)$	0.1 (0.1)	0.0-0.3
λ	$U(0, 50)$	16.8 (12.3)	0.6-45.1
τ	$U(0, 10,000)$	751.1 (131.4)	549.3-1,019.8
ξ	$U(0, 10)$	6.4 (1.8)	3.4-9.8
ψ	$U(0, 10)$	2.8 (0.3)	2.3-3.4
χ	$B(1, 1)$	0.5 (0.1)	0.4-0.6
cv_l	$B(1, 1)$	0.1 (0.0)	0.1-0.1
τ_C	$G(0.01, 0.01)$	0.0 (0.0)	0.0-0.0

Table 4: Calculation of phenotypic response to selection for smolting and resulting sockeye returns in mixed kokanee-sockeye system.

Phenotypic expression	
T4.1	$Sm^H = p_{(smolt)t-4} + h^2(1 - p_{(smolt)t-4})$
T4.2	$Sm^K = p_{(smolt)t-4} + h^2\left(\frac{E_{(S)t-2}}{E_{t-2}} - p_{(smolt)t-4}\right)$
Smolts and sockeye returns	
T4.3	$\hat{s}_{t,1} = (Sm^H x_t + Sm^K N_{(sp)t,1}) \frac{N_{(sp)t,1}}{N_{t-1,1}}$
T4.4	$\hat{N}_{(S)t} = \hat{s}_{t-2,1} S v_m$

Draft