DNA barcodes reveal deeply neglected diversity and numerous invasions of micromoths in Madagascar

Journal:	<i>Genome</i>
Manuscript ID	gen-2018-0065.R2
Manuscript Type:	Article
Date Submitted by the Author:	17-Jul-2018
Complete List of Authors:	Lopez-Vaamonde, Carlos; Institut National de la Recherche Agronomique (INRA), ; Institut de Recherche sur la Biologie de l'Insecte (IRBI), Sire, Lucas; Institut de Recherche sur la Biologie de l'Insecte Rasmussen, Bruno; Institut de Recherche sur la Biologie de l'Insecte Rougerie, Rodolphe; Institut Systématique, Evolution, Biodiversité (ISYEB), Wieser, Christian; Landesmuseum für Kärnten Ahamadi, Allaoui; University of Antananarivo, Department Entomology Minet, Joël; Institut de Systematique Evolution Biodiversite deWaard, Jeremy; Biodiversity Institute of Ontario, University of Guelph, Decaëns, Thibaud; Centre d'Ecologie Fonctionnelle et Evolutive (CEFE UMR 5175, CNRS–Université de Montpellier–Université Paul-Valéry Montpellier–EPHE), , CEFE UMR 5175 CNRS Lees, David; Natural History Museum London
Keyword:	Africa, invasive alien species, Lepidoptera, Malaise trap, plant pests
Is the invited manuscript for consideration in a Special Issue? :	7th International Barcode of Life

DNA barcodes reveal deeply neglected diversity and numerous invasions of micromoths in
Madagascar

Carlos Lopez-Vaamonde^{1,2}, Lucas Sire², Bruno Rasmussen², Rodolphe Rougerie³,
Christian Wieser⁴, Allaoui Ahamadi Allaoui ⁵, Joël Minet³, Jeremy R. deWaard⁶, Thibaud
Decaëns⁷, David C. Lees⁸

¹ INRA, UR633, Zoologie Forestière, F- 45075 Orléans, France.

² Institut de Recherche sur la Biologie de l’Insecte, UMR 7261 CNRS Université de Tours, UFR
Sciences et Techniques, Tours, France.

³Institut de Systématique Evolution Biodiversité (ISYEB), Muséum national d'Histoire naturelle,
CNRS, Sorbonne Université, EPHE, 57 rue Cuvier, CP 50, 75005 Paris, France.

⁴Landesmuseum für Kärnten, Abteilung Zoologie, Museumgasse 2, 9020 Klagenfurt, Austria

⁵ Department of Entomology, University of Antananarivo, Antananarivo 101, Madagascar

⁶ Centre for Biodiversity Genomics, University of Guelph, 50 Stone Road E., Guelph, ON
N1G2W1, Canada

⁷Centre d'Ecologie Fonctionnelle et Evolutive (CEFE UMR 5175, CNRS–Université de
Montpellier–Université Paul-Valéry Montpellier–EPHE), 1919 Route de Mende, F-34293
Montpellier, France.

⁸Department of Life Sciences, Natural History Museum, Cromwell Road, SW7 5BD, UK.

Email for correspondence: carlos.lopezvaamonde@inra.fr

Abstract

Madagascar is a prime evolutionary hotspot globally, but its unique biodiversity is under threat, essentially from anthropogenic disturbance. There is a race against time to describe and protect the Madagascan endangered biota. Here we present a first molecular characterization of the micromoth fauna of Madagascar. We collected 1572 micromoths mainly using light traps in both natural and anthropogenically disturbed habitats in 24 localities across eastern and northwest Madagascar. We also collected 1384 specimens using a Malaise trap in a primary rain forest at Andasibe. In total, we DNA barcoded 2956 specimens belonging to 1537 Barcode Index Numbers (BINs), 88.4% of which are new to BOLD. Only 1.7% of new BINs were assigned to species. Of 47 different families found, Dryadaulidae, Bucculatricidae, Bedelliidae, Batrachedridae and Blastobasidae are newly reported for Madagascar and the recently recognized Tonzidae is confirmed. For test faunas of Canada and Australia, 98.9-99.4% of Macroheterocera BINs exhibited the molecular synapomorphy of a Phenylalanine in the 177th complete DNA barcode codon. Non-macroheteroceran BINs could thus be sifted out efficiently in the Malaise sample. The Madagascar micromoth fauna shows highest affinity with the Afrotropics (146 BINs also occur in the African continent). We found 22 recognised pests or invasive species, mostly occurring in disturbed habitats. Malaise trap samples show high temporal turnover and alpha diversity with as many as 507 BINs collected; of these, astonishingly, 499 (98.4%) were novel to BOLD and 292 (57.6%) were singletons. Our results provide a baseline for future surveys across the island.

Key words

Africa, invasive alien species, Lepidoptera, Malaise trap, plant pests

Introduction

Madagascar is one of the top priority global hotspots for biodiversity conservation with high endemicity and under large anthropogenic pressure (Vences et al. 2009). There is an urgent need to describe what remains of the unique biota of Madagascar so as to locate hotspots of biodiversity and endemism and protect them. Conservation efforts in Madagascar are mainly focused on vertebrates (Herrera 2017; Jenkins et al. 2014) and plants (Royal Botanic Gardens Kew 2016). Arthropods are rarely taken into account in conservation in Madagascar, despite the fact that many species are micro-endemics at greatest risk of extinction (Danieleczak et al. 2017; Wesener & Rudolf 2017; Wesener et al. 2014).

With up to 4900 described species currently listed from Madagascar (Viette 1990; Krüger 2007; Lees & Minet 2003; Libert 2014; Lees 2016; De Prins & De Prins 2018), the order Lepidoptera (moths and butterflies) is a significant component of the arthropod biota. Since lepidopterans have been widely used as bioindicators of habitat disturbance (Kremen 1994; Enkhtur et al. 2017, Hawes et al. 2009), they could provide a strong signal for conservation efforts and priorities. Unfortunately, Madagascan Lepidoptera are relatively poorly known, particularly the 'micromoths', a polyphyletic group excluding Macroheterocera and butterflies (Lees et al. 2003) of about 1600 described species (Viette 1990; De Prins & De Prins 2018), with many species yet to be described (Lees & Minet 2003). Biodiversity assessment studies rarely take into account micromoths because of the difficulty in identifying them, for a general lack of taxonomic expertise, and the need for specialised technical skills for specimen mounting and dissecting. The use of DNA barcoding, however, has proved an efficient and affordable method to alleviate this taxonomic impediment. Operational taxonomic units derived from DNA barcodes can accurately and objectively represent species diversity and then be used to survey micromoth diversity in poorly known and hyperdiverse areas of the World (Lees et al. 2013; Miller et al. 2016).

The Barcode of Life Datasystem (BOLD; www.boldsystems.org; Ratnasingham and Hebert 2007) now contains over six million DNA barcodes and represents a huge resource to accelerate identification and quantify biodiversity. However, the coverage for the Madagascan lepidopteran fauna is very sparse. Nevertheless, the use of Barcode Index Numbers (BINs) (Ratnasingham and Hebert 2013) as proxies for species allows the assessment of hyperdiverse groups that are taxonomically poorly known, such as micromoths (Schmidt et al. 2017; Aagaard et al. 2016; Lees et al. 2013; Lopez-Vaamonde et al. 2012). As of 29th June 2018 (including the current study), there were 2852 DNA barcode BINs for Madagascar out of a total of 113,161 lepidopteran BINS, according to a search of the BIN Database in the public portal of BOLD. Nieuwerkerken et al. (2011) estimated 157,424 described species of Lepidoptera, and the upper bounds for true richness may be as much as half a million species (Solis and Pogue 1999). Very few of all these BINs representing Madagascan Lepidoptera are yet publicly identified on BOLD to described species. As of 30th March 2018 there were only 287 publicly released species names according to the BIN portal of BOLD, of which only 277 had correctly composed names; 173 represented Macroheterocera, 77 represented butterflies and only 27 represented micromoths, 24 of which were Tortricoidea and Pyraloidea. – Furthermore, only 201 of these species had BIN numbers allocated. Progress in DNA barcoding the described fauna of Madagascan Lepidoptera lags thus far behind most countries. The first implementation of the Global Malaise Program in Madagascar (Bio-Inventory and Collections Unit, Biodiversity Institute of Ontario, 2015) provides a local instance where identification of Lepidoptera samples below Order level is problematic by external morphology (Lepidoptera wings being poorly preserved) or very time consuming by individual sequence queries. We asked if a previously observed simple molecular synapomorphy in the DNA barcode (Lees et al. 2011) was reliable enough to filter out the clade Macroheterocera from such samples.

97 From a biogeographic point of view, Madagascar has a very unbalanced or disharmonic fauna,
98 with some taxa overrepresented and some underrepresented relative to the mainland source area
99 (Briggs 1987). Indeed, the Madagascan fauna is characterised by a significant number of large
100 endemic radiations such as lemurs and tenrecs now extinct on mainlands (Poux et al. 2005) and a
101 large number of major continental lineages that appear not to have established at all on the island
102 (the lack of poritiine lycaenids which are highly diverse in Africa is evident: Lees et al. 2003).
103 The lepidopteran fauna of Madagascar is, in particular, quite dissimilar to that of southern Africa,
104 much more so than the relatively more harmonic fauna of the neighbouring island fauna of La
105 Réunion (Krüger 2007). Southern Africa has twice as many described lepidopteran species
106 described as Madagascar, while Noctuoidea is overrepresented in Madagascar. By contrast,
107 “primitive” Lepidoptera (defined as consisting of the non-ditrysian grade of micromoths that
108 includes groups from Micropterigoidea to Tischerioidea; Krüger 2007), as well as Tineoidea and
109 Gelechioidea are, in particular, underrepresented in Madagascar. However, these general
110 faunistic patterns are based on current checklists, which are particularly incomplete for the
111 Madagascan lepidopteran fauna and also biased towards the best-studied families (for example,
112 Viette specialized on the noctuid fauna of both Madagascar and La Réunion: Viette, 1963, Viette
113 1965 and Viette 1967).

114 Finally, many microlepidopteran species are highly invasive and serious pests of agricultural and
115 ornamental plants (Lopez-Vaamonde et al. 2010). Despite their potential economic and
116 ecological impact, there is limited information available on invasive insects in Madagascar
117 (Fisher et al. 1998; Kull et al. 2014; Irwin et al. 2010).

118 The main aims of our study were: 1) to carry out a survey of micromoth diversity using DNA
119 barcodes across several sites in Madagascar from disturbed to primary rainforests using DNA
120 barcodes; 2) to identify any molecular synapomorphy(ies) within the DNA barcode fragment that

would allow us to more accurately identify samples and to better evaluate sequence queries where external morphology was problematic; 3) to characterize as far as possible the biogeographic origins of the Madagascan microlepidopteran (based purely on proximity, a predominantly African mainland affinity would be expected); 4) to identify the presence of any cosmopolitan, invasive, agricultural and forestry pest species, which should be more prevalent in disturbed habitats than in well preserved ones.

Material and Methods

Specimen collection

Micromoths were collected in non-protected areas by one of us (CW) between October-November 2013 and March 2015. CW used two to three light towers with 15W ultraviolet fluorescence actinic tubes (www.bioform.de) operated with lithium batteries (Li-Ion Akku HELLPOWER 12V/10.5Ah 116.60Wh). Micromoths were sampled from nine collecting sites in disturbed habitats around the Nosy Be area (northwestern Madagascar) (Table 1). All these specimens are deposited at the Natural History Museum of Carinthia (Austria).

Specimens were also collected largely within protected areas across eastern Madagascar by another of us (DCL) with 160W blended tungsten/mercury-vapour lamps or 15W actinic lights (Bioquip) powered with a generator (Honda EX350) (lights suspended on a white sheet with a protective transparent tarpaulin), sampled in November-December 2011, January-February 2014, and November 2014. All specimens collected by DCL are deposited at the Natural History Museum in London.

One Townes-style Malaise trap (standard for the Global Malaise Trap Program, Geiger et al. 2016) was set up by another of us (AA) in two sites of PN Andasibe-Mantadia, specifically the forest originally designated as the Réserve Spéciale d'Analamazaotra (for short, we refer

hereafter to this reserve as its current popular name “Andasibe”; it was also popularly known as Perinet). This is a c. 810-hectare fragment of the once far larger Analamazaotra rainforest (Table 1). One site was sampled during 65 days at the end of the wet to beginning of dry season (from April 1st until 28th May 2014) (M1) at 1000 m elevation and a second site, 0.8 km away from the first site at 1050 m, was sampled during 67 days at the end of dry to beginning of wet season (from September 1st until 6th November 2014) (M2) (elevations adjusted for coordinates in Table 1 using Google Earth). Each sample was collected in a 500 ml plastic Nalgene bottle that was filled with 375 ml of 95% ethanol and then attached to the trap head. The catch was harvested weekly by AA and brought to the University of Antananarivo where the bulk ethanol was replaced with fresh 95% ethanol before storage at -4°C until samples were drained and sent to the Centre for Biodiversity Genomics in Canada (CBG; www.biodiversitygenomics.net).

DNA barcoding

We DNA barcoded in most cases only one specimen per morphospecies for light-trapped and day-netted specimens. Morphospecies were defined using external morphology, mainly wing pattern. DNA was extracted using hind legs of pinned specimens or entire body extracts in the case of smaller Malaise-trapped Lepidoptera. DNA barcodes (658 bp of the COI mitochondrial gene) were generated using traditional Sanger sequencing at the CBG using standard high-throughput protocols (Ivanova et al. 2006).

Malaise trap samples were also processed at CBG as part of the Global Malaise Program (<http://biodiversitygenomics.net/projects/gmp/>) following the protocol described in deWaard et al. (2017), which involves unidirectional sequencing, so those sequences are usually shorter than 658 bp. Larger moths were pinned, smaller ones kept in their original wells. A randomly selected

example of each BIN was imaged at Guelph; as usual for the order Lepidoptera captured via this method, these samples tend not to be in good condition for external morphological analysis.

DNA sequences, along with the voucher data, images, and trace files, are deposited in the Barcode of Life Data Systems (BOLD v4) (Ratnasingham and Hebert 2007, www.barcodinglife.org), and the sequences were deposited in GenBank. All data are available in BOLD through the public dataset: DS-MICROMA (dx.doi.org/10.5883/DS-MICROMA).

To aggregate barcodes of the polyphyletic group micromoths (which includes some larger moths such as thyridids) from the Malaise trap data set, we asked if a previously noticed molecular synapomorphy for the clade Macroheterocera (Lees et al. 2011) was reliable enough to partition out all non-macroheterocerans. To do this we used a test dataset of two well-identified lepidopteran faunas, namely that of Australia (n=14965 BINs analyzed) and Canada (4684 BINs analyzed).

In the case of the Malaise trap sample, which had been predetermined to Lepidoptera before sequencing, we first filtered out all Papilionoidea (butterflies), which could be verified by batch queries on BOLD because all genera and most species had already been DNA barcoded.

To determine the number of BINs novel for this study for BOLD, we derived the number of uniques and non-uniques from the dataset front page “Data Summary”. However, we subtracted 36 BINs that were currently reported as private data to the CBG from the reported list of non-uniques. These data we inferred to be additional members from the Malaise trap not integral to our project as was derived from project container GMTAD (Global Malaise Programme Madagascar Malaise 2014).

Data analyses

Diversity analyses were carried out on both Malaise and light trap samples. Community analyses were performed only on Malaise samples from Andasibe since it is the only site for which we have abundance data.

Data analyses were done with R ver.3.4.3 (R Development Core Team 2004) using different packages for community and species richness analyses.

iNEXT (Chao et al., 2014, Hsieh et al. 2016) allowed us to calculate α -diversity and generate accumulation curves using 50 resampling replicates with replacement (Chao et al 2014). We used BINs as species proxy (Ratnasingham and Hebert 2013) and plotted them against both the sampling coverage (measure of sample completeness that estimates the proportion of the total number of individuals in a community that belong to the species represented in the sample) and the total number of caught individuals taken as a measure of sampling intensity. We ran the analyses for late wet to early dry (M1) and late dry to early wet (M2) seasons, both covered by the sampling at Andasibe.

Abundance Coverage Estimator (ACE) (Gotelli & Colwell 2010) and Chao1 (Chao et al. 2009) are two other diversity indices that were calculated with the package *Vegan* ver. 2.4-6 (Oksanen et al. 2016) in order to estimate the potential species richness in accordance with the sampling intensity.

We carried out a distributional data analysis by extracting from BOLD a list of all countries for which each BIN has been barcoded. Each appearance of a BIN per country was assigned to a biogeographical region (Afrotropical, Australasian, Nearctic, Neotropical, Oriental and Palearctic) by looking at the corresponding countries associated to the records in BOLD. Each BIN was counted only once per region but might be spread over multiple ones.

Specimen identification

Specimens were identified by both external morphology (without dissection) and by using DNA barcode queries using all data present in BOLD. For each sequence we queried, we used the “Current Database” and the “Search Database” query option “All Barcode Records” on the Identification Engine of BOLD and then we built a NJ tree in BOLD (=“Tree-based Identification”) to find the nearest neighbour. Then we searched for the minimum corresponding p-distance(s) in the list of 99 top hits. We looked in particular for interspecific query tails among the hit list that seemed informative, i.e. with the hit(s) showing potential signal standing proud of the noise of background hits (often evident as the sequence Similarity value directly preceding the sharpest inflexion in the Similarity Score graph before it starts to asymptote), or otherwise stated 'Non-informative' under Taxonomy Notes. We took particular note when nearest hits derived from apparent local radiations. We also considered amino acid information, in particular ignoring Similarity values for irrelevant hits inside or outside of the Macroheterocera (see below), and also looking qualitatively at unusual codon changes shared between taxa as shown in Conservation plot mode against a reference sequence in Bioedit v7. In most cases, the sequence divergence(s), to the nearest identified BINs on BOLD, expressed as 100-Similarity, are noted under "Taxonomy notes", particularly for records from the BOLD projects MADAM and MIMAD. In that field, we were often able to specify closely related BIN numbers by building a corresponding Image database for the Tree Based Identification query. Where relevant hits existed, we assigned species-level identifications for low pairwise divergences expressed as 100-Similarity (<approx. 2%) but if not, increasing taxonomic ranks where further hits showed taxonomic consistency in the NJ tree. Since application of strict thresholds may generally be misleading particularly for supraspecific ranks, and since no support levels are specified on BOLD NJ trees, we also used independent ML analyses in PHYML 3.0 (Guindon et al. 2010) to identify barcoded specimens by examining their phylogenetic position within a clade containing

identified individuals, some of which were downloaded from BOLD, at the best justifiable taxonomic rank. In Phyml 3.0, we used default options except: GTR (or automatic model selection), all parameters estimated, and SPR. In general, we looked for ABayes support levels >0.94 to assume nestedness within a clade. Identifications from the light-trapped and day-netted samples run alongside the malaise samples in an ML analysis helped the identifications of Malaise samples. We specified the identification method(s) or combination thereof (e.g. External morphology, COI-5P (NJ), COI-5P (ML), COI-5P (codons) i.e. amino-acid based identification) under the field Identification Method.

We compared the 1572 light-trapped moths and day-netted moths with specimens, including where possible, accessible types, deposited in the two most important reference collections of Madagascan Lepidoptera, namely the Muséum national d'Histoire naturelle (MNHN, Paris) and the Natural History Museum (NHMUK, London), and to illustrations in reference works. We have not attempted an exhaustive type comparison with our specimens and anticipate that more matches will come to light as the collections are digitised and/or as DNA sequencing of the types is attempted.

Results

DNA barcodes and identification rates

We successfully barcoded 2956 micromoth specimens (1572 light-trapped and day-netted moths and 1384 micromoths collected with the one Malaise trap) belonging to 1537 BINs (six of 2956 samples do not qualify as full barcodes and so lack BIN numbers). Those 1537 BINs belonged to 44 families as currently classified in BOLD (see Table 2, where 47 family-level groupings are specified; these include families currently lumped on BOLD). 32.7% of BINs (503 out of 1537 BINs) were identified to genus level and 6.2% of BINs (95 out of 1537 BINs) were identified to

species level. Many of those identified BINs correspond to well-known cosmopolitan species more likely to have been DNA barcoded elsewhere (Table 3).

88.4% of BINs (1358 out of 1537 BINs) obtained are new to BOLD and only 179 BINs (13.2%) were already in the BOLD database.

By analysing two barcoded lepidoptera faunas from Canada and Australia we found that almost all Macroheterocera indeed show a Phenylalanine rather than Leucine or other character state in the 177th complete codon (5'→3') of the (up to) 658 bp nucleotide sequence.

For the Australian fauna 4093 BINs (99.4%) exhibiting a Phenylalanine in the 177th position pertain to sequences identified as macroheteroceran families, while 11 exceptions belong to the genus *Aristeis* (Oecophoridae) and one to another Oecophoridae genus. Five others belong to Crambidae: Acentropinae, one to Crambinae, two to Lecithoceridae, two to Gelechiidae, two to Tineidae: Harmacloninae and one to Heliozelidae. Exceptions to the reliability of this synapomorphy (total n=26, discounting an apparently contaminated Lycaenidae) are not only rare in general, but phylogenetically also very narrowly represented. Also, true conversely for this dataset, 99.5% of 9070 BINs exhibiting another state than a Phenylalanine in that position (usually Leucine) are identified as belonging to non-macroheteroceran families, including those of butterflies. Of the exceptions (n=48), 21 belong to Oenosandridae and three to Nolidae, while three of seven Geometridae, three of eight Erebiidae and three of seven to Noctuidae seem correctly identified (the rest are micromoths from images), while one imaged “Saturniidae” also represents a micromoth. For the Canadian Lepidoptera fauna (4684 BINs analysed), the presence of a Phenylalanine in this position is 98.9% reliable as a surrogate for Macroheterocera (99.75% reliable when excluding Crambidae: Acentropinae and Tineidae: Meessiinae), while presence of other character states is 99.8% reliable for non-macroheteroceran Lepidoptera (exceptions one

geometrid, one nolid and two noctuids). We did not detect any cases of such parallelisms in our Madagascan dataset, suggesting that the synapomorphy was fully reliable for this fauna, but in the case of filtering of the malaise sample, Macroheterocera were only represented by 170 BINs (whereas Papilionoidea by 20 BINs).

Within the 507 BINs collected with the Malaise trap, 50 BINs (9.9%) have been identified to genus and only three BINs (0.59%) have been identified to species level (*Angustalius malacellus*, *Bradina admixtalis* and *Lobesia aeolopa*). The only other five BINs already on BOLD were a cosmopterigid *Stilbosis* sp. (BOLD:ABY7721, Kenya), a spilomeline (BOLD:ACT8113, South Africa), two tortricids and another spilomeline from Ranomafana (*Pandemis* sp. BOLD:ACO0519; *Olethreutes* sp. BOLD:ACS0054 and *Herpetogramma* sp. BOLD:ACT6691). All remaining 499 BINs, 270 of which singletons and 109 doubletons, are at present only known as endemic to Andasibe. Two BINs (BOLD:ACS0229, an *Elachista* and BOLD:ACS1392, Tineidae: Hieroxestinae) have more than 70 individuals ($n=75$ and 91 , respectively) but even these abundant taxa have as yet no species name (Fig. 3).

A total of 113 BINs were not identified to family level. According to NJ building on BOLD and/or external morphology of pinned specimens, these 113 BINs were overwhelmingly dominated by possible or probable Gelechioidea (>77%) which could not be reliably assigned at present to family. Around 16% may represent tineoids while superfamily was unassigned even tentatively for 6%. Maximum divergences among all of those unidentified BINs to any other BIN was no smaller than 14.3%. Over 40% of those unknown BINs were more closely related to one or more unidentified Madagascan BINs than to BINs outside Madagascar.

Only 23 BINs were shared between the samples collected with Malaise (507 BINs) and those collected with light trapping + netting by day (1053 BINs) (Table 2). Of the non Malaise-trapped material, approximately 92% were light-trapped and the remainder netted by day, so there was a

strong bias towards nocturnal activity. The low number of shared BINs is particularly striking for tineids, considering the high diversity of this family (95 BINs collected with Malaise and 65 BINs collected by other methods), with only one BIN shared (Table 2).

Some groups were much more strongly represented in Malaise samples such as Nepticulidae, Tineidae, Immidae, Lecithoceridae and Elachistidae. Other families were much better represented in the mainly light-trapped samples than in the Malaise, notably Gracillariidae, Tortricidae, Cosmopterigidae, Gelechiidae, Pyralidae, and Crambidae (Table 2).

Taxonomic composition and biogeographical distribution

Fig. 1 shows the difference in distribution of BINs per family between light trap and Malaise trap samples. The three families with highest number of BINs within the 507 BINs collected with Malaise traps are: Tineidae (95 BINs, 18.7%), Depressariidae s.l. ('Peleopodidae': Oditinae) (54 BINs, 10.7%), and Lecithoceridae (51 BINs, 10.1%). Within the 1053 BINs collected with light-traps and netted by day the three most representative families are: Gelechiidae (145 BINs, 13.8%), Crambidae (139 BINs, 13.2%) and Tortricidae (132 BINs, 12.5%).

Of the up to 47 different micromoth families found, Dryadaulidae, Bucculatricidae, Bedelliidae, Batrachedridae and Blastobasidae are newly reported for the island (Table 2). Other families, namely Micropterigidae, Opostegidae, Tonzidae and Eriocottidae, have been previously reported from Madagascar, but have no described species there (Krüger 1997; Lees & Minet 2003; Davis & Stonis 2007; Gibbs 2016; Kobayashi et al. 2018).

The analysis revealed that 55 BINs show a widespread distribution over more than one biogeographical region (Table 3). Out of the 162 BINs shared between Madagascar and other biogeographical regions, 146 BINs (90.1%) occur in Africa and 105 are found only in the Afrotropical region. More surprisingly, 49 BINs (30.3%) detected in Madagascar also occur in

Australasia, 29 BINs (17.9%) in the Oriental region, 27 BINs (16.7%) in the Palearctic, 18 BINs (11.1%) in the Neotropics, and 17 BINs (10.5%) in the Nearctic (Fig. 2).

Invasive and pest species

Of the above 55 BINs that show a widespread distribution occurring outside the Afrotropical region, at least 40% (22 out of 55) are known to be pests and/or invasive somewhere in their distribution range, while at least an additional five species occasionally feed on crops or may be minor pests. At least 50.9% (28 out of 55) are recorded for the first time in Madagascar (Table 3).

All widespread BINs are identified to species level except nine. These included a tineid (BOLD:ACS7592); a glyphipterigid (BOLD:AAY2216) previously barcoded from Australia but 1.7% divergent; a tortricid unidentified to genus (BOLD:ACS7628), one cosmopterigid of the genus *Gisilia* (BOLD:ACS6187), one *Ascalenia* (BOLD:AAG0134), two crambids of the genus *Herpetogramma* (BOLD:ACD5135 and BOLD:AAB6841), a gracillariid of the genus *Stomphastis* (BOLD:AAM6667) with barcodes from Australia (also currently without associated species name) and a pterophorid of the genus *Stenoptilia* (BOLD:AAD0716) with barcodes from Africa, Asia and Australia (Table 3).

38.2% (21 out of 55 BINs) of widespread BINs belong to the family Crambidae, a family known for many highly dispersive species (Lopez -Vaamonde et al. 2010).

Species richness and turnover

The analysis of 1384 microlepidopterans (three of which are without BIN allocations) collected over 16 weeks of Malaise trapping revealed a total of 507 BINs. Astonishingly, nearly all (499 BINs or 98.4%) were novel to BOLD given an also surprisingly small overlap (4.5% of the Malaise sample, 2.2% of others and 1.5% of the total sample) with the principally light-trapped

samples in this study. Moreover, 57.6% (292 out of the 507 Malaise BINs) are represented by singletons (i.e. by single individual) in our data set (Fig. 3). The high number of singletons demonstrates that even 16 weeks and two seasons are entirely inadequate to sample with Malaise traps most of the species that must be present in the studied area.

The Malaise trap automatically collected a total of 335 micromoths (representing 165 BINs) at the end of wet to beginning of dry season (M1), whereas 1046 individuals (representing 404 BINs) were collected at the end of dry to beginning of wet season (M2).

The BINs shared between Malaise trap samples collected at M1 and M2 are only 12.2% (62 BINs out of 507 BINs) but with many individuals (528 specimens out of 1384 individuals collected, 38.1%). Therefore many of the species collected during the two periods belong to relatively common species.

Fig. 4 shows a clear temporal turnover with a strong relationship between temporal distance of samples and amount of species overlap. For each site (M1 and M2) taken separately and compared, a clear decline in sample overlap with temporal distance is evident. The intercept for M2 is higher than M1, but slopes, p-values, and R-squared values are very similar.

Rarefaction curves show that both species diversity and sampling coverage indices were perhaps surprisingly higher at end of wet towards early dry season (M1) than at the end of the dry towards early wet season (M2) (Figs. 5a and 5b). They also show that 16 weeks of Malaise sampling is not nearly enough to capture all the Lepidoptera diversity in the studied area (Fig. 5c). We collected with one Malaise trap 507 BINs at Andasibe, whereas both non parametric indices, Chao 1 index and ACE suggest that at least twice as many species could occur in the studied area (Fig 6).

Discussion

383 *Massive 'Linnean shortfall' of micromoths in Madagascar*

384 The majority of BINs (1358 of 1537) found in our study are new to BOLD and most of them
385 remain unidentified to species level, as we only recovered 1.7% (23 out of 1358 BINs) of
386 corresponding taxonomic assignation (and several of these were assigned using external
387 morphology only since new to BOLD). Remarkably, despite the impressive lepidopteran
388 coverage on BOLD, we do not know the names of some of the most abundant species in
389 Madagascan ecosystems, and without comprehensive barcoding of museum collections, it is
390 difficult to be sure what names may already be available for them.

391 115/507 BINs (22.7%) in the Malaise trap were identified to five families within Tineoidea, one
392 of which is newly reported for Madagascar (Fig. 1). Among the non-tineoid Ditrysia, exceptional
393 diversity was found among Gelechioidea identified to families, which with 181 BINs (208
394 including Gelechioidea *incertae sedis*) form 35.7% (41%) of the entire Malaise sample. Of these,
395 Lecithoceridae (with 51 BINs by Malaise) was the richest family, while elsewhere in the
396 Depressariidae assemblage (Depressariinae s.l. on BOLD), the Malaise trap sampled a large
397 diversity of the local radiation of "Oditinae" (54 BINs) and Oecophoridae only had nine BINs,
398 most of these in *Metachanda*. This depressariid assemblage (Sohn et al. 2015), not yet adequately
399 sorted at family level but probably including numerous Peleopodidae, comprise a high proportion
400 of leaf litter detritivores (this provisional classification, including Gelechioidea *incertae sedis*, as
401 well as 'Stenomatidae' and Lecithoceridae, is included in Table 2). In the Malaise trap, we also
402 found eight BINs of Elachistinae (Elachistidae), a leaf mining group reported by Lees & Minet
403 (2003) but with only one reported (Parenti, 2006) and one undescribed (Lees and Minet 2003:
404 751) Madagascan species (De Prins and De Prins 2018 duplicate *Pauroptila* in Parametrioninae,
405 but it is here placed in Cosmopterigidae; see also Koster and van Nieukerken, 2018). In the
406 Malaise trapped Gelechiidae, Dichomeridinae with 27 BINs clearly form another significant local

radiation. The Malaise trap evidently captures diurnal as well as nocturnal species, and the use of such a passive and stationary sampling method allowed us to recover three families not detected by light trapping and a much better diversity of some local radiations. The Malaise sampling, however, was clearly limited in finding 27 rather than 45 families (using the expanded Depressariidae classification in Table 2), but the light trapping and day netting encompassed a greater geographic range and number of sampling sites.

The large differences in taxonomic composition between the two main collecting methods could be explained by differing geography, sampling times and human vs malaise collecting bias. Indeed, each method, of course, has its own inherent strong taxonomic biases, but the Malaise trap was largely free of human bias and its proportions reflect its passively sampled abundances. Indeed, Malaise trapping is likely to be the most unbiased method, since we did not control for all the possible biases (location, time of day, weather conditions, local flora, wind, etc.) that may have affected our acquired sample composition, in particular for light trapping.

The low number of “primitive” Lepidoptera in Madagascar reported by Krüger (2007) is probably an artefact of insufficient sampling. The only primitive non-ditrysians found in the Malaise trap samples were Nepticulidae (21 BINs with an additional seven BINs, all at light), a family with only one described species (*Fomoria scobleella* (Minet, 1990)) in Madagascar. Other non-ditrysians found by other methods (such as adelids) were hand netted by day (apart from one *Nemophora* sampled at light). The micropterigids (two BINs), a group also known to enter Malaise traps, were also hand netted by day by one of us (DCL), but it is likely that the Malaise trap did not intersect with their particularly narrow flight phenology. It may be that such lineages as adeloids actually need special sampling techniques and habitat surveys for their detection. Interestingly, we also noticed the absence of heliozelids, which does not fit field observations of their leaf-mines in Madagascar, and the fact that one species, *Antispila merinaella* Viette, [1956],

has been already described. The diverse primitive moth group Exoporia, that includes the Hepialidae, has not yet been detected in Madagascar, but are exceptionally depauperate in tropical Africa (in fact unknown there in tropical rainforests) and so might be really absent in Madagascar or present in poorly sampled habitats. Targeted samplings, such as plant internal feeder rearings, leaf mine and gall collections, soil and periphyton layer analysis, as well as vigorous netting for moths by daytime during peak emergence months synchronized to rains may increase the probability of discovering such primitive groups on the island.

These observations, reinforced by the apparent high seasonal turnover found in our study, demonstrate the paucity of knowledge regarding Madagascan lepidopteran diversity, notably for micromoths, and its true taxonomic makeup. They also highlight the lack of progress in sequencing identified and unidentified museum collections, as well as the extreme shortfall in documentation of true diversity of the fauna resulting from what remains to be found in the wild and then described. The actual number of public BINs on BOLD for Madagascan Lepidoptera (2852 BINs) relative to all described Madagascan species (~4900 species) is about 58.2%. This may seem like substantial progress, but a high percentage of those BINs are unlikely to correspond to checklists of Madagascan Lepidoptera. Most micromoths in Madagascar must be undescribed: the ratio of micromoths to Macroheterocera in the French checklist, for example, is about 2.1 and if such a ratio were to hold for Madagascar too, where ~ 3000 described Macroheteroceran species are currently known, one would expect to find at least 6300 micromoths. The paucity of well identified moth DNA barcode clusters highlights the gap in reference barcode libraries for Madagascar and the urgency of this task in regard to conservation of this biodiversity hotspot. Indeed, a comprehensive DNA barcoding library of Madagascan moths is needed, and that could be achieved by rapid digitization and sequencing of specimens deposited at the two main collections in the Natural History Museums of Paris and London. A

good example of how that could be achieved is the DNA barcoding of the Australian National Insect Collection (Hebert et al. 2013).

For micromoths, the numbers of BINs from our study (1537 BINs) equals approximately the number of non-macroheterocerans in the current list as updated in Table 2 (based on a resolution of Viette 1990 and De Prins and De Prins 2018). There are around 1510 described species for the micromoth families detected here [1539 including *Gelechioidea incertae sedis*] out of a tally today for all micromoth families of 1598 species (Table 2). Ultimately, we would anticipate a very low overlap between the current checklist of described species and the list of BINs in our samples, in the hypothetical case that the types in museums were successfully DNA barcoded. We were able to identify relatively few (162) specimens down to species level (94 species representing 95 BINs), either using DNA barcode searches on BOLD (for the Malaise trap, just three species) or for pinned material, by two weeks working in the Paris Museum (MNHN). Comparative analysis of the external morphology of our pinned material with museum reference collections suggests that a large percentage of our barcoded material are likely to represent undescribed species. Moreover, some higher taxa have so few described species that we can be almost certain (given a likely very high species endemism rate in these groups) that the undescribed rate is also very high in those groups (for example, only one nepticulid and eight hieroxestine tineids are described). For the 27 families that show more BINs than described species (notable among which are Gelechiidae and Tineidae), the number of BINs (889) exceeds the number of described species (270) by 619; only 15 of these BINs are identified to species. For the remaining 34 families (with 1298 described species), a minor proportion of their 482 BINs are likely to intersect greatly with their described species, considering only 79 of those BINs could be identified, 70 of which represent just four families (Crambidae, Pyralidae, Tortricidae and Limacodidae). These figures alone allow a range of 45-93% undescribed species, with a

tendency towards the upper figure, among the 1371 BINs identified to family (or family-level grouping). An additional 166 BINs were not even identified to family. It is of paramount importance to DNA barcode the reference collections deposited in both Paris and London in particular, using new barcoding technologies (Zuccon et al. 2012) and with a particular focus on types (Hausmann et al. 2016) in order to more precisely estimate the Linnean shortfall (Cardoso et al 2011) in Madagascan moths. Micromoths will be more challenging in this respect due to the need to minimise tissue removal on holotypes, but as an alternative, morphologically linkable non-primary type material is frequently available. This need for reference libraries from collection types also echoes the call for the barcoding effort to be extended to local metabarcoding studies, that all need to be linked into the BOLD system in order to improve database comprehensiveness (Porter and Hajibabaei 2017).

Molecular synapomorphy to identify Macroheterocera

We found that the presence of a Phenylalanine in the 177th position of the barcode fragment is shared by almost all Macroheterocera analysed from the two test data sets. We have determined that the Phenylalanine state is a shared character state of the clade Macroheterocera which is very seldom reversed or paralleled within the Lepidoptera. Its precise reliability is hard to gauge due to the possibility of false positive and false negative identifications, but appears to be of the order of 99.5%. This molecular synapomorphy allowed us to filter out reliably non-macroheteroceran barcodes from data sets where external morphology of voucher specimens is poorly preserved (e.g. Malaise trap samples), although butterflies, which share a Leucine with most micromoths in the homologous position, need to be independently removed. Use of this character should be very useful to barcoders of Lepidoptera and also provides a means when

using the identification engine to evaluate nearest neighbours (e.g. from irrelevant families) that are spuriously close to the sequence being queried.

Higher taxonomic diversity of micromoths in Madagascar

Our survey detected 38 micromoth families previously recognized for Madagascar (including confirmation of the newly recognized Tonzidae) and added five new ones (Dryadaulidae, Bucculatricidae, Bedellidae, Batrachedridae, Blastobasidae). It also added four higher taxa which may be valid at family level but are currently included on BOLD as subfamilies, respectively, of Lyonetiidae (s.l.) (Cemiostomidae) and Depressariidae (s.l.) (Ethmiidae, Peleopodidae, Stenomatidae; *Orygocera* is *incertae sedis* and needs to be excluded from the latter). Seventeen micromoth families previously listed for Madagascar (Viette, 1990; Lees and Minet, 2003; Sohn, 2015; De Prins and de Prins, 2018) that we did not detect in this survey were Heliozelidae, Tischeriidae, Lyonetiidae (s. str.), Ypsolophidae, Plutellidae, Dudgeoneidae, Metarbelidae, Sesiidae, Zygaenidae, Somabrachyidae, Xyloryctidae, Autostichidae, Momphidae, Coleophoridae, Hyblaeidae, Callidulidae, and Whalleyanidae. Whalleyanidae is included in Thyrididae and *Boisduvalodes tamatavana* in Limacodinae in De Prins and de Prins, 2018 but here we follow Lees and Minet 2003 (see p. 758, note 22) in treating the former as a valid family and the latter as a representative of Somabrachyidae. That brings the Madagascan micromoth fauna to as much as 64 families. It is not at all surprising that we did not find the other undetected families since they are essentially diurnal (Heliozelidae, Ethmiidae, Sesiidae, Zygaenidae, Hyblaeidae, Callidulidae), or are also rare and represented by only one or two described species (Tischeriidae, Lyonetiidae s.s., Ypsolophidae, Dudgeoneidae, Metarbelidae, Somabrachyidae, Autostichidae, Momphidae) or occur outside the sampled region (Whalleyanidae). The main

surprise is the absence of Sesiidae and Zygaenidae, known to occur in Malaise traps, and Heliozelidae, mentioned earlier.

It is always an exciting possibility that one or more new families could be represented in our dataset, considering that the 113 unknowns to family level include a number of local still unidentified radiations and exhibit sometimes striking divergences to any other BIN from Madagascar or outside it.

Invasive and pest species

We found 55 species with a widespread distribution range mostly outside the Afrotropical region. Of those, 28 species appear to be new to Madagascar as they have not been recorded previously by three works: Viette (1990) checklist, Martiré and Rochat, (2008), and the T@RTS database (Gilligan et al. 2014). Therefore, it is unknown whether these newly recorded species are established in the island or represent interceptions of new arrivals.

Most of the widespread and the 22 pest species recorded in Table 3 were detected in disturbed habitats of the Nosy Be area. However, we found some pest species even in primary habitats. For example, in primary forest in Andringitra we found both *Prays citri* and *Diasemiopsis ramburialis* for the first time although the former is not clearly distinguishable from the sympatric *P. oleaeoides* Gibeaux, 1985 (B. Heckford, pers. comm.), raising the possibility *P. citri* has been present there for decades.

These results show the impoverishment and homogenization of the micromoth fauna in disturbed areas and the importance of preserving intact primary forests (Watson et al. 2018). In addition, they highlight the importance of DNA barcoding as a bio-surveillance tool to facilitate the identification and detection of plant pests (Frewin et al. 2013).

549 *Biodiversity assessments*

550 DNA barcoding has facilitated the use of hyperdiverse groups such as micromoths in biodiversity
551 studies (Miller et al. 2016). Traditionally, especially in the tropics, micromoths have been largely
552 ignored in biodiversity assessment. This study adds much motivation to this type of effort,
553 considering also that they are so straightforward to distinguish with DNA barcodes (e.g., to
554 separate from very small erebids, or within types of sampling like Malaise trapping, where wing
555 pattern identification is rendered for the most part impractical). Identifying micromoths to family
556 level or below, however, still requires a large effort and integrated morphological and molecular
557 analysis. Here, building a comprehensive DNA barcode reference library on BOLD with as
558 complete taxonomic information as possible alongside lists of BINs will prove indispensable for
559 assisting future identification, surveys and comparisons of poorly known faunas such as that of
560 Madagascar. Hopefully such efforts will stimulate a new wave of species description while time
561 is left to highlight disappearing forest regions, with slash and burn for hill rice cultivation now
562 exacerbated by downwards spiralling poverty and the rosewood logging crisis. They will also
563 assist agriculturalists and horticulturalists to identify threats to plants via documentation of plant
564 pests and invasive species.

565

566

567 **Acknowledgements**

568 We would like to thank Sarah Adamowicz, Lori Cayer and two anonymous referees for their
569 careful reading of our manuscript. Monica Young helped by making the shared species decay
570 plot. Olivier Bouteleux photographed and processed some of the samples for DNA barcoding.
571 John Brown and Alecia Timm helped greatly with tortricid identifications. Sjaak Koster is also
572 thanked for identification of cosmopterigids. Joaquin Baixeras kindly assisted with information

573 about Madagascan species from the T@RTS database and checked information used in Table 3.
574 Peter Huemer kindly allowed us to use some unpublished barcodes from project CWLMA. Klaus
575 Sattler and Bob Heckford are thanked for discussion on Gelechiidae systematics and the genus
576 *Prays*, respectively. MICET and their staff are thanked for facilitating logistics and export.
577 Ravomiarana Ranaivosolo, Andrianjaka Ravelomanana, and Tahina Rajao helped in the field. We
578 are also grateful to Paul Hebert, Kate Perez, Claudia Steinke, Jayme Sones and other staff at the
579 Centre for Biodiversity Genomics. Funding for fieldwork came from from National Geographic
580 Society grant # 8316-07 (to DCL) and ERC grant EMARES #250325 (to Paul Brakefield). Lab
581 work was partly funded by grants from the Ontario Ministry of Research and Innovation and the
582 Canada Foundation for Innovation to the Centre for Biodiversity Genomics.
583
584 Relevant collecting permits are: 57/11/MEF/SG/DGF/DCB.SAP/SCB,
585 019/14/MEF/SG/DGF/DCB.SAP/SCB and 277/14/MEEF/SG/DGF/DCB.SAP/SCB (to DCL)
586 enabling collections in reserves and other protected areas. The director of MICET and their staff
587 including drivers are thanked for facilitating these permits, export and logistics. One of us (AA)
588 also worked in Andasibe under the authorization of a student program at Department of
589 Entomology (University of Antananarivo). Ravomiarana Ranaivosolo, Andrianjaka
590 Ravelomanana, and Tahina Rajao assisted DCL in the field.

References

- Aagaard, K., Berggren, K., Hebert, P.D.N., Sones, J., McClenaghan, B., and Ekrem, T. 2017. Investigating suburban micromoth diversity using DNA barcoding of Malaise trap samples. *Urban Ecosystems*, **20**(2): 353–361. doi:10.1007/s11252-016-0597-2.
- Baker R.H.A., Eyre D., and Brunel, S. 2013. Matching methods to produce maps for pest risk analysis to resources. *NeoBiota*, **18**: 25–40.
- Barrion, A., Litsinger, J., Medina, E.B., Aguda, R.M., Bandong, J.P., Pantua P.C., et al. 1991. The rice *Cnaphalocrocis* and *Marasmia* (Lepidoptera: Pyralidae) leafroller complex in the Philippines: taxonomy, bionomics and control. *Philippine Entomologist*, **8**: 987–1074.
- Bio-Inventory and Collections Unit, Biodiversity Institute of Ontario. 2015. Global Malaise Trap Program Progress Report 2015. <http://biodiversitygenomics.net/site/wp-content/uploads/2015/12/GMP-Progress-Report-2015.pdf>. Accessed 16 July 2018.
- Briggs, J.C. 1987. *Biogeography and plate tectonics*. Elsevier, Amsterdam 203 pp.
- Brown, J., Copeland, R., Aarvik, L., Miller, S., Rosati, M.E., and Luke, Q. 2014. Host Records for fruit-feeding Afrotropical Tortricidae (Lepidoptera). *African Entomology*, **22**: 343–376.

- 614 Buthelezi, N.M., Conlong, D.E., and Zharare, G.E. 2013. A comparison of the infestation of
615 *Aproaerema simplexella* (Walker) on groundnut and other known hosts for *Aproaerema*
616 *modicella* (Deventer) (Lepidoptera: Gelechiidae). African Entomology, **21**(2): 183–195 .
617
- 618 Cardoso, P., Erwin, T.L., Borges, P.A.V., and New, T.R. 2011. The seven impediments in
619 invertebrate conservation and how to overcome them. Biological Conservation, **144**(11): 2647–
620 2655. doi: 10.1016/j.biocon.2011.07.024.
621
- 622 Chao A., Colwell R.K., Lin C.W., and Gotelli N.J. 2009. Sufficient sampling for asymptotic
623 minimum species richness estimators. Ecology, **90**: 1125–1133.
624
- 625 Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., et al. 2014.
626 Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in
627 species diversity studies. Ecological Monographs, **84**: 45–67. doi: [https://doi.org/10.1890/13-](https://doi.org/10.1890/13-0133.1)
628 0133.1.
629
- 630 Danielczak, A., Devriese, H., and Hochkirch, A. 2017. *Agkistropleuron simplex*.
631 <<http://dx.doi.org/10.2305/IUCN.UK.2017-2.RLTS.T103896223A103900575.en>> Accessed
632 31/03/2018.
633
- 634 Davis, D.R., and Stonis, J.R. 2007. A revision of the new world plant-mining moths of the family
635 Opostegidae. Smithsonian Contributions to Zoology, 625: 1–212.
636 doi:<https://doi.org/10.5479/si.00810282.625>.
637

De Prins, J., and De Prins, W. 2018. Afromoths, online database of Afrotropical moth species (Lepidoptera). World Wide Web electronic publication (www.afromoths.net<<http://www.afromoths.net>>) [7-14 May, 2018].

deWaard, J.R., Levesque-Beaudin, V., deWaard, S.L., Ivanova, N.V., McKeown, J.T.A., Miskie, R., et al. 2017. Expedited assessment of terrestrial arthropod diversity by coupling malaise traps with DNA barcoding. bioRxiv doi:<http://dx.doi.org/10.1101/192732>.

Evans, D.E. 1970. The parasites of three Kenya coffee tortricids (Lepidoptera: Tortricidae). Journal of the Entomological Society of South Africa, **33**(2): 349–350.

Enkhtur, K., Pfeiffer, M., Lkhagva, A., and Boldgiv, B. 2017. Response of moths (Lepidoptera: Heterocera) to livestock grazing in Mongolian rangelands. Ecological Indicators, 72 : 667–674. doi:<https://doi.org/10.1016/j.ecolind.2016.08.053>.

Favetti, B.M., Catoia, B., Thais G. G., and Bueno, C.O.F. 2018. Population dynamics of *Omiodes indicata* (Fabricius) (Lepidoptera: Pyralidae) on Soybean in Brazil. Journal of Agricultural Science, **10**(1). DOI: 10.5539/jas.v10n1p245

Fisher, B.L., Ratsirarson, H., and Razafimandimby, S. 1998. Inventaire biologique de la forêt littorale de Tampolo (Fenoarivo Atsinanana). In Les Fourmis (Hymenoptera: Formicidae). Edited by J.Ratsirarson, and S.M. Goodman. CIDST, Antananarivo. Ppp. 107–131.

- 661 Frewin, A. Scott-Dupree, C., and Hanner, R. 2013. DNA barcoding for plant protection:
662 applications and summary of available data for arthropod pests. *CAB Reviews*, **8**(18) : 1–13.
663 doi:10.1079/PAVSNNR20138018.
- 664
- 665 Geiger, M. F., Moriniere, J., Hausmann, A., Haszprunar, G., Wagele, W., Hebert, P. D., et al.
666 2016. Testing the Global Malaise Trap Program - how well does the current barcode reference
667 library identify flying insects in Germany? *Biodiversity Data Journal*, **4**: e10671.
- 668
- 669 Gibbs, G.W. 2016 *Ghosts of Gondwana*. Revised Edition. Potton and Burton, New Zealand. 367
670 pp.
- 671
- 672 Gilligan, T.M., Baixeras, J., Brown, J.W., and Tuck, K.R. 2014. T@RTS: Online World
673 Catalogue of the Tortricidae (Ver. 3.0). <<http://www.tortricid.net/catalogue.asp>>. Accessed
674 31/03/2018.
- 675
- 676 Gotelli, N.J., and Colwell, R.K. 2010. Estimating species richness. *In* *Frontiers in Measuring*
677 *Biodiversity*. Edited by A.E. Magurran and B.J. McGill., Oxford University Press, New York. Pp.
678 39–54.
- 679
- 680 Guindon S., Dufayard J.F., Lefort V., Anisimova M., Hordijk W., and Gascuel O. 2010. New
681 algorithms and methods to estimate Maximum-Likelihood phylogenies: assessing the
682 performance of PhyML 3.0. *Systematic Biology*, **59**(3): 307–321.
- 683

- 684 Hausmann, A., Miller, S.E., Holloway, J.D., deWaard, J., Pollock, D., Prosser, S., et al. 2016.
685 Calibrating the taxonomy of a megadiverse insect family: 3000 DNA barcodes from geometrid
686 type specimens (Lepidoptera, Geometridae). *Genome*, **59**(9): 671–684.
687 doi:<http://dx.doi.org/10.1139/gen-2015-0197>.
688
- 689 Hayden, J.E., and Buss, L.J. 2013. Olive Shootworm, *Palpita persimilis* Munroe (Insecta:
690 Lepidoptera: Crambidae). Document EENY556, Department of Entomology and Nematology
691 series. <https://edis.ifas.ufl.edu/pdf/files/IN/IN99500.pdf>
692
- 693 Hawes, J., da Silva Motta, C., Overal, W.L., Barlow, J., Gardner, T.A., and Peres, C.A. 2009.
694 *Journal of Tropical Ecology*, **25**(3) : 281–300. doi:<https://doi.org/10.1017/S0266467409006038>.
695
- 696 Hebert, P.D.N., deWaard, J.R., Zakharov, E.V., Prosser, S.W.J., Sones, J.E., McKeown, J.T.A.,
697 et al. 2013. A DNA ‘Barcode Blitz’: rapid digitization and sequencing of a natural history
698 collection. *PLoS ONE*, **8**(7): e68535. doi:10.1371/journal.pone.0068535
699
- 700 Heckford, R.J. 2004. *Anatrachyntis simplex* (Walsingham, 1891) (Lepidoptera:
701 Cosmopterigidae), an adventive species new to the British Isles and a larval description.
702 *Entomologist’s Gazette*, **55**: 95–101.
703
- 704 Herrera, J.P. 2017. Prioritizing protected areas in Madagascar for lemur diversity using a
705 multidimensional perspective. *Biological Conservation*, **207**: 1–8. doi:
706 <https://doi.org/10.1016/j.biocon.2016.12.028>.
707

- 708 Hsieh, T.C., Ma, K.H., and Chao, A. 2016. iNEXT: An R package for interpolation and
709 extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, **7**(12) :
710 1451–1456. doi:<https://doi.org/10.1111/2041-210X.12613>.
711
- 712 Irwin, M.T., Wright, P.C., Birkinshaw, C., Fisher, B.L., Gardner, C.J., Glos, J., et al. 2010.
713 Patterns of species change in anthropogenically disturbed forests of Madagascar. *Biological*
714 *Conservation*, **143**(10) : 2351–2362. doi:[10.1016/j.biocon.2010.01.023](https://doi.org/10.1016/j.biocon.2010.01.023).
715
- 716 Ivanova, N.V., deWaard, J.R., and Hebert, P.D.N. 2006. An inexpensive, automation- friendly
717 protocol for recovering high-quality DNA. *Molecular Ecology Notes*, **6**(4) : 998–1002. doi :
718 <https://doi.org/10.1111/j.1471-8286.2006.01428.x>.
719
- 720 Jamieson, L.E., Suckling, D.M., and Ramankutty, P. 2008. Mass Trapping of *Prays nephelomima*
721 (Lepidoptera: Yponomeutidae) in *Citrus* orchards: optimizing trap design and density. *Journal of*
722 *Economic Entomology*, **101**(4): 1295–1301. <https://doi.org/10.1093>.
723
- 724 Jenkins, R.K.B., Tognelli, M.F., Bowles, P., Cox, N., Brown, J.L., Chan, L., et al. 2014.
725 Extinction risks and the conservation of Madagascar’s reptiles. *PLoS ONE*, **9**(8). doi:e100173.
726 doi:[10.1371/journal.pone.0100173](https://doi.org/10.1371/journal.pone.0100173).
727
- 728 Kremen, C. 1994. Biological inventory using target taxa: a case study of the butterflies of
729 Madagascar. *Ecological Applications*, **4**(3) : 407–422. <https://doi.org/10.2307/1941946>.
730

- 731 Kobayashi, S., Matsuoka, H., Kimura, M., Sohn, J-C., Yoshiyasu, Y., and Lees, D.C. 2018.
732 Designation of a new family group name, Tonzidae fam. nov., for the genus *Tonza* (Lepidoptera,
733 Yponomeutoidea), based on immature stages of *Tonza citrorrhoa*. European Journal of
734 Taxonomy, **443** : 1–32. <https://doi.org/10.5852/ejt.2018.443>
735
- 736 Koster, J.C.(S.), and van Nieukerken, E.J. 2018. *Gielisella* gen. n., a new genus and two new
737 species from southern Spain (Lepidoptera: Elachistidae: Parametriotinae) with a catalogue of
738 parametriotine genera. Nota Lepidopterologica, 40 (20): 163–202.
739 <https://doi.org/10.3897/nl.40.14528>
740
- 741 Krüger, M. 2007. Composition and origin of the Lepidoptera faunas of southern Africa,
742 Madagascar and Réunion (Insecta: Lepidoptera). Annals of the Transvaal Museum, **44**(1): 123–
743 178.
744
- 745 Kull, C., Tassin, J., and Carrière, S.M., 2014. Approaching invasive species in Madagascar.
746 Madagascar Conservation and Development, **9**(2) : 60–70.
747 [doi:http://dx.doi.org/10.4314/mcd.v9i2.2](http://dx.doi.org/10.4314/mcd.v9i2.2).
748
- 749 Kwadha, C.A, Ong'amo, G.O., Ndegwa, P.N., Raina, S.K., and Fombong, A.T. 2017. The
750 biology and control of the Greater Wax Moth, *Galleria mellonella*. Insects, **8**(2): E61. doi:
751 [10.3390/insects8020061](https://doi.org/10.3390/insects8020061).
752
- 753 Lees, D.C., Kremen, C., and Raharitsimba, H., 2003. Classification, diversity and endemism of
754 the butterflies (Papilionoidea and Hesperioidea): a revised species checklist. In The Natural

- 755 History of Madagascar. *Edited by* S.M. Goodman, S.M. and J.P. Benstead), , University of
756 Chicago Press, Chicago. pp. 762–793.
- 757
- 758 Lees, D.C., and Minet, J., 2003. Lepidoptera: systematics and diversity. *In* The Natural History of
759 Madagascar. *Edited by* S.M. Goodman, S.M. and J.P. Benstead). University of Chicago Press,
760 Chicago. pp. 748–761.and
- 761
- 762 Lees, D.C., Ohshima, I., Kawakita, A., Kawahara A., Rougerie, R., Adamski, D., et al. 2011.
763 Rapid inventory via DNA barcoding: cross-lepidopteran diversity survey of Nouragues
764 inselberg, French Guiana. Published abstract, XVIIth European Congress of Lepidopterology,
765 Luxembourg, 9-14.V.2011.
- 766
- 767 Lees, D.C., Kawahara, A. Y., Bouteleux, O., Ohshima, I., Kawakita, A., Rougerie, R., et al. 2013.
768 DNA barcoding reveals a largely unknown fauna of Gracillariidae leaf-mining moths in the
769 Neotropics. *Molecular Ecology Resources*, **14**(2): 286–296. doi: 10.1111/1755-0998.12178
- 770
- 771 Lees, D.C. 2016. *Heteropsis* (Nymphalidae: Satyrinae: Satyrini: Mycalesina): 19 new species
772 from Madagascar and interim revision. *Zootaxa*, **4118** : 1–97.
- 773
- 774 Libert, M. 2014. Sur la taxonomie du genre *Celaenorrhinus* Hübner en Afrique. Rouen : M.
775 Libert. 272 pp. 26 col plates, 54 text figs, 53 maps.
- 776
- 777 Lopez-Vaamonde, C., Agassiz, D.V.L., Augustin, S., De Prins, J., De Prins, W., Gomboc, S., et
778 al. 2010. Lepidoptera. Chapter 11. *In* Alien terrestrial arthropods of Europe. (*Edited by* A.

- 779 Roques, M. Kenis, D. Lees, C. Lopez-Vaamonde,, W. Rabitsch, J.Y. Rasplus and D.B. Roy. :
780 Pensoft, Sofia. pp. 603–668.
781
- 782 Lopez-Vaamonde, C., Bremn, F., Lees, D.C., van Houdt, J., and de Prins, J. 2012. Analysis of
783 tissue dependent DNA yield for optimal sampling of micro-moths in large-scale biodiversity
784 surveys. *European Journal of Entomology*, **109**(1): 1–6. <https://doi.org/10.14411/eje.2012.001>.
785
- 786 Martiré, D., and Rochat, J. 2008. Les papillons de la Réunion et leurs chenilles. Mèze, France,
787 Biotope, 496 pp.
788
- 789 Miller, S.E., Hausmann, A., Hallwachs, W., and Janzen, D.H. 2016. Advancing taxonomy
790 andbioinventories with DNA barcodes. *Philosophical Transactions of the Royal Society B*,
791 **371**(1702) : 20150339. doi:<https://doi.org/10.1098/rstb.2015.0339>.
792
- 793 Morland, G. 2015. The morphology and ecology of the Carob moth (*Ectomyelois ceratoniae*)
794 (Zeller) in citrus orchards of the Western Cape, South Africa. MSc. Thesis, Stellenbosch
795 University. 122 pp.
796
- 797 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. 2016.
798 Package ‘vegan’. Available at <https://cran.r-project.org>.
799
- 800 Parenti, U. 2006. *Elachista bassii* Parenti, a new species of Elachistidae from Nepal and
801 *Elachista crocogastra* Meyrick in Madagascar (Lepidoptera: Elachistidae). *SHILAP Revista de*
802 *Lepidopterologia*, **34**: 141-144.

803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826

Paulian, R., andand Viette, P. 1955. Essai d'un catalogue biologique des Lepidoptères Heterocères de Tananarive. Mémoires de l'Institut Scientifique de Madagascar, **6**(E): 141–281.

Porter, T.M., and Hajibabaei M. 2017. Automated High Throughput Animal DNA Metabarcoding Classification. bioRxiv. Springer US, 219675. doi :<https://doi.org/10.1101/219675>.

Poux, C., Madsen, O., Marquard, E., Vieites, D.R., de Jong, W.W., and Vences, M. 2005. Asynchronous colonization of Madagascar by the four endemic clades of primates, tenrecs, carnivores, and rodents as inferred from nuclear genes. Systematic Biology, **54**(5) : 719–730. doi:[10.1080/10635150500234534](https://doi.org/10.1080/10635150500234534).

R Development Core Team. 2004. R: a language and environment for statistical computing. Foundation for Statistical Computing, Vienna. Available at: <http://www.r-project.org>.

Ratnasingham, S., and Hebert, P.D.N. 2007. BOLD: the barcode of life data system (<http://www.barcodinglife.org>). Molecular Ecology Notes, **7**:355–364.

Ratnasingham, S., and Hebert, P.D.N. 2013. A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. PLoS ONE, **8**(7). doi:<https://doi.org/10.1371/journal.pone.0066213>.

Regier, J.C., Mitter, C., Davis, D.R., Harrison, T.L., Sohn J.-C., Cummings, M.P., et al. 2014. A molecular phylogeny and revised classification for the oldest ditrysian moth lineages

- 827 (Lepidoptera: Tineoidea), with implications for ancestral feeding habits of the mega-diverse
828 Ditrysia. *Systematic Entomology*, **40**(2) : 409–432.
- 829
- 830 Shankara Murthy, M., and Nagaraj, S. K. 2014. Outbreak of maize leaf roller, *Cnaphalocrocis*
831 *trapezalis* Guenee (= *Marasmia trapezalis*) (Crambidae: Lepidoptera) on sorghum in Karnataka,
832 south India. *Insect Environment*, **19**(4): 250–252
- 833
- 834 Schmidt, O., Hausmann, A., Cancian de Araujo, B., Sutrisno, H., Peggie, D., and Schmidt, S.
835 2017. A streamlined collecting and preparation protocol for DNA barcoding of Lepidoptera as
836 part of large-scale rapid biodiversity assessment projects, exemplified by the Indonesian
837 Biodiversity Discovery and Information System (IndoBioSys). *Biodiversity Data Journal*, **5**:
838 e20006. doi:<https://doi.org/10.3897/BDJ.5.e2000>.
- 839
- 840 Sharma, H.C., Saxena, K.B., and Bhagwat, V.R. 1999. The legume pod borer, *Maruca vitrata*:
841 bionomics and management. Information Bulletin no. 55 Patancheru 502 324, Andhra Pradesh,
842 India: International Crops Research Institute for the Semi-Arid Tropics. 42 pp.
- 843
- 844 Shimizu K. 2000. The biology of the cotton caterpillar (*Diaphania indica*) and the resistance to
845 insecticides. *Plant Protection*, **54**: 97–103.
- 846
- 847 Silva, E.B., and Mexia, A. 1999. The pest complex *Cryptoblabes gnidiella* (Millière)
848 (Lepidoptera: Pyralidae) and *Planococcus citri* (Risso) (Homoptera: Pseudococcidae) on sweet
849 orange groves (*Citrus sinensis* (L.) Osbeck) in Portugal: interspecific association. *Boletín de*
850 *sanidad vegetal Plagas*, **25**: 89–98.

- 851
- 852 Solis, M.A., and Pogue, M.G. 1999. Lepidopteran biodiversity: patterns and estimators.
- 853 American Entomologist, **45**(4) : 206–212. doi:<https://doi.org/10.1093/ae/45.4.206>.
- 854
- 855 Sohn J.-C., Regier, J.C., Mitter, C., Davis, D., Landry, J.-F., Zwick, A., et al. 2013. A molecular
- 856 phylogeny for Yponomeutoidea (Insecta, Lepidoptera, Ditrysia) and its implications for
- 857 classification, biogeography and the evolution of host plant use. PLoS ONE, **8**(1): e55066.
- 858 <https://doi.org/10.1371/journal.pone.0055066>
- 859
- 860 Sohn, J.-C. 2015. A taxonomic review of the genus *Phrealcia* Chrétien, 1900 with description of
- 861 a new species from Korea (Lepidoptera: Ypsolophidae). SHILAP Revista de Lepidopterología,
- 862 **43**: 385–423.
- 863
- 864 Sohn, J.-C., Regier, T.C., Mitter, C., Adamski, D., Landry, J.-F., Heikkilä, M., et al. 2015.
- 865 Phylogeny and feeding trait evolution of the mega-diverse Gelechioidea (Lepidoptera:
- 866 Obtectomera): new insight from 19 nuclear genes. Systematic Entomology, **41**(1) : 112–132.
- 867 doi:<https://doi.org/10.1111/syen.12143>.
- 868
- 869 van Nieukerken, E.J., Kaila, L., Kitching, I.J., Kristensen, N.P., Lees, D.C., Minet, J., et al. 2011.
- 870 Order Lepidoptera Linnaeus, 1758. Zootaxa **3148**: 212–221.
- 871
- 872 Vences, M., Wollenberg, K. C., Vieites, D.R., and Lees, D.C. 2009. Madagascar as a model
- 873 region of species diversification. Trends in Ecology and Evolution, **24**(8): 456–465.
- 874 doi:[10.1016/j.tree.2009.03.011](https://doi.org/10.1016/j.tree.2009.03.011).

875

876 Van Den Berg, H. Shepard, B.M., and Nasikin, 2010. Damage incidence by *Etiella zinckenella* in
877 soybean in East Java, Indonesia. *International Journal of Pest Management*, **44**(3): 153–159.

878

879 Viette, P. 1963. Noctuelles trifides de Madagascar, écologie, biogéographie, morphologie et
880 taxonomie (lep.). *Annales de la Société entomologique de France*, **131** (1): i–vi, 1–294 + pls 1–
881 10.

882

883 Viette, P. 1965. Insectes. Lépidoptères Noctuidae Amphipyridae (part.). *Faune de Madagascar*,
884 **20**(1): i–vi, 295–490 + pls 11–12.

885

886 Viette, P. 1967. Insectes. Lépidoptères Noctuidae Amphipyridae (part.) et Meliclyptidae. *Faune*
887 *de Madagascar*, **20** (2): i–iv, 491–825 + pls 13–14.

888

889 Viette, P. 1990. Liste récapitulative des Lépidoptères Hétérocères de Madagascar (A provisional
890 check-list of the Lepidoptera Heterocera of Madagascar). *Faune de Madagascar* **Suppl. 1**: 1–
891 264.

892

893 Watson, J.E.M., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C., et al. 2018. The
894 exceptional value of intact forest ecosystems. *Nature Ecology and Evolution*, **2**: 599–610.
895 doi:<https://doi.org/10.1038/s41559-018-0490-x>.

896

- 897 Wesener, T., and Rudolf, E. 2017. *Sphaeromimus splendidus*.
898 <<http://dx.doi.org/10.2305/IUCN.UK.2017-1.RLTS.T65526540A65527790.en>> 2017. Accessed
899 31/03/2018.
900
- 901 Wesener, T., Le, D.M.T., and Loria, S.F. 2014. Integrative revision of the giant pill-millipede
902 genus *Sphaeromimus* from Madagascar, with the description of seven new species (Diplopoda,
903 Sphaerotheriida, Arthrosphaeridae). ZooKeys, **414**: 67–107.
904 doi:<https://doi.org/10.3897/zookeys.414.7730>.
905
- 906 Yamada, H. 1979. Bionomics of the perilla leaf roller, *Pyrausta phoenicealis* Hübner. Annual
907 Report of The Kansai Plant Protection Society, **21**: 8–11. DOI: 10.4165/kapps1958.21.0_8
908
- 909 Yen, S.H. 2014. Insecta: Lepidoptera, Crambidae, Acentropinae. In Freshwater Invertebrates of
910 the Malaysian Region. Edited by C.M. Yule, C.M and H.S. Yong Academy of Sciences,
911 Malaysia. pp. 545–554.
912
- 913 Zharare, G.W. 2013. A Comparison of the infestation of *Aproaerema simplexella* (Walker) on
914 Groundnut and other known hosts for *Aproaerema modicella* (Deventer) (Lepidoptera:
915 Gelechiidae). African Entomology, **21**(2):183–195.
916
- 917 Zuccon D., Brisset J., Corbari L., Puillandre N., Utge J., and Samadi S. 2012. An optimised
918 protocol for barcoding museum collections of decapod crustaceans: a case-study for a 10-40-
919 years-old collection. Invertebrate Systematics, **26**(6):592–600.
920 doi:<https://doi.org/10.1071/IS12027>.

921

Draft

Figure Legends

Figure 1. Distribution of BINs over 47 different families collected with light traps or day-netted at 24 sites across Madagascar (light grey) and Malaise trap at Andasibe (dark grey). Unknown refers to individuals that could not be identified at family level. See text regarding Depressariidae s.l. (Gelechioidea i.s. are only those currently under Depressariidae in BOLD that do not fit in the expanded categories).

Figure 2. Distribution of BINs over biogeographic regions. Notice one BIN can appear in several biogeographic regions.

Figure 3. Abundance data for the 507 BINs detected in the Malaise trap samples. Notice that 57.6% (292 out of 507) BINs are singletons. Two BINs (BOLD:ACS0229, *Elachista* and BOLD:ACS1392, Tineidae: Hieroxestinae, exemplars illustrated left to right) have more than 70 individuals but no species name.

Figure 4. Shared species decay plot for Malaise trap samples collected at two sites in Andasibe: Malaise trap 1 (sampled from April 1st until 28th May 2014) in black; Malaise trap 2 (sampled from September 1st until 6th November 2014) in red.

Figure 5. Accumulation curves for Malaise trap samples over the two periods corresponding to end of the wet to beginning of dry season (M1) and end of dry to beginning of wet season (M2),

946 with: **A.** Species diversity (BINs) per number of individuals. **B.** Sample coverage per number of
947 individuals. **C.** Species diversity (BINs) per sample coverage.

948

949 **Figure 6.** Species richness observed and estimated, based both on Chao1 and ACE analyses for
950 the sites sampled with a Malaise trap at Andasibe.

951

Draft

953 Table 1. Study Sites.

Locality	Region	Habitat	Latitude & Longitude (decimal degrees)	Elevation (m)
Ambilobe	Mainland	Scrubland	-13.108 to -13.163, 49.097	25-40
Mont Passot	Nosy Be	Degraded forest	-13.282, 48.259	25
Ambaro	Nosy Be	Degraded forest	-13.31, 48.187	20
Dzamandzar	Nosy Be	open fields	-13.333, 48.196	25
Fascene	Nosy Be	Open field surrounded by degraded forest	-13.344, 48.299	75
Hell-Ville	Nosy Be	Degraded forest	-13.367, 48.283	15
Lac Ampobilava	Nosy Be	Degraded forest	-13.395, 48.241	40
Lac Djabala	Nosy Be	Degraded forest	-13.386, 48.244	40
Ambanoro	Nosy Be	Degraded forest	-13.389, 48.3	75
Ambondro	Nosy Be	Gardens	-13.382, 48.197	10
Ambanja	Mainland	Scrubland	-13.701, 48.464	40

Manongarivo Réserve Spéciale, Antsatrotro Mt	Mainland	Protected forest	-14.082, 48.366	1235
Marojejy National Park	Mainland	Protected forest	-14.433, 49.761 -14.44, 49.74	700 1540
Anjanaharibe Sud Réserve Spéciale, below Anjividibe summit	Mainland	Protected forest above dry stream bed	-14.739, 49.462	1540
Anjanaharibe Sud Réserve Spéciale, Indri Camp	Mainland	protected forest	-14.741, 49.497	960
Anjanaharibe Sud Réserve Spéciale	Mainland	Protected forest	-14.743, 49.464	1450
Anjozorobe Mananara Lodge	Mainland	Degraded forest	18.436, 47.942	1300
Feo-ny-ala, Andasibe	Mainland	Hotel near protected forest	-18.947, 48.419	945
Mantadia National Park, Belakato trail	Mainland	Protected forest	-18.82, 48.436	1000
Andasibe, Malaise trap M1	Mainland	Protected forest	-18.9484, 48.4256	1000
Andasibe, Malaise trap M2	Mainland	Protected forest	-18.9438, 48.4316	1050
Ankazomivady	Mainland	Degraded high	-20.778, 47.178	1710

		plateau forest	-20.7948, 47.1773	1830
Andringitra National Park, camp	Mainland	Protected forest	-22.147, 46.946	1570
Andringitra National Park	Mainland	Protected forest	-22.1504, 46.9487	1625
Sahavondronina, 7 km W Vohiparara, Community forest at Ranomafana National Park	Mainland	Open field surrounded by protected forest	-21.278, 47.331	1230

954

Draft

Table 2. Number of BINs per family and method of collection (Malaise, light trapping and netting by day). Families are ordered systematically and those recorded for the first time for Madagascar are highlighted in bold. All micromoth families are shown for completeness of the total microlepidopteran count (species included in Viette 1990 that are not synonyms of other species, although not listed for Madagascar in Afromoths, have been included in the count). Some other amendments have been made such as there is only one Elachistidae described from Madagascar (Parenti, 2006; Koster & van Nieukerken, 2017). See text regarding Depressariidae s.l. Families largely ordered following Nieukerken et al. 2011, Regier et al. (2014) for Tineoidea and Sohn et al. (2013, 2015) for Yponomeutoidea and Gelechioidea. *Placement in this family based entirely on COI data, specimens in poor condition. ** An additional six barcodes are too short to be allocated BINs so a total of 2950 barcodes have BINS and are subject to analysis. *** "*Iridostoma*" *catatella* Viette, 1956 is misplaced in *Iridostoma* Meyrick, 1909; this species is here transferred from Plutellidae to the Glyphipterigidae: Acrolepiinae (provisionally as *Acrolepia catatella* comb. nov.). ****Currently included in Lecithoceridae is an *Epichostis*-like species, *Lecithocera ojeiyella* Viette, 1958. In Gelechioidea i.s. we include 18 sequences representing 9 BINs of *Epichostis*-like moths as indicated in the field Extra info. There are currently no *Epichostis* barcodes on BOLD and we are currently uncertain if *L. ojeiyella* together with these BINs might represent true Xyloryctidae.

Family	Records	BINS	BINS Malaise	BINS Light trap + day-netted	BINS shared to BINs	BINS identified to Species	#Described species
--------	---------	------	--------------	------------------------------	---------------------	----------------------------	--------------------

Micropterigidae	2	2	0	2	0	0	0
Opostegidae	6	5	0	5	0	0	0
Nepticulidae	50	28	21	7	0	0	1
Heliozelidae	0	0	0	0	0	0	1
Adelidae	7	7	0	7	0	0	1
Tischeriidae	0	0	0	0	0	0	1
Psychidae	41	23	15	8	0	0	19
Eriocottidae	45	12	2	10	0	0	0
Dryadaulidae*	6	3	3	0	0	0	0
Tineidae	450	159	95	65	1	1	40
Lyonetiidae s.auct. ('Cemiostomidae')	2	2	1	1	0	0	2
Bucculatricidae	2	2	1	1	0	0	0
Gracillariidae	73	55	17	40	2	2	22
Bedelliidae	8	4	1	3	0	0	0
Praydidae	2	2	0	2	0	2	2
Lyonetiidae s.str.	0	0	0	0	0	0	1
Argyresthiidae	29	5	0	5	0	0	9
Yponomeutidae	5	5	2	3	0	0	6
Ypsolophidae	0	0	0	0	0	0	1
Plutellidae***	0	0	0	1	0	0	1
Tonzidae	1	1	0	1	0	0	0
Glyphipterigidae***	29	20	11	10	1	0	7
Alucitidae	3	3	0	3	0	0	4
Pterophoridae	31	19	1	19	1	4	65
Copromorphidae	9	5	0	5	0	0	1
Carposinidae	8	3	0	3	0	0	2
Epermeniidae	30	12	0	12	0	0	7
Immidae	34	21	17	5	1	0	1
Choreutidae	4	3	0	3	0	1	1
Galacticidae	3	1	0	1	0	0	1
Tortricidae	260	150	19	132	1	19	342
Brachodidae	1	1	0	1	0	1	8
Cossidae	1	1	0	1	0	0	26
Dudgeonidae	0	0	0	0	0	0	1
Metarbelidae	0	0	0	0	0	0	2
Sesiidae	0	0	0	0	0	0	32
Epipyropidae	2	2	1	1	0	0	3
Lacturidae	2	1	0	1	0	0	9
Limacodidae incl. Chrysopolominae	11	10	1	9	0	7	70
Somabrachyidae	0	0	0	0	0	0	1
Zygaenidae	0	0	0	0	0	0	5

Gelechioidea i.s., includes (<i>Orygocera</i> , <i>Prothamnoides</i> , " <i>Trichocirca</i> " <i>decaryanum</i>)	87	53	27	26	0	0	27
Depressariidae s.l. ('Stenomatidae': <i>Herbulotiana</i> , <i>Amontes</i>)	6	4	2	3	1	0	18
Depressariidae s.l. ('Peleopodidae': Oditinae)	230	131	54	81	4	0	61
Depressariidae s.l. ('Ethmiidae')	1	1	0	1	0	0	19
Depressariidae s.s. (Depressariidae: Depressariinae, Cryptolechiinae)	2	2	0	2	0	0	6
Oecophoridae	62	28	9	20	1	1	23
Lecithoceridae	219	79	51	31	3	0	28
Xyloryctidae****	0	0	0	0	0	0	1
Autostichidae	0	0	0	0	0	0	2
Elachistidae s.s.	85	8	8	0	0	0	1
Momphidae	0	0	0	0	0	0	1
Batrachedridae	4	4	0	4	0	0	0
Coleophoridae	0	0	0	0	0	0	1
Blastobasidae	13	7	1	6	0	1	0
Scythrididae	21	14	7	7	0	1	5
Stathmopodidae	27	17	0	17	0	0	4
Cosmopterigidae	105	59	13	47	1	2	14
Gelechiidae	313	178	36	145	3	6	32
Whalleyanidae	0	0	0	0	0	0	2
Thyrididae	4	3	0	3	0	2	32
Hyblacidae	0	0	0	0	0	0	4
Callidulidae	0	0	0	0	0	0	4
Pyalidae	171	107	14	93	0	10	271
Crambidae	253	162	25	139	2	34	346
Unknown/i.s. Lepidoptera	190	113	52	62	1	0	1
TOTAL	2950**	1537	507	1053	23	94	1598

978 Table 3. Species detected during this study in Madagascar that are known to occur outside the
 979 Afrotropical region. We also indicate those species that are known to be pests and/or invasive.
 980 Those known to feed on crops but not widely acknowledged pests are indicated with an asterisk.

Family/subfamily	Species name	BIN	Distribution	New record for Madagascar	Pest	Ref.
Tineidae: Erechthiinae	<i>Erechthias minutalis</i>	BOLD:ABW6327	Cosmopolitan	Yes	No	
Tineidae: Hieroxestinae	<i>Opogona</i> sp.	BOLD:ACS7592	Madagascar & Oriental	Yes	No	
Gracillariidae: Acrocercopinae	<i>Dialectica scalariella</i>	BOLD:AAL3278	Palaearctic, Afrotropics and Australia	Yes	No	
Gracillariidae: Ornixolinae	<i>Stomphastis</i> sp.	BOLD:AAM6667	Madagascar & Australia	Yes	No	
Praydidae	<i>Prays nephelomima</i>	BOLD:AAM9790	Madagascar & Australia	Yes	Yes	Jamieson et al. 2008
Praydidae	<i>Prays citri</i>	BOLD:AAW5122	Palaearctic, Afrotropics and Australia	?see text	Yes	Lopez-Vaamonde et al. 2010
Glyphipterigidae	<i>Glyphipterix</i> sp.	BOLD:AAY2216	Australia & Madagascar	Yes	No	
Pterophoridae: Platyptiliinae	<i>Hepalastis pumilio</i>	BOLD:AAD4253	Cosmopolitan	No	No	
Pterophoridae: Platyptiliinae	<i>Stenoptilia</i> sp.	BOLD:AAD0716	Cosmopolitan	Yes	No	
Pterophoridae: Platyptiliinae	<i>Sphenarches anisodactylus</i>	BOLD:AAD0725	Cosmopolitan	No	No	
Tortricidae: Olethreutinae	Genus sp.	BOLD:ACS7628	Palaearctic, Afrotropics, Oriental	Yes	No	
Tortricidae: Olethreutinae	<i>Bactra venosana</i>	BOLD:ABZ1079	Afrotropics, Oriental, Australia	Yes	No	
Tortricidae: Olethreutinae	<i>Crociosema lantana</i>	BOLD:AAH5763	Nearctic, Neotropics, Afrotropics and Australia.	Yes	No	
Tortricidae: Olethreutinae	<i>Cydia choleropa</i>	BOLD:ABW2540	Afrotropics and Oriental	Yes	No	
Tortricidae: Olethreutinae	<i>Dudua aprobola</i>	BOLD:AAT9574	Cosmopolitan	Yes	No*	
Tortricidae: Olethreutinae	<i>Lobesia aeolopa</i>	BOLD:AAJ2244	Afrotropics and Oriental	No	Yes	Evans 1970
Tortricidae: Olethreutinae	<i>Lobesia vanillana</i>	BOLD:ABV8007	Réunion, Madagascar	No	Yes	Brown et al. 2014
Tortricidae: Olethreutinae	<i>Thaumotobia leucotreta</i>	BOLD:AAE7729	Afrotropics, intercepted in Palaearctic and Nearctic	No	Yes	Baker et al. 2013
Cosmopterigidae: Cosmopteriginae	<i>Anatrachyntis simplex</i>	BOLD:ABX3349	Cosmopolitan	Yes	Yes	Heckford, 2004
Cosmopterigidae: Cosmopteriginae	<i>Cosmopterix athesiae</i>	BOLD:AAE4001	Palaearctic and Afrotropics	Yes	No	
Cosmopterigidae: Cosmopteriginae	<i>Cosmopterix</i> sp. cf. <i>attenuatella</i>	BOLD:AAC1744	Cosmopolitan	Yes	No	

Cosmopterigidae: Chrysopeliinae	<i>Ascalenia</i> sp.	BOLD:AAG0134	Cosmopolitan	Yes	No	
Cosmopterigidae: Chrysopeliinae	<i>Gisilia</i> sp.	BOLD:ACS6187	Bangladesh & Madagascar	Yes	No	
Gelechiidae: Anacampsinae	<i>Aproaerema simplexella</i>	BOLD:ACK6985	Cosmopolitan, invasive in Afrotropics	Yes	Yes	Zharare, 2013.
Gelechiidae: Dichomeridinae	<i>Dichomeris acuminatus</i>	BOLD:AAB6409	Cosmopolitan	Yes	No	
Choreutidae	<i>Tebenna micalis</i>	BOLD:AAH9855	Cosmopolitan	Yes	No	
Pyralidae: Galleriinae	<i>Achroia grisella</i>	BOLD:ACO9701	Cosmopolitan	No	No	
Pyralidae: Galleriinae	<i>Galleria mellonella</i>	BOLD:AAA0965	Cosmopolitan	No	Yes	Kwadha et al. 2017
Pyralidae: Pyralinae	<i>Hypsopygia nostralis</i>	BOLD:AAI3521	Nearctic, Neotropics and Afrotropics	Yes	No	
Pyralidae: Phycitinae	<i>Cadra cautella</i>	BOLD:AAB9605	Cosmopolitan	Yes	Yes	Paulian & Viette 1955
Pyralidae: Phycitinae	<i>Cryptoblabes gnidiella</i>	BOLD:AAW5129	Cosmopolitan	Yes	Yes	da Silva & Mexia 1999.
Pyralidae: Phycitinae	<i>Ectomyelois ceratoniae</i>	BOLD:AAU4812	Cosmopolitan	No	Yes	Morland, 2015
Pyralidae: Phycitinae	<i>Etiella zinckenella</i>	BOLD:AAB7420	Cosmopolitan	No	Yes	Van Den Berg et al. 2010
Pyralidae: Phycitinae	<i>Thylacoptila paurosema</i>	BOLD:AAV8326	Afrotropical, Oriental and Australia	No	No*	
Crambidae: Acentropinae	<i>Parapoynx fluctuosalis</i>	BOLD:AAA0473	Cosmopolitan	No	Yes	Yen, 2014
Crambidae: Crambinae	<i>Angustalius malacellus</i>	BOLD:AAV9127	Palaearctic & Afrotropics	No	No*	
Crambidae: Pyraustinae	<i>Isocentris filalis</i> (=Hyalobathra retinalis)	BOLD:AAL8896	Afrotropical, Oriental	No	No	
Crambidae: Spilomelinae	<i>Bocchoris inpersalis</i>	BOLD:AAC5466	Cosmopolitan	No	No	
Crambidae: Spilomelinae	<i>Cnaphalocrocis trapezalis</i>	BOLD:AAC0297	Cosmopolitan	No	Yes	Shankara Murthy & Nagaraj 2014
Crambidae: Spilomelinae	<i>Cnaphalocrocis exigua</i>	BOLD:AAO9362	Afrotropics and Oriental and Oceania	Yes	Yes	Barrión et al. 1991
Crambidae: Spilomelinae	<i>Diasemiopsis ramburialis</i>	BOLD:AAD0296	Old World (Africa, Oriental Australia)	No	No*	
Crambidae: Spilomelinae	<i>Diaphania indica</i>	BOLD:AAB1719	Cosmopolitan	No	Yes	Paulian & Viette 1955 ; Shimizu 2000
Crambidae: Spilomelinae	<i>Eurrhyarodes bracteolalis</i>	BOLD:AAD1173	Cosmopolitan	No	No*	
Crambidae: Spilomelinae	<i>Herpetogramma licarsisalis</i>	BOLD:AAA3965	Palaeotropics, Australasia, Hawaii, Canaries	No	Yes	Lopez-Vaamonde et al. 2010
Crambidae: Spilomelinae	<i>Herpetogramma</i> sp.	BOLD:AAB6841	Afrotropics and Oriental	No	No	
Crambidae: Spilomelinae	<i>Herpetogramma</i> sp.	BOLD:ACD5135	Oriental	Yes	No	
Crambidae: Spilomelinae	<i>Hymenoptychis sordida</i>	BOLD:AAF8520	Old World (Africa, Oriental Australia)	No	No	

Crambidae: Spilomelinae	<i>Hyalobathra olesialis</i>	BOLD:ACN7820	Afrotropical, India and Australia	Yes	No	
Crambidae: Spilomelinae	<i>Maruca fuscalis</i>	BOLD:AAD9057	Australia	Yes	No	
Crambidae: Spilomelinae	<i>Maruca vitrata</i>	BOLD:AAB2756	Pantropical	No	Yes	Sharma et al. 1999
Crambidae: Spilomelinae	<i>Omiodes indicata</i>	BOLD:AAB5389	Cosmopolitan	No	Yes	Favetti et al., 2018
Crambidae: Spilomelinae	<i>Palpita vitrealis</i>	BOLD:AAC1043	Cosmopolitan	No	Yes	Hayden & Buss, 2013
Crambidae: Spilomelinae	<i>Pyrausta phoenicealis</i>	BOLD:AAF5760	Cosmopolitan	No	Yes	Yamada 1979
Crambidae: Spilomelinae	<i>Salbia haemorrhoidalis</i>	BOLD:AAD3428	Cosmopolitan	Yes	No	
Crambidae: Spilomelinae	<i>Spoladea recurvalis</i>	BOLD:AAA3666	Cosmopolitan	No	Yes	Paulian & Viette 1955

981

982

Draft

Figure 2. Distribution of BINs over biogeographic regions. Notice one BIN can appear in several biogeographic regions.

408x267mm (72 x 72 DPI)

A **B** **C**

Figure 6. Species richness observed and estimates based both on Chao1 and ACE analyses for the site sampled with Malaise trap at Andasibe

408x267mm (72 x 72 DPI)