Bacterial diversity in the waterholes of the Kruger National Park: an eDNA metabarcoding approach

Journal:	<i>Genome</i>
Manuscript ID	gen-2018-0064.R1
Manuscript Type:	Article
Date Submitted by the Author:	08-Jul-2018
Complete List of Authors:	Farrell, Maxwell; McGill University, Department of Biology Govender, Danny; Scientific Services, Kruger National Park, SANParks; University of Pretoria, Department of Paraclinical Sciences, Faculty of Veterinary Science Hajibabaei, Mehrdad; University of Guelph VAN DER BANK, Michelle; University of Johannesburg, African Centre for DNA Barcoding Davies, T; University of British Columbia, Botany, Forestry & Conservation Sciences; University of Johannesburg, African Centre for DNA Barcoding
Keyword:	microbiome, conservation, biomonitoring, 16S, watering holes
Is the invited manuscript for consideration in a Special Issue? :	7th International Barcode of Life

SCHOLARONE™
Manuscripts

1
2
3
4
5

Bacterial diversity in the waterholes
of the Kruger National Park:
an eDNA metabarcoding approach

Maxwell J. Farrell^{1*}, D. Govender^{2,3},
M. Hajibabaei⁴, M. van der Bank⁵, T. J. Davies^{5,6}

¹Department of Biology, McGill University

²Scientific Services, Kruger National Park, SANParks

³Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria

⁴Integrative Biology & Biodiversity Institute of Ontario, University of Guelph

⁵African Centre for DNA Barcoding, University of Johannesburg

⁶Botany, Forest & Conservation Sciences, University of British Columbia

*To whom correspondence should be addressed; e-mail: maxwell.farrell@mail.mcgill.ca

September 21, 2018

6 **Abstract**

7 Bacteria are essential components of natural environments. They contribute to ecosystem functioning
8 through roles as mutualists and pathogens for larger species, and as key components of food webs
9 and nutrient cycles. Bacterial communities respond to environmental disturbances, and the tracking
10 of these communities across space and time may serve as indicators of ecosystem health in areas
11 of conservation concern. Recent advances in DNA sequencing of environmental samples allow for
12 rapid and culture-free characterization of bacterial communities. Here we conduct the first metabar-
13 coding survey of bacterial diversity in the waterholes of the Kruger National Park, South Africa.
14 We show that eDNA can be amplified from waterholes and find strongly structured microbial com-
15 munities, likely reflecting local abiotic conditions, animal ecology, and anthropogenic disturbance.
16 Over timescales from days to weeks we find increased turnover in community composition, indicating
17 bacteria may represent host-associated taxa of large vertebrates visiting the waterholes. Through taxo-
18 nomic annotation we also identify pathogenic taxa, demonstrating the utility of eDNA metabarcoding
19 for surveillance of infectious diseases. These samples serve as a baseline survey of bacterial diversity
20 in the Kruger, and in the future, spatially distinct microbial communities may be used as markers of
21 ecosystem disturbance, or biotic homogenization across the park.

22 **Introduction**

23 Traditional programs that monitor for early signs of ecosystem degradation require baseline data on the
24 distributions and ecology of species in an ecosystem. DNA barcoding uses differences in conserved
25 regions of genomes to classify sequences as belonging to particular taxonomic units, regardless of
26 whether or not they have been described formally by taxonomists (Hebert et al. 2003; Blaxter et al.
27 2005; Ratnasingham and Hebert 2013). DNA barcoding is thus a particularly powerful tool for explor-
28 ing microbial diversity, where there are many undescribed taxa that cannot be cultured using traditional
29 methods (Rappé and Giovannoni 2003). Molecular barcoding coupled with recent advances in genetic
30 sequencing have allowed for unprecedented exploration of microbial communities and the ability to
31 characterize organisms of interest from environmental samples with great sensitivity (Shokralla et al.
32 2012). In particular, sequencing of cellular and extracellular DNA that can be extracted from envi-
33 ronmental samples, collectively known as environmental DNA (eDNA) (Taberlet et al. 2012), is an
34 emerging approach for exploring diversity in aquatic ecosystems (Rees et al. 2014; Lodge et al. 2012).

35 Microbial diversity in freshwater systems responds to environmental conditions (Lozupone and
36 Knight 2007), and perturbations (Zeglin 2015) including multiple anthropogenic impacts such as ur-
37 banization (Fisher et al. 2015) and pollution (Bouskill et al. 2010). In addition to acting as indicators of
38 ecosystem health, changes in microbial diversity may be important in themselves. Bacteria are essen-
39 tial components of ecosystems and play important roles in food webs, nutrient recycling, disease, and
40 as important mutualists for larger multicellular species. Bacteria are thus integral to maintaining the
41 natural balance of ecosystems and shifts in taxonomic composition due to environmental change may
42 severely impact connectivity, functioning (Laforest-Lapointe et al. 2017; Delgado-Baquerizo et al.
43 2016) and increase exposure to pathogens (Cabral 2010). However, in most ecosystems of conserva-
44 tion priority, microbial diversity is poorly described.

45 Surface waters are a vital resource for savannah ecosystems (Redfern et al. 2005; Owen-Smith
46 1996), but frequent use by a large variety of species means they can also be a source of cross-species
47 infection and spread of harmful pathogens (Bengis and Erasmus 1988). These ecosystems provide
48 an ideal context for refining eDNA metabarcoding approaches as they act as sources and sinks of
49 microbial species for larger animals, however baseline information about microbial diversity in these

50 systems is lacking. Here we conduct a survey of bacterial diversity among watering holes of Kruger
51 National Park, South Africa (KNP) through spatio-temporal sampling and sequencing of the V3-V4
52 region of 16S genes present in water. Water can be scarce in the KNP throughout the dry season and
53 periods of drought (Redfern et al. 2005), and the park has a long history of water provisioning that
54 included the construction of a series of more than 300 artificial waterholes beginning in the 1930s
55 (Smit et al. 2007). These waterholes were intended to increase game numbers by stabilizing water
56 availability year-round and are frequently visited by a diversity of birds and mammals (Smit and
57 Grant 2009). However, they have proven to alter the distributions of wildlife, which in turn have
58 negative impacts on vegetation dynamics and the park-wide ecosystem (Smit et al. 2007; Smit and
59 Grant 2009). As a result, a number of artificial waterpoints have been closed since 1994 as the park
60 began reverting to a more natural cycle of water availability (Smit et al. 2007; Van Wyk 2011). A
61 subset of the waterholes still open are small concrete troughs which are well mixed, largely mud and
62 silt-free, and experience limited inflow from nearby surface waters. This means that eDNA samples
63 will largely represent microbes in sourcewater and those dispersed by air and animals, allowing us to
64 capture snapshots of local bacterial diversity across the park.

65 This study provides the first survey of bacterial diversity in the waterholes of the KNP, and is
66 among the first studies using next-generation sequencing to describe aquatic microbial diversity in
67 Africa (see Jordaan and Bezuidenhout (2016; 2013); Tekere, M., Prinsloo, A., Olivier, J., Jonker, N.,
68 Venter, S. (2012); Mwirichia, R., Cousin, S., Muigai, A. W., Boga, H. I., Stackebrandt, E. (2011);
69 Tekere, M., Lötter, A., Olivier, J., Jonker, N., Venter, S. (2011)). Here we explore bacterial diversity
70 across the southern half of the park and describe variation across, space, time, sample volume, and
71 abiotic influences.

72 **Methods**

73 **Study Site**

74 Waterholes were sampled in June and July of 2015 in the Kruger National Park, South Africa (KNP), a
75 large protected savannah ecosystem and a global diversity hotspot (Lahaye et al. 2008). Sampling was

76 conducted during the dry season when natural sources of surface water are largely dry and watering
77 hole visitation rates by medium and large vertebrates are highest. The park is divided into twenty-two
78 ranger sections, which range in size from roughly 520 to 1,170 square kilometers. Across the southern
79 half of the reserve below the Olifants river, ten concrete bottom artificial waterholes were selected
80 from five of these sections (Table 1, Fig. 1). The waterholes varied in shape with some mimicking the
81 contours of natural pans, making volume estimations difficult. However, the generic design included
82 longer and shorter axes, with comparable dimensions across waterholes. Each waterhole is equipped
83 with a ball-valve, which regulates water levels and re-fills the trough from nearby reservoirs when
84 water levels drop. Water is sourced predominantly from groundwater via boreholes, but three sites use
85 pipeline troughs filled with diverted river water.

86 **Water Sampling and Processing**

87 At each site, samples were taken once per week for three weeks. For one site (NWA), samples were
88 also taken every day for five consecutive days. Sampling consisted of two 1L water samples collected
89 in autoclaved, UV sterilized glass jars from opposite ends of the waterhole, approximately one foot
90 from the nearest edge. These two within-waterhole samples (A/B) were taken along the waterhole's
91 longest axis that maximized the distance and upwind-downwind gradient between them, if a strong
92 wind was present. Water samples were placed on icepacks in a cooler and kept between 4-8°C until
93 returning to the laboratory, where they were placed in the fridge.

94 Water quality parameters were taken during each sampling period using a YSI 650QS multi-
95 parameter sonde. Temperature (°C), conductivity (mS/cm), dissolved oxygen (in mg/L and % sat-
96 uration), and pH were recorded. Three measurements were taken along the same axis that the A/B
97 water samples were drawn, and then averaged to measure quality per sample-time.

98 In the lab, the outside of water sample collection bottles were washed with ELIMINase (Decon
99 Labs) and rinsed with deionized (DI) water to limit contamination. For each A/B sample, 150 mL of
100 water was sub-sampled and filtered through gamma-irradiated 0.2 μ m Supor hydrophilic polyether-
101 sulfone membranes (Pall no. 66234). The filtration apparatus consisted of three 300 mL Advantec
102 polysulfone 47mm filter funnels fitted to a Pall vacuum manifold with vacuum pressure maintained by

103 a Pall filtration vacuum/pressure pump (model no. 13158). After filtration, filters were stored in sterile
104 15 mL Falcon tubes and placed in a freezer at -60°C. On one sampling date for six sites, additional
105 volumes of 50 ml and 15 ml were filtered from each 1L sample to assess the impact of sample volume
106 filtered. Twice throughout sampling, BLANK samples were generated by filtering 1L of deionized
107 water used in the laboratory.

108 Prior to and between filtrations, all funnel components and tweezers used to manipulate the filters
109 were sterilized by soaking with 10% bleach for 10 minutes, rinsing with DI water, washing with
110 ELIMINase, rinsing with DI water, and subsequent exposure to UV radiation for a minimum of 30
111 minutes. Gloves were worn at all times and changed between samples to minimize cross-sample
112 contamination. To avoid sample freezing and bacterial growth in collection jars, all samples were
113 processed within 12 hours of collection. Frozen 50 mL unfiltered voucher samples were kept and
114 placed at -80°C at the University of Johannesburg's African Centre for DNA Barcoding for long term
115 storage.

116 DNA Extraction, Amplification, and Sequencing

117 DNA was isolated from filter papers using MO BIO PowerWater DNA Isolation Kits. Universal bac-
118 terial primer sets designed by Sundquist et al. (2007) (V3-F: 5'ACTCCTACGGGAGGCAGCAG 3';
119 V4-R: 5'GGACTACARGGTATCTAAT 3') tagged with an Illumina adapter sequence were used to
120 amplify the V3-V4 hypervariable region of the 16S ribosomal RNA gene through polymerase chain
121 reaction (PCR). The PCR used a standard mix of 17.8µL molecular grade water, 2.5µL 10 reaction
122 buffer (200mM Tris HCl, 500mM KCl, pH 8.4), 1µL MgCl₂ (50mM), 0.5µL dNTP (10mM), 0.5µL
123 forward primer (10mM), 0.5µL reverse primer (10mM), 0.2µL Platinum Taq DNA polymerase (Invit-
124 rogen), and 2µL DNA as template for a total volume of 25µL. PCRs underwent the following cyclor
125 conditions: initial 94°C for 5 minutes, then 30 cycles of 94°C for 40 seconds, 46°C for 1 minute, 72°C
126 for 30 seconds, and a final temperature of 72°C for 2 minutes. Amplification success was confirmed
127 through gel electrophoresis, using a 1.5% agarose gel. PCR products were purified using MinElute
128 PCR purification kit (Qiagen), and quantified through fluometry using a Quant-iT PicoGreen dsDNA
129 assay kit (Invitrogen). Samples were normalized, then multiplexed with the Nextera XT Index kit (96

indexes) (Illumina) and sequenced on an Illumina MiSeq flowcell using a V2 sequencing chemistry kit (2 x 250) making up approximately 1/8th of the run.

Sequence Processing, Taxonomy Assignment, and Phylogeny Construction

Across all samples, we generated a total of 2,164,262 Illumina reads. Primer sequences were removed using the trim.seqs function in mothur (Schloss et al. 2009). Reads were then processed in R (version 3.4.3) (R Development Core Team 2008) using the package *dada2* version 1.6.0 (Callahan et al. 2016) following a modified version of the DADA2 Bioconductor workflow (Callahan et al. 2017) and online tutorials v1.6 and workflow for big data v1.4 (benjjneb.github.io/dada2/tutorial.html). Reads were filtered by quality, removing sequences with maximum expected error (maxEE) greater than 6 for both forward and reverse reads, and reads with any base pair having Q of 6 or lower. Reads were truncated to a length of 230bp and 220bp for forward and reverse reads respectively, consistent with dropoffs in quality profiles, and reads shorter than this were removed. Since the samples were sequenced across four different runs, subsequent steps of learning error rates, dereplication, denoising and Amplicon Sequence Variant (ASV) calling (Callahan et al. 2017) using pooled samples, and merging of paired reads were performed separately for each run. Tables of ASV sequences per sample within each run were then combined and chimera detection using all pooled samples was performed (see SM Table S2 for the number of reads retained across each step). In total 1,184,831 reads were retained, representing 3533 ASVs.

Taxonomy assignment from Kingdom to Genus was performed using the RDP classifier and SILVA nr v128 reference database (Quast et al. 2013) formatted for DADA2 (benjjneb.github.io/dada2/training.html), using the assignTaxonomy function (Fig. 5). ASVs assigned as Archaea, Eukarya, Chloroplast, or Mitochondria were removed. Species level assignments were added by exact sequence matching using the addSpecies function. ASV sequences were aligned with the pynast algorithm via align_seqs.py in QIIME (Caporaso et al. 2010) and sequences with poor alignment automatically removed. A phylogenetic tree was constructed using the GTRCAT model in FastTree version 2.1.3 (Price et al. 2010) after filtering nucleotides with greater than 90% gap fraction and removing the 5% highest entropy positions with filter_alignment.py in QIIME (Caporaso et al. 2010). This resulted in a phylogenetic

157 tree of 3393 ASVs which were used in subsequent community analyses.

158 **Community Analyses**

159 The ASV sequence table was merged with the phylogeny and sample metadata using the R package
160 *phyloseq* version 1.22.3 (McMurdie and Holmes 2013). Negative controls (BLANK samples) used to
161 investigate contamination during sample filtration contained 43 ASVs collectively (with 9 ASVs found
162 in both samples). The sequence reads in each filtration blank were both dominated by the same ASV
163 (53% and 86% respectively), however none of the 43 ASVs identified in the blanks were identified in
164 any of the other samples. These control samples were removed prior to community analyses.

165 A subset of core samples was created by removing the first four daily NWA samples and samples
166 of differential volume (S & XS samples), resulting in 54 samples of 150 mL each (Table 2). An ASV
167 accumulation curve for core samples was generated using the *specaccum* function in the R package
168 *vegan* version 2.4.6 (Oksanen et al. 2018) using the “exact” method, and extrapolated to total ASV
169 richness using the Chao and Bootstrap methods in *vegan*’s *specpool* function. Alpha diversity was
170 calculated for the core samples as observed ASV richness and Shannon diversity using the *phyloseq*
171 package. Additive partitioning of Shannon diversity across core samples was investigated using the
172 *adipart* function in *vegan* (Table 3).

173 Taxonomic composition was assessed by merging core samples at each site, and plotting rela-
174 tive abundances of reads for the most common taxa at levels of phylum, class, and order (Fig. 6).
175 Temporal variation in taxonomic composition across core samples was assessed by merging A and B
176 samples and plotting relative abundances of reads for sites with two or more weekly samples, for the
177 levels of phylum (Fig. 7), and class (Fig. S6). To further investigate fine-scale temporal variation in
178 taxonomic composition (phylum, class, and order), relative abundance of reads were plotted across
179 the daily samples at site NWA (Fig. 8). We also explored temporal turnover among samples with
180 Sorensen’s dissimilarity calculated using the *beta.pair* function from the *betapart* package (Baselga
181 et al. 2018) (Fig. S9) and significant differences among daily and weekly samples was assessed using
182 permutational multivariate analysis of variance (*adonis* in *vegan*) with 999 permutations each.

183 Community composition across sites in the core samples was described with nonmetric multidimensional

184 mentional scaling (NMDS) ordinations on relative ASV abundances per sample using the Bray-Curtis
185 dissimilarity, and the abundance weighted Unifrac dissimilarity (Figs. 9, S10 & S11). Statistically,
186 associations between dissimilarities and both water quality properties and common taxonomic groups
187 were assessed using the envfit function in *vegan* for bacterial classes (Fig. 10) and orders (Fig. S12).

188 Phylogenetic community structure across core samples was calculated using standardized effect
189 sizes of mean pairwise phylogenetic distances (MPD) and mean nearest taxon distances (MNTD) in
190 the R package *picante* version 1.6.2 (Kembel et al. 2010) using the abundance weighted “richness”
191 null model and 999 randomizations in the *ses.mpd* and *ses.mntd* functions (Fig. S13). For a given sam-
192 ple, MPD calculates the mean phylogenetic distance among each pair of taxa present, while MNTD
193 calculates the mean phylogenetic distance from each taxa to its closest relative. These raw metrics
194 give an estimate of how closely related community members are to each other, and are then compared
195 to randomized communities to determine whether the observed metrics are different than what would
196 expected if communities were assembled at random from taxa pooled across all samples.

197 To assess the effect of differential sample volumes, S (50 mL) and XS (15 mL) samples were subset
198 along with their corresponding full volume samples (150 mL). Alpha diversity, calculated as observed
199 ASV richness and Shannon diversity were calculated as described above (Fig. S14). Variation in
200 taxonomic composition was investigated by comparing relative read abundances of bacterial phyla in
201 A/B samples across sites and differential volumes (Fig. S15).

202 Raw reads with primers removed are available via the NCBI Sequence Read Archive BioProject
203 PRJNA490450 (accession numbers SRR7822814 to SRR7822901). The ASV table, taxonomic as-
204 signments, phylogenetic tree, sample metadata, and scripts necessary to reproduce the results can be
205 found at 10.6084/m9.figshare.7119668.

206 Results

207 Our sampling design aimed to sequence a core set of samples with all ten sites being sampled once
208 per week for three weeks. Due to logistic constraints of sample storage and extremely low water
209 levels from drawdown by animals, we were only able to process 27 of the planned 30 weekly samples
210 (Table 2). In addition to this core sampling, we sequenced differential volumes for six samples, and

an additional four daily samples from Nwaswitshaka (site NWA). A/B samples were taken at each site-time, resulting in a total of 88 sequenced samples, including the two filtration blanks. Across all 88 samples, we identified a total of 3393 ASVs. Roughly 15% of ASVs (n=524) were represented by a single read, together comprising fewer than 0.05% of all reads. The DADA2 approach infers the biological sequences in the sample prior to the introduction of amplification and sequencing errors, and can distinguish sequence variants differing by as little as one nucleotide. As such, we included all ASVs, including those represented by single reads, in subsequent analyses of biodiversity.

The ASV accumulation curve generated for the core sample set (2603 ASVs) does not appear to saturate (Fig. 2). Estimates of total richness using the Chao estimator predicts 6164 ASVs (+/- 262 SE) among the core samples, indicating we may be capturing less than half of the total bacterial diversity present among our sites. However, estimates of total diversity using the Bootstrap method were more conservative, with 3260 (+/- 146 SE) estimated ASVs. ASV diversity varied across sites (Figs. 3, Fig. 4), but the largest turnover (β diversity) was observed among park sections (Table 3). Variation among A/B samples contributed very small amounts to β diversity, indicating that at a particular time, microbial diversity within each waterhole was fairly well mixed.

In terms of taxonomic composition, 99.2% of ASVs were assigned to a known phylum, with the proportion of assignments decreasing at lower taxonomic levels (Fig. 5). The majority of bacterial ASVs were classified as Proteobacteria (~ 59%), followed by Bacteroidetes (~ 14%), Firmicutes (~ 9%), Actinobacteria (~ 6%), and Verrucomicrobia (~ 2%). For bacterial classes, ASVs were largely classified as Betaproteobacteria (~ 34%), Alphaproteobacteria (~ 11%), Gammaproteobacteria (~ 10%), and Sphingobacteriia (~ 6%). Among core samples, relative abundances of phyla, classes, and orders varied across sites (Figs. 6, S4, S5). Across weeks, relative abundances of phyla varied within each site (Fig. 7), with some sites displaying more stability (IMB & HOY) compared to others (NYA & NGO). Patterns among bacterial classes (Fig. S6) largely reflected variation among phyla, though one site (HOY) displayed much more variation in relative abundances among classes, reflecting substantial turnover within Proteobacteria. Comparing weekly turnover with the five daily samples taken at Nwaswitshaka (NWA) (Fig. 8), taxonomic composition appeared more stable across days than weeks. Using hierarchical clustering of Sorensen's dissimilarity, we find that samples taken within

239 a single week cluster together (Fig. S9). Permutational multivariate ANOVAs on these distances
240 revealed a significant difference in beta diversity among weekly samples (NWA 2,7,8; $p = 0.02$), with
241 49% of the variance explained by sample date, but no significant difference among the additional daily
242 samples (NWA 3,4,5,6; $p = 0.54$), with 14% of the variance explained by sample date.

243 Community composition visualized through NMDS ordinations reflected results from the additive
244 partitioning of diversity, with core samples clustering by site (Fig. 9) and section (Fig. S10) for both
245 Bray-Curtis and abundance weighted UniFrac dissimilarities. Interestingly, waterholes filled by water
246 from pipeline troughs (NGO, NYA, WIT) grouped together (Fig. S10), although these three sites are
247 situated on a different geological type than sites fed by boreholes, making us unable to differentiate
248 the effects of each factor (Fig. S11). Bacterial community composition was significantly structured
249 by conductivity and pH for both Bray-Curtis and UniFrac dissimilarities, and dissolved oxygen also
250 had an influence on UniFrac dissimilarity (Figs. 10 & S12). The dissimilarity of high conductivity
251 sites (particularly HOY & IMB in the Kingfisherspruit section) was associated with high abundances
252 of Clostridia, Gammaproteobacteria, and Bacteroidia, while sites with high pH and dissolved oxygen
253 were positively related to the abundances of Actinobacteria and Alphaproteobacteria (Fig. 10).

254 Reflecting the NMDS structure of the abundance weighted UniFrac dissimilarities, MNTD, which
255 is most sensitive to phylogenetic structure towards the tips of the tree (Mazel et al. 2015), indicated
256 strong phylogenetic clustering within the majority of samples (Fig. S13). The strength of clustering
257 was weaker for MPD, which is more sensitive to phylogenetic structure deeper in the tree (Mazel et al.
258 2015).

259 We did not find any clear decrease in alpha diversity with smaller sample volumes (Fig. S14),
260 and one of the 15 mL samples returned the largest richness of ASVs, though the median value for
261 15 mL samples was lower and had a larger interquartile range than the 50 mL and 150 mL samples.
262 The major phyla detected within samples was also relatively consistent, with most groups represented
263 across different sample volumes, though not always in the same proportions (Fig. S15).

Discussion

Biological monitoring is an essential aspect of conservation for tracking contemporary changes in ecosystems as well as providing a historical baseline for making management decisions. The Kruger National Park, established in 1898, has a long history of management practices revolving around maintenance of large mammals (Venter et al. 2008). While bacterial diversity has been explored for important infectious agents in the system (Michel et al. 2007; Bengis and Erasmus 1988; Smith et al. 2000), recent advances in next generation sequencing methods now allow for the rapid and culture-free description of bacterial diversity throughout the park.

Here we present the first description of bacterial diversity in the waterholes of the Kruger National Park. In total we identified over 3000 unique taxa (referred to as amplicon sequence variants, or ASVs), only about half of which could be assigned to a previously described genus. The relative dominance of bacterial phyla was consistent with bacterial surveys of the Vaal River in central South Africa (Jordaan and Bezuidenhout 2016). However, bacterial diversity was strongly structured across space, with the largest turnover in diversity occurring among park sections. This is not surprising considering the distances from site to site range from 3km to 115km and represent a gradient in large animal density, rainfall, vegetation, and major subsurface geology (Chirima et al. 2012; Van Wilgen et al. 2000; Smit and Grant 2009; Smit et al. 2013). Samples also clustered by site, displaying substantial variation in taxonomic composition across sites. This variation was associated with physico-chemical properties of the water, with conductivity and pH being important explanatory variables. In addition to water quality, variability in taxonomic composition is likely influenced by the origin of the water used to fill each waterhole, differences in the surrounding soil and vegetation types, and the particular species and populations of animals using the waterholes.

We assessed daily turnover in composition at Nwaswitshaka, which appeared to be more stable over this shorter timescale when compared to turnover across weeks. However, Nwaswitshaka was less variable across weeks than other sites, indicating that daily variation in bacterial communities could be greater in other locations. Important water quality variables (conductivity and pH) were largely consistent across weeks (Table S1), suggesting that the observed temporal heterogeneity may be driven by differences in external factors influencing bacterial input and removal from the system,

such as variation in animal visitation throughout the sampling period. Between sampling events, water levels would sometimes drop substantially, indicating major drawdown by animals and likely removing bacteria deposited by animals visiting earlier in the week. Large mammal communities vary across the sampled regions of the park (Chirima et al. 2012), which may contribute to observed spatial variation in bacterial communities. However, different species also differ in their dependence on water, which is reflected in their rates of visitation to water points (Redfern et al. 2005). Variation in samples taken across subsequent weeks may therefore reflect different components of local animal communities, each with their unique host-associated bacterial taxa (Ley et al. 2008). By pairing bacterial composition with animal visitation prior to sampling (either through direct observation, or presence of genetic material), it may be feasible to build statistical probabilities of associations using co-occurrences of microbial and animal signatures.

Across samples, patterns of phylogenetic clustering were consistent with observed taxonomic variation. Multiple phyla were present in all samples, consistent with an even representation of deep bacterial lineages. However, turnover at lower taxonomic levels shown by significantly low mean nearest taxon distances indicate that there are distinct subsets of closely related taxa present at each site. This structuring may reflect filtering of bacterial communities by local environmental conditions, or the deposition of microbes by particular animal populations or individuals. Many vertebrate species have expansive home ranges, but during the dry season drought-intolerant animals will restrict their movement so as to stay close to permanent water bodies (Redfern et al. 2005). Thus the maintenance of major bacterial taxa may reflect both free-living environmental bacteria, and the core microbiome of water-dependent species. By taking repeat temporal samples, it may be possible to build association networks between bacterial taxa and host species, or even their local populations, solely from environmental DNA.

Through examination of taxonomic assignments we identified taxa belonging to genera that include important pathogens (*Arcobacter*, *Bacillus*, *Burkholderia*, *Coxiella*, *Legionella*, *Neisseria*, *Pasteurella*, *Rickettsia*, and *Yersinia*). While many of these genera include both pathogenic as well as benign species found in environmental samples, some of these genera are comprised solely of pathogenic species. For example, the genus *Coxiella* is represented by one species, *Coxiella burnetii*, the causative

agent of Q fever, which has previously been documented as causing disease in the park (Van Heerden et al. 1995). Additionally, taxa in the order *Chlamydiales* are all obligate intracellular pathogens of eukaryotes (Ball et al. 2015), and taxa in the genus *Neisseria* colonize mucosal surfaces of animals, some of which are pathogenic in humans (Liu et al. 2015). Interestingly, we also identified sequences classified as *Streptococcus urinalis*, a recently described species linked to urinary tract infections in humans (Peltroche-Llacsahuanga et al. 2012). While hypervariable regions of the 16S gene may not be the optimal genomic regions for detecting the presence of particular pathogenic species or strains, our findings indicate that broad scale surveys of microbial diversity may be useful in determining the presence of potential pathogens across vastly divergent groups of bacteria. This can in turn guide more targeted sampling of both pathogenic and commensal bacteria across the park.

In addition to the presence of pathogens, surveys of bacterial diversity may be used to detect anthropogenic influences in the park. For example, we found that bacterial diversity was quite different for the two sites sampled in the Kingfisherspruit section (Hoyo Hoyo and Imbali). These sites were dominated by an ASV assigned to the genus *Arcobacter*, but which did not exactly match any sequence in the SILVA reference database. Three of the five described members of this genus are known to be pathogenic (Fera et al. 2004) and include *A. butzleri*, which can cause severe diarrhea (Lerner et al. 1994) and was detected in two of the weekly samples at Hoyo Hoyo. The two waterholes in Kingfisherspruit are fed in part with greywater that is passed through reed beds. Greywater is untreated household wastewater that typically has not been contaminated by toilet waste and is often used as year-round sources of water, especially in water scarce areas (Nganga et al. 2012). Compared to source water, kitchen and laundry greywater can have elevated conductivity (Nganga et al. 2012), which may explain the high conductivity of water at these sites, and present a strong selective environment driving their unique bacterial communities.

We did not have sufficient sampling of sites to explore all possible drivers of differences in bacterial communities. Nonetheless, some features differed obviously among sites. For example, Witpens is a heavily vulture-dominated site. Bathing by vultures likely results in large influxes of nutrients such as blood, and vulture feces has been found to alter soil bacterial communities through elevated nitrogen and decreased pH (Ganz et al. 2012). Visually, water from Witpens was bright green, indi-

348 cating high abundance of photosynthetic species and consistent with the large variations observed in
349 dissolved oxygen (Table S1). However, we found no evidence of elevated abundances of *Microcystis*
350 or other Cyanobacteria, though samples from Witpens strongly clustered together and had relatively
351 high abundances of Rhizobiales and Rhodocyclales, both of which include species known to fix nitro-
352 gen (Carvalho et al. 2010; Loy et al. 2005).

353 Witpens, along with Nyamarhi and Ngotso North are filled by pipeline troughs that divert river
354 water to waterholes many kilometers away. While pipeline troughs are likely to reflect a subsample of
355 the diversity found in river water, the acts of pipeline transport and storage themselves may have strong
356 filtering effects on bacterial communities. Our results indicate that waterholes represent locally unique
357 bacterial communities, thus the practice of diverting river water to waterholes kilometers away may
358 homogenize microbial diversity across the landscape, ultimately disrupting local communities. The
359 consequences of such shifts in community structure are difficult to assess without comparing pipeline
360 troughs with their source waters, but the diversion of river water may have unintended impacts on
361 microbial diversity. For example, genes conferring antimicrobial resistance have been shown to spread
362 from river water to impala in the Kruger National Park (Mariano et al. 2009).

363 Here we show that eDNA can be amplified from waterholes in the Kruger National Park, and find
364 strongly structured microbial communities, likely reflecting local abiotic conditions, animal ecology,
365 and anthropogenic disturbance. We suggest that disruption of spatially distinct microbial communi-
366 ties may be used as a marker of ecosystem disturbance, or biotic homogenization across the park.
367 We find that for artificial waterholes, bacterial diversity is surprisingly insensitive to sample volume,
368 with even small volumes useful for capturing major components of bacterial communities, though
369 larger volumes are necessary to detect rare taxa. Replicating this study across different seasons, and
370 expanding sampling to include natural waterpoints will provide improved understanding of the roles
371 micro-organisms play in ecosystem stability and resilience, and offer an effective method for mon-
372 itoring of shifting species interactions in the face of environmental change. Just as studies of the
373 microbiome have revolutionized our understanding of human health, metagenomic analysis of envi-
374 ronmental DNA have the potential to revolutionize our understanding of ecosystem health. Tracking
375 of bacterial communities can provide a template for monitoring ecosystem disturbance through their

376 response to biological contaminants, documenting the spread of invasive species or infectious organ-
377 isms, and better understanding the impacts ecological disturbances have on the composition of native
378 communities.

379 **Acknowledgements**

380 This work would not have been possible without the logistical support of the African Centre for DNA
381 Barcoding and Purvance Shikwambana, molecular work by Shannon Eagle, encouragement and sup-
382 port from Lea Berrang-Ford, and the incredible knowledge and protection provided by Kruger game
383 guards Thomas, Velly, and Martin. We also thank Steven Kembel, his lab, Alexis Carteron, and
384 Benjamin Callahan for constructive feedback on bioinformatic analyses. Special thanks are due to
385 B.S. Bezeng for volunteering his friendship, dedication, and long days in the field. This work was
386 performed with the support and permission of SANParks (project DAVIJ1256). MJF was supported
387 by a Vanier NSERC CGS and the CIHR Systems Biology Training Program, with project funding
388 supported by the Quebec Centre for Biodiversity Science, the McGill Biology Department, and an
389 NSERC Discovery Grant awarded to TJD.

390 **References**

- 391 Ball, S. G., C. Colleoni, D. Kadouche, M. Ducatez, M. C. Arias, and C. Tirtiaux. 2015. Toward an
392 understanding of the function of Chlamydiales in plastid endosymbiosis. *Biochimica et Biophysica*
393 *Acta - Bioenergetics* 1847(6-7), 495–504.
- 394 Baselga, A., D. Orme, S. Villeger, J. De Bortoli, and F. Leprieur. 2018. betapart: Partitioning Beta
395 Diversity into Turnover and Nestedness Components. R package version 1.5.0.
- 396 Bengis, R. G. and J. M. Erasmus. 1988. Wildlife diseases in South Africa: a review. *Revue Scientifique*
397 *et Technique, (OIE)* 7(4), 807–821.
- 398 Blaxter, M., J. Mann, T. Chapman, F. Thomas, C. Whitton, R. Floyd, et al.. 2005. Defining operational

- 399 taxonomic units using DNA barcode data. *Philosophical Transactions of the Royal Society B:*
400 *Biological Sciences* 360(1462), 1935–1943.
- 401 Bouskill, N. J., J. Barker-Finkel, T. S. Galloway, R. D. Handy, and T. E. Ford. 2010. Temporal bacterial
402 diversity associated with metal-contaminated river sediments. *Ecotoxicology* 19(2), 317–328.
- 403 Cabral, J. P.. 2010. Water microbiology. Bacterial pathogens and water. *International Journal of*
404 *Environmental Research and Public Health* 7(10), 3657–3703.
- 405 Callahan, B. J., P. J. McMurdie, and S. P. Holmes. 2017. Exact sequence variants should replace
406 operational taxonomic units in marker-gene data analysis. *ISME Journal* 11(12), 2639–2643.
- 407 Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016.
408 DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13(7),
409 581–583.
- 410 Callahan, B. J., K. Sankaran, J. A. Fukuyama, P. J. McMurdie, and S. P. Holmes. 2017. Bio-
411 conductor Workflow for Microbiome Data Analysis: from raw reads to community analyses.
412 *F1000Research* 5(3), 1492.
- 413 Caporaso, J. G., J. Kuczynski, J. Stombaugh, K. Bittinger, F. D. Bushman, E. K. Costello, et al.. 2010.
414 QIIME allows analysis of high- throughput community sequencing data Intensity normalization
415 improves color calling in SOLiD sequencing. *Nature Publishing Group* 7(5), 335–336.
- 416 Carvalho, F. M., R. C. Souza, F. G. Barcellos, M. Hungria, and A. T. R. Vasconcelos. 2010. Genomic
417 and evolutionary comparisons of diazotrophic and pathogenic bacteria of the order Rhizobiales.
418 *BMC Microbiology* 10.
- 419 Chirima, G. J., N. Owen-Smith, and B. F. Erasmus. 2012. Changing distributions of larger ungulates
420 in the Kruger National Park from ecological aerial survey data. *Koedoe* 54(1), 1–11.
- 421 Delgado-Baquerizo, M., F. T. Maestre, P. B. Reich, T. C. Jeffries, J. J. Gaitan, D. Encinar, et al.. 2016.
422 Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature Communications* 7,
423 1–8.

- 424 Fera, M. T., T. L. Maugeri, C. Gugliandolo, C. Beninati, E. L. Camera, M. Carbone, et al.. 2004.
425 Detection of *Arcobacter* spp . in the Coastal Environment of the Mediterranean Sea Detection of
426 *Arcobacter* spp . in the Coastal Environment of the Mediterranean Sea. *Applied and Environmental*
427 *Microbiology* 70(3), 1271–1276.
- 428 Fisher, J. C., R. J. Newton, D. K. Dila, and S. L. McLellan. 2015. Urban microbial ecology of a
429 freshwater estuary of Lake Michigan. *Elementa: Science of the Anthropocene* 3, 000064.
- 430 Ganz, H. H., U. Karaoz, W. M. Getz, W. Versfeld, and E. L. Brodie. 2012. Diversity and structure of
431 soil bacterial communities associated with vultures in an African savanna. *Ecosphere* 3(6), art47.
- 432 Hebert, P. D. N., A. Cywinska, S. L. Ball, and J. R. DeWaard. 2003. Biological identifications through
433 DNA barcodes. *Proceedings. Biological sciences / The Royal Society* 270(1512), 313–21.
- 434 Jordaan, K. and C. C. Bezuidenhout. 2013. The impact of physico-chemical water quality parameters
435 on bacterial diversity in the Vaal River, South Africa. *Water SA* 39(3), 385–396.
- 436 Jordaan, K. and C. C. Bezuidenhout. 2016. Bacterial community composition of an urban river in the
437 North West Province, South Africa, in relation to physico-chemical water quality. *Environmental*
438 *Science and Pollution Research* 23(6), 5868–5880.
- 439 Kembel, S., P. Cowan, M. Helmus, W. Cornwell, H. Morlon, D. Ackerly, et al.. 2010. Picante: R tools
440 for integrating phylogenies and ecology. *Bioinformatics* 26, 1463–1464.
- 441 Laforest-Lapointe, I., A. Paquette, C. Messier, and S. W. Kembel. 2017. Leaf bacterial diversity
442 mediates plant diversity and ecosystem function relationships. *Nature* 546(7656), 145–147.
- 443 Lahaye, R., M. van der Bank, D. Bogarin, J. Warner, F. Pupulin, G. Gigot, et al.. 2008. DNA barcoding
444 the floras of biodiversity hotspots. *Proceedings of the National Academy of Sciences* 105(8), 2923–
445 2928.
- 446 Lerner, J., V. Brumberger, and V. Preac-Mursic. 1994. Severe diarrhea associated with *Arcobacter*
447 *butzleri*. *European Journal of Clinical Microbiology & Infectious Diseases* 13(8), 660–662.

- 448 Ley, R. E., M. Hamady, C. Lozupone, P. J. Turnbaugh, R. R. Ramey, J. S. Bircher, et al.. 2008.
449 Evolution of mammals and their gut microbes. *Science* 320(5883), 1647–1651.
- 450 Liu, G., C. M. Tang, and R. M. Exley. 2015. Non-pathogenic *Neisseria*: members of an abundant,
451 multi-habitat, diverse genus. *Microbiology (Reading, England)* 161(7), 1297–1312.
- 452 Lodge, D., C. Turner, and C. Jerde. 2012. Conservation in a cup of water: estimating biodiversity and
453 population abundance from environmental DNA. *Molecular Ecology* 21(11), 2555–8.
- 454 Loy, A., C. Schulz, S. Lückner, A. Schöpfer-Wendels, K. Stoecker, C. Baranyi, et al.. 2005. 16S rRNA
455 gene-based oligonucleotide microarray for environmental monitoring of the betaproteobacterial or-
456 der "Rhodocyclales". *Applied and Environmental Microbiology* 71(3), 1373–1386.
- 457 Lozupone, C. A. and R. Knight. 2007. Global patterns in bacterial diversity. *Proceedings of the*
458 *National Academy of Sciences of the United States of America* 104(27), 11436–11440.
- 459 Mariano, V., C. M. E. McCrindle, B. Cenci-Goga, and J. A. Picard. 2009. Case-control study to deter-
460 mine whether river water can spread tetracycline resistance to unexposed impala (*aepyceros melam-*
461 *pus*) in Kruger National Park (South Africa). *Applied and Environmental Microbiology* 75(1),
462 113–118.
- 463 Mazel, F., T. J. Davies, L. Gallien, J. Renaud, M. Groussin, T. Munkemuller, et al.. 2015. Influence of
464 tree shape and evolutionary time-scale on phylogenetic diversity metrics. *Ecography*.
- 465 McMurdie, P. J. and S. Holmes. 2013. Phyloseq: An R Package for Reproducible Interactive Analysis
466 and Graphics of Microbiome Census Data. *PLoS ONE* 8(4).
- 467 Michel, A. L., L.-M. de Klerk, N. van Pittius, R. M. Warren, and P. D. van Helden. 2007. Bovine
468 tuberculosis in African buffaloes: observations regarding *Mycobacterium bovis* shedding into water
469 and exposure to environmental mycobacteria. *BMC Veterinary Research* 3(1), 23.
- 470 Mwirichia, R., Cousin, S., Muigai, A. W., Boga, H. I., Stackebrandt, E.. 2011. Bacterial diversity in
471 the haloalkaline lake elmenteita, Kenya. *Current Microbiology* 62(1), 209–221.

- 472 Nganga, V. G., F. W. Kariuki, and K. Kotut. 2012. A comparison of the physico-chemical and bac-
473 teriological quality of greywater from water deficient households in Homabay town and Githurai
474 estates in Kenya. *Open Environmental Engineering Journal* 5, 110–118.
- 475 Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, et al.. 2018. *vegan*:
476 *Community Ecology Package*. R package version 2.4-6.
- 477 Owen-Smith, N.. 1996. Ecological guidelines for waterpoints. *South African Journal of Wildlife*
478 *Research* 26(4), 107–112.
- 479 Peltroche-Llacsahuanga, H., B. Frye, and G. Haase. 2012. Isolation of *Streptococcus urinalis* from a
480 human blood culture. *Journal of Medical Microbiology* 61(5), 740–742.
- 481 Price, M. N., P. S. Dehal, and A. P. Arkin. 2010. FastTree 2 - Approximately maximum-likelihood
482 trees for large alignments. *PLoS ONE* 5(3).
- 483 Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, et al.. 2013. The SILVA riboso-
484 mal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids*
485 *Research* 41(D1), 590–596.
- 486 R Development Core Team. 2008. *R: A Language and Environment for Statistical Computing*. Vienna,
487 Austria: R Foundation for Statistical Computing. ISBN 3-900051-07-0.
- 488 Rappé, M. S. and S. J. Giovannoni. 2003. The Uncultured Microbial Majority. *Annual Review of*
489 *Microbiology* 57(1), 369–394.
- 490 Ratnasingham, S. and P. D. N. Hebert. 2013. A DNA-Based Registry for All Animal Species: The
491 Barcode Index Number (BIN) System. *PLoS ONE* 8(7).
- 492 Redfern, J., C. Grant, A. Gaylard, and W. Getz. 2005. Surface water availability and the management
493 of herbivore distributions in an African savanna ecosystem. *Journal of Arid Environments* 63(2),
494 406–424.
- 495 Rees, H. C., B. C. Maddison, D. J. Middleditch, J. R. M. Patmore, and K. C. Gough. 2014. The

- 496 detection of aquatic animal species using environmental DNA - a review of eDNA as a survey tool
497 in ecology. *Journal of Applied Ecology*.
- 498 Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, et al.. 2009. Intro-
499 ducing mothur: Open-source, platform-independent, community-supported software for describing
500 and comparing microbial communities. *Applied and Environmental Microbiology* 75(23), 7537–
501 7541.
- 502 Shokralla, S., J. L. Spall, J. F. Gibson, and M. Hajibabaei. 2012. Next-generation sequencing tech-
503 nologies for environmental DNA research. *Molecular ecology* 21(8), 1794–805.
- 504 Smit, I. and C. Grant. 2009. Managing surface-water in a large semi-arid savanna park: Effects on
505 grazer distribution patterns. *Journal for Nature Conservation* 17(2), 61–71.
- 506 Smit, I. P., C. C. Grant, and B. J. Devereux. 2007. Do artificial waterholes influence the way herbivores
507 use the landscape? Herbivore distribution patterns around rivers and artificial surface water sources
508 in a large African savanna park. *Biological Conservation* 136(1), 85–99.
- 509 Smit, I. P., E. S. Riddell, C. Cullum, and R. Petersen. 2013. Kruger National Park research supersites:
510 Establishing long-term research sites for cross-disciplinary, multiscaled learning. *Koedoe* 55(1),
511 1–7.
- 512 Smith, K. L., V. DeVos, H. Bryden, L. B. Price, M. E. Hugh-Jones, and P. Keim. 2000. *Bacillus*
513 *anthracis* diversity in Kruger National Park. *Journal of Clinical Microbiology* 38(10), 3780–3784.
- 514 Sundquist, A., S. Bigdeli, R. Jalili, M. L. Druzin, S. Waller, K. M. Pullen, et al.. 2007. Bacterial
515 flora-typing with targeted, chip-based Pyrosequencing. *BMC Microbiology* 7(1), 108.
- 516 Taberlet, P., E. Coissac, M. Hajibabaei, and L. H. Rieseberg. 2012. Environmental DNA. *Molecular*
517 *ecology* 21(8), 1789–93.
- 518 Tekere, M., Lötter, A., Olivier, J., Jonker, N., Venter, S.. 2011. Metagenomic analysis of bacte-
519 rial diversity of Siloam hot water spring, Limpopo, South Africa. *African Journal of Biotechnol-*
520 *ogy* 10(78), 18005–18012.

- 521 Tekere, M., Prinsloo, A., Olivier, J., Jonker, N., Venter, S.. 2012. An evaluation of the bacterial
522 diversity at Tshipise, Mphephu and Sagole hot water springs, Limpopo Province, South Africa.
523 African Journal of Microbiology Research 6(23), 4993–5004.
- 524 Van Heerden, J., M. G. Mills, M. J. Van Vuuren, P. J. Kelly, and M. J. Dreyer. 1995. An investigation
525 into the health status and diseases of wild dogs (*Lycaon pictus*) in the Kruger National Park. Journal
526 of the South African Veterinary Association 66(1), 18–27.
- 527 Van Wilgen, B. W., H. C. Biggs, S. P. O'Regan, and N. Mare. 2000. A fire history of the savanna
528 ecosystems in the Kruger National Park, South Africa, between 1941 and 1996.
- 529 Van Wyk, L.. 2011. Review of the ecological implications of artificial waterhole closures in the Kruger
530 National Park and the effect thereof on tourism. pp. 1–80.
- 531 Venter, F. J., R. J. Naiman, H. C. Biggs, and D. J. Pienaar. 2008. The evolution of conservation
532 management philosophy: Science, environmental change and social adjustments in Kruger National
533 Park. Ecosystems 11(2), 173–192.
- 534 Zeglin, L. H.. 2015. Stream microbial diversity in response to environmental changes: Review and
535 synthesis of existing research. Frontiers in Microbiology 6(MAY), 1–15.

536 **Figures & Tables**

Section	Site	Type	Geology
Tshokwane (TSH)	Nhlanguleni (NHL)	Borehole	Granite
Skukuza (SKZ)	Nwaswitshaka (NWA)	Borehole	Granite
Skukuza (SKZ)	De LaPorte (DLP)	Borehole	Granite
Skukuza (SKZ)	Kwaggas Pan (KWA)	Borehole	Granite
Satara (SAT)	Girivana (GIR)	Borehole	Granite
SaTara (SAT)	Witpens (WIT)	Pipeline trough	Basalt
Kingfisherpruit (KFI)	Imbali (IMB)	Borehole	Granite
Kingfisherpruit (KFI)	Hoyo Hoyo (HOY)	Borehole	Granite
Houtboschrand (HOU)	Nyamarhi (NYA)	Pipeline trough	Basalt
Houtboschrand (HOU)	Ngosto North (NGO)	Pipeline trough	Basalt

Table 1: Sample locations

Figure 1: Map of site locations with park boundary indicated by dashed line. Circles represent sites filled by boreholes while triangles represent sites filled by river water via pipeline troughs.

Site	Weeks	S	XS	Daily	A/B	Total
Nhlanguleni (NHL)	3	0	0	0	Yes	6
Nwaswitshaka (NWA)	3	1	1	4	Yes	18
De LaPorte (DLP)	1	1	1	0	Yes	6
Kwaggas Pan (KWA)	2	1	1	0	Yes	8
Girivana (GIR)	3	0	0	0	Yes	6
Witpens (WIT)	3	0	0	0	Yes	6
Imbali (IMB)	3	0	0	0	Yes	6
Hoyo Hoyo (HOY)	3	1	1	0	Yes	10
Nyamarhi (NYA)	3	1	1	0	Yes	10
Ngosto North (NGO)	3	1	1	0	Yes	10
BLANK	2	0	0	0	No	2
	29	6	6	4		88

Table 2: Samples sequences, broken down by number of weekly samples, number of site-times for which S (50 mL) and XS (15 mL) samples were filtered, additional daily samples taken, whether A/B samples were taken, and the resulting total number of samples sequenced per site.

Figure 2: ASV accumulation curve of bacterial ASV richness using the “exact” method. Bars represent two standard deviations around mean estimates.

Figure 3: Phylogenetic tree of 16S ASV sequences in the core samples, paired with their relative abundances at each site. Sites are ordered by section.

Diversity	Level	Shannon	SES	2.5%	97.5%	Pr(sim.)
α	A/B samples	2.93	-9335.9	5.10	5.10	0.01
α	Temporal samples	2.99	-12535.9	5.13	5.13	0.01
α	Sites	3.39	-16873.6	5.15	5.15	0.01
α	Sections	3.94	-14919.9	5.16	5.16	0.01
γ	(Total)	5.17	0.0	5.17	5.17	1.00
β	A/B samples	0.05	153.3	0.027	0.027	0.01
β	Temporal samples	0.40	2688.2	0.022	0.022	0.01
β	Sites	0.55	6528.7	0.008	0.008	0.01
β	Sections	1.23	14919.9	0.008	0.008	0.01

Table 3: Additive partitioning of Shannon diversity into α , β , and γ diversities across sections, sites, temporal samples, and within site-time samples (A/B) as components of the total diversity observed across all core samples. Observed diversity is compared to 99 simulations and the standardized effect size (SES) using “r2dtable” null model with the *adipart* function in the R package *vegan*.

Figure 4: Plots of observed ASV richness and Shannon diversity across samples. Samples are grouped and coloured by park section and with shape indicating waterhole type.

Figure 5: The proportion of ASVs assigned a given taxonomic level using the SILVA database v128 and a pre-trained RDP classifier with minimum 50% bootstrap support.

Figure 6: Relative abundances of bacterial phyla across sites. Sites are ordered by relative abundance of phylum Proteobacteria.

Figure 7: Relative abundances of bacterial phyla across weekly samples.

Figure 8: Relative abundances of bacterial phyla across five days at a single site (NWA).

Figure 9: NMDS plots of a) Bray-Curtis and b) abundance-weighted UniFrac distances. Colours represent site.

Figure 10: Nonmetric multidimensional scaling (NMDS) ordination of variation in bacterial community structure across 54 samples based on a) Bray-Curtis and b) abundance-weighted UniFrac distances. Arrows indicate the direction of significant ($p < 0.05$) correlations among variables and the NMDS axes, with arrow length indicating the strength of the correlation. Blue arrows indicate environmental variables, while black arrows indicate relative abundances of sequences from different microbial classes. The ordination axes explain 96.8% (a) and 98.1% (b) of the variance in the dissimilarities (Fig. S17).