Canadian Journal of Fisheries and Aquatic Sciences

Long-term studies of lobster abundance at a salmon aquaculture site, eastern Canada

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2017-0547.R1
Manuscript Type:	Article
Date Submitted by the Author:	21-Apr-2018
Complete List of Authors:	Grant, Jon; Dalhousie University Simone, Michelle; Southern Cross University Daggett, Tara; Simcorp
Keyword:	AQUACULTURE < General, COASTAL WATERS < Environment/Habitat, SALMON < Organisms, LOBSTERS < Organisms, fish farming
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Long-term studies of lobster abundance at a salmon aquaculture site, eastern Canada

Jon Grant¹, Michelle Simone^{1,2} Tara Daggett³

¹Dept. of Oceanography, Dalhousie University, Halifax, NS B3H 4R2

jon.grant@dal.ca

^{1,2}Centre for Coastal Biogeochemistry, School of Environment, Science and Engineering,
Southern Cross University, Lismore, NSW 2480, Australia

mnhsimone@gmail.com

³Sweeney International Marine Corp., 46 Milltown Blvd., St. Stephen, NB E3L 1G3

tdaggett@simcorp.ca

Keywords: Lobster fisheries; aquaculture; fish farming; salmon; *Homarus americanus*

Abstract

Wild lobster (*Homarus americanus*) abundance was monitored before, during, and after salmon aquaculture production in a bay on Grand Manan Island, New Brunswick, Canada in an 8-year survey, 2008 to 2015. Diver transects and free-area spot-dives were used to measure the carapace length (CL) and determine sex (including berried state) of each lobster encountered both inside (farm) and outside (reference) the lease boundaries. In pairwise comparisons of each sampling date, there was no significant difference between the number of lobsters inside the salmon farming area vs. a nearby reference site, and no significant difference in the number of berried females in or out of the farm lease area. Combining data from all lobster surveys (farm and reference sites) indicated a increase over 8 years, similar in slope to the increase of the trap fishery in Lobster Fishing Area (LFA) 38. These results indicate that the fish farm had no obvious impact on lobster density at any point in the salmon production cycle, and that inshore lobster abundance followed trends similar to those of the general fishery of LFA 38.

47 **Introduction**

48 The sustainability of coastal aquaculture is in part dependent on its fit with other coastal zone
49 activities, including commercial fisheries, recreational use, resource extraction and conservation.
50 Fish farming in net-pens or cages occurs in discrete well-defined areas, but two concerns of
51 farm siting are its boundaries which may exclude other activities, and its ecological footprint
52 including solid and dissolved waste dispersion. In Atlantic Canada net-pen culture of Atlantic
53 salmon is a major economic activity. The lobster (*Homarus americanus*) fishery takes place in
54 the same coastal environments as fish farming and is among the most valuable of commercial
55 species. Designated Lobster Fishing Areas (LFA) in the Bay of Fundy are the most prolific of *H.*
56 *americanus* fisheries (Serdynska and Coffen-Smout 2017). Landings have shown huge increases
57 over the last 10 years (DFO 2015), but conflicts between lobster fishing and fish farming remain
58 for several reasons. For example, the availability of space for deployment of lobster traps is
59 suggested to be compromised by space occupied by fish farms. However, stronger interactions
60 have been suggested regarding the waste stream from the farm. First, organic loading of
61 sediments due to deposition of waste food and feces in some cases produces hypoxic sediments
62 beneath fish pens (Chang et al. 2014). Lobsters occupy a variety of substrates from mud to rock,
63 with a preference for gravel in the early benthic phase (Pottle and Elner 1982). Fish farms are
64 also sited on many bottom types, and locations with dispersive flows are less likely to
65 accumulate waste and alter bottom habitat (Keeley et al. 2013). Second, some pesticide
66 treatments specifically for sea lice, applied either in feed or in bath treatments (tarps surrounding
67 pens or in well boats) may be harmful to crustaceans (Burridge et al. 2010), and there have been
68 lobster mortality events due to pesticide dispersion (Wiber et al. 2012).

69 Grand Manan Island, a large Canadian island in the Gulf of Maine off New Brunswick is home

to multiple salmon farming sites and to a lucrative lobster fishery (Campbell 2011). The island is part of LFA 38 in which the fishery is conducted from about mid-November to the end of June via 136 trap licenses. According to DFO (2015, page 1), ‘Landings in LFAs 35-38 began a long-term increase in the mid-1990s and current landings are at record highs’. Grand Manan is characterized by typically large regional tides (5m) and is also the site of documented conflicts between these industries (Walters 2007).

In the 1980s, salmon aquaculture was active on the northeast coast of Grand Manan in Flagg’s Cove and Dark Harbour. As a result of originally unregulated farming practices, the seafloor experienced benthic degradation (Rosenthal and Rangeley 1989), reducing lobster habitat quality in Flagg’s Cove. Lobster abundance recovered after the site was removed. This failed attempt to sustainably culture salmon was a result of suboptimal practices, which are now monitored and regulated with government involvement through New Brunswick’s aquaculture environmental monitoring program (EMP), via their Department of Environment and Local Government (NBDELG 2018). Regardless of the development of best management practices in aquaculture, when a new salmon aquaculture lease was approved in 2007, 12 km south of Flagg’s Cove on the northeast coast of Cheney Head Island (Duck Island Sound), surveys of lobster abundance became a condition of the lease. The goal of the surveys was to track the occurrence of lobster inside and outside the lease area, before, during, and after production. The following study presents the findings of the mandated surveys to determine how aquaculture practices might affect lobster populations in the immediate vicinity (within lease boundaries) and at the scale of a portion of Duck Island Sound.

In the following study, we undertook diver-based lobster surveys over an 8-year survey in a BACI design experiment (Underwood 1991). We hypothesize (1) trends in lobster abundance at

a salmon aquaculture site, including pre-aquaculture and into production cycles, are identical to trends at a reference site of similar depth and bottom type, and (2) both sites follow a trend similar to the larger regional fishery of LFA 38.

Methods

Lobster surveys over 8 years were conducted via diver transects and free-area spot-dives during the summer/fall months of August and September, 2008 to 2015. The timing of the surveys targeted the first two weeks in each of those months to capture summer lobster populations before their fall migration to deeper waters; however, in 2008 only one survey took place, August 20 – 21. The August 2011 survey was delayed until the third week of the month due to weather. No aquaculture activities took place on the site in preproduction years 2008 and 2009. Within the remaining six years there were two production cycles, 2010-2011 and 2013-2014, where Atlantic salmon (*Salmo salar*) were grown on site and two fallow periods, 2012 and 2015, where all production stopped.

Study Site

All surveys were conducted in and around lease area MF-0503 at Cheney Head (26.51 ha) located on the northeast side of Cheney Island, off Grand Manan Island, New Brunswick (Fig. 1). Water depth in the area ranges from approximately 5 – 15 m, while the center of the Cheney Head site sits in approximately 14 - 16 m of water at low tide. Location in the Bay of Fundy signifies large semi-diurnal tidal currents ($>0.5\text{ m sec}^{-1}$) with a range of 6m. The general direction of flow is North-South, however direction and speed are influenced by rocks and ledges. The

Cheney Head site is typical for a salmon farm off Grand Manan, employing plastic circular cages and standard husbandry methods. The Cheney Head site started off at 100,000 fish, which is small in comparison to other farms in the area of SWNB. For the second production cycle, the number of fish increased to ~336,000. Other farms in the area were stocked at 380,000 to 490,000 fish during that time.

Transect Dives

A total of six diver transects were established off the coast of Cheney Island based on the transect locations from a previous lobster survey conducted in the same area by the Department of Fisheries and Oceans Canada in 2007 (Robichaud 2007), with three transects inside (farm) and three outside the lease boundaries (reference) (Fig. 1A). Location of transects at the farm were constrained by the farm boundaries and avoidance of net-pen infrastructure. Reference transects were along similar depths and substrate as the farm. Geo-referenced coordinates were used to position the transect lines in approximately the same locations each year. However, after the grid system and cages were placed on site in 2010, positions for transects 1, 2 and 3 were altered somewhat to avoid the cage array. For instance, transect 2 was broken into two segments as a result of cage interference with the original transect line. Additional adjustments to transects 1 and 2 were required in 2013 when a larger cage array was installed.

Each transect was 300 m long and was marked every 25 m, yielding 12 sections per line. A weighted sinking rope was used to position and anchor the line on the bottom during diver surveys. Lines were marked with floats at each end and at the 150-m mark to facilitate line retrieval and to aid in diver orientation. Each survey consisted of two divers swimming the length of the transect, and for each of the 12 sections observations within 1 m on either side of

the line were recorded. One diver captured lobsters while the second measured the carapace length (CL) with stainless steel calipers (scale in 1-mm increments) and determined the sex of each lobster, including the egg stage of berried females. All lobsters < 50 mm-CL were considered juvenile due to the difficulty in accurately sexing small lobsters underwater.

Spot-Dives

Two free-area dive searches were also included in each survey period; one just inside the northwest corner of the lease area and the second 1.1 km outside the lease to the southwest (Fig. 1A). The starting coordinates of the spot-dives were the same for each survey; divers then swam in an haphazard manner and searched for lobsters for approximately 30 minutes. The same carapace length and lobster sex data were recorded during the spot-dives as the transect dives.

Benthic Conditions

Bottom sediment type was defined using video footage collected during each of the transect and spot-dives. Videos were recorded with a SONY Handycam camcorder housed in an Amphibico Dive Buddy or a GoPro Hero (2, 3 or 3+) with GoPro dive camera housing case. Battery time constraints limited video collection to approximately 15 minutes of each dive. The provincially mandated environmental monitoring program (EMP) that classifies the level of benthic health using the concentration of dissolved sulfide in undisturbed surface sediments coincided with the survey dates and was used to further characterize the substrate. EMPs were conducted within one week of the September lobster survey each year.

LFA 38 data

Aquaculture lease boundaries and the larger bay fall within Lobster Fishing Area 38 (LFA 38; Fig. 1B) and in Grid Group 6 (GG-6; Fig. 1C) as defined by the Department of Fisheries and Oceans Canada (DFO, 2013). In order to compare lobster abundance trends in the study area to the greater Lobster Fishing Area, aggregated landings from fisher logbooks and the corresponding effort data spanning from the 2007/08 to the 2014/15 season were collated. Data were provided directly by DFO (2016). We found no difference in fishery statistics by analysing GG-6 separately from LFA 38 and consider only the latter which is the more commonly known management unit.

Results

Benthic Conditions

A mixture of bottom types ranging from coarse sand to gravel and cobble, characterized the sedimentary environment of the study area. No changes were apparent over the 8 years. This bottom type is typical of most salmon farms on Grand Manan Island. Among 20 farms on Grand Manan, only one has sediments categorized as mud, the rest being sand, gravel, and shell debris (personal observations). At Cheney Head, there were increases in farm production in the second production cycle (2013-14) compared to the first production cycle (2010-11) and changes to the provincial EMP standard operating procedures in 2013 and 2014. Regardless, the Cheney Head lease site was consistently classified as Oxid-A (<750 μM of dissolved sediment sulfide, which indicates normal/oxid sediment) from 2010-2015 (Table 1). Prior to production, EMP monitoring was not required, but it is a condition of the government-issued Approval to Operate certificate. When an aquaculture site is fallow (i.e. not under production), this certificate is not required.

184 Survey data

185 Lobsters were numerous in all the survey areas and a total of 1255 individuals were
186 sampled inside the lease (farm area) and 1171 outside the lease (reference area) during the study
187 period. The aggregate population was dominated by individuals of commercial size ($>82.5\text{mm}$
188 carapace length CL). Over 50% of the lobsters observed during the eight-year survey were
189 within the 61-100mm CL size range (Fig. 2). This mean size range was smaller than that found
190 off Flagg's Cove on Grand Manan Island by Tremblay et al. (2006) in 2001-2002 where most
191 lobsters fell in the 100-130 mm CL range.

192 Sex ratio (male:female) was usually >1 with a maximum ~ 1.6 and no significant difference in
193 numbers of either sex in comparisons in the farm area vs. the reference (paired Wilcoxon ranked
194 sign test, $p = 0.097$ males; $p = 0.416$ females, $n = 15$ pairs; data were non-normal; $\alpha = 0.05$ here
195 and following). The % of berried females ranged from 0-38% over all the sampling, with about
196 half of the values $\leq 15\%$. There was no significant difference in the number of berried females in
197 the farm vs. reference areas (paired Wilcoxon ranked sign test, $p = 0.071$, $n = 15$ pairs). However,
198 on 7 of 15 comparison dates, there were 3 or fewer berried lobsters at both reference and farm
199 sites (Fig. 3). Numbers of berried females were correlated in the farm and reference areas
200 (Pearson $r = 0.797$, $p < 0.001$, $n = 15$).

201 The mean lobster density calculated from all transect dives during the study ($600\text{ m}^2/\text{transect}$)
202 was 0.035 ind m^{-2} inside the lease (farm) and 0.028 ind m^{-2} outside the lease (reference). Lower
203 densities were found at farm and reference areas by Robichaud (2007) in the year prior to our
204 starting date, 0.016 and 0.01 ind m^{-2} respectively. Robichaud's (2007) data did however fall
205 within the min and max densities observed during the time of the current study (farm range =
206 $0.009 - 0.079\text{ ind m}^{-2}$; reference range = $0.08 - 0.062$).

The total number of lobsters (transects and spot dives) were summed for each sampling date, separately at farm and reference areas. There was no significant difference between the number of lobsters in comparisons of farm vs. reference totals, paired at each sampling date (paired t-test, $p = 0.79$; D'Agostino & Pearson normality test, $p = 0.371$ and 0.227 respectively, $n = 15$).

Mean numbers of lobsters shown for each sampling date indicate that both farm and reference locations had increasing trends in lobster density during the study period (Fig. 4). In the reference area, this trend was significant through time (Fig. 4). Within the farm, the trend was not significant due to the generally greater spatial variation in lobster density found there (Fig. 4). Part of the variance is due to an increase in numbers consistently occurring from August to September in each of the sampling years. We emphasize that these trends were independent of production phases of the farm, and that lowest lobster density occurred before the farm was established. It is apparent from Fig. 4 that the trajectories of increase in lobster abundance converged on similar density at the farm and reference sites. The ranges of density overlapped in every comparison between farm and reference site, reinforcing the results of the paired t-test for total abundance which indicate no significant location difference. In summary, these data provide no statistical evidence of an effect of the fish farm location on lobster density. However, it is important to interpret these results in the context of the regional lobster fishery in LFA 38.

Comparison to LFA Data

The landings from LFA 38 were standardized by effort (trap hauls) to derive catch per unit effort (CPUE) and expressed as % increase from the starting value in 2008. In order to compare the Cheney Head survey data to the regional abundance in LFA38, the farm and reference data were summed on each sampling date to produce a single estimate of total lobster density for the

entire sampling area. Lack of significant differences between these samples justify their summation. The August and September survey dates were then averaged per year (see August-September variation in Fig. 4) and compared to the DFO lobster fishery season that had just ended. In LFA38 the fishery is conducted from the 2nd Tuesday in November to June 29th. Because each data set produced a single annual value, we made the date consistent between them.

LFA38 displayed significant increases in lobster abundance (estimated as CPUE) over the 8 seasons (Fig. 5). Cheney Head (farm+reference) also showed a significant increase in abundance (Fig. 5). An F-test of the slopes indicated no significant difference between them ($F = 3.608$; $p=0.0818$). However, the intercepts were significantly different ($F = 23.33$; $P=0.0003$) indicating that although similar in slope, the regression lines were displaced in elevation. Variation in the regressions is obvious, each with a single substantial outlier beyond 95% CI (Fig. 5). Differences in intercept between the lines is not surprising since the data are from very different sources. Beyond their location difference, CPUE data are a cumulative total over the fishing season, while the Cheney data are from averaged August and September samples outside of the fishing season. Despite the difference in data sources, LFA 38 and Cheney show a similar temporal trend, suggesting that over this multi-year time series inshore lobsters reflect the abundance of the regional (LFA 38) population.

Discussion

Our results show that the salmon farming cycle at Cheney Head, including fallow periods, had no obvious influence on lobster abundance. It is important however, to examine the applicability of this finding to other fish farming sites, the modes by which lobster populations could be affected by salmon farming, and the extent to which they apply to the present study.

The farm site and reference area transects are ideal gravel-cobble habitats for lobster (Pottle and Elmer 1982). Density values as much as 0.1 ind m⁻² are typical for Grand Manan shallow waters (Tremblay et al. 2006), but higher densities (1 ind m⁻²) have been observed in Maine (Steneck and Wilson 2001). Coarse sediments are often non-depositional, and it is obvious that there is not local accumulation of farm waste in the sediment, since regulatory monitoring indicates a consistent classification of Oxidic A, the least impacted category (NBDELG 2018). Using available provincial EMP data for the 20 salmon farms on Grand Manan, 11 had sulfides exceeding 750 µm (limit of Oxidic A classification) in EMP years 2012-14, but none in more than one of those years. Collectively, these are largely non-depositional sites at similar depths with few near-field eutrophication impacts, and our results should be widely applicable to Grand Manan. Although a similar lobster study would be of interest in depositional soft sediments, such conditions are not typical lobster habitat and substrate suitability for lobsters might confound fish farm effects in interpretation of results. The Cheney Head farm is an example of a well-managed, well-sited fish farm and as such stands as an example that fish farms can have no apparent negative impact on local lobster abundance.

The topic of far-field effects often arises in consideration of organic deposition dispersing away from fish farms. The literature has limited documentation of far-field impacts (Grant 2010), and typically involves small inlets with closely spaced fish farms (Wildish 2005), a situation that is avoided in modern husbandry. In the highly dispersive macrotidal environment of the Cheney Head salmon farm, it is unlikely that there are regional (farm + reference) impacts of organic deposition in the benthos, confirmed by the EMP data which includes reference sites. Trace metals from either food waste or antifouling treatments (e.g. Zn, Cu) have also been associated with sediments near salmon farms (Chou et al. 2002), but in non-depositional environments we

277 expect lack of accumulation as with organic matter.

278 The risk to lobster populations from pesticide exposure applied to treat sea lice is difficult to
279 assess under field conditions. Our limited data for the Cheney Head farm indicate that bath (well
280 boat and tarp) and in-feed treatments were used for sea lice. Exposure to water-borne
281 therapeutants studied via brief exposure in laboratory experiments indicates variable toxicity
282 (Burridge et al. 2014), depending on the which product is applied. Recent studies by Ernst et al.
283 (2014) in the Bay of Fundy examined the scope for pesticide dispersion in a series of dye
284 experiments simulating tarp treatments. Their results suggest that potentially deleterious
285 pesticide concentrations could occur up to several hundred meters from a treated cage, but this is
286 dependent on the specific pesticide applied, exposure time, non-target species- and life-stage-
287 specific toxicity, and hydrodynamic regime. Although in-feed treatments might be expected to
288 provide a direct route for pesticide exposure since lobsters could ingest waste feed beneath
289 cages, laboratory studies suggest that the dose in feed pellets is not necessarily harmful to adult
290 lobsters (Burridge et al. 2004). In addition, salmon food pellets are not a preferred food source
291 for *H. americanus* (Waddy et al. 2007). In-feed treatments may persist in sediments for long
292 periods (Benskin et al. 2016), but we would not expect this mode of exposure at a non-
293 depositional site like Cheney Head.

294 Pesticides used for sea lice have also been shown to impact lobster spawning (Burridge et al.
295 2008). Our data suggest that gravid females on the farm follow a similar temporal pattern
296 compared to the reference site but this trend was not obviously related to the farm
297 production/fallow cycle, and the highest number of gravid females occurred at the end of the
298 time series. Correlation between the dynamics of farm and reference sites indicates that
299 application of sea lice treatments on the farm is not decoupling these patterns. In describing the

preliminary results of a field study, Maillet et al. (2017) suggest that gravid females avoid farms undergoing pesticide treatment. Laboratory experiments on female lobster behavior relative to pesticides either in sediments or in water-borne application would be fruitful in addressing this question (see also Abgrall et al. (2000)).

Our second hypothesis was that lobster abundance at the fish farm/reference site followed the temporal pattern shown in the wider population. This is demonstrated in the similarity of slope in trends at the combined farm/reference site and LFA 38. If the fish farm affected lobster abundance, either positively or negatively, the trend demonstrated in the LFA38 fishery would not be observed at the 1-10km scale of our sampling (farm and reference). Due to seasonal migrations (see below), inshore-offshore lobsters may be viewed as effectively the same populations. In an investigation of indices used for assessing lobster populations, research trawl data demonstrated the same large increase in abundance as fishery statistics (DFO 2015). Our diver surveys indicate similar agreement with fishery data providing confidence that the temporal trends documented for the fish farm and reference area accurately capture dynamics of local abundance.

There are clearly ecosystem-level factors affecting increased lobster abundance in eastern Canada, and our results suggest that they are reflected in the nearshore as well. The loss of groundfish due to overfishing in continental shelf ecosystems in parts of eastern Canada is a major alteration to these systems, including a shift from a demersal to a pelagic food chain (Bundy 2011). Among other observed effects, large increases in lobster abundance in recent decades have not been easy to relate to apparent regime shifts occurring from fishing down the food web. It has been suggested that increased lobster abundance is due to release from predation pressure by the decline in demersal fish (Boudreau and Worm 2010). Wahle et al. (2013) suggest

323 that overfishing of larger predatory fish in the Gulf of Maine has given lobsters a size refuge
324 from predation, contributes to their increasing numbers. If there is any local impact of fish
325 farming on lobster populations at Cheney Island, it is obscured by the larger trend of increasing
326 abundance.

327 Lobsters are known to be attracted to food resources and shelter (Karnovsky 1989), these
328 attractants being the basis of the trap fishery. Lobster fishers on Grand Manan have in the past
329 trapped close to salmon cages to take advantage of potentially higher abundance (Walters 2007).
330 However, our data do not indicate increased abundance on the farm site compared to reference
331 areas. Moreover, lobster abundance at the farm site continues to increase even through fallow
332 periods.

333 It has been suggested that lobsters seek warmer water to enhance temperature-dependent
334 processes such as molting and egg development. Campbell (1986) conducted mark-recapture
335 studies of lobsters off Grand Manan and confirmed that they undertake two seasonal migrations,
336 entering shallow waters in summer and retreating to deeper colder waters in fall. This might lead
337 to higher densities in August compared to September if fall migrations to deeper water occurred
338 during the study period. However, we usually saw higher densities in September than in August,
339 and water temperatures are similar between these months (Campbell 1986), confirmed in our
340 data collected during this study (data not shown).

341 There are few studies which specifically consider the interaction between lobster fisheries and
342 fish farming. Loucks et al (2015) suggested that a steelhead farm in coastal Nova Scotia caused
343 baywide reduction in lobster catches. The methods and results of this research were subsequently
344 called into question (Grant et al. 2016). Trapping locations by fishers were estimated in their
345 study rather than documented by GPS and the designated spatial sampling regions for lobster

overlapped in their distances from the farm. Moreover, sampling involving only the final two weeks on the fishing season was potentially influenced by the prior months of catches (see also Loucks et al. 2016). We consider our study to be a comprehensive work on lobster-fish farm interactions because it contained a multi-year time series, a repeated spatial sampling plan, includes all aspects of the salmon production cycle, and comparison to wild fishery dynamics. The timing of our annual sampling timing pre-dates the start of the annual lobster fishing season in LFA38, so we anticipate that our survey results were not influenced by trapping pressures.

It is likely that potential conflicts between lobster fishing and fish farming will persist. Conflict resolution in coastal resource use is managed via marine spatial planning (MSP), an approach previously applied to aquaculture (Filgueira et al. 2014). Although scientific studies are essential in this process, consultation among stakeholders is key to decision making. Mapping of fishing areas and lobster habitat is important in this planning as demonstrated for lobsters by Serdynska and Coffen-Smout (2017). Initial efforts at MSP were promoted for Grand Manan in a study by Lane et al. (2008). An ecosystem approach to aquaculture requires an assessment of zones of impact surrounding coastal activities and an understanding far-field effects. Continued dialog in the framework of a structured planning process among coastal resource industries, coastal residents, and regulators is key to avoiding conflict and pursuit of sustainability in shared use of the ocean.

References

- Abgrall, P., Rangeley, R.W., Burrige, L.E. and Lawton, P. 2000. Sublethal effects of azamethiphos on shelter use by juvenile lobsters (*Homarus americanus*). *Aquaculture* **181**: 1–10.
- Benskin, J. P., Ikonomou, M. G., Surridge, B. D., Dubetz, C., and Klaassen, E. 2016. Biodegradation potential of aquaculture chemotherapeutants in marine sediments. *Aquacult. Res.* **47**: 482–497.
- Boudreau, S. A., and Worm, B. 2010. Top-down control of lobster in the Gulf of Maine: insights from local ecological knowledge and research surveys. *Mar. Ecol. Prog. Ser.* **403**: 181–191.
- Bundy, A. 2011. Structure and functioning of the eastern Scotian Shelf ecosystem before and after the collapse of groundfish stocks in the early 1990s. *Can. J. Fish. Aquat. Sci.* **62**: 1453–1473.
- Burrige, L. E., Hamilton, N., Waddy, S. L., Haya, K., Mercer, S. M., Greenhalgh, R., Tauber, R., Radecki, S. V., Crouch, L.S., Wislocki, P. G., and Endris R.G. 2004. Acute toxicity of emamectin benzoate (SLICETM) in fish feed to American lobster, *Homarus americanus*. *Aquacult. Res.* **35**: 713–722.
- Burrige L.E., Haya K., Waddy S.L. 2008. The effect of repeated exposure to azamethiphos on survival and spawning in the American lobster (*Homarus americanus*). *Ecotoxicol. Environ. Safety* **69**: 411-415.

- 393 Burridge, L.E., Weis, J. S., Cabello, F., Pizarro, J., and Bostick, K. 2010. Chemical use in salmon
394 aquaculture: A review of current practices and possible environmental effects. *Aquaculture*
395 **306**: 7–23.
- 396 Burridge, L. E., Lyons, M. C., Wong, D. K. H., MacKeigan, K., and Van Geest, J. L. 2014. The
397 acute lethality of three anti-sea lice formulations: AlphaMax®, Salmosan®, and
398 Interlox®Paramove™50 to lobster and shrimp. *Aquaculture* **420-421**: 180–186.
- 399 Campbell, A. 1986. Migratory movements of ovigerous lobsters, *Homarus americanus*, tagged
400 off Grand Manan, Eastern Canada. *Can. J. Fish. Aquat. Sci.* **43**: 2197–2205.
- 401 Campbell, A. 2011. Characteristics of the American lobster fishery of Grand Manan, New
402 Brunswick, Canada. *North American Journal of Fisheries Management*, **12**: 139–150.
- 403 Chang, B. D., Page, F. H., Losier, R. J., and McCurdy, E. P. 2014. Organic enrichment at salmon
404 farms in the Bay of Fundy, Canada: DEPOMOD predictions versus observed sediment sulfide
405 concentrations. *Aquacult. Environ. Interact.* **5**: 185–208.
- 406 Chou, C.L., Haya, K., Paon, L.A., Burridge, L., and Moffatt, J.D. 2002. Aquaculture-related trace
407 metals in sediments and lobsters and relevance to environmental monitoring program ratings
408 for near-field effects. *Mar. Poll. Bull.* **44**: 1259–1268.
- 409 DFO. 2013. Assessment of Lobster (*Homarus americanus*) in lobster fishing areas (LFA) 35-38.
410 Maritimes Region: Canadian Science Advisory Report.
- 411 DFO. 2015. 2015 Stock Status Update of Lobster (*Homarus americanus*) in the Bay of Fundy.
412 (Lobster Fishing Areas 35-38). DFO Can. Sci. Advis. Sec. Sci. Resp. 2015/030.
- 413 DFO. 2016. LFA38 lobster fishery data from 2007-2015, Adam Cook, Bedford Institute of
414 Oceanography, Dartmouth Nova Scotia, Canada, 2016-04-21.
- 415 Ernst, W., Doe, K., Cook, A., Burridge, L., Lalonde, B., Jackman, P., Aube, J.G., and Page F.

- 416 2014. Dispersion and toxicity to non-target crustaceans of azamethiphos and deltamethrin after
417 sea lice treatments on farmed salmon, *Salmo salar*. *Aquaculture* **424-425**: 104–112.
- 418 Filgueira, R., Grant, J., and Strand, Ø. 2014. Implementation of marine spatial planning in
419 shellfish aquaculture management: modelling studies in a Norwegian fjord. *Ecol. Applic.* **24**:
420 832-843.
- 421 Grant, J. 2010. Coastal communities, participatory research, and far-field effects of aquaculture.
422 *Aquacult. Environ. Interact.* **1**: 85-93.
- 423 Grant, J., Barrell, J. and Filgueira, R. 2016. Lack of interaction between finfish aquaculture and
424 lobster catch in coastal Nova Scotia. *Mar. Poll. Bull.* **110**: 613-615.
- 425 Keeley, N.B. 2013. Novel observations of benthic enrichment in contrasting flow regimes with
426 implications for marine farm monitoring and management. *Mar. Poll. Bull.* **66**: 105-116.
- 427 Karnofsky, E.B., Atema, J., and Elgin, R.H. 1989. Field observations of social behavior, shelter
428 use, and foraging in the lobster, *Homarus americanus*. *Biol. Bull.* **176**: 239–246.
- 429 Lane, D., Michalowski, W., Stephenson, R., and Page, F. 2008. Integrated Systems Analysis for
430 Marine Site Evaluations and Multicriteria Decision Support for Coastal Aquaculture. In
431 *Aquaculture, Innovation and Social Transformation* (Vol. 17, pp. 255–264). Dordrecht:
432 Springer Netherlands.
- 433 Loucks, R. H., Smith, R. E., and Fisher, E. B. 2014. Interactions between finfish aquaculture and
434 lobster catches in a sheltered bay. *Mar. Poll. Bull.* **88**: 255-259.
- 435 Loucks, R. H., Smith, R. E., and Fisher, E. B. 2016. A response to the letter to the editor ‘Lack of
436 interaction between finfish aquaculture and lobster catches in coastal Nova Scotia.’ *Marine*
437 *Pollution Bulletin* **110**: 616–618.
- 438 Maillet, D. G. C., Wiber, M. G., and Barnett, A. 2017. Actions towards the joint production of

- 439 knowledge: the risk of salmon aquaculture on American Lobster. *Journal of Risk Research* **55**:
440 1–14.
- 441 NBDELG. 2018.
442 [http://www2.gnb.ca/content/gnb/en/departments/elg/environment/content/water/content/marin](http://www2.gnb.ca/content/gnb/en/departments/elg/environment/content/water/content/marine_aquaculture.html)
443 [e_aquaculture.html](http://www2.gnb.ca/content/gnb/en/departments/elg/environment/content/water/content/marine_aquaculture.html)
- 444 Pottle, R. A., and R. W. Elner. 1982. Substrate preference behavior of juvenile American lobsters.
445 *Homarus americanus*, in gravel and silt clay sediments. *Can. J. Fish. Aquatic. Sci.* **39**: 928-
446 932.
- 447 Robichaud, D., 2007. Preliminary summary of 2007 lobster dive survey off Cheney Island,
448 Grand Manan. DFO Science, St. Andrews Biological Station, 6 pp.
- 449 Rosenthal, H., and R.W. Rangeley. 1989. The effect of a salmon cage culture on the benthic
450 community in a largely enclosed bay (Dark Harbour, Grand Manan Island, N.B., Canada). *Int.*
451 *Counc. Explor. Sea C.M.* 1989/F-23 Mariculture Committee.
- 452 Serdynska, A. and S. Coffen-Smout. 2017. Mapping inshore lobster landings and fishing effort
453 on a Maritimes region statistical Grid (2012–2014). *Can. Tech. Rep. Fish. Aquat. Sci.* 3177.
- 454 Wilson, C. J., Steneck, R. S., and Wilson, C. J. 2002. Large-scale and long-term, spatial and
455 temporal patterns in demography and landings of the American lobster, *Homarus americanus*,
456 in Maine. *Mar. Fresh. Res.* **52**: 1303–1319.
- 457 Tremblay, M. J., Smith, S. J., Robichaud, D. A., and Lawton, P. 2006. The catchability of large
458 American lobsters (*Homarus americanus*) from diving and trapping studies off Grand Manan
459 Island, Canadian Maritimes. *Can. J. Fish. Aquat.* **63**: 1925–1933.
- 460 Underwood, A.J. 1991. Beyond BACI: experimental designs for detecting human environmental
461 impacts on temporal variations in natural populations. *Austral. J. Mar. Fresh. Res.* **42**: 569 –

462 587.

463 Waddy, S.L., Merritt, V.A, Hamilton-Gibson, M.N., Aiken, D.E., and Burrige, L.E.. 2007.

464 Relationship between dose of emamectin benzoate and molting response of ovigerous

465 American lobsters (*Homarus americanus*). Ecotoxicol. Environ. Safety **67**: 95-99

466 Wahle, R. A., Brown, C., & Hovel, K. (2013). The geography and body-size dependence of top-

467 down forcing in New England's lobster-groundfish Interaction. Bull. Mar. Sci. **89**: 189–212.

468 Walters, B. B. 2007. Competing use of marine space in a modernizing fishery: salmon farming

469 meets lobster fishing on the Bay of Fundy. Can. Geogr. **51**: 139–159.

470 Wiber, M. G., Young, S., and Wilson, L. 2012. Impact of aquaculture on commercial fisheries:

471 fishermen's local ecological knowledge. Human Ecol. **40**: 29-40.

472 Wildish, D.J. 2005. Benthic macrofaunal changes resulting from finfish mariculture. Benthic

473 macrofaunal changes resulting from finfish mariculture. Handbook Env. Chem. Vol. 5, Part M:

474 275–304, Springer-Verlag, Berlin.

475

476 **Acknowledgements**

477 This study was funded by the NSERC-Cooke Industrial Research Chair in Sustainable

478 Aquaculture awarded to Jon Grant and the Atlantic Canada Fish Farmers Association. We thank

479 Amanda Martin and Bob Sweeney of SIMCorp and Benson Aquaculture Ltd.

480

481

482

Figure 1. A. Cheney Head bathymetry, salmon farm lease boundaries, sampling transects and free dive locations. Farm transects are inside the lease polygon, reference transects are outside. White areas are >10m depth. Map was created with Garmin MapSource software using BlueChart Americas v9.5. B. Lobster Fishing Areas (LFAs) 35-38. C. Grid Groups (GGs) from 10 minute grids in LFA 35-38 logbooks (bottom-right). The red circle in GG-6 and LFA 38 is the location of the Cheney Head lease area on Grand Manan Island. B and C from DFO (2013).

MF-0503 Cheney Head	Sulphide (µM)	Classification
2015	4	Oxic A
2014	160	Oxic A
2013	144	Oxic A
2012	0	Oxic A
2011	705	Oxic A
2010	349	Oxic A

Table 1. New Brunswick environmental monitoring program results for the Cheney Head salmon farm. Free dissolved sulfide determined by ion-specific electrode. Source: produced from data on NBDELG 2018.

Draft

Figure 2. Numbers of lobsters from all sites in 10mm carapace length bins. Vertical lines represent the upper and lower quartiles.

Figure. 3. Numbers of berried female lobsters during the study period at the fish farm site vs. the reference site.

Figure 4. Mean number of lobsters observed per dive at the farm site ($n=4$; 3 transects + 1 spot dive) and reference site ($n=4$; 3 transects + 1 spot dive) during each survey from August 2008 (Day 0) to September 2015 (Day 2571). Every year except 2008 has both an August and September sampling date. Error-bars represent standard deviation of each mean. Linear regression farm: $y = 0.0065x + 12.028$, $R^2=0.252$, $p=0.0567$; reference, $y = 0.0077x + 9.0074$, $R^2=0.572$, $p = 0.001$.

Figure 5. Seasonal catch per unit effort data (lbs/haul) for LFA38 (Fishery) and annual August-September mean of total number of lobsters observed in combined farm and reference surveys (Cheney) from 2007/08 to 2014/15. Both data sets have been normalized as % increases from the initial value. Cheney regression is $y = 0.083x + 17.27$, $R^2 = 0.81$, $p = 0.002$; and Fishery: $y = 0.043x - 18.31$, $R^2 = 0.878$, $p < 0.001$.

Draft