Canadian Journal of Fisheries and Aquatic Sciences

Predicting species distribution from fishers' local ecological knowledge: an alternative for data-poor management

Journal:	<i>Canadian Journal of Fisheries and Aquatic Sciences</i>
Manuscript ID	cjfas-2018-0148.R1
Manuscript Type:	Article
Date Submitted by the Author:	06-Sep-2018
Complete List of Authors:	Lopes, Priscila; Universidade Federal do Rio Grande do Norte, Ecology Verba, Júlia; Universidade Federal do Rio Grande do Norte, Program in Ecology Begossi, Alpina; Universidade Estadual de Campinas, CAPESCA/NEPA Pennino, Maria Grazia; Instituto Espanol de Oceanografia
Keyword:	data-poor fisheries, Brazil, ethnobiology, species distribution model, <i>Epinephelus marginatus</i>
Is the invited manuscript for consideration in a Special Issue? :	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Running head: LEK to predict fish distribution

**Predicting species distribution from fishers’ local ecological knowledge: a new
alternative for data-poor management**

Priscila F. M. Lopes^{1,2,3,*}, Júlia T. Verba^{1,3}, Alpina Begossi^{3,4} & Maria Grazia Pennino^{1,5}

¹Fisheries Ecology, Management and Economics Unit – FEME, Natal, Brazil; Department
of Ecology, Federal University of Rio Grande do Norte, Natal, Brazil, 59078-900; +55 84
3342-2334, Route 423; priscila@cb.ufrn.br

²Fisheries and Food Institute, UNISANTA, Santos, Brazil;

³Graduate Program in Ecology, Federal University of Rio Grande do Norte, Natal, Brazil.
julitovarv@gmail.com

⁴Capesca/Nepa/Unicamp, Rua Albert Einstein s/n, Campinas, 13083-852, SP, Brazil and
Ecomar/Unisanta (Rua Oswaldo Cruz, 277, Santos, 11045-907, SP, Brazil).
albegossi@gmail.com

⁵Instituto Español de Oceanografía. Centro Oceanográfico de Murcia, Murcia, Spain;
graziapennino@yahoo.it

*Corresponding author

Abstract

Many developing countries lack information to manage their endangered species, urging the need for affordable and reliable information. We used Bayesian hierarchical spatial models, with oceanographic variables, to predict the distribution range of *Epinephelus marginatus*, the dusky grouper, for the entire Southwest Atlantic. We ran a model using scientific information gathered from the literature, and another using information gathered from fishers on species presence or absence. In both models, temperature was an important determinant of species occurrence. The predicted occurrence of the dusky grouper overlapped widely (Schoener's $D = 0.71$; Warren's $I = 0.91$) between the models, despite small differences on the southern and northern extremes of the distribution. These results suggest that basic information provided by fishers on species occurrence in their area can be reliable enough to predict species occurrence over large scales and can be potentially useful for marine spatial planning. Fishers' knowledge may be an even more viable alternative to data collection than what was previously thought, for countries that both struggle with financial limitations and have urgent conservation needs.

Keywords: Brazil; data-poor fisheries; dusky grouper; ethnobiology; species distribution model; *Epinephelus marginatus*

Introduction

One of the main obstacles that developing countries face to manage fisheries or assign a conservation status to target species is a lack of basic and reliable information on species, habitats and ecosystems (Kaufman et al. 2004; Mora et al. 2009; Houk et al. 2012; Worm and Branch 2012; Pauly and Zeller 2016). Without this information, species management is just a guessing game subject to failure (Hilborn 2007; Cinti et al. 2010; Finkbeiner and Basurto 2015). Lack of reliable information also opens the possibility for criticism, with resource users, for instance, opposing measures because they are not convinced that they are necessary or legitimate (Jentoft 2000; Lopes et al. 2013b; Di Dario et al. 2015). Precautionary approaches, which are usually advisable in such cases (Johannes 1998; Kaufman et al. 2004), are rarely implemented because of the high immediate political, social and economic costs associated with management (Marshall and Marshall 2007; Gaines et al. 2010). While the “burden of proof” remains unjustly on science rather than on users (Dayton 1998), managers and researchers need basic supporting data to justify the need for immediate management (Houk et al. 2012; Roos et al. 2015, 2016).

However, acquiring species, ecosystem or fisheries data in developing countries is proportionally costly, not only because funding for science and research is limited (science usually ranks low in government priorities) (Chao et al. 2015; Pinheiro et al. 2015), but also because such countries are burdened with bureaucracy, fear of biopiracy, lack of specific tax incentives and access to technologies, and neglect for resource use statistics (Barber et al. 2014; Pauly and Zeller 2016). In such a context, developing country researchers face the constant need to seek innovative affordable solutions to data acquisition and effective management implementation (Berkes 2003).

64 In fact, globally, two of the most successful management systems seem to be well-
65 established practices in the developing tropics: co-management and customary management
66 (Berkes 2003; Mora et al. 2009; Lopes et al. 2011; Cinner and Huchery 2014). These
67 solutions include users in both the decision-making and surveillance processes (Castello et
68 al. 2009) and rely on very little data (Castello, 2004).

69 Researchers have proposed that such knowledge and data gaps be filled with
70 information provided by fishers, based on their knowledge and experience, which is known
71 in the literature as fishers' local ecological knowledge - FEK or LEK (Johannes 1981;
72 Moller et al. 2004; Silvano and Valbo-Jorgensen 2008; Silvano and Begossi 2010). Since
73 the beginning, these studies have shown that fishers retain an amazing amount of accurate
74 information on fish biology and ecology, such as diet, reproduction, habitat preference, and
75 schooling habit of their target species (Neis et al. 1999; Silvano and Begossi 2010). More
76 recently, studies using LEK have demonstrated the potential to tap into fishers' knowledge
77 to reconstruct information on fish stocks and to detect stock declines unknown to science,
78 given the lack of past scientific data. With regards to the latter, older fishers provide
79 evidence for stock decline by reporting on their past catches (Tesfamichael et al. 2014;
80 Bender et al. 2014; Damasio et al. 2015).

81 Clearly, fishers' LEK is not taken for granted, as it indeed should not be (Davis and
82 Wagner 2003). Fishers' information, similar to any other information provided by humans,
83 can be distorted by their cultural and socioeconomic background and by their perception
84 and interpretation of reality (Jones et al. 2008; Ruddle and Davis 2011). For instance,
85 fishers may attribute a strong predatory role to a given species if they think it competes
86 with them for valuable prey (Davis et al. 2004). They can also overemphasize the role of
87 certain food items on fish diets, if the food item is used as bait (Begossi et al. 2016).

Moreover, fishers may recollect their most positive memories (largest catch or largest fish) much better than their average day-to-day fishing success (Damasio et al. 2015). The role of research is to identify the information that is most applicable to management (Silvano and Valbo-Jorgensen 2008).

Taking this a step further, recent studies have started to test the usefulness of LEK data in ecosystems modeling (Anadón et al. 2010; Eddy et al. 2010). Although less detailed than traditional scientific modeling, LEK-based models can provide some of the required information for proper ecosystem management (Bevilacqua et al. 2016). It is known that fishers can identify areas in their own region where target species are, however, it remains to be tested whether fishers' LEK can be used to predict species occurrence (niche distribution). (Begossi et al. 2013). Although basic, the prediction of species occurrence may represent non-existent information for certain areas and for certain species (Roos et al. 2015). Knowing species–environment relationships and their suitable habitats could contribute to the implementation of marine spatial planning and, if necessary, the protection of relevant habitats for the conservation of some species (Agardy et al. 2011; Pennino et al. 2016). Habitat and species mapping are especially relevant for conservation programs for threatened or endangered species because they provide clear pictures of their distribution, and thus facilitate management of the marine environment (Roos et al. 2015).

Within this context, we modeled the distribution of *Epinephelus maginatus* (Lowe 1834), the dusky grouper, along the Southern Atlantic coast (South American coast). This species is economically relevant, has been widely exploited in Europe, Africa and South America, has been extensively studied in the Mediterranean (e.g., Bouchereau et al. 1999; Gilles et al. 2000; Andrello et al. 2013), and is considered endangered by the IUCN Red List. In the South Atlantic the species is much less known (Machado et al. 2003, 2008;

Andrade et al. 2003; Irigoyen et al. 2005). However, some studies have already suggested that in the Southern Atlantic *E. marginatus* has a smaller effective population size compared to its Mediterranean counterpart (Priolli et al. 2016), and that fishers have noticed significant declines in their catches over time (Lopes et al. 2013a). We also compared the distribution prediction generated from traditional scientific data to the one generated from information provided by fishers. Our intention was to provide alternative basic, reliable and cheap information for spatial planning and management of fish species that can be applied worldwide, particularly in areas where there is no other source of information available.

2. Material and methods

Scientific data

The scientific data used here to establish where *E. marginatus* was present or absent along the south Atlantic coast came from three different sources: 1) our own data collected through multiple projects, 2) bibliographic sources, and 3) online databases that provide data on species presence and absence (all sources are in Supplement S1). With respect to our own data, we used information present in our databases for the period 1988-2016, combined with geographical location of fishing grounds, which are published elsewhere (Begossi et al. 2013). With the exception of the period 2013-2016, when our research was specifically designed to sample and study *E. marginatus*, the remaining studies concerned small-scale fisheries in general, and we simply registered the presence of the dusky grouper whenever available in the landings (sometimes in numbers, sometimes in kilograms, sometimes just whether or not it was present). This means that the occurrence of *E. marginatus* may have been registered in a given site (fishing grounds) multiple times, as

our data is fishing data and not scientific sampling data. As most of our research to date has covered a specific part of the extensive Brazilian coast, we also carried out an exhaustive bibliographic review of previous research to account for existing information about geo-referenced occurrences or absences of *E. marginatus* in all South American waters (Supplement S1). The studies we identified were mostly scientific samplings that registered the presence of the species in a given site, using methods such as transects; where the researcher samples multiples areas near one another. For those cases, we randomly took one of the sampled locations and assigned the species presence (or absence) to it. To make sure we included all available information, we extracted presence data from online databases (Kaschner et al. 2013) and excluded duplicate observations. We then combined the three data sources pertaining to the presence or absence of *E. marginatus* into a single presence-absence dataset to perform a Bayesian spatial model with a binomial distribution.

Overall, we registered 826 suitable observations to run in the model (Figure 1a): 578 presences (356 from our own data, six presences from the AQUAMAPS database, and 216 presences from the bibliographic sources) and 26 absences (all from the bibliographic sources). Of the presence data, 43 were unique presences (registered just once).

Ethnobiological data

Following the same procedure used for the scientific datasets, an exhaustive literature review of previous studies was performed to collect occurrence information for all South American waters that were provided by fishers (Supplement S1). This information was usually obtained through interviews when fishers were asked if *E. marginatus* was present in their region. As these studies relied on interviews, no geo-referenced information was available. To account for an approximate location, we estimated the presence or

absence of the species in a coastal point close to the village where the interview took place (average of 120 meters away from the shore).

We recorded a total of 60 observations, of which 37 referred to presences and 23 to absences of *E. marginatus* (Figure 1b).

Environmental data

Overall, six environmental variables were considered to be potential predictors of *E. marginatus* distribution, both for the scientific and for the ethnobiological data: sea surface temperature (SST in °C), sea surface salinity (SSS in PSU), net primary productivity (NPP in mg C m⁻² d⁻¹), depth (in meters), slope (% grade) and concavity of the seabed. The latter variable is not a physical unit, but the second derivative of the bathymetry layer (or the slope of the slope) and represents whether a raster cell is on a hill (negative values) or in a valley (positive values). These variables were chosen as they had previously been associated with the presence of either the same species or a related species (Harmelin and Harmelin-Vivien 1999; Andreello et al. 2013).

SST and SSS were both extracted with a spatial resolution of 1° x 1° (~ 100 km) from NODS_WOA09 as long-term monthly climatological means provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their web site at <http://www.esrl.noaa.gov/psd/>.

Net primary productivity (NPP) was retrieved from a 920 x 1680 global grid of NPP and calculated as a function of chlorophyll, available light, and photosynthetic efficiency using the entire SeaWIFS chlorophyll record (1998-2016) on the Ocean Productivity website with a spatial resolution of 1° x 1° (<http://www.science.oregonstate.edu/ocean.productivity/index.php>).

Bathymetric features were derived from the MARSPEC database, (<http://www.marspec.org>), which is a world ocean dataset with a spatial resolution of 1 km x 1 km (30 arc-second) developed for marine spatial ecology (Sbrocco and Barber 2013).

To ensure the same spatial resolution, all environmental data were aggregated to the lowest spatial resolution common to them ($1^{\circ} \times 1^{\circ}$), using the “raster” package in R (Hijmans et al. 2016) to avoid downscaling. This means that more detailed information, when available (e.g., slope), was retained. All environmental variables were standardized in order to enable relative weight comparisons between variables (Kinas and Andrade 2014). Multicollinearity was checked using Pearson’s correlation index. Other typical data exploration procedures were routinely done according to Zuur et al. (2010).

High correlations were found between SSS and SST ($r = 0.79$), NNP and SSS ($r = -0.91$) and NPP and SST ($r = 0.69$). Consequently, these variables were introduced separately in the different models.

Sampling uncertainties

As data were collected from different sources, a potential additional origin of variation on the species occurrence could be due to the observer effect, especially for the observations collected from the literature, from AQUAMAPS, and from fishers for whom we do not have any additional sampling information. The behavior of individual observers caused by random aspects, and/or unobserved sampling characteristics could have caused some of the variation in the data. Ignoring such non-independence of the data can lead to an invalid statistical inference (Roos et al. 2015; Costa et al. 2017). To remove any bias caused by observer-specific differences in the sampling process, an observer effect was added to the models (both for the scientific and ethnobiological datasets) as a random effect because there was no interest in knowing the specific nature of the observers.

Statistical models

The probability of *E. marginatus* occurrence (for both the scientific and ethnobiological data) was modeled using Bayesian Gaussian iCAR (intrinsic conditional autoregressive model) to take into account spatial autocorrelation (Latimer et al. 2006) and different sources of uncertainties. Bayesian species distribution models are especially suitable for data limited situations (Costa et al. 2017). We used a Gaussian iCAR (Besag 1974) to account for any eventual spatial autocorrelation between observations, assuming that the probability of species presence at one site depends on the probability of species presence in neighboring sites. Assuming the random variable Y_i follows a Bernoulli distribution, it can take on a value of 1 or 0 depending on whether the habitat is suitable ($Y_i = 1$) or not ($Y_i = 0$), then:

$$Y_i \sim \text{Bernoulli}(\pi_i)$$

$$\text{logit}(\pi_i) = X_i\beta + Z_i + \rho_{j(i)}$$

where β represents the vector of the regression coefficients, X is the matrix of covariates, Z_i is the observer random effect, and ρ represents the spatial random effect for any given cell j . The relation between π_i , the covariates of interest and spatial effect is the usual logit link.

We used uninformative priors centered at zero with a fixed large variance of 100 for all parameters involved in the model and adopted a uniform distribution for the variance of the spatial effect.

We fit these models using the “hSDM” package (Vieilledent et al. 2014) of the R statistical environment (R Development Team 2017).

We fit the resulting models obtained by combining the mentioned variables and the respective interactions and compared them using both backwards and forwards approaches

using the standard Deviance Information Criterion (DIC) (Spiegelhalter et al. 2002). Lower values of DIC represent better compromises between fit and parsimony.

Finally, in order to plot the functional response between the selected environmental variables and the predicted values, we used the “ggplot” package of the R software to apply a smoothing function to attempt to capture the general patterns in the spatial trend of the species, while also reducing the noise. This technique uses locally weighted scatterplot smoothing (lowess), which is an outlier-resistant method that estimates a polynomial regression curve by using local and bootstrap techniques with the percentile method in order to account for the variability in the original lowess fit. With this methodology, each functional response shows a 95% confidence interval for the original lowess.

Model validation for occurrence

We validated the best selected model by using an internal 10-fold cross validation based on randomly selected training and test datasets (created by a random selection of 75% and 25% of the data respectively) (Fielding and Bell 1997), with the “*PresenceAbsence*” package in R (Freeman and Moisen 2008). We assessed the model’s performance by using the area under the receiver-operating characteristic curve (AUC) (Fielding and Bell 1997) and the “True Skill Statistic” (TSS) (Allouche et al. 2006).

Comparison of model predictions

We compared the predictions of *E. marginatus* distribution from the scientific and ethno datasets using the similarity statistics Shoener’s D and Warren’s I (Warren et al. 2008). These statistics range from 0 (no overlap between areas) to 1 (distributions are identical) (Pennino et al. 2016; 2017). We carried out these analyses using the *nicheOverlap* function of the *dismo* package (Hijmans et al. 2011) in R software.

Results

Scientific data

The occurrence of *E. marginatus* seems to be primarily determined by the depth, slope, and SST of the habitat, in addition to the spatial component that accounted for the residual spatial autocorrelation and the observer random effect (Table 1). The species is more likely to be present in shallow (≈ 300 -400 m, posterior mean = 3.71; 95% CI = [1.99, 5.64]) and cooler waters (11-25° C max, posterior mean = -4.63; 95% CI = [-8.10, -1.51]) (Figure 2a - c). Although the species seems to be present in areas ranging from sandy (low slope values) to rocky bottoms (high slope values), it showed a marked preference for a structured (rocky in most cases of the study area) seabed (slope value preference = 75-90; posterior mean = 0.18; 95% CI = [0.10, 0.99]), demonstrated by the highest occurrence probability at higher slope values (Figure 2b).

The best conditions for species occurrence in South America were met in shallow and structured grounds, with slightly cooler waters than those found in the north (observed by plotting the median posterior occurrence probability), specifically between the southeastern coast of Brazil (state of Espírito Santo) and Rio de Plata in Uruguay (Figure 3). The expected occurrence of *E. marginatus* in the north of Argentinian Patagonia was driven by three reports of its presence in the literature, but there are few observations here and the presences could be rare events.

Ethnobiological data

Based on the ethnobiological dataset, the main predictors of *E. marginatus* habitats were depth and SST, along with the spatial component that accounted for the residual spatial autocorrelation and the observer random effect (Table 2). In this specific case, slope

was not relevant. This was expected given that the occurrence locations were assigned to the ocean grid point closest to the reporting village, therefore disregarding the local type of seabed.

According to the presence data credited to the fishers, *E. marginatus* was also expected to occur in shallow (posterior mean = 3.49; 95% CI = [2.35, 7.10]) and cooler waters (posterior mean = -5.24; 95% CI = [-7.48, -0.22]). However, given that species presence or absence was assigned to the ocean grid point closest to the reporting village in our study, i.e., areas which are usually shallow, the actual most relevant information is the effect of water temperature (Figure 4). Higher occurrence probabilities were found at a depth range of ≈ 250 -450 m (Figure 4a). Interestingly, the species occurrence probability was shown to be most likely at a temperature range of between 11° C and 25° C, which is the exact range predicted from the scientific dataset (Figure 4 b).

The expected geographical distribution of *E. marginatus* in South America suggests a wider distribution pattern along the Brazilian coast than the one we observed in the scientific database (based on its median posterior probability). For instance, based on the ethnobiological data, the species is expected to occur further up north, especially between the northeastern Brazilian state of Sergipe and the southern state of Santa Catarina (Figure 5).

Model performance and comparisons

For both datasets, all the best models achieved AUC values of >0.65 , which indicates a good degree of discrimination between the locations where a species is present and where it is absent. For the scientific dataset the AUC value was 0.83, whereas this value was slightly lower for the ethnobiological dataset (0.73). Similarly, a higher TSS

value was achieved using the scientific dataset (0.74 versus 0.67). However, again both datasets achieved values higher than 0.60, indicating a good ability for the models to predict real absence and presence records for *E. marginatus*.

Interestingly, the predictions generated from both models showed an excellent degree of overlap, suggesting high similarity between them (Schoener's $D = 0.71$; Warren's $I = 0.91$).

Discussion

This study highlights two major points of relevance: (i) it is the first reliable occurrence prediction of the dusky grouper *E. marginatus* carried out using a state-of-the-art species distribution model (Costa et al. 2017), (ii) and it provides evidence that such models can be built from data-poor situations, such as when the only information available is that provided by fishers.

Similar to what has been shown in the Mediterranean, where *E. marginatus* has been widely studied, in our study we also expected that the species would have a strong preference for sub-tropical shallower waters, and for habitats with complex topography (Mesa et al. 2002). Whereas in well studied parts of the Mediterranean the species has been shown to prefer areas with a depth of up to 50 m (Heemstra and Randall 1993), our model of the Southwest Atlantic waters suggested a higher likelihood of its occurrence in areas with a depth of between 300 - 400 m. Although different *E. marginatus* populations (Mediterranean x Southwest Atlantic) might be suited to different habitats, we cannot rule out the possibility that overfishing or coastal degradation in the Southwest Atlantic plays a role (Bruslé 1985; Morris et al. 2000; Begossi et al. 2012), thus making it harder to find adults in shallower waters. Even in the Mediterranean the species can be found in deeper

waters (up to 250 m), especially adults (Harmelin and Harmelin-Vivien 1999). We did not have information on individual size, which may have supported the depth pattern observed in the Mediterranean. Still, the habitat suitability comparison between regions is a rough attempt, since this is the first prediction ever made for this species using similar models.

What is mainly innovative here is the suggestion that mere information provided by geographically dispersed fishers on whether they catch this species in their customary fishing grounds or not is enough to predict species distribution and achieve results similar to those made with scientific data. This achievement is especially relevant given our sampling limitations. For instance, our technique of allocating a random spot right across from each fishing village where species occurrence was mentioned, to represent where the species is caught, is an important caveat, because we did not consider ground specificities. Clearly fishers go to specific grounds, such as rocky outcrops, to catch the dusky grouper (Begossi and Silvano 2008; Begossi et al. 2012), but this information was not available in most of the papers we reviewed. Nevertheless, even considering this limitation, the predictive model was accurate enough to show between 71% and 91% of overlap with its counterpart scientific model, in addition to performing well as a reliable predictive model (see values of AUC and TSS). Similar to the scientific model, the ethnobiological model also predicted *E. marginatus* occurrences in cooler and shallower waters. However, depth in this case was just an artifact of our method of assigning a random point in the water to represent its occurrence. On the other hand, for a species that seems to have a well-established temperature range (Hereu et al. 2006), the simple information of village location works as an accurate predictor of expected species occurrence, especially because small-scale fishers do not venture to distant grounds with their small vessels.

Although overall the expected distribution was the same for both methods, there are

some differences worth investigating. The first is the extended expected distribution down south, towards Argentinian Patagonia, which is observed only in the scientific modeling approach. We considered species occurrence along the southern coast of Brazil, close to the mouth of Rio de la Plata in Uruguay, and in Argentinian Patagonian due to three reports that expanded the geographical limit of the species occurrence in the last decades (Irigoyen et al. 2005; Condini et al. 2016). The species is by no means common in those waters, although the existence of suitable habitats in the region might represent a future geographical range in a warming world (Perry et al. 2005). However, the ethnobiological model could not predict *E. marginatus* occurrence south of Brazil because no study has ever approached Uruguayan and Argentinian fishers about species occurrence in that region.

Another striking difference between the two models was that the ethnobiological model predicted the expected occurrence of the dusky grouper to be at the southern tip of the northeastern Brazilian coast, which is north of what the scientific model predicted. With the exception of three identical reports from the online databases that attributed the presence of *E. marginatus* to the southern part of the Brazilian northeast, the scientific model did not predict the northeast region to house the species (although there were some reports of its presence in an offshore location in the north). In specific ethnobiological studies designed to confirm its existence in the north with fishers, no such confirmation was obtained above -12° degrees of latitude (Ribeiro 2016). However, the region between -15° and -12° degrees seems to be a grayish zone, with some fishers reporting its occurrence (Silvano et al. 2006). The northern limit of the *E. marginatus* distribution established from information given by fishers seems to be close to a geographical barrier for some species, such as some of the Dasyatidae rays (Costa et al. 2017). One possible explanation has to do

with the San Francisco river; a large warm river that could be working as a barrier for multiple species for various reasons, such as changes in temperature (Costa et al. 2017). For *E. marginatus*, for instance, the warm river waters may make the northern coast unfit for it. This break also broadly coincides with the breaks between the ecoregions of Spalding et al. (2007).

However, it is not possible to rule out the possibility that fishers may have been mistaken, confounding *E. marginatus* with another similar grouper that does occur in the region, *E. morio* (Valenciennes, 1828) (Costa et al. 2003; Begossi et al. 2012). Another possibility is that the fishers that mentioned the occurrence of *E. marginatus* north of its known distribution are those relatively close to the expected distribution limit of this species. They could be fishers that encounter the grouper when fishing slightly down south. Although small-scale fishers in Brazil are not really mobile, performing mostly daily fishing trips close to their villages (Lopes et al. 2009), some cross state borders to fish in other regions (Silva, pers. comm.). This could represent an opportunity to come across species not usually found close to their home villages. Finally, we should not disregard the species' past presence in the area either, perhaps in lower abundances that were either overfished or dislodged with warmer waters over the last decades.

It is also important to take into account larval dispersal distances where predictions differed slightly between models. For the Mediterranean it is estimated that the median distance for larval dispersal is 120km, but a maximum distance of 906 km is also possible (Andrello et al. 2013). If larvae reach such distances, we could expect some sporadic occurrences of the species outside its standard geographical limits when the larvae eventually find some intermittent favorable conditions to settle. That would possibly allow the maturation of a few extralimital adults beyond the species range, although not enough

395 to form a viable fishing population.

396 Overall, this study points towards a promising low-cost approach (LEK) to provide
397 baseline data and to fill knowledge gaps to support not only fisheries management but also
398 marine conservation initiatives, such as marine spatial planning. Regardless if it is to be
399 used alone or as part of an integrative approach, we advise future studies on fishers' local
400 ecological knowledge to compile data on species occurrence per fishing ground with their
401 respective locations (e.g., Begossi et al. 2013). We also recommend future studies to be
402 specifically designed to collect LEK fish abundance data to support its reliability as a
403 conservation tool (Anadón et al. 2010). Finally, it is worth mentioning that this study also
404 shows that the integration of multiple sources of information into science and management
405 is a potentially invaluable tool. Such tool should not be overlooked, despite the limitations
406 of integrating information obtained through different sampling designs, which limit the
407 statistical methods that can be used. Indeed, further statistical advancements should work
408 towards developing single models that can account for all the variability that emerges
409 from combining data collected with different sampling techniques over distinct temporal
410 and spatial coverage. The Bayesian approach may provide the answer to combine datasets
411 from different sampling schemes while maintaining a sound statistical framework.

412 In conclusion, this is the first study to predict *E. marginatus* occurrence along the South
413 American coast, which is highly relevant given the threatened state of the species and its
414 importance as a fishing target (Sadovy de Mitcheson et al. 2013). Our study could also possibly be
415 hinting at an expected future distribution towards the southern cooler waters in the future (Perry
416 et al. 2005), by confirming the existence of suitable habitats close to Patagonia. Moreover, our
417 study demonstrated that with little data, provided by fishers, it is possible to get accurate species

distributions at least for species limited by more general aspects such as temperature and salinity, which are not expected to vary greatly between fishing grounds. For species whose occurrence are affected by different requirements, more specific ethnobiological knowledge might be necessary, whereas fishers may not be able to provide accurate information for species they do not target regularly. Nevertheless, sampling ethnobiological data to predict the distribution of target species (or species with which people interact regularly) is affordable and allows the coverage of large geographical areas, which may be a considerable advancement for data-poor countries with urgent needs for proper fisheries management. Ethnobiological data can also be incorporated into integrative and potentially more informative approaches to advance marine spatial planning and conservation in general.

Acknowledgements

We are deeply grateful to all the fishers who took part in our research over the last 30 years. AB and PFML thank CNPq for a productivity grant. This research was indirectly funded by different institutions over the last 30 years, specifically FAPESP (1992-2017), CNPq (1986-1987), FAPERJ (1991-1992) and Alfred Sloan Foundation (1986-1987).

References

- Agardy, T., di Sciara, G. N., and Christie, P. 2011. Mind the gap: Addressing the shortcomings of marine protected areas through large scale marine spatial planning. *Marine Policy* 35:226–232.
- Allouche, O., Tsoar, A., and Kadmon, R. 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology* 43:1223–1232.
- Anadón, J. D., Giménez, A., and Ballestar, R. 2010. Linking local ecological knowledge and habitat modelling to predict absolute species abundance on large scales. *Biodiversity and Conservation* 19:1443–1454.

- 444 Andrade, Á. B., Machado, L. F., Hostim-Silva, M., and Barreiros, J. P. 2003. Reproductive biology of
445 the dusky grouper *Epinephelus marginatus* (Lowe, 1834). Brazilian Archives of Biology and
446 Technology 46:373–382.
- 447 Andrello, M., Mouillot, D., Beuvier, J., Albouy, C., Thuiller, W., and Manel, S. 2013. Low
448 Connectivity between Mediterranean Marine Protected Areas: A Biophysical Modeling Approach
449 for the Dusky Grouper *Epinephelus marginatus*. PLOS ONE 8:e68564.
- 450 Barber, P. H., Ablan-Lagman, M. C. A., Ambariyanto, Berlinck, R. G. , Cahyani, D., Crandall, E.D.,
451 Ravago-Gotanco, R., Juinio-Meñez, M.A., Mahardika, I. N., Shanker, K., Starger, C.J, Toha, A. H. A.,
452 Anggoro, A. W., and Willette, D. A. 2014. Advancing biodiversity research in developing countries:
453 the need for changing paradigms. Bulletin of Marine Science 90:187–210.
- 454 Begossi, A., Camargo, E., and Carpi Jr, S. 2013. Os mapas da pesca artesanal: pescueiros e
455 pescadores na costa do Brasil. Editora Rima, São Carlos.
- 456 Begossi, A., Lopes, P. F. M., and Silvano, R. A. M. 2012. Co-Management of Reef Fisheries of the
457 Snapper-Grouper Complex in a Human Ecological Context in Brazil. Pages 1–22 Global Progress in
458 Ecosystem-Based Fisheries Management. Alaska Sea Grant, University of Alaska Fairbanks, Alaska.
- 459 Begossi, A., Salivonchyk, S., Lopes, P. F. M., and Silvano, R. A. M. 2016. Fishers' knowledge on the
460 coast of Brazil. Journal of Ethnobiology and Ethnomedicine 12:20.
- 461 Begossi, A., and Silvano, R. A. M. 2008. Ecology and Ethnoecology of dusky grouper, garoupa,
462 [*Epinephelus marginatus* (Lowe, 1834)] along the coast of Brazil. Journal of Ethnobiology and
463 Ethnomedicine 4:1–20.
- 464 Bender, M. G., Machado, G. R., Silva, P. J. de A., Floeter, S. R., Monteiro-Netto, C., Luiz, O. J., and
465 Ferreira, C. E. L. 2014. Local Ecological Knowledge and Scientific Data Reveal Overexploitation by
466 Multigear Artisanal Fisheries in the Southwestern Atlantic. PLoS ONE 9:e110332.
- 467 Berkes, F. 2003. Alternatives to conventional management: lessons from small-scale fisheries.
468 Environments 31:5–19.
- 469 Besag, L. 1974. Spatial interaction and the statistical analysis of Lattice systems. Journal of the
470 Royal Statistical Society. Series B (Methodological) 36:192–236.
- 471 Bevilacqua, A. H. V., Carvalho, A. R., Angelini, R., and Christensen, V. 2016. More than anecdotes:
472 fishers' ecological knowledge can fill gaps for ecosystem modeling. PLOS ONE 11:e0155655.
- 473 Bouchereau, J. L., Body, P., and Chauvet, C. 1999. Growth of the dusky grouper *Epinephelus*
474 *marginatus* (Linnaeus, 1758) (Teleostei, Serranidae), in the Natural Marine Reserve of Lavezzi
475 Islands, Corsica, France. Scientia Marina 63:71–77.

- 476 Bruslé, J. 1985. Exposé synoptique des données biologiques sur les mérous *Epinephelus aeneus*
477 (Geoffroy Saint Hilaire, 1809) et *Epinephelus guaza* (Linnaeus, 1758) de l'Océan Atlantique et de
478 la Méditerranée. *FAO Syn. Pêches* 129:1–64.
- 479 Castello, L., Viana, J. P., Watkins, G., Pinedo-Vasquez, M., and Luzadis, V. A. 2009. Lessons from
480 integrating fishers of arapaima in small-scale fisheries management at the Mamirauá Reserve,
481 Amazon. *Environmental Management* 43:197–209.
- 482 Chao, N. L., Frédou, F. L., Haimovici, M., Peres, M. B., Polidoro, B., Raseira, M., Subirá, R., and
483 Carpenter, K. 2015. A popular and potentially sustainable fishery resource under pressure—
484 extinction risk and conservation of Brazilian Sciaenidae (Teleostei: Perciformes). *Global Ecology*
485 *and Conservation* 4:117–126.
- 486 Cinner, J., and Huchery, C. 2014. A comparison of social outcomes associated with different
487 fisheries co-management institutions. *Conservation Letters* 7:224–232.
- 488 Cinti, A., Shaw, W., Cudney-Bueno, R., and Rojo, M. 2010. The unintended consequences of formal
489 fisheries policies: Social disparities and resource overuse in a major fishing community in the Gulf
490 of California, Mexico. *Marine Policy* 34:328–339.
- 491 Condini, M. V., Velez-Rubio, G. M., Fallabrino, A., and Garcia, A. M. 2016. First occurrence of dusky
492 grouper *Epinephelus marginatus* (Lowe, 1834) in a Marine Protected Area on the Uruguayan coast.
493 *Cah. Biol. Mar.*:57.
- 494 Costa, P. A. S., Braga, A. D., and da Rocha, L. O. F. 2003. Reef fisheries in Porto Seguro, eastern
495 Brazilian coast. *Fisheries Research* 60:577–583.
- 496 Costa, T. L. A., Pennino, M. G., and Mendes, L. F. 2017. Identifying ecological barriers in marine
497 environment: The case study of *Dasyatis marianae*. *Marine Environmental Research* 125:1–9.
- 498 Damasio, L. de M. A., Lopes, P. F. M., Guariento, R. D., and Carvalho, A. R. 2015. Matching Fishers'
499 Knowledge and Landing Data to Overcome Data Missing in Small-Scale Fisheries. *PLoS ONE*
500 10:e0133122.
- 501 Davis, A., Hanson, J. M., Watts, H., and MacPherson, H. 2004. Local ecological knowledge and
502 marine fisheries research: the case of white hake (*Urophycis tenuis*) predation on juvenile
503 American lobster (*Homarus americanus*). *Canadian Journal of Fisheries and Aquatic Sciences*
504 61:1191–1201.
- 505 Davis, A., and Wagner, J. R. 2003. Who Knows? On the Importance of Identifying “Experts” When
506 Researching Local Ecological Knowledge. *Human Ecology* 31:463–489.
- 507 Dayton, P. K. 1998. Reversal of the burden of proof in fisheries management. *Science* 279:821–
508 822.

- 509 Di Dario, F., Alves, C. B. M., Boos, H., Frédou, F. L., Lessa, R. P. T., Mincarone, M. M., Pinheiro, M. A.
510 A., Polaz, C. N. M., Reis, R. E., Rocha, L. A., Santana, F. M., Santos, R. A., Santos, S. B., Vianna, M.,
511 and Vieira, F. 2015. A better way forward for Brazil's fisheries. *Science* 347:1079.
- 512 Eddy, T. D., Gardner, J. P. A., and Pérez-Matus, A. 2010. Applying fishers' ecological knowledge to
513 construct past and future lobster stocks in the Juan Fernández archipelago, Chile. *PLoS ONE* 5:1–
514 12.
- 515 Fielding, A. H., and Bell, J. F. 1997. A review of methods for the assessment of prediction errors in
516 conservation presence/absence models. *Environmental Conservation* 24:38–49.
- 517 Finkbeiner, E. M., and Basurto, X. 2015. Re-defining co-management to facilitate small-scale
518 fisheries reform: An illustration from northwest Mexico. *Marine Policy* 51:433–441.
- 519 Freeman, E. A., and Moisen, G. 2008. PresenceAbsence: An R Package for Presence Absence
520 Analysis. *Journal of Statistical Software* 23.
- 521 Gaines, S. D., White, C., Carr, M. H., and Palumbi, S. R. 2010. Designing marine reserve networks
522 for both conservation and fisheries management. *Proceedings of the National Academy of*
523 *Sciences* 107:18286–18293.
- 524 Gilles, A., Miquelis, A., Quignard, J.-P., and Faure, É. 2000. Molecular phylogeography of western
525 Mediterranean dusky grouper *Epinephelus marginatus*. *Comptes Rendus de l'Académie des*
526 *Sciences - Series III - Sciences de la Vie* 323:195–205.
- 527 Harmelin, J.-G., and Harmelin-Vivien, M. 1999. A review on habitat, diet and growth of the dusky
528 grouper *Epinephelus marginatus* (Lowe, 1834). Pages 11–20 *Marine life*. Institut océanographique
529 Paul Ricard, Les Embiez.
- 530 Heemstra, P. C., and Randall, J. E. 1993. *FAO Fisheries Catalogue: An annotated and illustrated*
531 *catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date*. FAO,
532 Rome.
- 533 Hereu, B., Diaz, D., Pasqual, J., Zabala, M., and Sala, E. 2006. Temporal patterns of spawning of the
534 dusky grouper *Epinephelus marginatus* in relation to environmental factors. *Marine Ecology*
535 *Progress Series* 325:187–194.
- 536 Hijmans, R. J., van Etten, J., Cheng, J., Mattiuzzi, M., Summer, M., Greenberg, J. A., and Ghosh, A.
537 2016. Package "raster". R package. R package.
- 538 Hijmans, R. J., Phillips, S., Leathwick, L., and Elith, J. 2011. Package "dismo."
- 539 Hilborn, R. 2007. Defining success in fisheries and conflicts in objectives. *Marine Policy* 31:153–
540 158.

- 541 Houk, P., Rhodes, K., Cuetos-Bueno, J., Lindfield, S., Fread, V., and McIlwain, J. L. 2012. Commercial
542 coral-reef fisheries across Micronesia: A need for improving management. *Coral Reefs* 31:13–26.
- 543 Irigoyen, A. J., Galván, D. E., and Venerus, L. A. 2005. Occurrence of dusky grouper *Epinephelus*
544 *marginatus* (Lowe, 1834) in gulfs of northern Patagonia, Argentina. *Journal of Fish Biology*
545 67:1741–1745.
- 546 Jentoft, S. 2000. Legitimacy and disappointment in fisheries management. *Marine Policy* 24:141–
547 148.
- 548 Johannes, R. 1998. The case for data-less marine resource management: examples from tropical
549 nearshore finfisheries. *Trends in Ecology & Evolution* 13:243–246.
- 550 Johannes, R. E. 1981. Words of the lagoon: fishing and marine lore in the Palau district of
551 Micronesia. University of California Press, Berkeley.
- 552 Jones, J. P. G., Andriamarovololona, M. M., Hockley, N., Gibbons, J. M., and Milner-Gulland, E. J.
553 2008. Testing the use of interviews as a tool for monitoring trends in the harvesting of wild
554 species. *Journal of Applied Ecology* 45:1205–1212.
- 555 Kaufman, L., Heneman, B., Barnes, T. J., and Fujita, R. 2004. Transition from low to high data
556 richness: an experiment in ecosystem-based fishery management from California. *Bulletin of*
557 *Marine Science* 74:693–708.
- 558 Kinas, P. G., and Andrade, H. A. 2014. *Introdução à Análise Bayesiana (com R)*. Buqui Livros
559 Digitais, Porto Alegre.
- 560 Latimer, A. M., Wu, S., Gelfand, A. E., and Silander, J. A. 2006. Building statistical models to
561 analyze species distributions. *Ecological Applications: A Publication of the Ecological Society of*
562 *America* 16:33–50.
- 563 Lopes, P. F. M., Francisco, A. S., and Begossi, A. 2009. Artisanal commercial fisheries at the
564 southern coast of São Paulo state, Brazil: Ecological, social and economic structures. *Interciencia*
565 34:536–542.
- 566 Lopes, P. F. M., Rosa, E. M., Salyvonchik, S., Nora, V., and Begossi, A. 2013a. Suggestions for fixing
567 top-down coastal fisheries management through participatory approaches. *Marine Policy* 40:100–
568 110.
- 569 Lopes, P. F. M., Silvano, R. A. M., and Begossi, A. 2011. Extractive and Sustainable Development
570 Reserves in Brazil: resilient alternatives to fisheries? *Journal of Environmental Planning and*
571 *Management* 54:421–443.
- 572 Lopes, P. F. M., Silvano, R. A. M., Nora, V. A. and Begossi, A. 2013b. Transboundary Socio-Ecological
573 Effects of a Marine Protected Area in the Southwest Atlantic. *AMBIO* 42:963–974.

- 574 Machado, L. F., Andrade, A. B., Hostim-Silva, M., and Barreiros, J. P. 2003. Habitat use by the
575 juvenile dusky grouper *Epinephelus marginatus* and its relative abundance, in Santa Catarina,
576 Brazil. *Journal of Ichthyology and Aquatic Biology* 6:133–138.
- 577 Machado, L. F., Daros, F. A. M. L., Andrade Bertoncini, A., Hostim-Silva, M., and Barreiros, J. P.
578 2008. Feeding strategy and trophic ontogeny in *Epinephelus marginatus* (Serranidae) from
579 Southern Brazil. *Cybio* 32:33–41.
- 580 Marshall, N. A., and Marshall, P. A. 2007. Conceptualizing and operationalizing social resilience
581 within commercial fisheries in Northern Australia. *Ecology And Society* 12:1.
- 582 Mesa, G. L., Louisy, P., and Vacchi, M. 2002. Assessment of microhabitat preferences in juvenile
583 dusky grouper (*Epinephelus marginatus*) by visual sampling. *Marine Biology* 140:175–185.
- 584 Moller, H., Berkes, F., Lyver, P. O., and Kislalioglu, M. 2004. Combining Science and Traditional
585 Ecological Knowledge: Monitoring Populations for Co-Management. *Ecology and Society* 9.
- 586 Mora, C., Myers, R. A., Coll, M., Libralato, S., Pitcher, T. J., Sumaila, R. U., Zeller, D., Watson, R.,
587 Gaston, K. J., and Worm, B. 2009. Management effectiveness of the world's marine fisheries. *PLOS*
588 *Biology* 7:e1000131.
- 589 Morris, A. V., Roberts, C. M., and Hawkins, J. P. 2000. The threatened status of groupers
590 (*Epinephelinae*). *Biodiversity & Conservation* 9:919–942.
- 591 Neis, B., Schneider, D. C., Felt, L., Haedrich, R. L., Fischer, J., and Hutchings, J. A. 1999. Fisheries
592 assessment: what can be learned from interviewing resource users? *Canadian Journal of Fisheries*
593 *and Aquatic Sciences* 56:1949–1963.
- 594 Pauly, D., and Zeller, D. 2016. Catch reconstructions reveal that global marine fisheries catches are
595 higher than reported and declining. *Nature Communications* 7:10244.
- 596 Pennino, M. G., Arcangeli, A., Fonseca, V. P., Campana, I., Pierce, G. J., Rotta, A., and Bellido, J. M.
597 2017. A spatially explicit risk assessment approach: Cetaceans and marine traffic in the Pelagos
598 Sanctuary (Mediterranean Sea). *PLOS ONE* 12:e0179686.
- 599 Pennino, M. G., Conesa, D., López-Quílez, A., Muñoz, F., Fernández, A., and Bellido, J. M. 2016.
600 Fishery-dependent and -independent data lead to consistent estimations of essential habitats.
601 *ICES Journal of Marine Science* 73:2302–2310.
- 602 Perry, A. L., Low, P. J., Ellis, J.R., and Reynolds, J. D. 2005. Climate change and distribution shifts in
603 marine fishes. *Science* 308:1912–1915.
- 604 Pinheiro, H. T., Dario, F. D., Gerhardinger, L. C., de Melo, M. R. S., Moura, R. L. de, Reis, R. E.,
605 Vieira, F., Zuanon, J., and Rocha, L. A. 2015. Brazilian aquatic biodiversity in peril. *Science*
606 350:1043–1044.

- 607 Priolli, R. H. G., Bajay, M. M., Silvano, R. A. M., and Begossi, A. 2016. Population genetic structure
608 of an estuarine and a reef fish species exploited by Brazilian artisanal fishing. *Scientia Marina*
609 80:467–477.
- 610 R Development Team. 2017. R: A language and environment for statistical computing. R
611 Foundation for Statistical Computing, Vienna, Austria.
- 612 Ribeiro, A. R. 2016. Conhecimento ecológico local de pescadores sobre peixes recifais
613 (Epinephelidae) no litoral nordeste do Brasil. Master's, Universidade Federal do Rio Grande do Sul,
614 Porto Alegre.
- 615 Roos, N. C., Carvalho, A. R., Lopes, P. F. M., and Pennino, M. G. 2015. Modeling sensitive parrotfish
616 (Labridae: Scarini) habitats along the Brazilian coast. *Marine Environmental Research* 110:92–100.
- 617 Roos, N. C., Pennino, M. G., Lopes, P. F. M., and Carvalho, A. R. 2016. Multiple management
618 strategies to control selectivity on parrotfishes harvesting. *Ocean & Coastal Management* 134:20–
619 29.
- 620 Ruddle, K., and Davis, A. 2011. What is “Ecological” in Local Ecological Knowledge? Lessons from
621 Canada and Vietnam. *Society & Natural Resources* 24:887–901.
- 622 Sadovy de Mitcheson, Y., Craig, M.T., Bertoncini, A.A., Carpenter, K. E., Cheung, W. W. L., Choat, J.
623 H., Cornish, A. S., Fennessy, S. T., Ferreira, B. P., Heemstra, P. C., Liu, M., Myers, R. F., Pollard, D. A.,
624 Rhodes, K. L., Rocha, L. A., Russell, B. C., Samoily, M. A., and Sanciangco, J. 2013. Fishing groupers
625 towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery.
626 *Fish and Fisheries* 14:119–136.
- 627 Sbrocco, E. J., and Barber, P. H. 2013. MARSPEC: ocean climate layers for marine spatial ecology.
628 *Ecology* 94:979–979.
- 629 Silvano, R. A. M., and Begossi, A. 2010. What can be learned from fishers? An integrated survey of
630 fishers' local ecological knowledge and bluefish (*Pomatomus saltatrix*) biology on the Brazilian
631 coast. *Hydrobiologia* 637:3–18.
- 632 Silvano, R. A. M., MacCord, P. F. L., Lima, R.V., and Begossi, A. 2006. When does this fish spawn?
633 Fishermen's local knowledge of migration and reproduction of Brazilian coastal fishes.
634 *Environmental Biology of Fishes* 76:371–386.
- 635 Silvano, R., and Valbo-Jorgensen, J. 2008. Beyond fishermen's tales: contribution of fishers' local
636 ecological knowledge to fish ecology and fisheries management. *Environment Development and*
637 *Sustainability* 10:657–675.
- 638 Spalding, M. D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern, B.S.,
639 Jorge, M.A., Lombana, A., Lourie, S.A., Martin, K.D., Mcmanus, E., Molnar, J., Recchia, C.A., and
640 Robertson, J. 2007. Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf
641 Areas. *BioScience* 57:573–583.

- 642 Spiegelhalter, D. J., Best, N.G., Carlin, B. P., and Van Der Linde, A. 2002. Bayesian measures of
643 model complexity and fit. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*
644 64:583–639.
- 645 Tesfamichael, D., Pitcher, T. J., and Pauly, D. 2014. Assessing Changes in Fisheries Using Fishers’
646 Knowledge to Generate Long Time Series of Catch Rates: a Case Study from the Red Sea. *Ecology*
647 and Society 19.
- 648 Vieilledent, G., Merow, C., Guélat, J., Latimer, A. M., Kéry, M., Gelfand, A. E., Wilson, A. M.,
649 Mortier, F., and Silander Jr, J.A. 2014. hSDM: hierarchical Bayesian species distribution models.
- 650 Warren, D. L., Glor, R. E., and Turelli, M. 2008. Environmental niche equivalency versus
651 conservatism: quantitative approaches to niche evolution. *Evolution* 62:2868–2883.
- 652 Worm, B., and Branch, T. A. 2012. The future of fish. *Trends in Ecology & Evolution* 27:594–599.
- 653 Zuur, A. F., Leno, E.N., and Elphick, C. S. 2010. A protocol for data exploration to avoid common
654 statistical problems. *Methods in Ecology and Evolution* 1:3–14.

655

Draft

Table 1: Comparison of the most relevant *E. marginatus* scientific dataset models. Deviance Information Criterion (DIC) scores measure the goodness-of-fit of each model. Predictor acronyms are: sea surface temperature (SST), sea surface salinity (SSS), net primary productivity (NPP), depth (D), slope (S) concavity of the seabed (C), spatial effect (ρ), observer effect (Z). Relevant environmental predictors are indicated by *, i.e., those predictors with 95% credibility intervals not covering zero. The best model is highlighted in bold.

Models	DIC
$1 + \text{SST}^* + \text{D}^* + \text{S} + \text{C} + \text{Z}^* + \rho^*$	47.43
$1 + \text{SST}^* + \text{D}^* + \text{S}^* + \text{Z}^* + \rho^*$	45.62
$1 + \text{D}^* + \text{S} + \text{Z}^* + \rho^*$	50.58
$1 + \text{D}^* + \text{C} + \text{Z}^* + \rho^*$	52.36
$1 + \text{SST}^* + \text{D}^* + \text{Z}^* + \rho^*$	47.74
$1 + \text{SSS} + \text{D}^* + \text{S} + \text{Z}^* + \rho^*$	50.38
$1 + \text{SSS} + \text{D}^* + \text{Z}^* + \rho^*$	50.17
$1 + \text{NPP} + \text{D}^* + \text{S} + \text{Z}^* + \rho^*$	51.77
$1 + \text{NPP} + \text{D}^* + \text{Z}^* + \rho^*$	57.97
$1 + \text{NPP}^* + \text{Z}^* + \rho^*$	57.36
$1 + \text{S} + \text{Z}^* + \rho^*$	61.08
$1 + \text{D}^* + \text{Z}^* + \rho^*$	51.44
$1 + \text{SST}^* + \text{Z}^* + \rho^*$	57.19
$1 + \text{SSS}^* + \text{Z}^* + \rho^*$	57.66

Table 2: Comparison of the most relevant *E. marginatus* ethnobiological dataset models. Deviance Information Criterion (DIC) scores measure goodness-of-fit. Predictor acronyms are: sea surface temperature (SST), sea surface salinity (SSS), net primary productivity (NPP), depth (D), slope (S) concavity of the seabed (C), spatial effect (ρ), observer effect (Z). Relevant environmental predictors are indicated by *, i.e., those predictors with 95% credibility intervals not covering zero. The best model is highlighted in bold.

Models	DIC
$1 + \text{SST}^* + \text{D}^* + \text{S} + \text{C}^* + \text{Z}^* + \rho^*$	21.54
$1 + \text{SST} + \text{D} + \text{S} + \text{Z}^* + \rho^*$	19.61
$1 + \text{D} + \text{S} + \text{Z}^* + \rho^*$	18.57
$1 + \text{D}^* + \text{C}^* + \text{Z}^* + \rho^*$	17.12
$1 + \text{SST}^* + \text{D}^* + \text{Z}^* + \rho^*$	14.89
$1 + \text{SSS} + \text{D} + \text{C} + \text{Z} + \rho^*$	15.13
$1 + \text{SSS} + \text{C}^* + \text{Z}^* + \rho^*$	16.39
$1 + \text{SSS} + \text{D}^* + \text{Z}^* + \rho^*$	18.70
$1 + \text{NPP} + \text{D}^* + \text{Z}^* + \rho^*$	19.23
$1 + \text{NPP} + \text{Z}^* + \rho^*$	21.31
$1 + \text{C}^* + \text{Z}^* + \rho^*$	16.53
$1 + \text{D}^* + \text{Z}^* + \rho^*$	18.26
$1 + \text{SST}^* + \text{Z}^* + \rho^*$	19.24
$1 + \text{SSS} + \text{Z}^* + \rho^*$	21.48

Draft

Figure legends

Figure 1: Sampling locations for the scientific (a) and ethnobiological (b) datasets. Red dots indicate presences, while black dots reflect absences.

Figure 2. Smooth function of the predicted occurrence probability for depth (a), slope (b) and SST (c), using the scientific dataset. The solid line is the smooth function estimate and shaded regions represent the approximate 95% credibility interval (CI).

Figure 3. Mean occurrence probability of the *E. marginatus* species along South American waters, obtained using the scientific dataset on its presence/absence.

Figure 4. Smooth function of the predicted occurrence probability of depth (a) and SST (b), using the ethnobiological dataset. The solid line is the smooth function estimate. Shaded regions represent the approximate 95% credibility interval (CI).

Figure 5. Mean occurrence probability of the *E. marginatus* species along South American waters obtained using the ethnobiological dataset.

334x206mm (300 x 300 DPI)

369x127mm (300 x 300 DPI)

186x219mm (300 x 300 DPI)

243x117mm (300 x 300 DPI)

186x219mm (300 x 300 DPI)