Canadian Journal of Zoology

Experimental assessments of marine species sensitivities to ocean acidification and co-stressors: how far have we come?

Journal:	<i>Canadian Journal of Zoology</i>
Manuscript ID	cjz-2018-0198.R1
Manuscript Type:	Review
Date Submitted by the Author:	30-Oct-2018
Complete List of Authors:	Baumann, Hannes; University of Connecticut, Department of Marine Sciences
Is your manuscript invited for consideration in a Special Issue?:	Canadian Society of Zoologists 2018
Keyword:	FISH < Taxon, Ocean Variability Hypothesis, ocean warming, CO2 sensitivity, oxygen decline

SCHOLARONE™
Manuscripts

1 PERSPECTIVE

2 **Experimental assessments of marine species sensitivities to ocean acidification and**
3 **co-stressors: how far have we come?**¹

4 Hannes Baumann

5

6 University of Connecticut, Department of Marine Sciences, 1080 Shennecossett Rd,
7 Groton, CT 06340 | hannes.baumann@uconn.edu, 860-405-9297

8

9 Running title: CO₂ experiments – progress and challenges

10

11

12

13

14

15

16

17

18

19

20

21

22

¹ This review is one of a series of invited papers arising from the symposium “Aquatic environments and organisms in transition” that was co-sponsored by the Canadian Society of Zoologists and the Canadian Journal of Zoology and held during the Annual Meeting of the Canadian Society of Zoologists at Memorial University of Newfoundland, St. John’s, Newfoundland, 7–11 May 2018.

Abstract

Experimental studies assessing the potential impacts of ocean acidification on marine organisms have rapidly expanded and produced a wealth of empirical data over the past decade. This perspective examines four key areas of transformative developments in experimental approaches: **(1)** methodological advances; **(2)** advances in elucidating physiological and molecular mechanisms behind observed CO₂ effects; **(3)** recognition of short-term CO₂ variability as a likely modifier of species sensitivities (**Ocean Variability Hypothesis**) and **(4)** consensus on the multistressor nature of marine climate change where effect interactions are still challenging to anticipate. No single experiment allows predicting the fate of future populations. But sustaining the accumulation of empirical evidence is critical for more robust estimates of species reaction norms and thus for enabling better modeling approaches. Moreover, advanced experimental approaches are needed to address knowledge gaps including changes in species interactions, intra-specific variability in sensitivity and its importance for the adaptation potential of marine organisms to a high CO₂ world.

Keywords: Ocean Variability Hypothesis, ocean warming, acidification, oxygen decline

The challenge of experimentally addressing marine climate change

Anthropogenic climate change has been a scientific consensus for decades, but its reality is coming into ever sharper focus with each new data trove we collect on earth's climate and biosphere (IPCC 2014; Shepherd et al. 2018). Excessive carbon dioxide emissions (CO₂) have intensified earth's greenhouse effect, resulting in land and ocean warming (Mitchell 1989). Warming oceans decline in oxygen content and stratify more (Levin and Breitburg 2015; Breitburg et al. 2018), while increased CO₂ dissolution causes ocean acidification (Doney et al. 2009). The problem is regionally exacerbated by anthropogenic nutrient loads (eutrophication) stimulating primary production and successive microbial respiration (Diaz and Rosenberg 2008; Cloern et al. 2016), causing metabolic oxygen depletion and acidification in coastal habitats (Cai et al. 2011; Wallace et al. 2014; Gobler and Baumann 2016). Ocean warming, acidification and oxygen decline therefore comprise the main concurrent symptoms of marine climate change (Pörtner 2012; Baumann 2016).

There is no doubt that marine climate change will profoundly impact marine life in ways that are of concern to human food security, well-being, and places to inhabit. But which species exactly will thrive and which will decline? Which will genetically adapt? Will ecosystems become less productive or resilient? None of these questions can presently be answered with confidence, given the large inherent uncertainties in forecasts of non-linear population and ecosystem responses and the systemic lack of controls, replication, and the immediate ability to falsify predictions, which challenge the scientific method. The difficulties at times elicit extreme notions – fatalism, naiveté, or cynicism – three fallacies that must be avoided. Fatalism can emerge when pioneering research implies the *potential* for broad, devastating impacts (“All is lost!”) and such messages then reinforce themselves in press releases and public commentary (Browman 2016). This has arguably happened for some groundbreaking experimental assessments on coral, mollusk, and larval fish CO₂ sensitivities (e.g., Orr et al. 2005; Fine and Tchernov 2007; Baumann et al. 2012). However, subsequent studies have since broadened the empirical database, and when more complex and often contradictory patterns emerged (Busch et al. 2015) some have questioned the threat altogether (“Don’t worry!”). Yet naiveté is as unhelpful as the cynic notion that all approaches have inherent shortcomings and answers are therefore

elusive (“Nothing we can do!”). In reality, not all is lost, but there is reason to worry, and science has tools to address the problem. As the past decade of experimental, observational, and modeling approaches has demonstrated, by avoiding extremes science has advanced our understanding of potential species and ecosystem responses to marine climate change. While considerable knowledge gaps remain, this perspective focuses on the progress made, specifically in the field of experimental ocean acidification research.

Most studies addressing acidification, warming, and low oxygen effects in marine organisms have taken an experimental approach (Browman 2016). Its central premise is to trade environmental realism (e.g., CO₂ variability, species interactions) for environmental control (Sommer 2012) to obtain basal information on CO₂-sensitive vs. CO₂-tolerant traits, life-stages, and species and to elucidate mechanisms that cause observed effects (Boyd et al. 2018; Snyder et al. 2018). Experiments are therefore critical stepping-stones on the way to higher order questions such as community effects and adaptive evolution (Pfister et al. 2014; Sunday et al. 2014; Gaylord et al. 2015). The considerable progress made by experiments is evident from the large number of recent reviews and meta-analyses, e.g., on multistressor experiments (Byrne and Przeslawski 2013; Przeslawski et al. 2015; Boyd et al. 2018), acidification effects on fish (Ashur et al. 2017; Esbaugh 2017; Cattano et al. 2018), CO₂ × temperature interactions (Harvey et al. 2013; Kroeker et al. 2013; Lefevre 2016), and CO₂ × O₂ effects (Gobler and Baumann 2016). Thus, past decades have amassed empirical datasets that can now be mined for overarching patterns, however incomplete yet. Importantly, this work in progress needs to be sustained, based on the lessons learned.

I suggest that transformative progress over the past decade has occurred in four key areas (Fig.1). A first area contains methodological advances that include standardization and identification of confounding factors. A second area encompasses advances in understanding physiological and molecular mechanisms of CO₂ sensitivity in marine organisms. The third area is comprised by the recognition that short-term variability in environmental CO₂ conditions likely modifies species sensitivities to acidification. This involves intriguing plasticity in responses to highly dynamic environments and the general expectation of changing vulnerabilities across spatial CO₂ variability gradients (*Ocean Variability Hypothesis*). The fourth area contains the broad consensus on marine

climate change as a multistressor problem and the progress in assessing stressor interactions.

(Key area 1) Methodological advances

CO₂ manipulation & reporting. To ensure comparability of studies, experimental researchers have strived to standardize the terminology and practices of CO₂ manipulation, measurement, and reporting, while further improving husbandry methods of model organisms (e.g., Gattuso and Hansson 2011; Riebesell et al. 2011). It is now standard that experiments manipulate seawater pH only by CO₂ dissolution (not acid) and that the full experimental water chemistry is assessed, calculated (e.g., via CO2SYS) and reported in the primary literature. These developments have been aided by parallel refinements of carbon chemistry kinetics (Dickson and Millero 1987) particularly for brackish water systems (Millero 2010), and recent inter-laboratory calibration studies for measurements of alkalinity and pCO₂ (Bockmon and Dickson 2015). Better consensus also emerged on terminology, e.g., by more clearly distinguishing the long-term process of ocean acidification from experimental assessments of species sensitivities to elevated CO₂ conditions (Browman 2016; McElhany 2017). Regarding CO₂ levels, there is now principal agreement that treatments should reflect plausible future levels in aquatic habitats. For example, aquaculture experiments over the past 50 years often tested pCO₂ levels of 10,000 – 50,000 µatm on juvenile and adult fishes (Ishimatsu et al. 2008), which is unrealistic in the context of ocean acidification. Rather, pCO₂ levels of ~ 2,000 µatm have emerged as a common upper benchmark, based largely on Caldeira and Wickett's (2003) seminal estimate of worst-case pCO₂ concentrations in the average surface ocean by the year 2300 (Riebesell et al. 2011). End-of-this-century treatments of ~ 1,000 µatm are also commonly employed. However, one issue with such open ocean levels is that most marine organisms do not occur there; instead, they mostly inhabit coastal shelf and nearshore environments (Costanza et al. 1997; Hendriks et al. 2010) with profoundly different CO₂ baselines and fluctuations (key area 3, Vargas et al. 2017). Due to the declining buffering capacity of ocean (Revelle factor), these fluctuations will further intensify and thus expose many organisms to critical CO₂ levels earlier than previously

predicted (Shaw et al. 2013; McNeil and Sasse 2016). Hence, there is now consensus that upper realistic pCO₂ levels in experiments need to be flexible and cognizant of the differing acidification trajectories in marine habitats (Salisbury and Jönsson ; Busch et al. 2015; Vargas et al. 2017).

Factors confounding CO₂ experiments. Experimenters have also identified factors that can bias CO₂ sensitivity assessments in marine organisms. Adverse CO₂ effects on coral calcification, for example, appear to depend on experimental light levels that thus need to be controlled and reported (Suggett et al. 2013). Food availability has been shown to affect CO₂ sensitivities across taxa including pteropods (Seibel et al. 2012), bivalves (Thomsen et al. 2013), and fish (Gobler et al. 2018; Stiasny et al. 2018). In the latter, high CO₂ is often believed to increase metabolic costs of acid-base regulation and therefore to reduce growth (key area 2, Heuer and Grosell 2016), but firm empirical support for this hypothesis is lacking (Esbaugh 2017; Hurst et al. 2017; Cattano et al. 2018). It is possible that these costs are too small to be detectable in short-term exposures (Baumann et al. 2018a). Another reason could be that fish larvae are often fed in excess (*ad libitum*), which reduces their overall mortality but also allows survivors to compensate for increased metabolic costs by increased food consumption. Hence controlling or accounting for experimental feeding conditions seems critical for comparable estimates of CO₂ effects. Inadvertent buildup of nitrogenous waste in rearing containers can also confound estimates, because ammonia toxicity to fish and other organisms generally increases with pH (Randall and Tsui 2002), thus affecting predominantly the control over high CO₂ (low pH) treatments. Largely unresolved, however, are frequent concerns about pseudoreplication in CO₂ exposure experiments (e.g., shared water sources, enclosures, founding populations). How exactly this issue confounds empirical estimates remains poorly quantified, and there is currently no clear consensus on acceptable compromises between the most rigorous experimental designs and those that are logistically feasible.

Experiments across multiple life stages and generations. One of the most robust patterns across taxa is the general decrease in CO₂ sensitivity from early life to juvenile and adult ontogenetic stages (Harvey et al. 2013; Przeslawski et al. 2015; Cattano et al. 2018). This may explain why a majority of published experiments have focused on short-term exposures and the most vulnerable early life stages (embryos, larvae). However,

evolving CO₂ rearing systems and better husbandry methods now also allow longer-term experiments that span multiple life stages, whole life cycles, and even generations. While these approaches are still scarce, logistically challenging, and feasible only for select model species, they have clearly highlighted the potential for carry-over effects and transgenerational acclimation, both of which alter our overall assessment of species CO₂ sensitivities. For example, work on the Atlantic silverside (*Menidia menidia* Linnaeus, 1766), an important nearshore forage fish along the North-American Atlantic coast, demonstrated that high CO₂ environments can indeed incur growth costs (Fig.2), but this effect was only detectable after rearing large numbers of individuals from fertilization to late juvenile or adult stages under non-excess food conditions (Murray et al. 2017).

Conversely, when CO₂ experiments have included parental generations, offspring CO₂ tolerance frequently increased (Munday 2014); a form of transgenerational plasticity (TGP) via maternal provisioning or epigenetic modifications that is surprisingly well documented for many stressors and taxa (Jablonka and Raz 2009; Salinas and Munch 2012; Salinas et al. 2013; Munday 2014). TGP may allow species in highly dynamic coastal environments to precondition their offspring to seasonally changing CO₂ conditions (key area 3, Murray et al. 2014). Moreover, it may provide a short-term refuge to the increasing stressors of marine climate change. Unfortunately, the ubiquity of TGP may also comprise yet another confounding factor for experiments, which by definition seek to assess CO₂ sensitivities of species as they currently exist in the wild. But when experimental organisms are obtained from commercial or laboratory brood stocks (Baumann et al. 2012; Chambers et al. 2014; Bromhead et al. 2015; Stiasny et al. 2016) they are likely genetically less diverse, and the parental generation at least partially experienced artificial temperature, CO₂, light, and feeding conditions (Baumann et al. 2018b). Via TGP, these parental conditions influence offspring CO₂ sensitivities, and this additional uncertainty needs to be more clearly acknowledged and then accounted for in scaled up models.

(Key area 2) Mechanisms underlying CO₂ effects in marine organisms

The question what exactly high pCO₂ exposure does to marine organisms is central to experiments, as only a firm mechanistic understanding allows correct interpretations of observed CO₂ effects, improves modeling approaches, or reveals mitigating pathways such as acclimation or evolutionary adaptation. Physiologists have long understood that marine metazoans – unlike single cell organisms – have extracellular fluids that they control to ensure the proper functioning of pH sensitive, metabolic enzymes (Fabry et al. 2008; Pörtner 2008). Regulatory capacity appears to increase from sessile to more mobile taxa (Melzner et al. 2009) and generally involves either responses to accelerate the off-gassing of excessive CO₂ (e.g., by increased ventilation) or responses to increase the buffering capacity of extracellular fluids (e.g., by increasing bicarbonate concentrations) (Esbaugh 2017). As active swimmers, juvenile and adult fishes are particularly effective at both CO₂ and acid/base regulation, so that even a doubling of environmental pCO₂ may only lead to a 10% pCO₂ increase in their blood (Melzner et al. 2009; Heuer and Grosell 2014, 2016). This ability is likely further enhanced in species occurring in highly CO₂ variable habitats (key area 3) and explains why most juvenile and adult fishes tolerate even extreme CO₂ levels on time scales of weeks to months (Ishimatsu et al. 2004), whereas early life stages that still develop acid-base competency may not. It also explains, why most observed CO₂ effects in fish involve sub-lethal, downstream responses related to physiological CO₂ or pH compensation (e.g., behavior, metabolism, growth, aerobic scope, overcalcification), not direct acidosis. For example, the consistently observed overcalcification of fish otoliths under high CO₂ conditions likely results from increased CO₃²⁻ concentrations in the endolymph fluid of the inner ear caused by increased bicarbonate (HCO₃⁻) pH buffering of the blood (Checkley et al. 2009; Hurst et al. 2012; Bignami et al. 2013; Schade et al. 2014).

Our mechanistic understanding of behavioral impairments in fish larvae has arguably progressed the most, thanks to sustained empirical assessments combined with neuro-physiological approaches (reviewed by Ashur et al. 2017). These revealed that impairments of chemical, auditory, and visual senses in fish can already manifest in response to small increases in ambient pCO₂ (from 400 to 700-800µatm) and often result in a large suite of maladaptive behaviors such as the inability to distinguish chemical cues

from predators or preferable settlement habitats (Munday et al. 2009; Munday et al. 2010), to hear nearby reefs (Simpson et al. 2011; Castro et al. 2017) or visually assess predation risk (Ferrari et al. 2012a). Increased anxiety (Hamilton et al. 2014), lateralization (Domenici et al. 2012) and reduced learning ability have also been observed (Ferrari et al. 2012b; Jutfelt et al. 2013). The underlying mechanism for these impairments is likely a malfunction of GABA_A, the primary inhibitory neurotransmitter receptor in vertebrate brains (Ashur et al. 2017). Its crucial, inhibitory function results from membrane hyperpolarization; however, when fish increase blood HCO₃⁻ to buffer against decreasing pH due to elevated pCO₂, the ionic shift results in depolarizing rather than hyperpolarizing neuron membranes when GABA binds to GABA_A, causing maladaptive behavior (Nilsson et al. 2012). However, alternative mechanisms almost certainly exist. Recently, behavioral abnormalities of European seabass (*Dicentrarchus labrax* Linnaeus, 1758) were shown to result from general impairments of the peripheral olfactory system after high CO₂ exposure, not GABA_A inhibition in the brain (Porteus et al. 2018). In both cases, the deleterious behavior appears to manifest only after prolonged, constant CO₂ exposure (>24h), whereas under more realistic diel CO₂ fluctuations most effects are mitigated (Jarrold et al. 2017). As with all other traits studied thus far, behavioral CO₂ sensitivities, too, vary even between taxonomically and ecologically similar fish species (Ferrari et al. 2011), and neutral responses are likely more common than currently acknowledged (Jutfelt and Hedgärde 2013).

As an alternative to species-by-species assessments of response traits, physiologists have suggested that experiments test bioenergetic frameworks (Pörtner 2012; Sokolova 2013), because they focus on e.g., aerobic scope as the affected, fundamental trait (i.e., a proxy for the *surplus* energy available for growth, reproduction, predator avoidance etc.). For example, the oxygen- and capacity-limited thermal tolerance (OCLTT) hypothesis stipulates that species- and life stage-specific thermal windows constrain an organism's capacity for aerobic metabolism (Pörtner 2010; Pörtner et al. 2017). In pH-conformers (some invertebrates), high CO₂ levels may further reduce these thermal windows by directly depressing the oxygen content of blood and other body fluids or by decreasing the functional capacity of pH-sensitive proteins (e.g., hemoglobin and hemocyanin). PH-regulators (e.g., fish), on the other hand, likely respond by increasing ventilation rates,

acid/base regulation, or activate mechanisms for protection and damage repair, which elevates basal metabolic rate and shrinks the aerobic scope. While intuitive, the OCLTT hypothesis is not universally supported by all available empirical evidence (Jutfelt et al. 2018), given that many ectotherms appear to lack clear thermal optima for aerobic scope (Lefevre 2016), or thermal optima are inconsistent with temperatures experienced in the wild and often fail to align with other thermal reaction norms, e.g., in growth and reproduction (Gräns et al. 2014; Norin et al. 2014). Hence, alternative physiological frameworks that focus instead on food assimilation and/or stress-induced impacts on the aerobic production of ATP (Sokolova 2013) and protein synthesis (Frieder et al. 2018) are worth exploring more broadly.

Finally, experiments have already begun to make progress in addressing questions beyond acute species CO₂ sensitivities, such as assessing *de novo* evolution in single-cell organisms (Lohbeck et al. 2012) or quantifying genotypic differences in longer-lived metazoans that will determine the evolutionary potential of these organisms. For example, an experiment on purple sea urchin larvae (*Strongylocentrotus purpuratus* Stimpson, 1857) demonstrated that high CO₂ conditions lead to allelic shifts in the surviving larval population (Pespeni et al. 2013). Others have quantified the heritability of traits such as growth or survival under acidified conditions; generally revealing that CO₂ tolerance has a small but significant genetic component and could therefore evolve (Parker et al. 2011; Malvezzi et al. 2015). In addition, progress with respect to transgenerational acclimation via epigenetic mechanisms or maternal provisioning (Snyder et al. 2018) has suggested that short-term pathways exist for many marine organisms to cope with acidification (Munday 2014; Murray et al. 2014; Thor and Dupont 2015). All recent syntheses agree that acclimation and adaptation represent important frontiers in experimental ocean acidification research, with the potential to change our perception of potentially vulnerable vs. resilient species in a high CO₂ ocean (Munday et al. 2013; Sunday et al. 2014; Boyd et al. 2018).

(Key area 3) **Spatial variability in short term CO₂ fluctuations**

Empirical assessments of CO₂ responses are necessarily limited to a very small fraction of existing marine species, e.g., in marine fish, where only 60 of approximately 20,000 species have been studied so far (~0.3%, Cattano et al. 2018). This means that experiments generally have a broader implicit objective, which is to reveal patterns applicable to other taxonomically or ecologically similar species and thus to provide a basis for inferring community and ecosystem responses to ocean acidification. Unfortunately, the empirical evidence so far suggests that CO₂ sensitivities in marine organisms are intriguingly complex, with examples for negative, neutral (likely underreported, Browman 2016), and positive responses to high CO₂ documented for a large number of traits, between ecologically similar or closely related species, populations, and even repeated experiments. For example, direct negative CO₂ effects on fish early life survival have been reported for some but not most studied species (Cattano et al. 2018). Traits such as metabolism, aerobic scope, swimming performance, and growth, which are largely controlled by other abiotic (e.g., temperature, salinity, oxygen) and biotic factors (food, genes) appear to be most heterogeneous in their CO₂ sensitivity (Lefevre 2016; Cattano et al. 2018). Even among the more consistent behavioral and otolith overcalcification responses in fish (Ashur et al. 2017; Cattano et al. 2018), unexplained differences between species exist (Ferrari et al. 2011; Munday et al. 2011; Jutfelt and Hedgärde 2013). While some variability in experimental outcomes may be attributed to methodological uncertainty (key area 1), most of it still deserves exploring beyond broad meta-analytical approaches. An oft-invoked claim is that responses may simply be species-specific, which is unsatisfactory for lacking causation and for its logical but impractical consequence that thousands of additional species would still require testing. Instead, further insights could still come from more explicitly considering species responses against the level of CO₂ variability in their natural environments, a framework related to the more general ‘Climate variability hypothesis’ (e.g., Compton et al. 2007; Markle and Kozak 2018; Ortega-Mayagoitia et al. 2018), but which in the specific context of CO₂ sensitivities I suggest be called “*Ocean Variability Hypothesis*” (OVH, Fig.3).

The OVH is based on the broad recognition that marine habitats differ vastly in the frequency and magnitude of short-term $p\text{CO}_2$ fluctuations (Hofmann et al. 2011; Duarte et al. 2013). These fluctuations can be ephemeral in upwelling regions or in the vicinity of volcanic CO_2 seeps (Hall-Spencer et al. 2008). But for most marine habitats they follow annual, diel, or tidal cycles caused by changes in net ecosystem metabolism (Odum et al. 1995; Gagliano et al. 2010; Baumann and Smith 2018), where net heterotrophy leads to metabolic acidification (pH decrease) and net autotrophy causes the reverse process, basification (pH increase). Importantly, this natural CO_2 variability follows a clear spatial pattern, with generally decreasing amplitudes from nearshore to open ocean environments (Hofmann et al. 2011). In nearshore habitats, where biological productivity is usually highest (Cloern et al. 2016), diel and seasonal CO_2 fluctuations between 200-5,000 μatm (more than one full pH unit in full strength seawater) are common and more than twice the worst-case, open ocean $p\text{CO}_2$ increase over the next three centuries (Caldeira and Wickett 2003). With increasing distance from shore, however, the seasonality and metabolic dominance of the marine carbon system dissipates until approaching near-stable $p\text{CO}_2$ conditions year-round in the open ocean (Hofmann et al. 2011; Duarte et al. 2013). This spatial gradient means, first, that the anthropogenic acidification signal strengthens from largely undetectable in nearshore habitats (Duarte et al. 2013) to clearly discernable over decadal time scales in the open ocean (Doney et al. 2009; Lauvset et al. 2015). It also implies that marine organisms are likely adapted to the level of contemporary CO_2 fluctuations in their natural habitats. Assuming that high CO_2 tolerance comes at a cost to organisms, one should therefore expect CO_2 sensitivities to change from tolerant nearshore to more sensitive open ocean species. Indeed, Vargas et al. (2017) recently showed that CO_2 sensitivities in several Chilean mollusk and copepod species decrease with increasing short-term CO_2 fluctuations in their natural habitat – a first explicit test of the OVH. For fish, this framework has not yet been explicitly tested, although a recent meta-analysis by Cattano et al. (2018) differentiated CO_2 responses between marine fishes from several climatic regions (polar, temperate, subtropical, tropical) or benthic vs. pelagic habitats. While valuable, such contrasts are still likely to group species from habitats with different baselines of CO_2 variability.

CO₂ sensitivities in marine fish illustrate the potential of the Ocean Variability Hypothesis (Fig.3). Early life stages of nearshore species such as sticklebacks (*Gasterosteus aculeatus* Linnaeus 1758, Schade et al. 2014), Atlantic silversides (*Menidia menidia*, Baumann et al. 2018b; Murray and Baumann 2018), red drum (*Sciaenops ocellatus* Linnaeus 1766, Lonthair et al. 2017), or Baltic spring-spawning herring (*Clupea harengus* Linnaeus 1758, Franke and Clemmesen 2011) appear generally tolerant of CO₂ levels of 2,000 µatm and above with no or very small effects on survival or growth. In Atlantic silversides, years of repeated experimentation documented average reductions in embryo survival by only 10% in response to ~2,000 µatm CO₂, and these effects mainly occurred in offspring produced at the beginning or end of the species spawning season (Baumann et al. 2018b). Marine medaka (*Oryzias melastigma* McClelland, 1839) and sheepshead minnow (*Cyprinodon variegatus* Lacepède, 1803) embryos and larvae appear to tolerate even the most extreme CO₂ levels tested so far (Depasquale et al. 2015; Mu et al. 2015). While taxonomically diverse, these examples all represent species that reproduce in very shallow, nearshore waters with likely large short-term CO₂ fluctuations to which they must be adapted.

Slightly greater CO₂ sensitivities, on the other hand, appear to exist for coastal shelf species including Atlantic cod (*Gadus morhua* Linnaeus 1758, Stiasny et al. 2016; Stiasny et al. 2018), summer flounder (*Paralichthys dentatus* Linnaeus 1766, Chambers et al. 2014) or Senegalese sole (*Solea senegalensis* Kaup 1858, Pimentel et al. 2014b), where significant reductions in embryo and larval survival have already been observed in response to 1,000 – 2,000 µatm pCO₂. Coral reef fish may fall under this category, too, given the diminished but still substantial CO₂ variability in their habitats (Gagliano et al. 2010; Hofmann et al. 2011) and the well-documented behavioral responses to CO₂ levels below 1,000 µatm (Ashur et al. 2017). However, direct CO₂-induced survival effects are conspicuously absent in this group (Munday et al. 2011; Welch and Munday 2016), which might be due interactive effects of development time on early life CO₂ sensitivity. In contrast to higher latitude species, most tropical fish hatch within few days post fertilization and become actively swimming larvae within as little as one week. It is possible that offspring in these cases acquire acid-base competency too fast for significant adverse survival effects to accrue. The same cave-at applies to the few tested

open ocean species such as yellowfin tuna (*Thunnus albacares* Bonnaterre 1788, Frommel et al. 2016), dolphinfish (*Coryphaena hippurus* Linnaeus 1758, Pimentel et al. 2014a) or kingfish (*Seriola lalandi* Valenciennes 1833, Munday et al. 2015), which reproduce in relatively CO₂ stable environments and should therefore be most sensitive to high CO₂ conditions. While there is evidence for lethal and sublethal effects of high CO₂ in these pelagic species (but see Watson et al. 2018), all have fast developing early life stages that become active piscivores within days post hatch, which requires well-developed acid-base competency (Melzner et al. 2009).

Stage duration may therefore comprise a second important dimension of the Ocean Variability Hypothesis (Fig.3). It suggests that the most CO₂ tolerant marine organisms are those that develop fast and/or in habitats with large contemporary CO₂ fluctuations, whereas the potentially most vulnerable species are those that develop slowly in relatively stable open ocean habitats. Unfortunately, at least in marine fish, this latter group remains almost unstudied, but likely includes offshore winter-spawners in temperate or high latitude environments that have extended embryonic and larval durations (Murray et al. in review). In addition, there are other spatial CO₂ variability gradients beyond the general nearshore-offshore axis that appear equally suitable to test the Ocean Variability Hypothesis. Such gradients likely exist with depth, as many marine species exhibit distinct daily or ontogenetic vertical migrations (Godø et al. 2009), or the level of eutrophication in coastal waters. In upwelling regions such as the California Current Large Marine Ecosystem (Kelly et al. 2013) or along the coast of Chile (Vargas et al. 2017) different regions exhibit different levels of ephemeral upwelling and therefore short-term CO₂ variability. Consistent with the hypothesis, a study on purple sea urchin (*S. purpuratus*) found that offspring from high CO₂ variability regions were less CO₂ sensitive than their conspecifics from low CO₂ variability areas (Kelly et al. 2013).

(Key area 4) Marine climate change is a multistressor problem

The concurrent rise in ocean temperature and CO₂ or the intensifying co-occurrence of high CO₂ and low oxygen conditions in urban estuaries or oxygen minimum zones (Cai et

al. 2011; O'Boyle et al. 2013; Breitburg et al. 2015; Baumann and Smith 2018) are just two of many examples that demand assessments of species sensitivities to stressor combinations. This is necessary given the likelihood that combined stressors elicit quantitatively or qualitatively different effects than the sum of each stressor alone (Breitburg et al. 2015). These stressor interactions may be antagonistic and thus mitigate the overall response; however, the potential for synergistic interactions causing disproportionately adverse effects is the most concerning. For example, early experiments on the hard coral *Stylophora pistillata* (Esper 1797) found that calcification rates increased with CO₂ under control temperatures but decreased at combined high CO₂ and high temperature treatments (Reynaud et al. 2003). A recent CO₂ × O₂ factorial experiment on Atlantic silversides (Depasquale et al. 2015) observed no effects of high CO₂ on early life survival at normoxic treatments, but synergistic survival reductions under low oxygen × high CO₂ treatments. Even in elasmobranchs such as juvenile bamboo sharks (*Chiloscyllium punctatum* Müller and Henle 1838), both additive and synergistically negative CO₂ × temperature interactions were evident in traits like survival and ventilation rate (Rosa et al. 2014).

The most extensive empirical evidence exists for CO₂ × temperature effects so far, particularly in mollusks, crustaceans, and echinoderms, while less so for marine fish. A pioneering meta-analysis thus revealed that high CO₂ × temperature conditions appear to have mostly additive effects (no interaction) on early life survival in marine crustaceans, whereas mollusk species are synergistically negatively affected (Harvey et al. 2013). However, a different pattern emerged for another trait, growth, where strong and synergistically negative CO₂ × temperature effects characterized crustaceans, only weakly negative additive effects were found for mollusks, and overall positive effects occurred in echinoderm species. Thus, multistressor responses vary across taxa, stages, and trophic levels, and our current predictive capacity is still in its infancy with respect to where and when they occur. Furthermore, while earlier meta-analyses found synergistic interactions to be less frequent than additive or antagonistic effects (Byrne and Przeslawski 2013; Harvey et al. 2013), a more recent synthesis concluded just the opposite (Przeslawski et al. 2015).

For the combined effects of high CO₂ and low dissolved oxygen (DO), the empirical evidence is much scarcer still and not yet sufficient for conclusive meta-analyses (Gobler and Baumann 2016), despite the universal recognition that these two stressors are tightly coupled in most marine habitats on short and long time scales (Baumann and Smith 2018). An initial review concluded that, individually, low oxygen conditions are more stressful to marine organisms than high CO₂. In combination, the two stressors appear to cause mostly additive negative effects; however, every study to date also found synergistic negative interactions in a least one examined trait (Gobler et al. 2014; Gobler and Baumann 2016; Miller et al. 2016). There is a clear and well-understood need to expand multiple stressor experiments in this area, including the important addition of testing fluctuating CO₂ × O₂ environments as they are most common in nature and may mitigate or further exacerbate effects.

Importantly, designing multiple stressor experiments also comes with new and complex challenges, which have recently been comprehensively reviewed by Boyd et al. (2018). The authors find that modern empirical assessments have begun to diverge in scope in two complementary directions. The first group comprises reductionist approaches that gain mechanistic insights into the nature of stressor interactions via full-factorial designs in well-controlled laboratory settings. This comes, however, at the expense of environmental realism both in terms of near-future projections and the lack indirect effects that will ultimately determine ecosystem responses and thus guide policy decisions. Furthermore, factorial designs become rapidly unwieldy if they try to include more than two stressors and more than two exposure levels at the time. Beyond the obvious logistical challenges, difficulties in interpretation and presentation of multistressor observations increase exponentially with design complexity, while making spurious patterns more likely. Suggested solutions include sequential testing and then combining of multiple stressors (Boyd et al. 2018) as well as repeating experiments themselves to increase statistical power (Baumann et al. 2018b; Murray and Baumann 2018). A second group of multistressor experiments encompasses mesocosms and field manipulations (Sswat et al. 2018), which trade some environmental control for greater realism to measure overall community effects in response to projections of ocean change (Alsterberg et al. 2013), while consciously ceding the ability to distinguish between the

exact mechanisms behind observed effects. These approaches are often even more logistically challenging and thus allow fewer replicates and reduced statistical power. However, both groups of multistressor approaches are important to sustain (Boyd et al. 2018).

Going forward, refined experiments will continue to contribute important data toward anticipating the impacts of marine climate change, particularly when they include multiple life stages, examine potential carryover effects and are focused on traits of known fitness-relevance.

References

- Alsterberg, C., Eklöf, J.S., Gamfeldt, L., Havenhand, J.N., and Sundbäck, K. 2013. Consumers mediate the effects of experimental ocean acidification and warming on primary producers. *Proc. Natl. Acad. Sci. U.S.A.* **110** (21): 8603-8608. doi: 10.1073/pnas.1303797110
- Ashur, M.M., Johnston, N.K., and Dixson, D.L. 2017. Impacts of ocean acidification on sensory function in marine organisms. *Integr. Comp. Biol.* **57**(1): 63-80.
- Baumann, H. 2016. Combined effects of ocean acidification, warming, and hypoxia on marine organisms. *Limnol. Oceanogr. e-lectures*. doi:10.1002/loe2.10002.
- Baumann, H., and Smith, E.M. 2018. Quantifying metabolically driven pH and oxygen fluctuations in US nearshore habitats at diel to interannual time scales. *Estuar. Coasts* **41**: 1102-1117.
- Baumann, H., Talmage, S.C., and Gobler, C.J. 2012. Reduced early life growth and survival in a fish in direct response to increased carbon dioxide [Letter]. *Nat. Clim. Change* **2**: 38-41. doi:10.1038/nclimate1291.
- Baumann, H., Parks, E.M., and Murray, C.S. 2018a. Starvation rates in larval and juvenile Atlantic silversides (*Menidia menidia*) are unaffected by high CO₂ conditions. *Mar. Biol.* **165**(4): 75. doi:10.1007/s00227-018-3335-x.
- Baumann, H., Cross, E.L., and Murray, C.S. 2018b. Robust quantification of fish early life CO₂ sensitivities via serial experimentation. *Biol. Lett.* **14**: 20180408. doi:10.1098/rsbl.2018.0408.
- Bignami, S., Enochs, I.C., Manzello, D.P., Sponaugle, S., and Cowen, R.K. 2013. Ocean acidification alters the otoliths of a pantropical fish species with implications for sensory function. *Proc. Natl. Acad. Sci. U.S.A.* **110**(18): 7366-7370. doi:10.1073/pnas.1301365110.
- Bockmon, E.E., and Dickson, A.G. 2015. An inter-laboratory comparison assessing the quality of seawater carbon dioxide measurements. *Mar. Chem.* **171**: 36-43.
- Boyd, P.W., Collins, S., Dupont, S., Fabricius, K., Gattuso, J.P., Havenhand, J., Hutchins, D.A., Riebesell, U., Rintoul, M.S., Vichi, M., Biswas, H., Ciotti, A., Gao, K., Gehlen, M., Hurd, C.L., Kurihara, H., McGraw, C.M., Navarro, J.M., Nilsson, G.E., Passow, U., and Pörtner, H.O. 2018. Experimental strategies to assess the

- biological ramifications of multiple drivers of global ocean change—A review. *Glob. Change Biol.* **24**: 2239-2261. doi:10.1111/gcb.14102.
- Breitbart, D., Levin, L.A., Oschlies, A., Grégoire, M., Chavez, F.P., Conley, D.J., Garçon, V., Gilbert, D., Gutiérrez, D., and Isensee, K. 2018. Declining oxygen in the global ocean and coastal waters. *Science*, **359**(6371): eaam7240.
- Breitbart, D.L., Salisbury, J., Bernhard, J.M., Cai, W.-J., Dupont, S., Doney, S.C., Kroeker, K.J., Levin, L., Long, W.C., Milke, L.M., Miller, S.H., Phelan, B., Passow, U., Seibel, B.A., Todgham, A.E., and Tarrant, A.M. 2015. And on top of all that... Coping with ocean acidification in the midst of many stressors. *Oceanography*, **28**: 48-61.
- Bromhead, D., Scholey, V., Nicol, S., Margulies, D., Wexler, J., Stein, M., Hoyle, S., Lennert-Cody, C., Williamson, J., Havenhand, J., Ilyina, T., and Lehodey, P. 2015. The potential impact of ocean acidification upon eggs and larvae of yellowfin tuna (*Thunnus albacares*). *Deep Sea Res. Part II: Top. Stud. Oceanogr.* **113**: 268–279. doi:10.1016/j.dsr2.2014.03.019.
- Browman, H.I. 2016. Applying organized scepticism to ocean acidification research. *ICES J. Mar. Sci.* **73**(3): 529-536. doi:10.1093/icesjms/fsw010.
- Busch, D.S., O'Donnell, M.J., Hauri, C., Mach, K.J., Poach, M., Doney, S.C., and Signorini, S.R. 2015. Understanding, characterizing, and communicating responses to ocean acidification: Challenges and uncertainties. *Oceanography*, **28**: 30-39.
- Byrne, M., and Przeslawski, R. 2013. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integr. Comp. Biol.* **53**(4): 582-596. doi:10.1093/icb/ict049.
- Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M.C., Lehrter, J.C., Lohrenz, S.E., Chou, W.-C., Zhai, W., Hollibaugh, J.T., Wang, Y., Zhao, P., Guo, X., Gundersen, K., Dai, M., and Gong, G.-C. 2011. Acidification of subsurface coastal waters enhanced by eutrophication [10.1038/ngeo1297]. *Nat. Geosci.* **4**(11): 766-770. doi:10.1038/ngeo1297.
- Caldeira, K., and Wickett, M.E. 2003. Anthropogenic carbon and ocean pH. *Nature*, **425**(6956): 365-365. doi:10.1038/425365a
- Castro, J.M., Amorim, M.C.P., Oliveira, A.P., Gonçalves, E.J., Munday, P.L., Simpson, S.D., and Faria, A.M. 2017. Painted goby larvae under high-CO₂ fail to recognize reef sounds. *PLoS One*, **12**(1): e0170838.
- Cattano, C., Claudet, J., Domenici, P., and Milazzo, M. 2018. Living in a high CO₂ world: a global meta - analysis shows multiple trait - mediated fish responses to ocean acidification. *Ecol. Monogr.* **88**: 320-335. doi:10.1002/ecm.1297.
- Chambers, R.C., Candelmo, A.C., Habeck, E.A., Poach, M.E., Wicczorek, D., Cooper, K.R., Greenfield, C.E., and Phelan, B.A. 2014. Effects of elevated CO₂ in the early life stages of summer flounder, *Paralichthys dentatus*, and potential consequences of ocean acidification. *Biogeosciences*, **11**: 1613-1626. doi:10.5194/bg-10-13897-2013.
- Checkley, D.M., Dickson, A.G., Takahashi, M., Radich, J.A., Eisenkolb, N., and Asch, R. 2009. Elevated CO₂ enhances otolith growth in young fish. *Science*, **324**(5935): 1683-1683. doi:10.1126/science.1169806.

- Cloern, J.E., Abreu, P.C., Carstensen, J., Chauvaud, L., Elmgren, R., Grall, J., Greening, H., Johansson, J.O.R., Kahru, M., and Sherwood, E.T. 2016. Human activities and climate variability drive fast - paced change across the world's estuarine-coastal ecosystems. *Glob. Change Biol.* **22**: 513-529.
- Compton, T.J., Rijkenberg, M.J., Drent, J., and Piersma, T. 2007. Thermal tolerance ranges and climate variability: a comparison between bivalves from differing climates. *J. Exp. Mar. Biol. Ecol.* **352**(1): 200-211.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., and van den Belt, M. 1997. The value of the world's ecosystem services and natural capital [10.1038/387253a0]. *Nature*, **387**: 253-260. doi:10.1038/387253a0.
- Depasquale, E., Baumann, H., and Gobler, C.J. 2015. Variation in early life stage vulnerability among Northwest Atlantic estuarine forage fish to ocean acidification and low oxygen. *Mar. Ecol. Prog. Ser.* **523**: 145-156.
- Diaz, R.J., and Rosenberg, R. 2008. Spreading dead zones and consequences for marine ecosystems. *Science* **321**(5891): 926-929. doi:10.1126/science.1156401.
- Dickson, A., and Millero, F. 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Res. Part 1 Oceanogr. Res. Pap.* **34**(10): 1733-1743.
- Domenici, P., Allan, B., McCormick, M.I., and Munday, P.L. 2012. Elevated carbon dioxide affects behavioural lateralization in a coral reef fish. *Biol. Lett.* **8**(1): 78-81.
- Doney, S.C., Fabry, V.J., R.A., F., and Kleypas, J.A. 2009. Ocean acidification: the other CO₂ problem. *Annu. Rev. Mar. Sci.* **1**: 169-192.
- Duarte, C.M., Hendriks, I.E., Moore, T.S., Olsen, Y.S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J.A., and McCulloch, M. 2013. Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH. *Estuar. Coasts*, **36**(2): 221-236. doi:10.1007/s12237-013-9594-3.
- Esbaugh, A.J. 2017. Physiological implications of ocean acidification for marine fish: emerging patterns and new insights. *J. Comp. Physiol. B* **188**: 1-13.
- Fabry, V.J., Seibel, B.A., Feely, R.A., and Orr, J.C. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* **65**(3): 414-432. doi:10.1093/icesjms/fsn048.
- Ferrari, M.C., Dixon, D.L., Munday, P.L., McCormick, M.I., Meekan, M.G., Sih, A., and Chivers, D.P. 2011. Intrageneric variation in antipredator responses of coral reef fishes affected by ocean acidification: implications for climate change projections on marine communities. *Glob. Change Biol.* **17**(9): 2980-2986.
- Ferrari, M.C., McCormick, M.I., Munday, P.L., Meekan, M.G., Dixon, D.L., Lönnstedt, O., and Chivers, D.P. 2012a. Effects of ocean acidification on visual risk assessment in coral reef fishes. *Funct. Ecol.* **26**(3): 553-558.
- Ferrari, M.C., Manassa, R.P., Dixon, D.L., Munday, P.L., McCormick, M.I., Meekan, M.G., Sih, A., and Chivers, D.P. 2012b. Effects of ocean acidification on learning in coral reef fishes. *PLoS One*, **7**(2): e31478.
- Fine, M., and Tchernov, D. 2007. Scleractinian coral species survive and recover from decalcification. *Science*, **315**(5820): 1811. doi:10.1126/science.1137094.

- Franke, A., and Clemmesen, C. 2011. Effect of ocean acidification on early life stages of Atlantic herring (*Clupea harengus* L.). *Biogeosciences*, **8**: 3697-3707. doi:10.5194/bg-8-3697-2011.
- Frieder, C.A., Applebaum, S.L., Pan, T.-C.F., and Manahan, D.T. 2018. Shifting balance of protein synthesis and degradation sets a threshold for larval growth under environmental stress. *Biol. Bull.* **234**(1): 45-57. doi:10.1086/696830.
- Frommel, A.Y., Margulies, D., Wexler, J.B., Stein, M.S., Scholey, V.P., Williamson, J.E., Bromhead, D., Nicol, S., and Havenhand, J. 2016. Ocean acidification has lethal and sub-lethal effects on larval development of yellowfin tuna, *Thunnus albacares*. *J. Exp. Mar. Biol. Ecol.* **482**: 18-24. doi:10.1016/j.jembe.2016.04.008.
- Gagliano, M., McCormick, M.I., Moore, J.A., and Depczynski, M. 2010. The basics of acidification: baseline variability of pH on Australian coral reefs. *Mar. Biol.* **157**(8): 1849-1856.
- Gattuso, J.-P., and Hansson, L. 2011. Ocean acidification. Oxford University Press.
- Gaylord, B., Kroeker, K.J., Sunday, J.M., Anderson, K.M., Barry, J.P., Brown, N.E., Connell, S.D., Dupont, S., Fabricius, K.E., Hall-Spencer, J.M., Klinger, T., Milazzo, M., Munday, P.L., Russell, B.D., Sanford, E., Schreiber, S.J., Thiagarajan, V., Vaughan, M.L.H., Widdicombe, S., and Harley, C.D.G. 2015. Ocean acidification through the lens of ecological theory. *Ecology*, **96**(1): 3-15. doi:10.1890/14-0802.1.
- Gobler, C.J., and Baumann, H. 2016. Hypoxia and acidification in marine ecosystems: Coupled dynamics and effects on ocean life. *Biol. Lett.* **12**: 20150976.
- Gobler, C.J., Depasquale, E., Griffith, A., and Baumann, H. 2014. Hypoxia and acidification have additive and synergistic negative effects on the growth, survival, and metamorphosis of early life stage bivalves. *PLoS ONE*, **9**: e83648.
- Gobler, C.J., Merlo, L.R., Morrell, B.K., and Griffith, A.W. 2018. Temperature, acidification, and food supply interact to negatively affect the growth and survival of the forage fish, *Menidia beryllina* (Inland silverside), and *Cyprinodon variegatus* (Sheepshead minnow) [Original Research]. *Front. Mar. Sci.* **5**(86). doi:10.3389/fmars.2018.00086.
- Godø, O.R., Patel, R., and Pedersen, G. 2009. Diel migration and swimbladder resonance of small fish: some implications for analyses of multifrequency echo data. *ICES J. Mar. Sci.* **66**(6): 1143-1148.
- Gräns, A., Jutfelt, F., Sandblom, E., Jönsson, E., Wiklander, K., Seth, H., Olsson, C., Dupont, S., Ortega-Martinez, O., Einarsdottir, I., and Björnsson, B.T. 2014. Aerobic scope fails to explain the detrimental effects on growth resulting from warming and elevated CO₂ in Atlantic halibut. *J. Exp. Biol.* **217**(5): 711-717.
- Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S.M., Rowley, S.J., Tedesco, D., and Buia, M.-C. 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification [10.1038/nature07051]. *Nature*, **454**(7200): 96-99. doi:10.1038/nature07051.
- Hamilton, T.J., Holcombe, A., and Tresguerres, M. 2014. CO₂-induced ocean acidification increases anxiety in Rockfish via alteration of GABA_A receptor functioning. *Proc. R. Soc. Lond. B Biol. Sci.* **281**(1775): 20132509.

- Harvey, B.P., Gwynn-Jones, D., and Moore, P.J. 2013. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecol. Evol.* **3**(4): 1016-1030. doi:10.1002/ece3.516.
- Hendriks, I.E., Duarte, C.M., and Álvarez, M. 2010. Vulnerability of marine biodiversity to ocean acidification: A meta-analysis. *Estuar. Coast. Shelf Sci.* **86**(2): 157-164. doi:10.1016/j.ecss.2009.11.022.
- Heuer, R.M., and Grosell, M. 2014. Physiological impacts of elevated carbon dioxide and ocean acidification on fish. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **307**(9): R1061-R1084. doi:10.1152/ajpregu.00064.2014.
- Heuer, R.M., and Grosell, M. 2016. Elevated CO₂ increases energetic cost and ion movement in the marine fish intestine. *Sci. Rep.* **6**: 34480. doi:10.1038/srep34480.
- Hofmann, G.E., Smith, J.E., Johnson, K.S., Send, U., Levin, L.A., Micheli, F., Paytan, A., Price, N.N., Peterson, B., Takeshita, Y., Matson, P.G., Crook, E.D., Kroeker, K.J., Gambi, M.C., Rivest, E.B., Frieder, C.A., Yu, P.C., and Martz, T.R. 2011. High-frequency dynamics of ocean pH: A multi-ecosystem comparison. *PLoS ONE*, **6**(12): e28983. doi:10.1371/journal.pone.0028983.
- Hurst, T.P., Laurel, B.J., Hanneman, E., Haines, S.A., and Ottmar, M.L. 2017. Elevated CO₂ does not exacerbate nutritional stress in larvae of a Pacific flatfish. *Fish. Oceanogr.* **26**(3): 336-349.
- Hurst, T.P., Fernandez, E.R., Mathis, J.T., Miller, J.A., Stinson, C.M., and Ahgeak, E.F. 2012. Resiliency of juvenile walleye pollock to projected levels of ocean acidification. *Aquat. Biol.* **17**: 247-259.
- IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, (eds.)]. IPCC, Geneva, Switzerland, 151 pp. IPCC, Geneva, Switzerland.
- Ishimatsu, A., Hayashi, M., and Kikkawa, T. 2008. Fishes in high-CO₂, acidified oceans. *Mar. Ecol. Prog. Ser.* **373**: 295-302. doi:10.3354/meps07823.
- Ishimatsu, A., Kikkawa, T., Hayashi, M., Lee, K.-S., and Kita, J. 2004. Effects of CO₂ on marine fish: larvae and adults. *J. Oceanogr.* **60**(4): 731-741. doi:10.1007/s10872-004-5765-y.
- Jablonka, E., and Raz, G. 2009. Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. *Q. Rev. Biol.* **84**(2): 131-176. doi:10.1086/598822.
- Jarrold, M.D., Humphrey, C., McCormick, M.I., and Munday, P.L. 2017. Diel CO₂ cycles reduce severity of behavioural abnormalities in coral reef fish under ocean acidification. *Sci. Rep.* **7**(1): 10153.
- Jutfelt, F., and Hedgärde, M. 2013. Atlantic cod actively avoid CO₂ and predator odour, even after long-term CO₂ exposure. *Front. Zool.* **10**(1): 81.
- Jutfelt, F., de Souza, K.B., Vuylsteke, A., and Sturve, J. 2013. Behavioural disturbances in a temperate fish exposed to sustained high-CO₂ levels. *PLoS One*, **8**(6): e65825.
- Jutfelt, F., Norin, T., Ern, R., Overgaard, J., Wang, T., McKenzie, D.J., Lefevre, S., Nilsson, G.E., Metcalfe, N.B., Hickey, A.J.R., Brijs, J., Speers-Roesch, B., Roche, D.G., Gamperl, A.K., Raby, G.D., Morgan, R., Esbaugh, A.J., Gräns, A.,

- Axelsson, M., Ekström, A., Sandblom, E., Binning, S.A., Hicks, J.W., Seebacher, F., Jørgensen, C., Killen, S.S., Schulte, P.M., and Clark, T.D. 2018. Oxygen- and capacity-limited thermal tolerance: blurring ecology and physiology [10.1242/jeb.169615]. *J. Exp. Biol.* **221**(1). doi:10.1242/jeb.169615.
- Kelly, M.W., Padilla-Gamiño, J.L., and Hofmann, G.E. 2013. Natural variation and the capacity to adapt to ocean acidification in the keystone sea urchin *Strongylocentrotus purpuratus*. *Glob. Change Biol.* **19**(8): 2536-2546. doi:10.1111/gcb.12251.
- Kroeker, K.J., Kordas, R.L., Crim, R., Hendriks, I.E., Ramajo, L., Singh, G.S., Duarte, C.M., and Gattuso, J.P. 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Change Biol.* **19**(6): 1884-1896.
- Lauvset, S., Gruber, N., Landschützer, P., Olsen, A., and Tjiputra, J. 2015. Trends and drivers in global surface ocean pH over the past 3 decades. *Biogeosciences*, **12**(5): 1285.
- Lefevre, S. 2016. Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO₂ and their interaction. *Conserv. Physiol.* **4**(1): cow009.
- Levin, L.A., and Breitburg, D.L. 2015. Linking coasts and seas to address ocean deoxygenation. *Nat. Clim. Change*, **5**(5): 401-403.
- Lohbeck, K.T., Riebesell, U., and Reusch, T.B.H. 2012. Adaptive evolution of a key phytoplankton species to ocean acidification [10.1038/ngeo1441]. *Nat. Geosci.* **5**: 346-351. doi:10.1038/ngeo1441.
- Lonthair, J., Ern, R., and Esbaugh, A.J. 2017. The early life stages of an estuarine fish, the red drum (*Sciaenops ocellatus*), are tolerant to high pCO₂. *ICES J. Mar. Sci.* **74**(4): 1042-1050.
- Malvezzi, A., Murray, C.S., Feldheim, K.A., Dibattista, J.D., Garant, D., Gobler, C.J., Chapman, D.D., and Baumann, H. 2015. A quantitative genetic approach to assess the evolutionary potential of a coastal marine fish to ocean acidification. *Evol. Appl.* **8**: 352-362. doi:10.1111/eva.12248.
- Markle, T.M., and Kozak, K.H. 2018. Low acclimation capacity of narrow - ranging thermal specialists exposes susceptibility to global climate change. *Ecol. Evol.* **8**(9): 4644-4656.
- McElhany, P. 2017. CO₂ sensitivity experiments are not sufficient to show an effect of ocean acidification. *ICES J. Mar. Sci.* **74**(4): 926-928.
- McNeil, B.I., and Sasse, T.P. 2016. Future ocean hypercapnia driven by anthropogenic amplification of the natural CO₂ cycle. *Nature*, **529**(7586): 383-387.
- Melzner, F., Gutowska, M.A., Langenbuch, M., Dupont, S., Lucassen, M., Thorndyke, M.C., Bleich, M., and Pörtner, H.O. 2009. Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences*, **6**(10): 2313-2331. doi:10.5194/bg-6-2313-2009.
- Miller, S.H., Breitburg, D.L., Burrell, R.B., and Keppel, A.G. 2016. Acidification increases sensitivity to hypoxia in important forage fishes. *Mar. Ecol. Prog. Ser.* **549**: 1-8.
- Millero, F.J. 2010. Carbonate constants for estuarine waters. *Mar. Freshw. Res.* **61**(2): 139-142. doi:10.1071/MF09254.

- Mitchell, J.F. 1989. The “greenhouse” effect and climate change. *Rev. Geophys.* **27**(1): 115-139.
- Mu, J., Jin, F., Wang, J., Zheng, N., and Cong, Y. 2015. Effects of CO₂-driven ocean acidification on early life stages of marine medaka (*Oryzias melastigma*). *Biogeosciences*, **12**(12): 3861-3868.
- Munday, P.L. 2014. Transgenerational acclimation of fishes to climate change and ocean acidification. *F1000Prime Rep.* **6**(99). doi:10.12703/P6-99pmid:25580253 doi:10.12703/P6-99.
- Munday, P.L., Gagliano, M., Donelson, J.M., Dixon, D.L., and Thorrold, S.R. 2011. Ocean acidification does not affect the early life history development of a tropical marine fish. *Mar. Ecol. Prog. Ser.* **423**: 211-221.
- Munday, P.L., Warner, R.R., Monro, K., Pandolfi, J.M., and Marshall, D.J. 2013. Predicting evolutionary responses to climate change in the sea. *Ecol. Lett.* **16**(12): 1488-1500. doi:10.1111/ele.12185.
- Munday, P.L., Dixon, D.L., McCormick, M.I., Meekan, M., Ferrari, M.C.O., and Chivers, D.P. 2010. Replenishment of fish populations is threatened by ocean acidification. *Proc. Natl. Acad. Sci. U.S.A.* **107**: 12930-12934. doi:10.1073/pnas.1004519107.
- Munday, P.L., Dixon, D.L., Donelson, J.M., Jones, G.P., Pratchett, M.S., Devitsina, G.V., and Døving, K.B. 2009. Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proc. Natl. Acad. Sci. U.S.A.* **106**(6): 1848-1852. doi:10.1073/pnas.0809996106.
- Munday, P.L., Watson, S.-A., Parsons, D.M., King, A., Barr, N.G., Mcleod, I.M., Allan, B.J.M., and Pether, S.M.J. 2015. Effects of elevated CO₂ on early life history development of the yellowtail kingfish, *Seriola lalandi*, a large pelagic fish. *ICES J. Mar. Sci.* **73**: 641-649. doi:10.1093/icesjms/fsv210.
- Murray, C.S., and Baumann, H. 2018. You better repeat it: complex temperature × CO₂ effects in Atlantic silverside offspring revealed by serial experimentation. *Diversity*, **10**(69): 1-19. doi:10.3390/d10030069.
- Murray, C.S., Fuiman, L., and Baumann, H. 2017. Consequences of elevated CO₂ exposure across multiple life stages in a coastal forage fish. *ICES J. Mar. Sci.* **74**: 1051-1061. doi:10.1093/icesjms/fsw179.
- Murray, C.S., Malvezzi, A.J., Gobler, C.J., and Baumann, H. 2014. Offspring sensitivity to ocean acidification changes seasonally in a coastal marine fish. *Mar. Ecol. Prog. Ser.* **504**: 1-11.
- Nilsson, G.E., Dixon, D.L., Domenici, P., McCormick, M.I., Sorensen, C., Watson, S.-A., and Munday, P.L. 2012. Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function [10.1038/nclimate1352]. *Nat. Clim. Change*, **2**: 201-204. doi:10.1038/nclimate1352.
- Norin, T., Malte, H., and Clark, T.D. 2014. Aerobic scope does not predict the performance of a tropical eurythermal fish at elevated temperatures. *J. Exp. Biol.* **217**(2): 244-251.
- O’Boyle, S., McDermott, G., Noklegaard, T., and Wilkes, R. 2013. A simple index of trophic status in estuaries and coastal bays based on measurements of pH and dissolved oxygen. *Estuar. Coasts*, **36**: 158-173. doi:10.1007/s12237-012-9553-4.

- Odum, W.E., Odum, E.P., and Odum, H.T. 1995. Nature's Pulsing Paradigm. *Estuaries*, **18**(4): 547-555. doi:10.2307/1352375.
- Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner, G.-K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D., Totterdell, I.J., Weirig, M.-F., Yamanaka, Y., and Yool, A. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**(7059): 681-686. doi:10.1038/nature04095.
- Ortega-Mayagoitia, E., Hernández-Martínez, O., and Ciro-Pérez, J. 2018. Phenotypic plasticity of life-history traits of a calanoid copepod in a tropical lake: Is the magnitude of thermal plasticity related to thermal variability? *PLoS One*, **13**(4): e0196496.
- Parker, L., Ross, P., and O'Connor, W. 2011. Populations of the Sydney rock oyster, *Saccostrea glomerata*, vary in response to ocean acidification. *Mar. Biol.* **158**(3): 689-697. doi:10.1007/s00227-010-1592-4.
- Pespeni, M.H., Sanford, E., Gaylord, B., Hill, T.M., Hosfelt, J.D., Jaris, H.K., LaVigne, M., Lenz, E.A., Russell, A.D., Young, M.K., and Palumbi, S.R. 2013. Evolutionary change during experimental ocean acidification. *Proc. Natl. Acad. Sci. U.S.A.* **110**(17): 6937-6942. doi:10.1073/pnas.1220673110.
- Pfister, C.A., Esbaugh, A.J., Frieder, C.A., Baumann, H., Bockmon, E.E., White, M.M., Carter, B.R., Benway, H.M., Blanchette, C.A., Carrington, E., McClintock, J.B., McCorkle, D.C., McGillis, W.R., Mooney, T.A., and Ziveri, P. 2014. Detecting the unexpected: a research framework for ocean acidification. *Environ. Sci. Technol.* **48**: 9982-9994. doi:10.1021/es501936p.
- Pimentel, M., Pegado, M., Repolho, T., and Rosa, R. 2014a. Impact of ocean acidification in the metabolism and swimming behavior of the dolphinfish (*Coryphaena hippurus*) early larvae. *Mar. Biol.* **161**(3): 725-729. doi:10.1007/s00227-013-2365-7.
- Pimentel, M.S., Faleiro, F., Dionísio, G., Repolho, T., Pousão, P., Machado, J., and Rosa, R. 2014b. Defective skeletogenesis and oversized otoliths in fish early stages in a changing ocean. *J. Exp. Biol.* **217**: 2062-2070. doi:10.1242/jeb.092635.
- Porteus, C.S., Hubbard, P.C., Webster, T.M.U., van Aerle, R., Canário, A.V., Santos, E.M., and Wilson, R.W. 2018. Near-future CO₂ levels impair the olfactory system of a marine fish. *Nat. Clim. Change*, **8**(8): 737-746.
- Pörtner, H.-O. 2008. Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar. Ecol. Prog. Ser.* **373**: 203-217.
- Pörtner, H.-O. 2010. Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *J. Exp. Biol.* **213**(6): 881-893. doi:10.1242/jeb.037523.
- Pörtner, H.-O., Bock, C., and Mark, F.C. 2017. Oxygen-and capacity-limited thermal tolerance: bridging ecology and physiology. *J. Exp. Biol.* **220**(15): 2685-2696.
- Pörtner, H.O. 2012. Integrating climate-related stressor effects on marine organisms: unifying principles linking molecule to ecosystem-level changes. *Mar. Ecol. Prog. Ser.* **470**: 273-290.

- Przeslawski, R., Byrne, M., and Mellin, C. 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. *Glob. Change Biol.* **21**(6): 2122-2140. doi:10.1111/gcb.12833.
- Randall, D., and Tsui, T. 2002. Ammonia toxicity in fish. *Mar. Pollut. Bull.* **45**(1-12): 17-23.
- Reynaud, S., Leclercq, N., Romaine-Lioud, S., Ferrier-Pagés, C., Jaubert, J., and Gattuso, J.-P. 2003. Interacting effects of CO₂ partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Glob. Change Biol.* **9**(11): 1660-1668. doi:10.1046/j.1365-2486.2003.00678.x.
- Riebesell, U., Fabry, V.J., Hansson, L., and Gattuso, J.P. 2011. Guide to best practices for ocean acidification research and data reporting. Publications Office of the European Union: 260.
- Rosa, R., Baptista, M., Lopes, V.M., Pegado, M.R., Ricardo Paula, J., Trübenbach, K., Leal, M.C., Calado, R., and Repolho, T. 2014. Early-life exposure to climate change impairs tropical shark survival. *Proc. R. Soc. Lond. B Biol. Sci.* **281**(1793): 20141738. doi:10.1098/rspb.2014.1738.
- Salinas, S., and Munch, S.B. 2012. Thermal legacies: transgenerational effects of temperature on growth in a vertebrate. *Ecol. Lett.* **15**: 159-163. doi:10.1111/j.1461-0248.2011.01721.x.
- Salinas, S., Brown, S.C., Mangel, M., and Munch, S.B. 2013. Non-genetic inheritance and changing environments. *Non-Genetic Inheritance*, **1**: 38-50. doi:10.2478/ngi-2013-0005.
- Salisbury, J.E., and Jönsson, B.F. Rapid warming and salinity changes in the Gulf of Maine alter surface ocean carbonate parameters and hide ocean acidification. *Biogeochemistry*, **2018**: 1-18.
- Schade, F.M., Clemmesen, C., and Wegner, K.M. 2014. Within-and transgenerational effects of ocean acidification on life history of marine three-spined stickleback (*Gasterosteus aculeatus*). *Mar. Biol.* **161**(7): 1667-1676.
- Seibel, B.A., Maas, A.E., and Dierssen, H.M. 2012. Energetic plasticity underlies a variable response to ocean acidification in the pteropod, *Limacina helicina antarctica*. *PLoS ONE*, **7**(4): e30464. doi:10.1371/journal.pone.0030464.
- Shaw, E.C., Munday, P.L., and McNeil, B.I. 2013. The role of CO₂ variability and exposure time for biological impacts of ocean acidification. *Geophys. Res. Lett.* **40**(17): 4685-4688. doi:10.1002/grl.50883.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., and Krinner, G. 2018. Mass balance of the Antarctic ice sheet from 1992 to 2017. *Nature*, **558**: 219-222. doi:10.1038/s41586-018-0179-y.
- Simpson, S.D., Munday, P.L., Wittenrich, M.L., Manassa, R., Dixon, D.L., Gagliano, M., and Yan, H.Y. 2011. Ocean acidification erodes crucial auditory behaviour in a marine fish. *Biol. Lett.* **7**(6): 917-920. doi:10.1098/rsbl.2011.0293.
- Snyder, J.T., Murray, C.S., and Baumann, H. 2018. Potential for maternal effects on offspring CO₂ sensitivities in the Atlantic silverside (*Menidia menidia*). *J. Exp. Mar. Biol. Ecol.* **499**: 1-8. doi:10.1016/j.jembe.2017.11.002.

- Sokolova, I.M. 2013. Energy-limited tolerance to stress as a conceptual framework to integrate the effects of multiple stressors. *Integr. Comp. Biol.* **53** (4): 597-608. doi:10.1093/icb/ict028.
- Sommer, U. 2012. Experimental systems in aquatic ecology. *In* eLS (Ed.). doi:10.1002/9780470015902.a0003180.pub2.
- Sswat, M., Stiasny, M.H., Taucher, J., Algueró-Muñiz, M., Bach, L.T., Jutfelt, F., Riebesell, U., and Clemmesen, C. 2018. Food web changes under ocean acidification promote herring larvae survival. *Nat. Ecol. Evol.* **2**(5): 836.
- Stiasny, M., Mittermayer, F., Göttler, G., Bridges, C., Falk-Petersen, I.-B., Puvanendran, V., Mortensen, A., Reusch, T., and Clemmesen, C. 2018. Effects of parental acclimation and energy limitation in response to high CO₂ exposure in Atlantic cod. *Sci. Rep.* **8**(1): 8348.
- Stiasny, M.H., Mittermayer, F.H., Sswat, M., Voss, R., Jutfelt, F., Chierici, M., Puvanendran, V., Mortensen, A., Reusch, T.B.H., and Clemmesen, C. 2016. Ocean acidification effects on Atlantic cod larval survival and recruitment to the fished population. *PLoS ONE*, **11**(8): e0155448. doi:10.1371/journal.pone.0155448.
- Suggett, D.J., Dong, L.F., Lawson, T., Lawrenz, E., Torres, L., and Smith, D.J. 2013. Light availability determines susceptibility of reef building corals to ocean acidification. *Coral Reefs*, **32**(2): 327-337.
- Sunday, J.M., Calosi, P., Dupont, S., Munday, P.L., Stillman, J.H., and Reusch, T.B.H. 2014. Evolution in an acidifying ocean. *Trends Ecol. Evol.* **29**(2): 117-125. doi:10.1016/j.tree.2013.11.001.
- Thomsen, J., Casties, I., Pansch, C., Körtzinger, A., and Melzner, F. 2013. Food availability outweighs ocean acidification effects in juvenile *Mytilus edulis*: laboratory and field experiments. *Glob. Change Biol.* **19**(4): 1017-1027. doi:10.1111/gcb.12109.
- Thor, P., and Dupont, S. 2015. Transgenerational effects alleviate severe fecundity loss during ocean acidification in a ubiquitous planktonic copepod. *Glob. Change Biol.* **21**(6): 2261-2271.
- Vargas, C.A., Lagos, N.A., Lardies, M.A., Duarte, C., Manríquez, P.H., Aguilera, V.M., Broitman, B., Widdicombe, S., and Dupont, S. 2017. Species-specific responses to ocean acidification should account for local adaptation and adaptive plasticity. *Nat. Ecol. Evol.* **1**(4): 0084.
- Wallace, R.B., Baumann, H., Grear, J.S., Aller, R.C., and Gobler, C.J. 2014. Coastal ocean acidification: The other eutrophication problem. *Estuar. Coast. Shelf Sci.* **148**(0): 1-13. doi:10.1016/j.ecss.2014.05.027.
- Watson, S.A., Allan, B.J., McQueen, D.E., Nicol, S., Parsons, D.M., Pether, S.M., Pope, S., Setiawan, A.N., Smith, N., Wilson, C. 2018, and Munday, P.L. Ocean warming has a greater effect than acidification on the early life history development and swimming performance of a large circumglobal pelagic fish. *Glob. Change Biol.* **24** (9): 4368-4385.
- Welch, M.J., and Munday, P.L. 2016. Contrasting effects of ocean acidification on reproduction in reef fishes. *Coral Reefs*, **35**(2): 485-493. doi:10.1007/s00338-015-1385-9.

Figure captions

Figure 1: Overview of the discussed major advances and challenges in marine climate change experiments

Figure 2: Long-term growth costs of high CO₂ conditions in the Atlantic silverside (*M. menidia*). **A:** Violin plot of total length (TL) frequency distributions of juveniles reared from fertilization to 122 days post hatch under ambient (blue, ~500 µatm, n = 1,025) vs. high CO₂ conditions (red, ~2,300 µatm, n = 1,100). Symbols denote mean ± SD. Data source for **A** is Murray et al. 2017. **B:** Condition factors K ($10^5 \times \text{wet weight (g)}/\text{TL(mm)}^{2.944}$) of adults reared in duplicates from fertilization to 280 dph under ambient (blue, ~500 µatm, n = 155) vs. high CO₂ conditions (red, ~2,300 µatm, n = 132). Symbols denote mean ± 95% bootstrapped confidence intervals.

Figure 3: Illustration of the Ocean Variability Hypothesis positing that the CO₂ sensitivity of marine organisms is related to the magnitude of short-term CO₂ fluctuations in their habitat (e.g., from nearshore to open ocean) and length of their early life stage durations. It suggests that the most CO₂ tolerant marine organisms are those that develop fast and/or in habitats with large contemporary CO₂ fluctuations, whereas the potentially most vulnerable species are those that develop slowly in relatively stable open ocean habitats.

	Advances	Challenges
Methods	<p>Best practices for CO₂ manipulation, measurements, reporting</p> <p>Identification of confounding factors</p> <p>Multiple life-stages and generations important, carry-over effects</p>	<p>Scale and realism of experimental designs, pseudo-replication</p> <p>Multi-species experiments, indirect effects</p> <p>Field and long-term experiments</p>
Mechanisms	<p>Behavioral effects in fish due to GABAA receptor depolarization</p> <p>Metabolic costs of acid-base regulation to high CO₂ small</p> <p>Physiological frameworks, aerobic performance, ATP</p>	<p>Transcriptomic, proteomic up-/downregulation</p> <p>Trade-offs of CO₂ acclimation</p> <p>Selection under high CO₂</p>
Variability	<p>CO₂ varies naturally in time and space, largely due to ecosystem metabolism/upwelling</p> <p>Likely influences species CO₂ sensitivities</p> <p>Potential for local adaptation, transgenerational plasticity</p>	<p>Importance of extreme over average conditions, incorporating CO₂ fluctuations</p> <p>Use of laboratory/commercial broodstocks for sensitivity experiments</p>
Multistressors	<p>Major concurrent stressors = warming, acidification, oxygen loss</p> <p>Synergistic interactions are the norm, not the exception</p> <p>Early life stages are most vulnerable</p>	<p>Mechanistic experiments to study effect interactions vs. realistic experiments to study climate change scenarios</p> <p>Incorporating more than 3 stressors, sequential experimentation</p>

Overview of the discussed major advances and challenges in marine climate change experiments

127x143mm (300 x 300 DPI)

Long-term growth costs of high CO₂ conditions in the Atlantic silverside (*M. menidia*). A: Violin plot of total length (TL) frequency distributions of juveniles reared from fertilization to 122 days post hatch under ambient (blue, ~500 μ atm, n = 1,025) vs. high CO₂ conditions (red, ~2,300 μ atm, n = 1,100). Symbols denote mean $\pm$ SD (modified after Murray et al. 2017). B: Condition factors K ($105 \times \text{wet weight (g)} / \text{TL (mm)}^2.944$) of adults reared in duplicates from fertilization to 280 dph under ambient (blue, ~500 μ atm, n = 155) vs. high CO₂ conditions (red, ~2,300 μ atm, n = 132). Symbols denote mean $\pm$ 95% bootstrapped confidence intervals.

240x134mm (300 x 300 DPI)

Illustration of the Ocean Variability Hypothesis positing that the CO₂ sensitivity of marine organisms is related to the magnitude of short-term CO₂ fluctuations in their habitat (e.g., from nearshore to open ocean) and length of their early life stage durations. It suggests that the most CO₂ tolerant marine organisms are those that develop fast and/or in habitats with large contemporary CO₂ fluctuations, whereas the potentially most vulnerable species are those that develop slowly in relatively stable open ocean habitats.

174x134mm (300 x 300 DPI)