Canadian Journal of Plant Science

AAC Stronghold Durum Wheat

Journal:	<i>Canadian Journal of Plant Science</i>
Manuscript ID	CJPS-2018-0224.R1
Manuscript Type:	Cultivar description
Date Submitted by the Author:	15-Nov-2018
Complete List of Authors:	Ruan, Yuefeng; Swift-Current Research and Development Centre, AAFC Singh, Asheesh; Iowa State University, Agronomy DePauw, R.; Advancing Wheat Technologies, Knox, Ron; Agriculture & Agri-Food Canada, Swift Current Research and Development Centre Cuthbert, Richard; Swift Current Research & Development Centre, Agriculture & Agri-Food Canada McCallum, Brent; Morden Research and Development Centre, , AAFC Fetch, Thomas; Morden Research and Development Centre, , AAFC Beres, Brian; Agriculture and Agri-Food Canada, Sustainable Production Systems
Keywords:	Triticum turgidum,, grain yield, solid stem,, cadmium uptake
Is the invited manuscript for consideration in a Special Issue?:	Not applicable (regular submission)

SCHOLARONE™
Manuscripts

Cultivar Description

AAC Stronghold Durum Wheat

Short title: Ruan et al. – AAC Stronghold durum wheat

Y. Ruan¹, A.K. Singh^{1,2*}, R.M. DePauw^{3*}, R.E. Knox¹, R.D. Cuthbert¹, B. McCallum⁴, T. Fetch⁴, B.L. Beres⁵

¹Agriculture and Agri-Food Canada, Swift Current Research and Development Centre, Box 1030, Swift Current, SK, S9H 3X2; ²Present Address: Department of Agronomy, Iowa State University, Ames, IA, USA; ³Advancing Wheat Technologies, 870 Field Drive, Swift Current, SK, S9H 4N5; ⁴Agriculture and Agri-Food Canada, Morden Research and Development Centre, 101 Route 100, Morden, Manitoba; ⁵Agriculture and Agri-Food Canada, Lethbridge Research and Development Centre, Lethbridge, Alberta, Canada.

* Corresponding authors: A.K. Singh (email: singhak@iastate.edu) and R.M. DePauw (email: ron.depauw1944@gmail.com)

Y. Ruan, A.K. Singh, R.M. DePauw, R.E. Knox, R.D. Cuthbert, B. McCallum, T. Fetch, B. Beres. 20xx. AAC Stronghold durum wheat. Can. J. Plant Sci. xx:000-000. AAC Stronghold durum wheat (*Triticum turgidum* L. subsp. *durum* (Desf.) Husn.) is adapted to the durum production area of the Canadian prairies. Averaged over three years, AAC Stronghold yielded significantly more grain than AC Navigator. AAC Stronghold had protein concentration significantly less than Strongfield but significantly more than Brigade. AAC Stronghold had plant height significantly shorter than Brigade, Strongfield and AAC Cabri with lodging scores significantly less than Strongfield and AAC Cabri. AAC Stronghold has a solid stem which confers resistance to cutting by the wheat stem sawfly (*Cephus cinctus* Norton). AAC Stronghold is eligible for grades of Canada Western Amber Durum. AAC Stronghold had low grain cadmium concentration and stronger gluten than Strongfield.

Key words: *Triticum turgidum*, durum wheat, cultivar description, grain yield, solid stem, cadmium uptake

AAC Stronghold durum wheat (*Triticum turgidum* L. subsp. *durum* (Desf.) Husn.) was developed at the Swift Current Research and Development Centre (SCRDC), Agriculture and Agri-Food Canada (AAFC), Swift Current, SK. Plant Breeders' Rights, grant of rights certificate #5769 was granted on 11 May 2018, and AAC Stronghold received registration No. #8085 from the Variety Registration Office, Canadian Food Inspection Agency, on 8 July 2016.

Pedigree and Breeding Method

AAC Stronghold (experimental names: DT862, A0719-KZ06) was selected from the cross DT805/DT795 made in 2007 at the Swift Current Research and Development Centre, Swift

Current, SK. DT805 (A0138-KZ01) is a breeding line derived from a cross of 9667A-AV6/DT704//Strongfield. Strongfield (Clarke et al. 2005a) is a high yielding, high grain protein concentration, low cadmium uptake Canadian durum cultivar. DT795 (A0050-EC03) is a solid stem breeding line derived from a cross of 94D11-K*3B/Strongfield//9688A-245D2 where the solid stem line 9688A-245D2 derived from the cross of DT663/W9262-260D1//DT665/DT673. The solid stem line W9262-260D1 derived from the cross Kyle*2/Biodur, where Kyle (Townley-Smith et al. 1987) is a widely adapted high cadmium uptake, hollow stem Canadian cultivar and Biodur is a low cadmium uptake, solid stem cultivar from Germany. The parents were haplotyped with molecular markers linked to *Cdu1* controlling cadmium uptake, and *Sst1* controlling stem solidness (Houshmand et al. 2007, Knox et al. 2009).

In 2007, F₁ seeds were planted in the greenhouse to increase F₂ seed. In the spring of 2008, approximately 8000 seeds of the F₂ generation were space-planted at 14 cm intervals within a row in an irrigated epiphytotic field nursery near Swift Current. Genotypes susceptible to prevalent races of leaf rust (*Puccinia triticina* Eriks.) and stem rust (*P. graminis* Pers.:Pers. f. sp. *tritici* Eriks. and E. Henn.) were planted as disease spreaders every tenth row. Between the spreader rows, five rows of spring planted winter wheat were alternated with four rows of F₂ seed at a row spacing of 23 cm. The winter wheat cultivar CDC Kestrel (Fowler 1997), which is susceptible to leaf and stem rust, was used to contribute to the multiplication of rust inoculum. Spreader rows were inoculated by injecting, with a syringe and needle, a water suspension of leaf rust and stem rust spores into a sample of plants every 3 m. Representative leaf rust races found the previous year were applied (McCallum and Seto-Goh 2006). Stem rust races used were: QTHJF (C25), RHTSC (C20), RKQSC (C63), RTHJF (C57), TMRTF (C10), and TPMKC (C53) (Roelfs and Martens 1988; Fetch et al. 2015). Leaf spotting diseases developed through natural

infection. Individual plants were selected for plant height, straw strength, maturity, and resistance to leaf spotting diseases, leaf rust, and stem rust.

The F_3 seeds from individual spikes from 261 selected F_2 plants were grown in 2 m long rows in a contra season nursery near Lincoln, New Zealand, in 2008-2009. Based on plant height, days to maturity, and straw strength, 137 rows were selected, and the rows were harvested individually to produce the seed used for agronomic and disease trials in Canada. In 2009, the 137 F_4 lines, their parents, and other check cultivars were grown in unreplicated 2.74 m² four-row plot experiments near Swift Current and Regina, SK. The traits grain yield, plant height, time to maturity, straw strength, and leaf spotting infection were assessed. Seven spikes per F_4 line from within plots grown near Swift Current were selected for plant height, straw strength, and leaf spotting disease symptoms caused primarily by tan spot [*Pyrenophora tritici-repentis* (Died.) Drechs., anamorph *Drechslera tritici-repentis* (Died.) Shoemaker] and stagonospora nodorum blotch [*Phaeosphaeria nodorum* (E. Müll.) Hedjaroude, anamorph *Stagonospora nodorum* (Berk.) Castell. & E.G. Germano]. The grain quality traits protein concentration, yellow pigment concentration, gluten strength, and volume weight were assessed on grain harvested from field trials. Based on this suite of agronomic, disease, and quality traits, 21 F_4 lines were selected.

In 2009-10, 147 F_5 lines (from the 21 $F_{4:5}$ families at 7 spikes per F_4 line) were grown in 2 m long rows near Leeston, New Zealand and selected primarily on plant height, straw strength, and days to maturity. After selection, 86 $F_{4:6}$ lines were grown in 2010 under dryland conditions near Swift Current and Regina, SK, and under irrigation near Lethbridge, AB. Twenty-eight genotypes were selected based on agronomic performance, disease resistance, and quality traits assessed as described for the F_4 generation.

Four F₇ genotypes were grown in the 2011 Durum A-level tests as a two replicate lattice design with four-row plots planted near Swift Current, Regina, and Indian Head, SK, Lethbridge, AB, and Brandon, MB, to assess agronomic performance as described for the F₄ generation. Check cultivars in the Durum A tests were AC Avonlea (Clarke et al.1998), AC Morse, AC Navigator (Clarke et al. 2000), Brigade (Clarke et al. 2009), Commander (Clarke et al.2005b), and Strongfield. Remnant seed from the yield trials was bulked over replications within a location and a subsample from each location that graded better than #3 CWAD was used to assess end-use suitability by the Central Quality Lab, Cereal Research Centre, Winnipeg, MB. Assessment included grain protein concentration, yellow pigment concentration, milling properties, gluten strength, and Hagberg Falling Number. Response to loose smut [*Ustilago tritici* (Pers.) Rostr.] was tested with a mixture of races T26, T32, and T33 (Nielsen 1987) under field conditions near Swift Current. Response to leaf rust and stem rust were evaluated in hill plots in a rust nursery near Glenlea, MB, using a mixture of races similar to that in the F₂ rust nursery. Response to leaf spotting pathogens was assessed from within the yield plots under conditions of natural inoculum. Response to Fusarium head blight (FHB) caused by *Fusarium graminearum* Schwabe (teleomorph *Gibberella zeae* (Schwein. Petch) was assessed in nurseries near Portage la Prairie and Carman, MB (Bokore et al. 2017). Plots at Carman were scored for incidence (%) and severity (%), and at Portage la Prairie the plots were scored on a 1 to 9 scale on increments of 10% incidence and severity symptoms. Scoring for FHB was performed when a significant differential reaction was observed among checks. These procedures identified the line A0719-KZ06, which met all of the selection criteria at each stage of selection.

The four lines from the population designated A0719, including A0719-KZ06 were tested in the 2012 Durum-B test in an alpha-lattice design with two replications and grown near Swift

Current, Regina, Saskatoon, Floral, SK, Lethbridge, AB, and Brandon, MB, using the same check cultivars as in the 2011 Durum A tests. Response to diseases was measured using protocols similar to that for the A-level tests described above. Remnant seed from the yield trials was used to prepare a composite, using degrading factors as a consideration for suitability for inclusion in the composite, to assess the same end-use suitability parameters as in the Durum A-level test, by the Central Quality Lab, Cereal Research Centre, MB. This procedure identified the line A0719-KZ06, which met all of the selection criteria at each stage of selection.

A0719-KZ06 was advanced to the Durum Wheat Cooperative Test and evaluated as DT862 from 2013 to 2015. The Durum Wheat Cooperative Test was grown in four row plots at up to 12 locations annually in a two replicate 5 x 6 lattice design including four check cultivars in two repetitions and two replications. The check cultivars were AAC Cabri (Singh et al. 2017), AC Navigator, Brigade, and Strongfield. The Durum Wheat Cooperative Test operating protocols are described in the Prairie Recommending Committee for Wheat Rye and Triticale operating procedures (Anonymous 2015). The PROC MIXED procedure in SAS version 9 (SAS Institute Inc. 2003) was used to analyze the data annually and to perform a combined analysis over years, using a mixed model with sub-blocks, replications, zones, locations and years considered random effects, and genotypes considered fixed effects (Littell et al. 2006). The diff command was used to estimate the standard error of the difference between entries, which in turn was used to estimate an F-protected least significant difference (LSD) at a significance level of 5% ($LSD_{0.05}$).

The Durum Wheat Cooperative Test entries were evaluated in inoculated disease nurseries near Brandon, Portage la Prairie and Morden MB to determine the response to leaf rust, stem rust, and loose smut. Fusarium head blight was assessed in inoculated nurseries near

Carman, Glenlea, Portage la Prairie and Morden MB, Ottawa, ON, and Charlottetown, PEI.

Inoculum composition for leaf rust, stem rust, and loose smut was as described above. Response to common bunt caused by *Tilletia laevis* Kuhn in Rabenh., and *T. tritici* (Bjerk.) G. Wint. in Rabenh., was assessed in a nursery grown near Lethbridge, using a mixture of prevalent races: T-1, T-6, T-13, T-19, L-1, and L-16 (Hoffmann and Metzger 1976, Gaudet and Puchalski 1989). Leaf spotting reaction was determined based on natural infection at Swift Current.

A sample of grain of DT862 and the check cultivars from each location was submitted to the Canadian Grain Commission to determine grain grade and protein concentration. End-use suitability was determined on a composite sample made up from sites with grain samples representative only of the top durum wheat grades available. The quantity of grain from a location was adjusted to achieve a final composite protein concentration approximating that of the average for the crop that year. A consistent quantity of grain within a location for all experimental lines was used to make up the composite each year. All end-use suitability analyses were performed by personnel at the Grain Research Laboratory, Canadian Grain Commission, Winnipeg, MB following protocols of the AACC (American Association of Cereal Chemists, 2000).

Performance

In three years of cooperative testing, the grain yield of AAC Stronghold was significantly greater than AC Navigator in both zones (Table 1). Averaged over zones, AAC Stronghold was significantly later maturing than Strongfield but earlier than Brigade (Table 2). Test weight (kg hL⁻¹) of AAC Stronghold was significantly heavier than AC Navigator and Strongfield. The 1000-kernel weight (g) of AAC Stronghold was significantly heavier than Strongfield and AAC Cabri, but similar to the other checks. AAC Stronghold had plant height significantly shorter

than Brigade, and AAC Cabri while being significantly taller than AC Navigator. Lodging score was significantly less than Strongfield and AAC Cabri. Grain protein concentration of AAC Stronghold was similar to AC Navigator and AAC Cabri, significantly more than Brigade, and significantly less than Strongfield (Table 3).

AAC Stronghold was resistant to leaf rust, stem rust, and loose smut, moderately resistant to stripe rust and resistant to common bunt and intermediately resistant to leaf spots (Table 4). The FHB symptoms of disease incidence and severity of AAC Stronghold was rated as moderately susceptible while the deoxynivalenol (DON) accumulation was rated as susceptible (Table 5).

AAC Stronghold is a solid stemmed genotype (Tables 6) a trait that reduces cutting by the wheat stem sawfly (*Cephus cinctus* Norton) (Holmes and Peterson 1961a, 1961b). AAC Stronghold has low grain cadmium concentration similar to Strongfield (Table 7). The grain and milling attributes of AAC Stronghold were intermediate to the checks. AAC Stronghold had gluten index and alveograph W values significantly higher than Strongfield as well as pasta b* colour desirably higher. The semolina ash of AAC Stronghold was significantly higher than for Strongfield.

Other Characteristics

Straw (at maturity): thick pith in cross-section, no anthocyanin coloration.

Spikes: parallel-sided in profile, mid-dense to dense, erect; off-white at maturity; awns longer than spike, white at maturity.

Kernel: colour amber; kernel size large, elliptical, short brush hairs.

Lower glume: medium length and width; glabrous.

Lower glume shoulder: very narrow to medium width; slightly sloping shape.

Lower glume beak: narrow to medium length, straight to slightly curved shape.

End-use suitability: Eligible for the grades of Canada Western Amber Durum wheat market class.

Maintenance and Distribution of Pedigreed Seed

The 107 Breeder Lines originated from random $F_{4:9}$ single plants of A0719-KZ06 grown as 108 pre-Breeder Lines in 3 m long rows in isolation near Swift Current in 2014, and again as 15 m rows near Indian Head in 2015. Breeder Seed will be maintained by the Seed Increase Unit of the Research Farm, Indian Head, SK, S0G 2K0. Distribution and multiplication of pedigreed seed stocks will be handled by SeCan Association, 400 * 300 Terry Fox Drive, Ottawa, ON., K2K 0E3, Canada. www.secan.com

Acknowledgements

We gratefully acknowledge the financial support of the producer funded wheat research check-off (administered by the Western Grains Research Foundation); D. Niziol and J. Fehr of AAFC, Cereal Research Centre (CRC) (Winnipeg, MB) for providing end-use quality analyses; J. Gilbert of AAFC, CRC for leaf spotting and FHB reactions; J. Menzies of AAFC, CRC for loose smut evaluation; D. Gaudet and T. Despins of AAFC, Lethbridge Research Centre for providing reaction to common bunt; E. Johnson, Scott Research Farm, AAFC, for agronomic assessment at Scott; C.J. Pozniak, University of Saskatchewan, for agronomic assessment near Saskatoon, SK; Francis Kirigwi, Syngenta AgriPro, for agronomic assessment at Souris, MB and Pense, SK; Tim Ferguson, Viterra, for agronomic assessment at Moose Jaw, SK, and Vulcan, AB; Stephen Fox, Gavin Humphreys and Doug Brown of AAFC, CRC for agronomic assessment at Brandon, MB

and FHB nursery at Portage la Prairie, MB; Anita Brûlé-Babel of University of Manitoba for FHB evaluations at Carman, MB; B.X. Fu and L. Schlichting of the Grain Research Laboratory, Canadian Grain Commission, Winnipeg, MB, for end-use quality assessment; D. Gehl of AAFC Seed Increase Unit, Indian Head, SK for multiplication of Breeder seed; M. Olfert, J. Ross, L. Oakman, J. Powell, H. Campbell, S. Friesen, T. Greenwood, Brad Meyer, Alison Banman, Theresa Colenutt, and Cam Barlow of the wheat molecular genetics lab at SCRDC-AAFC for molecular marker work, and all members of the wheat genetic enhancement group at AAFC, Swift Current; M. Knelsen, AAFC Regina, SK.; O. Thompson, AAFC Indian Head, SK; and R. Dyck, AAFC, Lethbridge, AB, for their assistance in conducting field trials.

American Association of Cereal Chemists. 2000. Approved Methods of the AACC, 10th Ed. St. Paul, MN.

Anonymous. 2015. Operating Procedures, Prairie Recommending Committee for Wheat, Rye and Triticale. [Online Accessed 12 June 2018]. Available at:

<http://www.pgdc.ca/pdfs/wrt/Proposed%20PRCWRT%20OPS%20-20FINAL%20DRAFT%20-%2027%20Nov%202013%20Updated%205%20December%202015.pdf>

Bokore, F. E., Knox, R. E., DePauw, R. M., Clarke, F., Cuthbert, R. D., Campbell, H. L., Brûlé-Babel, A. L., Gilbert, J. and Ruan, Y. 2017. Validation of molecular markers for use with adapted sources of fusarium head blight resistance in wheat. *Plant Disease* **101**:1292–1299.

<http://dx.doi.org/10.1094/PDIS-10-16-1421-RE>.

Clarke, J. M., Knox, R. E., DePauw, R. M., Clarke, F. R., Fernandez, M. R., McCaig, T. N. and Singh, A. K. 2009. Brigade durum wheat. *Can. J. Plant Sci.* **89**: 505 – 509.

doi:[10.4141/CJPS08168](https://doi.org/10.4141/CJPS08168).

Clarke, J. M., McCaig, T. N., DePauw, R. M., Knox, R. E., Clarke, F. R., Fernandez, M. R. and

- Ames, N. P. 2005a. Strongfield durum wheat. *Can. J. Plant Sci.* **85**: 651 – 654. doi:[10.4141/P04-119](https://doi.org/10.4141/P04-119).
- Clarke, J. M., McCaig, T. N., DePauw, R. M., Knox, R. E., Ames, N.P., Clarke, F.R., Fernandez, M. R., Marchylo, B. A., and Dexter, J. A. 2005b. Commander durum wheat. *Can. J. Plant Sci.* **85**: 901 – 904. doi:[10.4141/P04-189](https://doi.org/10.4141/P04-189).
- Clarke, J.M., McLeod, J.G., DePauw, R.M., Marchylo, B.A., McCaig, T.N., Knox, R.E., Fernandez, M.R., and Ames, N. 2000. AC Navigator durum wheat. *Can. J. Plant Sci.* **80**:343-345. doi:[10.4141/P99-108](https://doi.org/10.4141/P99-108).
- Clarke, J. M., McLeod, J. G., McCaig, T. N., DePauw, R. M., Knox, R. E., and Fernandez, M. R. 1998. AC Avonlea durum wheat. *Can. J. Plant Sci.* **78**: 621-623. doi:[10.4141/P98-002](https://doi.org/10.4141/P98-002).
- Fetch, T., Mitchell Fetch, J. Zegeye, T. and Xue, A. 2015. Races of *Puccinia graminis* on wheat, oat, and barley in Canada in 2009 and 2010. *Can. J. Plant Pathol.* **37**:476-484. doi:[10.1080/07060661.2015.1119735](https://doi.org/10.1080/07060661.2015.1119735).
- Fowler, D.B. 1997. CDC Kestrel winter wheat. *Can. J. Plant Sci.* **77**: 673-675. doi:[10.4141/P96-193](https://doi.org/10.4141/P96-193).
- Gaudet, D.A. and Puchalski, B.L. 1989. Races of common bunt (*Tilletia caries* and *T. foetida*) of wheat in western Canada, *Can. J. Plant Pathol.* **11**: 415-418. doi:[10.1080/07060668909501089](https://doi.org/10.1080/07060668909501089).
- Gilbert, J. and Woods, S. 2006. Strategies and considerations for multi-location FHB screening nurseries. Pages 93-102 in T. Ban, J. M. Lewis, and E. E. Phipps, eds. The global Fusarium initiative for international collaboration: A strategic planning workshop held at CIMMYT, El Batàn, Mexico. 2006. Mar. 14-17. CIMMYT, Mexico, D.F.

- Hoffmann, J.A. and Metzger, R.J. 1976. Current status of virulence genes and pathogenic races of the wheat bunt fungi in the northwestern USA. *Phytopathology* **66**: 657-660.
doi:[10.1094/Phyto-66-657](https://doi.org/10.1094/Phyto-66-657).
- Holmes, N.D. and Peterson, L.K. 1961a. Resistance of spring wheats to the wheat stem sawfly, *Cephus cinctus* Nort. (*Hymenoptera: Cephidae*. I. Resistance to the egg. *Can. Entomol.* **94**:250-259. doi:[10.4039/Ent93250-4](https://doi.org/10.4039/Ent93250-4).
- Holmes, N.D. and Peterson, L.K. 1961b. Resistance of spring wheats to the wheat stem sawfly, *Cephus cinctus* Nort. (*Hymenoptera: Cephidae*. II. Resistance to the larvae. *Can. Entomol.* **94**: 348-365. doi:[10.4039/Ent94348-4](https://doi.org/10.4039/Ent94348-4).
- Houshmand, S., Knox, R.E., Clarke, F.R., Clarke, J.M. 2007. Microsatellite markers flanking a stem solidness gene on chromosome 3BL in durum wheat. *Mol Breeding*.**20**:261-70.
- Knox, R.E., Pozniak, C.J., Clarke, F.R., Clarke, J.M., Houshmand, S. and Singh, A.K. 2009. Chromosomal location of the cadmium uptake gene (*Cdu1*) in durum wheat. *Genome* **52**:741-7. doi:[10.1139/G09-042](https://doi.org/10.1139/G09-042). PMID:[19935921](https://pubmed.ncbi.nlm.nih.gov/19935921/).
- Littell, R. C., Milliken, G. A., Stroup, W. W. and Wolfinger, R. D. 2006. SAS® system for mixed models. 2nd ed. SAS Institute, Inc., Cary, NC.
- McCallum, B. D. and Seto-Goh, P. 2006. Physiologic specialization of *Puccinia triticina*, the causal agent of wheat leaf rust, in Canada in 2004. *Can. J. Plant Pathol.* **28**:566-576.
doi:[10.1080/07060660609507335](https://doi.org/10.1080/07060660609507335).
- McFadden, W. 1991. Etiology and epidemiology of leaf-spotting diseases in winter wheat in Saskatchewan. Ph.D. Thesis, University of Saskatchewan, Saskatoon, SK. 151 pp.
- Nielsen, J. 1987. Races of *Ustilago tritici* and techniques for their study. *Can. J. Plant Pathol.* **9**: 91-105. doi:[10.1080/07060668709501888](https://doi.org/10.1080/07060668709501888).

Roelfs, A.P. and Martens, J.W. 1988. An international system of nomenclature for *Puccinia graminis* f. sp. *tritici*. *Phytopathology* **78**: 525-533. doi:[10.1094/Phyto-78-526](https://doi.org/10.1094/Phyto-78-526).

SAS Institute Inc. 2003. SAS Software. Version 9. SAS Institute Inc. Cary, NC, USA.

Singh, A.K., R.M. DePauw, R.E. Knox, J.M. Clarke, T.N. McCaig, R.D. Cuthbert, Y. Ruan. 2017. AAC Cabri durum wheat. *Can. J. Plant Sci.* **97**: 135-143. [dx.doi.org/10.1139/cjps-2016-0044](https://doi.org/10.1139/cjps-2016-0044)

Townley-Smith, T. F., DePauw, R. M., Lendrum, C. W. B., McCrystal, G. E., and Patterson, L. A. 1987. Kyle durum wheat. *Can. J. of Plant Sci.* **67**: 225–227. doi:[10.4141/cjps87-026](https://doi.org/10.4141/cjps87-026).

Table 1. Grain yield (kg ha⁻¹) of AAC Stronghold and check cultivars in the Durum Cooperative Test, 2013 to 2015 in Zones 1 and 2^a.

	2013			2014			2015			2013 – 2015		
	Zone 1	Zone 2	Mean ^b	Zone 1	Zone 2	Mean	Zone 1	Zone 2	Mean	Zone 1	Zone 2	Mean
AC Navigator	3556	4540	4367	4045	3349	3490	3640	3099	3197	3733	3668	3688
Brigade	4891	5156	5108	5170	4173	4375	4062	3415	3540	4719	4254	4345
Strongfield	4174	4816	4701	4896	3748	3978	4160	3231	3391	4413	3926	4023
AAC Cabri	4619	4951	4891	5078	3874	4113	3734	3549	3594	4473	4130	4202
Mean of Checks	4310	4866	4767	4797	3786	3989	3899	3324	3431	4342	3995	4058
AAC Stronghold	4732	4825	4809	5874	3900	4286	4277	3452	3593	4930	4061	4225
LSD ^c _{0.05}	670	245	241	782	355	335	828	181	203	506	246	223
No. of tests	2	9	11	2	8	10	2	9	11	6	26	32

^a Zone 1 (Black Soils): Indian Head, Brandon; Zone 2 (Brown and Dark Brown Soils): Swift Current, Stewart Valley, Saskatoon, Lethbridge, Vulcan, Moose Jaw (2013, 15), Pense, Scott, Vanguard.

^b Means are LS means obtained from PROC MIXED procedure of SAS.

^c Least significant difference, $P \leq 0.05$, includes the genotype by environment interaction variation.

Table 2. Agronomic characteristics of AAC Stronghold and check cultivars in the Durum Wheat Cooperative Test, 2013 – 2015.

	Days to maturity ^{a,b}			Test weight ^a (kg hL ⁻¹)			1000-kernel wt ^a (g)	Height ^d (cm)	Lodging ^e (1-9)
	Zone 1	Zone 2	Mean ^c	Zone 1	Zone 2	Mean			
AC Navigator	98.7	102.6	102.1	73.2	79.0	78.0	44.3	77.2	2.2
Brigade	99.9	105.0	104.2	74.0	79.6	78.7	42.9	98.1	2.0
Strongfield	97.3	101.6	101.0	74.4	78.8	78.1	41.5	89.5	3.1
AAC Cabri	99.1	103.3	102.7	74.7	79.7	78.8	40.7	93.0	3.0
AAC Stronghold	100.2	103.6	103.1	75.1	79.7	79.0	44.0	87.4	1.4
LSD ^f _{0.05}	2.8	1.2	0.9	2.0	0.7	0.7	1.1	2.3	0.8
No. of tests	4	24	28	5	26	31	31	32	14

^a Zone 1 (Black Soils): Indian Head, Brandon (2013,15); Zone 2 (Brown and Dark Brown Soils): Swift Current, Stewart Valley, Saskatoon, Lethbridge, Vulcan, Moose Jaw (2013, 15), Pense, Scott, Vanguard.

^b All Zone 1 and Zone 2 locations except Stewart Valley (in Zone 2).

^c Means are LS means obtained from PROC MIXED procedure of SAS.

^d Includes Brandon 2014.

^e Swift Current (2014), Moose Jaw (2015), Saskatoon (2014), Stewart Valley, Brandon (2013), Pense, Vanguard, Indian Head (2014).

^f Least significant difference, $P \leq 0.05$, includes the genotype by environment interaction variation.

Table 3. Grain protein concentration (13.5% moisture basis) of AAC Stronghold and check cultivars measured on grain samples bulked across replications at each location of the Durum Cooperative Test, 2013–2015.

	2013			2014			2015			2013-15
	Zone 1 ^a	Zone 2	Mean ^b	Zone 1	Zone 2	Mean	Zone 1	Zone 2	Mean	Mean
AC Navigator	13.5	12.9	13.0	12.8	13.3	13.2	14.8	13.5	13.7	13.4
Brigade	12.7	12.7	12.7	12.8	13.0	12.9	14.1	12.8	13.0	13.0
Strongfield	14.7	13.6	13.8	13.5	13.9	13.8	15.2	13.5	13.8	13.9
AAC Cabri	13.9	13.1	13.3	13.5	13.9	13.8	16.3	13.1	13.7	13.6
AAC Stronghold	13.2	13.1	13.1	13.7	13.5	13.5	15.0	13.4	13.7	13.5
LSD ^c _{0.05}	1.2	0.4	0.4	0.6	0.6	0.6	0.9	0.6	0.5	0.3
No. of tests	2	9	11	1	8	9	2	9	11	31

^a Zone 1 (Black Soils): Indian Head, Brandon; Zone 2 (Brown and Dark Brown Soils): Swift Current, Stewart Valley, Saskatoon, Lethbridge, Vulcan, Moose Jaw (2013, 15), Pense, Scott, Vanguard.

^b Means are LS means obtained from PROC MIXED procedure of SAS.

^c Least significant difference, $P \leq 0.05$, includes the genotype by environment interaction variation.

Table 4. Summary of disease reactions to stem rust, leaf rust, stripe rust, common bunt, loose smut, and leaf spots of AAC Stronghold and check cultivars grown in the durum cooperative test, 2013-2015.

	Year	Stem rust		Leaf rust	Common bunt		Loose smut		Leaf spot		Stripe rust			
		Severity ^a (%)	Rating ^b	Rating ^b	Incidence (%) ^c	Rating ^b	Incidence (%) ^d	Rating ^b	SC ^e		LB ^e		CT ^e	
									Infection response ^f	Rating ^b	Severity ^g (%)	Infection response ^b	Severity (%)	Infection response
AC Navigator	2013	5 MR		R	1 R		35 MR		9.3 MS		60 S		15 R	
	2014	7 MR		R	1 R		7 R		9.8 S		5 R		15	
	2015	15 MR		R	0 R		0 R		-		15 I		-	
Brigade	2013	1 R		R	1 R		0 R		8.3 MS		15 R		15 R	
	2014	7 MR		R	1 R		3 R		8.5 MS		25 MR		25	
	2015	5 R		R	0 R		0 R		-		25 I		-	
Strongfield	2013	1 R		R	7 R		8 R		8.3 MS		15 R		5 R	
	2014	1 R		R	1 R		11 MR		8.8 MS		5 R		5	
	2015	2 R		R	1 R		0 R		-		10 MR		-	
AAC Cabri	2013	1 R		R	4 R		14 R		7.8 I		10 R		5 R	
	2014	10 MR		R	0 R		8 R		8.5 MS		20 MR		15	
	2015	5 R		R	1 R		0 R		-		5 R		-	
AAC Stronghold	2013	1 R		R	26 I		2 R		7.8 I		10 R		15 R	
	2014	1 R		R	0 R		6 R		7.3 I		20 MR		5	
	2015	5 MR		R	1 R		4 R		-		10 MR		-	

^a Severity is a percentage of the stem infected with stem rust using the Modified Cobb Scale.
^b Rating: VR, very resistant; R, resistant; MR, moderately resistant; I, intermediate; MS, moderately susceptible; S, susceptible.
^c Percentage of total spikes with common bunt symptoms.
^d Percentage of total plants with loose smut symptoms.
^e SC, Swift Current; LB, Lethbridge; CT, Creston.
^f Adult plant, rated mid-grainfill at Swift Current McFadden scale (0=no symptoms, 11=severe symptoms) (McFadden 1991).
^g Severity is a percentage of the leaf infected with stripe rust using the Modified Cobb Scale.

Table 5. Summary of Fusarium head blight disease reactions of AAC Stronghold and check cultivars grown in the Durum Cooperative Test, 2013-2015.

	Year	Fusarium Head Blight									DON (ppm) ^e				ISD ^f				FDK ^g	
		Carman		Portage		Morden		Ottawa	Glenlea	PEI ^d					MD		CM		CM	PEI
		Index ^a	Rating ^b	Index	Rating ^b	Index	Rating ^b	Score ^c (1 – 100)	Index	Index	CM ^d	MD ^d	OT ^d	PEI	Score	Rating ^b	Score	Rating	% ^g	1 – 9 ^h
AC Navigator	2013	51 MS		21 MS				73	9	73			17	17						9
	2014	56 S				89 S		90		49	42	118	19	87	74 S		28 S		31	9
	2015	56 MS				68 S					70	91	23	26	58 S		45		48	8
Brigade	2013	23 MR		17 I				48	7	48			12	15					32	9
	2014	19 I				16 R		37		46	30	101	5	47	62 S		20 S		23	9
	2015	30 MS				29 MR					45	20	8	19	14 MR		29		45	7
Strongfield	2013	30 I		17 I				90	10	72			9	22					19	9
	2014	40 MS				42 I		53		52	35	80	26	56	51 MS		24 S		24	9
	2015	39 MS				67 S					36	40	13	18	27 MS		24		39	8
AAC Cabri	2013	34 I		17 I				45	15	46			16	21					16	8
	2014	32 MS				41 I		27		47	32	72	14	44	46 I		22 S		15	8
	2015	41 MS				57 MS					41	37	13	17	25 I		27		41	7
AAC Stronghold	2013	43 MS		20 I				70	11	64			19	16					19	8
	2014	33 MS				43 MR		48		43	31	106	16	33	66 S		21 S		18	9
	2015	44 MS				64 MS					47	69	28	11	44 S		31		31	7

^a Fusarium head blight disease index = (percentage of infected heads x percentage of diseased florets on infected heads)/100.

^b Disease response category: R, resistant; MR, moderately resistant; I, intermediate resistant; MS, moderately susceptible; S, susceptible.

^c Score is based on a 1 to 100 scale of spikes infected with Fusarium head blight.

^d PEI, Prince Edward Island; CM, Carman; MD, Morden; OT, Ottawa.

^e DON, Deoxynivalenol in parts per million.

^f ISD, Incidence-Severity-DON index calculated as (0.2*mean incidence) + (0.2*mean severity) + (0.6* DON).

^g FDK, Fusarium damaged kernels on a weight of kernels with Fusarium symptoms as a percent of the total sample weight.

^h FDK, Fusarium damaged kernels on a 1 (low) to 9 (high) scale.

Table 6. Stem solidness^a rating of each of five internodes on plants grown near Swift Current, 2013 to 2015, of AAC Stronghold, durum checks and hexaploid wheat checks.

	Ploidy	Internode 1 ^b	Internode 2	Internode 3	Internode 4	Internode 5
AC Navigator	Tetraploid	2.8	2.6	2.0	1.5	1.3
Strongfield	Tetraploid	3.1	2.7	2.2	1.5	1.3
AAC Raymore	Tetraploid	4.8	4.7	4.6	4.5	4.3
AAC Cabri	Tetraploid	4.9	4.9	4.7	4.5	4.1
AAC Stronghold	Tetraploid	5.0	5.0	4.9	4.7	4.5
AC Abbey	Hexaploid	3.8	3.3	2.8	2.2	1.7
AC Eatonia	Hexaploid	4.4	4.2	3.6	2.5	1.8
Glenlea	Hexaploid	3.1	2.4	2.0	1.5	1.2
Lancer	Hexaploid	4.8	4.6	3.9	2.6	1.8
Lillian	Hexaploid	4.0	3.9	3.4	2.4	1.6
LSD ^c 0.05		1.0	0.9	0.9	0.7	0.5

^a Solidness rated on a 1 (stem cavity hollow and thin walled) to 5 (stem cavity completely filled with pith) scale. Mean of solidness rating based on 10 plants per entry per rep.

^b Internode 1 is basal internode and internode 5 is the neck.

^c Least significant difference, $P \leq 0.05$, includes the genotype by environment interaction variation.

Table 7. End-use suitability^{a,b,c} measured on yearly composites of AAC Stronghold and check cultivars from 2013 to 2015 in the Durum Cooperative Test.

	FN	HVK	Cd	Milling yld	Semo yld	Semo ash	Wht prot	Semo prot	GI	P/L	W	Semo YP	Pasta colour	
	(s)	(%)	(mg kg ⁻¹)	(%)	(%)	(%)	(%)	(%)	(%)		(ergs)	(mg kg ⁻¹)	b*	a*
AC Navigator	395	79.0	232	76.2	67.7	0.67	12.8	11.9	86.1	1.01	217	10.1	64.8	5.0
Brigade	345	73.7	74	74.8	66.2	0.66	12.6	11.6	95.8	0.92	256	10.2	63.7	4.5
Strongfield	345	77.0	78	74.8	66.1	0.61	13.3	12.3	76.0	0.79	194	9.1	63.3	3.7
AAC Cabri	360	81.7	68	75.4	66.7	0.64	13.0	12.0	78.0	0.56	171	10.4	65.7	4.4
AAC Stronghold	388	78.3	81	74.6	66.8	0.65	13.3	12.1	84.9	1.23	222	9.7	65.0	4.4
Std Dev ^d	5		0.001	0.4	0.4	0.006	0.06	0.05	3	0.04	6	0.04	0.3	0.1

^a American Association of Cereal Chemists methods were followed by the Grain Research Laboratory (GRL), Canadian Grain Commission (CGC) for determining the various end-use suitability traits on a composite of 8 to 9 locations each year.

^b FN, Hagberg falling number; HVK, hard vitreous kernel; Cd, grain cadmium; Semo yld, semolina yield; Wht prot, wheat protein; GI, gluten index; P/L and W values determined through Alveograph; Semo YP, semolina yellow pigment; spectrophotometer colour: b*, yellowness; a*, redness on the CIE scale.

^c Means are from 2013, 2014 and 2015 durum composites.

^d Std. Dev. is the standard deviation based on repeated testing of check samples with replicate tests carried out over an extended period of time each season, provided by GRL, CGC.