Canadian Journal of Earth Sciences

A new approach to the opening of the Eastern Mediterranean Sea and the origin of the Hellenic Subduction Zone

Part 2: The Hellenic subduction zone

Journal:	<i>Canadian Journal of Earth Sciences</i>
Manuscript ID	cjes-2018-0315.R1
Manuscript Type:	Article
Date Submitted by the Author:	18-Mar-2019
Complete List of Authors:	Le Pichon, Xavier; Collège de France, Şengör, Celâl; İstanbul Teknik Üniversitesi, İmren, Caner; İstanbul Teknik Üniversitesi,
Keyword:	Hellenic Subduction Zone, Eastern Mediterranean Sea, kinematics, evolution of Anatolia-Aegea
Is the invited manuscript for consideration in a Special Issue? :	Understanding tectonic processes and their consequences: A tribute to A.M. Celal Sengor

SCHOLARONE™
Manuscripts

**A new approach to the opening of the Eastern Mediterranean Sea
and the origin of the Hellenic Subduction Zone**

Part 2: The Hellenic Subduction Zone

Xavier Le Pichon¹, A.M. Celâl Şengör² and Caner İmren³

¹ Collège de France, Aix-en-Provence

² İstanbul Teknik Üniversitesi, Maden Fakültesi, Jeoloji Bölümü ve Avrasya Yerbilimleri Enstitüsü, Ayazağa 34469 İstanbul, TURKEY

³ İstanbul Teknik Üniversitesi, Maden Fakültesi, Jeofizik Bölümü, Ayazağa 34469 İstanbul, TURKEY

Abstract

We discuss the structure of the present Hellenic subduction zone. We show that the present Hellenic subduction zone was formed about 15 Ma when this subduction zone started to consume Mediterranean lithosphere and to form the huge accretionary wedge that covers a great part of the Eastern Mediterranean. We establish that there is independent evidence that the very large Hellenic Trough that it created was formed simultaneously. Shortly before, an 8-10 km thick backstop, that extends 200 km southward, where it presently abuts the African margin, was put into place. We reconstruct the northern margin of the Eastern Mediterranean Sea prior to the Hellenic subduction in a new and independent way. The faults recently identified by Sachpazi et al. (2016a and 2016b) within the Hellenic seismic slab are a key element of our reconstruction. This is because the slab, which is part of the Nubia plate, is rigid and the faults within it coincide with lines of slip congruent with the relative motion of the Aegean block over it. These faults demonstrate that about 400 to 500 kilometers of East Mediterranean lithosphere have been subducted with essentially the same southwestward direction of motion during the last 15 Myr. Our reconstruction shows that before the onset of the Hellenic subduction, the northern margin of the Eastern Mediterranean Sea coincided with a major Jurassic transform fault that limited the Eastern Mediterranean to the north during its formation in Jurassic and Lower Cretaceous time as proposed in part 1. We discuss the implications of this reconstruction on the Neogene evolution of the Anatolia-Aegea block and its geodynamics.

Keywords: Hellenic Subduction Zone, Eastern Mediterranean Sea, kinematics, evolution of Anatolia-Aegea

35 ***Introduction***

36

37 In part 1, we identified long transform faults that frame the Eastern Mediterranean Sea and
38 proposed that they were created during Jurassic and probably early Cretaceous time when
39 this sea was formed as a pull-apart (Figs. 1 and 2). We called the northern boundary the
40 Eastern Mediterranean North Transform Fault (EMNT). A large portion of this boundary was
41 destroyed during late Neogene by the Hellenic subduction which created the Hellenic
42 subduction trough. The subduction trough, best identified by the gravity low it creates, was
43 formed by the flexure of the Mediterranean lithosphere subducted below Aegea. This trough
44 now occupies the whole central part of the Eastern Mediterranean over a width of 1000 km
45 (Fig. 2). The purpose of this paper is to investigate the Hellenic subduction zone since its
46 origin and attempt to reconstruct the EMNT prior to the formation of the Hellenic subduction
47 zone using new data coming from the faults identified on the Hellenic slab by Sachpazi et al.
48 (2016a and 2016b). We establish that this subduction zone started to consume
49 Mediterranean lithosphere about 15 Ma and that at this time the northern limit of the
50 Eastern Mediterranean coincided with the trace of the EMNT that we have proposed in part
51 1.

52

53 Most of the recent interpretations of the tectonics of Aegea have been based upon "the
54 hypothesis that a single subduction zone has been active through most of the Mesozoic and
55 the entire Cenozoic" (Jolivet and Brun 2010). This hypothesis was suggested by tomographic
56 models (e.g. Bijwaard et al. 1998) that appeared to represent a single slab entering the
57 mantle below the Aegean along a total length of at least 1500 km (Jolivet and Brun 2010) or

up to 1500 to 2500 km (van Hinsbergen et al. 2005). Does this hypothesis invalidate the reconstruction of the Eastern Mediterranean we made in part 1?

As stated by Jolivet and Brun (2010), this hypothesis implies that "throughout the Cenozoic, Vardar ocean, Pelagonian microcontinent, Pindos ocean, Apulian continent and the Eastern Mediterranean domain were successively subducted and partly accreted to the southern margin of Eurasia". These domains are not small. The Vardar ocean may have been up to 1000 km wide according to Stampfli and Borel (2002) and the Pindos ocean at least 500 km wide according to van Hinsbergen et al. (2005), comparable to the present Eastern Mediterranean. In addition, the nappes that were previously lying above continental crust imply a total width of subducted continent of about 900 km (van Hinsbergen et al. 2005). These pieces of continent were above sea-level when this subduction occurred since what would become the Aegean Sea only subsided below sea-level in Middle Miocene time (Drooger and Meulenken 1973). Actually, it is most likely that, during Oligocene, the altitude may have been quite significant (see discussion for example in Le Pichon et al. 2002) and consequently the continental crust thickness quite large. This continuous subduction thus alternated oceanic subduction (Vardar, prior to 65 Ma and Pindos between 49 and 35 Ma) with true continental subduction, while the domains subducted had significantly different widths. Such an alternation of oceanic and continental subduction must have implied significant jumps of the actual subduction zone as new deep decollements were successively activated below the thick (of the order of 10 km) detached nappes within the continental crust entering the oceanic subduction zone.

This was certainly the case when the Eastern Mediterranean lithosphere first entered what would become the Hellenic subduction zone after the Apulian (Gavrovo-Tripolitza, Ionian and Plattenkalk nappes) subduction had ceased. The reconstructions made with the single plate hypothesis agree on the fact that the Eastern Mediterranean lithosphere entered the subduction zone about 15 Ma or slightly before and that at this time the southern limit of Aegea along which this subduction occurred was rectilinear (see Fig. 10 B of Jolivet and Brun 2010, Fig. 12 of van Hinsbergern and Schmidt 2012, Fig. 9 of Faccenna et al. 2014). That the southern limit of Aegea was rectilinear sometime after Lower Miocene had been established earlier on the basis of paleomagnetism (e.g. Kissel and Laj 1988). It had been proposed much before by McKenzie (1972) on the basis of seismological considerations and by Garfunkel (2004) on the basis of geological considerations.

This is a very important conclusion. Whether one believes in the single plate subduction hypothesis during the whole Cenozoic or even earlier, or whether one prefers the idea that there has been at least one detachment of slab prior to the formation of the Hellenic subduction zone, as we do and have defended earlier (Le Pichon et al. 2014, 2016), there was no subduction of Eastern Mediterranean lithosphere below Aegea prior to about 15 Ma and the boundary between the Eastern Mediterranean and Aegea was rectilinear. In other words, Aegea during this time belonged to Africa as recognized by paleomagneticians (e.g. Kissel and Laj 2008). Aegea was progressively incorporated in the accretionary wedge of Europe with a north to south migration of the accretion that finally reached southern Aegea in latest Oligocene with the overthrusting of the Tripolitza and Ionian zones. The formation of the huge Hellenic trough is a recent post-15 Ma geological event. The important Africa-

Europe shortening was absorbed to the north of the Eastern Mediterranean and its southern migration only reached northern Aegea after Paleocene.

Actually, recent seismological work appears to us to be much simpler to interpret in terms of detachments and southward jumps of new subduction than in terms of a continuous slab with tears. For example, Hansen et al (2019) show that the present slab below Peloponnesus is not in contact with the lower slab that dips below 200 km toward the northeast. In southernmost Peloponnesus, the present slab is slightly longer and may be in contact with the lower slab but its character is quite different. Continuing eastward the present slab begins a steep concave bend with a radius of curvature decreasing downslope. Bocchini et al. (2018) confirm the spectacular amphitheater shape of the slab below the Aegean that is impossible to put in continuity with a deeper slab below 200 km supposed to dip either to the northeast or to the north (e.g. Biryol et al. 2011). Further east, where the subduction velocity gets much lower, the situation is even more complex. The slabs entering the mantle are separated by important discontinuities opposite Isparta (between 28°E and 29°E) et east of Cyprus (between 32°E and 33°E) (Biryol et al. 2011). There are portions of nearly vertical slabs which join a reservoir of deeper lithosphere that appears to have accumulated in the transition zone, within the lower part of the upper mantle, between 400 km and 650 km. The recent seismological data demonstrate that the present extremely complex geometry of subduction and highly variable subduction rates that range between 35 mm/yr and less than 5 mm/yr are incompatible with a continuous slab at depth. Is there any reason to believe that it was different in earlier times? Would it not be simpler to assume that these highly diverse subductions piled portions of slabs that accumulated within the transition zone before leaking downward to 1000 km and finally deeper as they seem to do elsewhere in the

world. We wish to stress however once more that even if one believes that a single subduction zone has been active since at least upper Cretaceous time, there was no subduction of Eastern Mediterranean lithosphere below Aegea prior to about 15 Ma and the original northern boundary of the Eastern Mediterranean along Aegea at this time was rectilinear.

In this paper, we first discuss the structure of the present subduction zone. We show that the Hellenic subduction zone was formed about 15 Ma and that there is independent evidence that the Hellenic Trough was formed simultaneously. This is the time when the 8-10 km thick backstop, that extends 200 km southward where it presently abuts the African margin, was put into place, as will be discussed later in the backstop section. We discuss the geometry of the seismic slab with its peculiar amphitheater shape that leads to a decrease of the radius of the concavity of the slab as it sinks. The amphitheater shape results in shortening strain along strike with a P-axis parallel to the lines of equal depth. This demonstrates complete decoupling between the slab and the overlying Aegea that is in extension, instead of compression, parallel to the arc.

We then discuss the Sachpazi faults that are collinear with the direction of subduction and consequently coincide with lines of slip. We use these lines of slip to reconstruct in a new and independent way the geometry of Aegea during the last 15 Myr. The result we obtain is essentially similar to those mentioned above using completely different criteria. Finally, we discuss the post-15 Ma geodynamics of Anatolia-Aegea on the basis of these new results.

The Hellenic subduction trough within the EMSZ

152

153 **The geometry of the Hellenic subduction trough**

154 We first consider the Hellenic subduction trough that now occupies a large part of the EMSZ
155 (Figs. 1 and 2). It is best defined by the gravity trough due to the bending of the African
156 plate as it is subducted below Aegea. Although we call it the Hellenic subduction zone, this
157 should not be confused with the Hellenic trench. This so-called trench is a forearc structure,
158 as most of the basin to its south is covered by the huge Eastern Mediterranean Ridge which
159 is an accretionary complex (e.g. Le Pichon and Angelier 1979). Actually, the nose of the
160 subduction zone abuts at the Libyan continental margin, because this is how far the Eastern
161 Mediterranean Ridge extends.

162

163 Note first the striking difference between the northwestern and northeastern limits of the
164 gravity trough. The northwestern limit is abrupt with a large lateral 100 mgal gravity outer
165 high about 130 km wide next to a steep downward gravity gradient. The northwestern
166 extremity of the subducted oceanic plate is limited by the southeastern edge of the Apulian
167 platform. Further northwest, the 20 km thick Apulian continental platform is itself
168 subducted to the northeast but at a much slower rate, 4 mm/yr instead of 34 mm/yr for
169 the oceanic Mediterranean plate (Pérouse et al. 2012). The southeastern limit of the
170 subduction trough is much more subdued with a gravity gradient about five times smaller.
171 And there is no clear gravity outer high. The large gravity high opposite Egypt is due to the
172 uncompensated load of the Nile delta. The subduction decreases to about 12mm/yr east of
173 Antalya and less than 5 mm/yr east of western Cyprus. The decrease in subduction rate is
174 thus much less rapid than to the northwest. The Herodotus abyssal plain is consequently

175 subducted at the relatively low subduction rate of 12 mm/yr, much slower than the 35
176 mm/yr of the Ionian abyssal plain.

177

178 We conclude that the flexure of the part of the plate that is subducted is much steeper to
179 the northwest than to the southeast. This is in agreement with the fact that the
180 northwestern edge of the subducting plate is limited by a tear fault (Spakman et al. 1993),
181 interpreted as a STEP (Subduction-Transform Edge Propagator) fault by Govers and Wortel
182 (2005). The detailed seismological investigation of Pearce (2015) confirms that there is at
183 a minimum a steep ramp between the Hellenic slab and the Apulian slab to the northwest.
184 Royden and Papanikolaou (2011) have proposed that a tear between the two slabs is
185 geologically justified although others argue for a transition through a ramp (e.g. Hansen et
186 al. 2019). The presence of an abrupt transition along the northwestern limit is in contrast
187 with the southeastern limit that is much more gradual. This subduction is asymmetric. To
188 the southeast the Mediterranean plate is bending below the advancing Aegean landmass
189 whereas a tear fault to the northwest let the slab enter into the asthenosphere
190 unconstrained by lateral attachment. Note that the subduction occurs in such a way that
191 only the oceanic crust is being subducted, as the thickly covered Herodotus thinned
192 continental crust (see part 1) does not participate in the fast subduction.

193

194 To the south, the gravity outer high coincides with the EMST. This high is due to the flexure
195 of the subducted plate. It follows the EMST from the Cyrenaica continental margin with
196 which it coincides to the more subduced continent-ocean transition north of the Sirte basin.
197 As a consequence, the whole Cyrenaica margin has been uplifted and is part of this gravity

high. Of course, as Aegea moves southwestward, the Hellenic gravity trough should also move southwestward and one could consider that it just happens that the southern axial high just reached the Libyan margin. However, this is not so as the uplift dates from the Tortonian (between 12 and 7 Ma, Chamot-Rooke et al. 2005), shortly after the beginning of the present Hellenic subduction (Le Pichon et al. 2002, 2016). In addition, the lateral northwestern gravity high coincides with a large topographic high that extends from the Medina Mounts to the Ionian Islands (Fig. 2). This lateral high is also due to the flexure of the Mediterranean plate. The Medina Mounts are situated on the southern extremity of this lateral high (Fig. 2). A thrust fault system is situated on the lateral gravity high. It is remarkable that the thrust faulting also occurred during Tortonian (Gallais et al. 2011). Gallais et al. (2011) related the formation of the Medina high to the initiation of the Calabrian subduction zone. However, the spatial and temporal coincidence with the formation of the Hellenic outer rise leads us to attribute the deformation of both gravity highs during Tortonian to the formation of the Hellenic subduction trough in Serravalian-Tortonian time, although it is quite possible that the initiation of the Calabrian zone increased the amplitude of the flexure.

We conclude that the southern limit of the Hellenic subduction trough already coincided with the EMST during Tortonian. The northwestern limit of the trough, the gravity high that ends to the SSW within the Medina Mounts, was also already established along its present-day limit. This is rather surprising as the edge of the Aegean landmass and consequently the subduction zone was situated in Tortonian about 350 km farther away than it is today, that is about 550 km from the Libyan margin.

221

222 Do we have more information on the formation of the gravity high? Consider the shape of
223 the gravity outer high. The southern high has an amplitude of about 50 mgal and a width of
224 80 km. It appears to be symmetric (Fig. 2). Let us call x_m the abscissa of the maximum of the
225 bulge, and x_{10} the abscissa of the outer limit, x_{20} the abscissa of the inner limit. Using the
226 simplest theory for a thin elastic plate horizontally loaded, in the absence of horizontal
227 loading, the ratio r between the inner and outer lengths of the bulge $r = (x_{20} - x_m)/(x_m - x_{10})$
228 is about 3 (e.g. Le Pichon et al. 1973). The high is elongated away from the subduction zone.
229 Here r is about equal to 1. The bulge is symmetric which corresponds to the buckling case.
230 One would then expect a secondary bulge 160 km further south. There is indeed a secondary
231 high 140 km to the south. The corresponding flexural parameter is 35 km. With usual values
232 for the density of mantle, Young's modulus and Poisson's ratio, it corresponds to a low value
233 of 12 km for the elastic thickness. This suggests that the initiation of the Hellenic subduction
234 began with a strong shortening episode that led to buckling of the Sirte basin lithosphere
235 immediately south of the EMST, before the subduction zone was formed along the northern
236 boundary. This Sirte basin was presumably weaker than the Mediterranean oceanic
237 lithosphere as it had been tectonized during the Cenomanian rifting episode that opened the
238 Sirte basin to the north, as discussed above. This episode presumably preceded the actual
239 beginning of the Hellenic subduction after the northern Hellenic margin failed.

240

241 The larger lateral Medina-Ionian islands gravity high is clearly asymmetric with a ratio r of
242 1.6. Note that the long side of the bulge is to the northwest, indicating that the dominating
243 influence was the bending of the Mediterranean lithosphere to form the Hellenic Trough, not
244 the Calabrian one. The corresponding flexural parameter is equal to 35 km indicating a more

important elastic thickness of 20 km. This is to be expected as the bulge is now entirely within Mediterranean oceanic lithosphere that should be stronger. The ratio r would indicate that a significant horizontal compressive stress component has been acting perpendicular to the bulge, but was not sufficient to produce buckling. Of course, one should be aware of the important limitations related to the thin plate model used. These are simply qualitative indications on the state of stress. Yet, the indications we have suggest that, although both episodes of deformation occurred during Tortonian, the Medina deformation was posterior to the Sirte deformation, as this lateral bulge is best explained in the context of an active Hellenic subduction. When the buckling to the south failed to yield, the subduction zone was created along the Hellenic boundary and the lateral high was formed.

The geometry of the Hellenic margin

We now look at the geometry of the Hellenic margin (Fig. 1). Ryan (1970) proposed that the Hellenic arc and trench system is a consuming boundary. McKenzie (1970, 1972) then proposed that this subduction results from the westward motion of Anatolia-Aegea due to the northward motion of Arabia. The Hellenic scarp has been the leading edge of the Anatolia-Aegea platelet in its escape toward the Central Mediterranean since its initiation during Serravallian-Tortonian time (Le Pichon and Angelier 1979). The role of the Hellenic Scarp is beautifully illustrated in Figure 3 of Floyd et al. (2010). The overall eastward motion of Anatolia-Aegea is accounted for by the present dynamics of this whole block (Özeren and Holt 2010; Özbakır et al. 2017; England et al. 2016). England et al. (2016) in particular have demonstrated that the present dynamics of the Anatolia-Aegea system can be simply accounted for by the distribution of gravitational potential energy within it and by two

boundary forces: a pull applied to the Nubia-Aegea boundary and a push applied to the Arabia-Anatolia boundary. These two forces can be simply predicted by the contrast in lithostatic pressure between the Aegean mass and the adjacent Mediterranean lithosphere and between the high Anatolian Plateau and the adjacent Arabia. We will demonstrate that the Hellenic Scarp at the front of Aegea has been moving within the same kinematic system since the initiation of the motion. This 600 km long rectilinear main Hellenic scarp is oriented N40°W. It is approximately orthogonal to an irregular 400 km long southeast margin. We will see later that this latter margin has been created during Neogene. The subduction direction, as given by the subduction earthquakes, such as the large 2008 Methoni earthquake, is not strictly perpendicular to the Hellenic scarp but makes an angle of 70°-75° with it (Howell et al. 2017a, see Figure 1, see also Shaw and Jackson 2010). On the southeast margin, the subduction direction is much more oblique making an angle of 20-40° with the margin (Shaw and Jackson 2010, see Fig. 1).

In part 1, we presented the Hellenic scarp as a preserved remnant of the EMNT. The fact that it preserved its rectilinearity while the Aegean landmass was strongly deformed behind it by distension indicates that it played a key role in front of this advancing land mass. It plowed ahead while the Aegean basin adjusted behind. As proposed by Le Pichon (1982) and by England et al. (2016) the sharp gradient of potential energy due to the juxtaposition of the dense Mediterranean lithosphere to the Aegean thicker continental crust results in a stress system that pushes the Hellenic scarp outward. Thus, the present southwestward motion of the Hellenic scarp is consistent with the general geometry of the Hellenic trough as described above. The geometry of the gravity lateral high confirms that this southwestward motion has been going on since the initiation of the subduction zone some

time close to Tortonian. An important property of the gravity force acting on the Hellenic scarp is that it maintained its perpendicularity to it as the scarp was rotating clockwise. This is a major difference with the slab roll-back. How can roll-back be the main engine behind the clockwise rotation of the Hellenic scarp as the pull exerted by the long slab assumed to dip to the north should have remained N/S throughout its rollback?

The geometry of the slab and its state of stress

We now look more closely at the geometry of the present slab below the Aegean. Figure 1 compiles various data on the geometry of the slab coming from different sources identified in the caption. The isodepths of the western part of the slab are from Sachpazi et al. (2016a and 2016b). They are much more detailed than those reported in previous studies and show offsets with respect to active fractures that will be discussed later. The isodepths of the eastern part of the slab are from Shaw and Jackson (2010) for the deeper part and from Vernant et al. (2014) for the shallower part. We have made no attempt at reconciling these isodepths but there is broad agreement between them. A first general remark is that the general geometry of the subducted slab is governed by the geometry of the edge of the Aegean landmass below which it subducts. The 90° abrupt turn from the Hellenic scarp to the southeast margin is accommodated by a bend of the slab that gets steeper as the slab deepens.

The increasing curvature with depth should produce shortening strain along strike as pointed out by Shaw and Jackson (2010). As a consequence, the P-axes of the earthquakes within the slab should be parallel with the local strike of the subduction zone. In other words,

the P-axes are parallel to the lines of equal depth of the slab. This was already noted by Taymaz et al. (1990) and was confirmed by the compilation of fault plane solutions published by Shaw and Jackson (2010, see Figure 13). We have added lines of approximate direction of P axis within the slab. They are parallel with isodepth contours and show the same increasing curvature with depth as subduction changes from frontal to the southwest to nearly transform to the southeast. Note in particular the pure thrust earthquake just north of the margin of Africa between faults F10 and F11. This is where the backstop enters into collision with the margin. Yet, the Mediterranean lithosphere immediately below the 8-10 km thick backstop was affected by a large thrust earthquake with a WSW-ENE P-axis. The along-strike orientation of the P-axis indicates that this event was not related to the collision with Africa. Rather, it resulted from the initiation of the circular bending of the Mediterranean lithosphere to subduct below the southwestward advancing Aegea-Anatolia. This is a spectacular confirmation that the deformation of the subducting Mediterranean lithosphere is completely decoupled from the deformation of the overlying Aegean landmass and adjacent backstop. We relate the Tortonian shortening deformation of the Medina outer high to the same pattern of deformation as the one affecting the slab below the Hellenic trough.

The backstop

The backstop is easy to map because of its characteristic reflectivity when it is not covered by thick Mediterranean ridge sediments thrust over it from the south. Figures 1, 2, 3 and 4 show the limit of the backstop proposed by Huguen et al. (2006). Opposite Crete, the backstop is overthrust by the Mediterranean ridge and consequently, its outward limit

cannot be mapped accurately. However, the gravity gradient map suggests that it extends to the base of the African continental margin (dashed black lines). This outward limit has an unusual triangular shaped geometry, with the apex opposite Central Crete. There, its width reaches nearly 200 km. It consists of an 8-10 km thick high velocity (4.8 to 6.6 km/s, Sachpazi et al. 2016b) slab, probably made of limestone, cut by low-angle reflectors that may correspond to overthrusts (Le Pichon et al. 2002). DSDP 377 borehole within a trough near the backstop leading edge drilled through middle Miocene turbidites (between 19 and 15 Ma), probably of African origin, covered by 15 to 14 Ma pelagic marls (Kastens 1991). Kastens (1991) concluded that the accretionary wedge there was not present before 15 Ma, otherwise the turbidites would not have been able to reach this location. She concluded further that the volume and stratigraphy of the wedge which extends south of DSDP 377 and consequently south of the backstop was compatible with an age of 15 Ma. Since then numerous studies have shown that the backstop is overlain by an evaporite Messinian sequence and layered Plio-Quaternary sediments. A thin layer of pre-Messinian sediments underlies the backstop locally. Le Pichon et al. (2002) concluded that the backstop reached its present configuration in upper middle Miocene and that the adjacent accretionary wedge was initiated immediately after, near 15 Ma. Consequently, the subduction that resulted in the formation of this wedge was also initiated 15 Ma.

Le Pichon et al. (2002) noted the similarity of the backstop geometry with the pattern of extension at sea of the Hellenic nappe system proposed by Aubouin et al. (1976). Figure 3 compares the extension of the nappes at sea after Aubouin et al. (1976) with the limit of the backstop. These nappes traveled outward during Lower Middle Miocene. The proposal of Aubouin et al. (1976) implied that the outer nappes migrated beyond the continental margin

between 20 and 15 Ma to form the present backstop, just before the initiation of the present subduction zone. Figure 3 also shows the Aegean extensional areas and their development since the early Miocene after Chamot-Rooke et al. (2005). The general agreement in geometry and timing of this outward tectonic migration is striking and is in favor of the origin of the backstop by overboard migration of part of the Gavrovo-Tripolitza, Ionian and Plakenkalk nappes involved in the Apulian subduction during early to medial Miocene time. In the recent geological interpretations, the most recently exhumed unit is the Phyllite-Quartzite unit which was exhumed 15 Ma (e.g. Jolivet and Brun 2010).

There is no continuity between the northern edge of the backstop and the Hellenic Scarp. Shaw et al. (2008) and Howell et al. (2017b) show that a backthrust dipping northeastward 30° marks the contact between the backstop and the Hellenic Scarp and that rare very large earthquakes along this fault are responsible for rapid uplift of the southern coast of Crete. A similar process was proposed to explain the uplift of Rhodes. Howell et al. (2017b) generalize this proposal. They describe a band of crustal thickening inland of the Matapan, Hellenic and Strabo trenches that is produced by this backtrusting. Le Pichon and Angelier (1981) had earlier proposed that present crustal thickening below Crete was due to scraped sediments from the accretionary wedge.

The presence of the backstop might be expected to impose strong constraints on the geometry of subduction but the shape of the gravity trough shows that this is not so (Fig. 2) as the outward limit of the gravity trough is parallel with the Hellenic scarp, not with the backstop limit. This is because the thickness of the backstop is only 8-10 km whereas the

thickness of the Hellenic crust is at least three times more. Thus, the backstop only influences the iso-contours of gravity in the immediate vicinity of its outward limit.

The Sachpazi faults

We now consider in this context new enigmatic tectonic objects that have been discovered by Sachpazi et al. (2016a and 2016b). Using teleseismic receiver function on a dense 2D seismic array, Sachpazi et al. (2016a) showed that the slab below Greece, between Kythira island to the south and Corinth to the north, over a length of 300 km is segmented by nine along-dip faults (Figs. 1, 2 and 4) with an average separation of 35 km for F1 to F6, 50 km for F6 to F9. In addition, the isodepth lines are dextrally offset along the faults. Although there is significant scatter in the amount of horizontal offset, which is of the order of 10-30 km, it tends to decrease as the slab deepens. These faults are confined to the slab although there is one case where a fault broke within the overlying crust just over fault F2 during the 2008 Movri earthquake (Sachpazi et al. 2016a; Serpetsidaki et al. 2014). However, in this case, the slab is affected by scissor faulting, whereas the faulting in the overlying crust is right-lateral strike-slip. We propose that this is because the slab is affected by vertical shear strain as it subsides below the overriding lithosphere. The overriding lithosphere instead is affected by horizontal shear strain in its southwestward move. There is a complete disconnection between the two strain systems.

Sachpazi et al. (2016a) show three sections perpendicular to the fractures that intersect faults F5 to F8 at eight locations. The surface of the slab is vertically offset along the faults, the southeast side being about 8 ± 3 kilometers down with respect to the northwest one.

408 There is no detectable systematic difference in vertical offset between the fractures in this
409 area where the dip of slab is everywhere close to 13-15°. The vertical offset V_o is directly
410 related to the horizontal offset H_o and to the dip D ($V_o = H_o \tan D$). This relation predicts that
411 the average observed vertical offset of 8 km and dip of the slab of 15° should correspond to
412 a horizontal offset of 30 km. As the depth of the top of the slab increases to the northeast to
413 reach 70 km, the dip abruptly increases to 30-50° and the vertical offset would be expected
414 to increase again for the same offset. We propose that this is the cause of the increase in
415 seismic activity that Sachpazi et al. (2016a) observed along the faults as their dip increases
416 beyond 70 km depth. Sachpazi et al. (2016a) reported further three thrust earthquakes that
417 occurred on the fault interfaces beyond 60-70 km depth of the slab in the area of sharply
418 increasing dip from 15° to 30-50°. These earthquakes show that the southeast side
419 underthrusts the northwest one and they confirm the state of compressive stress along
420 strike within the slab described by Shaw and Jackson (2010).

421
422 Sachpazi et al. (2016b) using 4 OBS's to extend the land network at sea have shown that the
423 fractures extend offshore below the 8-10 km thick backstop to its outer limit. The addition
424 of OBS's enabled them to present one additional section perpendicular to the fractures over
425 the backstop. The vertical offsets along this section are three times smaller than on the
426 adjacent continent. The combined offset is 5 km over F5-F6 and also 5 km over F7-F8,
427 indicating an average 2.5 km vertical offset over this section below the backstop, compared
428 to an average 8 km offset below the continent to the northeast. This result implies that the
429 offset is progressively created as the slab increases its dip to underthrust the 40 km Hellenic
430 crust.

431

432 Sachpazi et al. (2016b) also suggested that the faults terminate seaward below a dextral
433 offset of the outer limit of the backstop. On the map of Huguen et al. (2006, see Figure 1), this
434 is true for fault F5, possibly so for faults F6 and F7, although the amount of offset and its
435 direction do not show an agreement with those on these faults. No offset can be detected for
436 the other faults. The resolution of mapping is not sufficient to exclude the presence of other
437 as yet undetected offsets. Yet, the available data do not prove that there is a systematic
438 correspondence between dextral offsets of the backstop outer limit with horizontal offsets
439 along the Sachpazi faults and we see no reason why it should be so. The increase in dip of
440 the slab occurs when it approaches the thicker crust of the Hellenic Scarp. Yet there is no
441 detectable offset of the Hellenic Scarp. The faults are confined to the slab. They only affect
442 the overlying crust indirectly through the vertical offset created.

443

444 Finally, another important property of these faults is that they are collinear with the
445 subduction direction as can be seen in the case of the large subduction 2008 Methoni
446 earthquake. Sachpazi et al. (2016b) showed that the Methoni earthquake was initiated along
447 fault F6 and that this fault exerted a major control on the delimitation of the zone of rupture
448 that it limits to the northeast. The Methoni slip vector on Figure 1 coincides with the trace
449 of fault F6. We conclude that the segmentation of the slab follows slip lines.

450

451 Let us recapitulate at this stage. The Sachpazi faults are scissor vertical faults along slip lines
452 offsetting vertically the slab but they do not extend within the overlying continental crust.
453 They are initiated below the backstop, in the general area of large inter-plate subduction

20

454 earthquakes and the vertical offset increases as the slab begins to dip to underthrust the
455 Hellenic continental margin. Below the Hellenic continental crust, with slab depth between
456 40 and 70 km, the dip of the slab stays close to 13-15° and the vertical offset stays close to 8
457 ± 3 km. The activity of the faults resumes sharply beyond 70 km depth, because of the rapid
458 increase there in the dip of the slab. The spacing between faults is quite regular, 35 km for
459 F1 to F6, 50 km for F7 to F9, suggesting a mechanical control of the spacing. The faults
460 segment the slab in slices that have the same geometry, but are offset geometrically. The
461 offsets compensate the differential advance of the Aegean lithosphere on top of it. The
462 velocity of the Aegean lithosphere increases toward the southeast whereas a rigid rotation
463 would imply a decrease. This results in right-lateral horizontal shear strain. Being rigid, the
464 African lithosphere is not submitted to horizontal shear strain but instead must adjust to the
465 vertical strain due to the increasing velocity of the Aegean lithosphere to the southeast.
466 Instead of adjusting continuously to the resulting gradient in vertical velocity, is cut by
467 regularly spaced vertical tear faults that appear whenever the vertical shear strain between
468 slices exceeds a certain value. This can be understood as a variation of compartmental
469 faulting. One can then understand why the crustal fault triggered over fault F2 during the
470 2008 Movri earthquake (Sachpazi et al. 2016a; Serpetsidaki et al. 2014) was right-lateral.
471 Whenever the vertical offset of the Sachpazi faults initiate a fault within the overlying crust,
472 they localize the horizontal shear strain within it and initiate right-lateral strike-slip faulting.
473 The faulting within the Sachpazi faults is similar to the one within the STEP (Subduction-
474 Transform Edge Propagator) faults of Govers and Wortel (2005), both are scissor faulting,
475 but they have a different origin.

476

477 **The kinematic significance of the Sachpazi faults**

The Sachpazi faults have an obvious kinematic significance, because they are planes of slip between the subducting Mediterranean slab and the overlying Aegean landmass. The strikes of these faults correspond with the direction of motion of the southwestward overthrusting landmass with respect to the slab. Similarly, the striations and grooves left by a glacier give the direction of motion but not the relative velocity, because the glacier significantly deforms during its motion. For example, the lateral parts of a glacier are moving slower than the medial part. In the case of the Aegean landmass, it is clear that the velocity of slip is not governed by a rigid rotation as it has been rapidly deforming in oblique transextension since the middle of the Neogene (e.g. Dewey and Şengör 1979; Le Pichon and Angelier 1979; Şengör et al. 1985; Şengör 1987; Chamot-Rooke et al. 2005; Jolivet and Brun 2010; Royden and Papanikolaou 2011; see the reconstruction of van Hinsbergen and Schmid 2012 and the discussion of Le Pichon et al. 2016).

We know further from paleomagnetic studies that the area between the Scutari-Pec fault near 42°N in northern Albania and northern Peloponnesus, Western Greece and Evia, north of faults F6-F7 (Fig. 1) was affected by clockwise rotations of about 40° between 15-13 and 8 Ma (Kissel et al. 2003; van Hinsbergen et al. 2005). The analysis of van Hinsbergen et al. (2005) excludes the possibility of significant paleomagnetic rotations there before 15 Ma. These rotations may have extended over southern Peloponnesus between faults F6-F7 and fault F9, but in the absence of suitable middle Miocene sediments this cannot be established (van Hinsbergen et al. 2005). The 15-13 to 8 Ma rotations occurred contemporaneously with motion along the detachment systems of the Cyclades and of the Rhodope (van Hinsbergen et al. 2005; Jolivet et al. 2015). An additional clockwise rotation of about 10° was detected by paleomagnetism during Plio-Pleistocene time (van Hinsbergen et al. 2005). Finally,

geodetic data show that the whole area north of fault F5 is now rotating clockwise with a rotation rate larger than $2^{\circ}/\text{Myr}$, the rotation rate increasing to a maximum along an axis corresponding to the prolongation of the North Anatolian Fault where it exceeds $5^{\circ}/\text{Myr}$ (Chousianitis et al. 2015; see also Le Pichon and Kreemer 2010; Floyd et al. 2010; also see Şengör and Zabcı in press). The rotation rate progressively decreases southward and becomes insignificant south of fault F8. Since 13-15 Ma, clockwise rotation of the whole area below which the Sachpazi faults are now situated, appears to have continuously operated, the present rotation rates being quite large north of fault F5 and decreasing progressively southward to disappear south of fault F9. Crete has not been affected by any clockwise rotation during the 15 Ma – Present interval (Van Hinsbergen et al. 2005). The present distribution of clockwise rotations within Aegea demonstrates that these do not correspond to a single rigid rotation. Yet, we are faced with this paradox that these clockwise rotations occur in the framework of a general anticlockwise rotation of Anatolia-Aegea. How is it possible to have simultaneous clockwise and anticlockwise rotations affecting Aegea?

Le Pichon et al. (2016) have pointed out there was no plate boundary north of Aegea during most of the Anatolia-Aegea motion. This is because the North Anatolian Fault (NAF) propagated westward since about 11 Ma (Şengör et al. 2005) and has not yet reached the northern extremity of the Hellenic subduction whereas the Hellenic subduction had been initiated approximately 15 Ma ago. Where the NAF was not yet present, the shear strain due to the motion of Anatolia-Aegea had to be accommodated over a wide strip of Western Anatolia-Aegea. As the NAF progressed westward, the zone of shear strain progressively became narrower and more concentrated. Two million years ago, the NAF reached the North Aegean Trough and a narrower $\sim 100\text{-}200$ km shear zone, called Grecian Shear Zone (GSZ)

by Şengör (1979), appeared along its prolongation across Greece in the area of the Gulf of Corinth, with the formation of the Gulf of Corinth and the Gulf of Evia. But the shear is not limited to the Grecian Shear zone as clockwise rotation extends as far south as Kythira as discussed above. Le Pichon et al. (2016) noted further that the ductile state of the lower crust and upper mantle of Western Anatolia-Aegea facilitated the distribution of shear strain.

We next investigate whether the necessary distribution of shear strain due to the absence of plate boundary is obtained through non-rigid clockwise rotation extending from the Scutari Pec fault, in Albania, near 42°N, to Fault F9, near Kythira island. This clockwise rotation would have to be superposed to the anticlockwise southwestward progression of Aegea-Anatolia with respect to Nubia with a pole of rotation situated to the southeast of the Mediterranean Sea. The anticlockwise Anatolia-Aegea rotation is governed by the traces of the Sachpazi faults, which determine a pole compatible with the Anatolian motion to the east. The superposed non-rigid clockwise rotation should increase the expected velocity southward, introducing a shear strain gradient. This gradient imposes to the rigid subducted plate a gradient of vertical subsidence that is absorbed along regularly spaced scissor faults. Thus, each slice of slab between faults would have above it Aegean crust with an overthrusting velocity with respect to the expected Anatolian-Aegean velocity, larger than the one immediately to the north. This northward decrease in velocity is the cause of the horizontal offset. We should then expect the horizontal offsets to increase as the slab gets deeper because the offsets are the product of the increase in velocity by the subduction time length.

In addition, the edge of the Greek continent has maintained its rigidity behind the Hellenic Scarp and the slab depth is in great part controlled by the part of the subduction path along which it was in contact with the continental crust, from the margin to the deepest Moho depth, before entering the asthenosphere. Consequently, one should expect the isodepth lines of the slab to keep memory of their passage below the 40 km deep continental crust and their orientations should reflect the clockwise rotation of the Hellenic scarp. We should therefore expect a clockwise rotation from the deeper isodepth lines (90-100 km) to the shallower one (40 km just below the Hellenic scarp area).

The Hellenic Scarp pole of rotation

We first test whether the geometry of the Sachpazi faults corresponds to the traces of a rigid rotation, in other words whether the traces of these faults are small circles about the same pole. We fit the faults as given by Sachpazi et al. (2016a, thick continuous white lines in Figs. 1 and 2) with their probable prolongations over the backstop as given by Sachpazi et al. (2016b, thick dashed white lines in same figures). We obtain a pole of rotation situated at 30.3°N, 31.4°E, near Cairo. The corresponding small circles are shown on the different faults. The agreement is good and confirms that these slip lines are governed in a first approximation by a rigid rotation. We call this pole the HS/NU pole because it governs the slip lines of the Hellenic Scarp with respect to Nubia (HS/NU).

Figure 4 investigates whether the Sachpazi faults can be independently detected below the backstop using the gravity gradient map of Sandwell (Sandwell et al. 2014). We show the mapped faults and the small circles fitted to them on each side of the backstop. Although one

is looking there at something that happens 8-10 kilometers below the sea-floor and corresponds to a relatively modest vertical offset of 2-3 km diminishing outward, there is clear indication of segmentation along a general N30°E direction that is the direction of the faults. This strongly strengthens the proposal of Sachpazi et al. (2016b) that the faults extend below the backstop. It also strengthens the confidence we can give to the determination of this pole of rotation, as the directions of the faults below the backstop are significantly different from the directions of the faults below the continent. The oblique Mercator projection of Figure 5 illustrates the fact that the faults are small circles about this pole as it shows that they do follow lines of latitude in this oblique projection. It is interesting that the prolongations of the faults within Aegea appear to follow discontinuities in the pattern of anomalies suggesting that the extension there occurred along the same direction. We will come back to this point later.

We have used the gravity maps (Figs. 2 and 4) to propose the possibility of three additional faults, that we named F10, F11 and F12, beyond the area investigated by Sachpazi et al. (2016a and 2016b). These are obviously quite hypothetical. The absence of significant clockwise rotations south of fault F9 indicated by paleomagnetic data as well as by GPS suggests that segmentation should not be present south of fault F9. As these hypothetical faults follow flow lines, confirming that the overall motion becomes mostly transform to the east, we believe that they exist as structural elements, although they are not the same type of faults as the Sachpazi faults. The exact nature of these faults cannot be solved with the information we have.

The HS/NU pole of rotation we just obtained is close to the Anatolia/Nubia (AN/NU) pole of rotation (32.71°N , 34.2°E , $1.29^{\circ}/\text{Ma}$, Le Pichon and Kreemer 2010). The slip lines do not give any information on the velocity. However, we know from GPS that the subduction of Nubia below Aegea along the Hellenic Scarp is close to 35 mm/yr. We attribute this velocity to the Methoni earthquake location, at the level of fault F6, sufficiently south not to be in the rapidly rotating area (Fig. 1). The corresponding angular velocity is $1.78^{\circ}/\text{Ma}$. Such a pole of rotation, 400 km to the southwest of the AN/NU pole reorients the Aegea motion to the southwest instead of southward if Aegea was strictly following the Anatolian motion. In addition, it increases the velocities. For example, at the Methoni earthquake location, the velocity predicted by the AN/NU Euler vector would be 27 mm/yr instead of 35. The differential motion between the Hellenic Scarp motion and the Anatolian motion produces an E/W extension of 12-13 mm/yr in Central Aegean. This extension is distributed in a complex way northeast of the scarp, all the way to West Anatolia. We conclude that this result is geologically quite reasonable.

The clockwise rotation

The first test is whether the geometry of the slab tells us something about the existence of a clockwise rotation that operates simultaneously with the counterclockwise HS/NU rotation. As stated above, the Hellenic scarp area keeps its rigidity as it rotates. As the slab depth should keep a memory of its subduction below it, one should expect the isodepth lines of the slab to show a clockwise rotation from the deeper ones (90-100 km) to the shallower one (40 km just below the Hellenic scarp area). Let us consider the isodepth lines of Sachpazi et al. (2016a) northeast of fault F6. The F6 limit is adopted because the isodepth lines are

rectilinear to the northeast of F6 whereas they are highly curved to the southwest of F7. In addition, we know from geodesy but also from paleomagnetism that the clockwise rotation is quite large northwest of fault F5 and inexistent southeast of F9.

There is obviously a lot of noise in the mapping data of Sachpazi et al. (2016a). However, one observes a southeastward fanning of the isodepth lines (Fig. 2b of Sachpazi et al. 2016a, see Fig. 1). The 90-100 km isodepth lines make an angle of about 15° with the perpendicular to the faults, whereas the 40 km isodepth line makes an angle of about 35° with it. We do find consequently the expected clockwise rotation instead of the anticlockwise one predicted by the HS/NU pole of rotation. This clockwise rotation from 90-100 km isodepth to 40 km isodepth is about 20° .

The length of slab between the 40 and 90-100 km isodepth lines is 180 km at the level of F6. At the present 35 mm/yr rate of subduction, it takes 5.15 Myr to produce 180 km of subduction. This is a minimum time interval because, as will be discussed later, the rate of subduction had to increase through time. Thus, the rate of rotation is about $3.9^\circ/\text{Ma}$. We can compare it with the clockwise rotation rate necessary to produce the present decrease of velocity from 35 mm/yr at F6 to 26 mm/yr at F2 (Pérouse et al. 2012) instead of the 38.5 mm/yr predicted by the HS/NU Euler vector. The decrease in velocity of 12.5 mm/yr requires a clockwise rotation rate of $5.7^\circ/\text{Myr}$, somewhat larger but of the same order of magnitude. We know from geodesy that at present the rotation rate progressively increases from about 2° at the level of F5 to more than 5° at the level of F2. This is consequently a reasonable approximation. We stress here that although we can combine the AN/NU

rotation with this clockwise rotation, the resulting instantaneous pole of rotation changes with time because the clockwise rotation is attached to the Hellenic Scarp and moves with it. Even with a single rotation for the whole area, the corresponding motion is non-rigid. And as we know that the rate of rotation decreases southward, this non-rigidity becomes evident. The combination of this clockwise rotation to the HS/NU rotation results in an instantaneous pole situated to the northwest that increases velocities southeastward along the subduction zone (25 mm/yr at F2, 34 mm/yr at F6).

The horizontal offsets

The second test is whether the horizontal offset between slices decreases with depth as it should. Again, there is a lot of noise in the Sachpazi data. But their mapping of the offset does show a progressive decrease from about 20 km at the 40 km isodepth line to a value of 10 km or less at the 80-90 km isodepth lines. Another way to look at it is to use the obliquity of the isodepth lines with the perpendiculars to the faults obtained earlier to predict the total offset between F6 and F1. We find a total horizontal offset of 85 km at the level of the 40 km isodepth line (average offset 17 km) and 32 km at the level of the 90-100 km isodepth lines (average offset 6 km). The increase in total offset of 53 km was obtained in 5.1 Myr. This gives a decrease in velocity between F6 and F1 of 10 mm/yr compared to the decrease of 12.5 mm/yr that we obtained above for the present situation. This is again an encouraging result.

A schematic reconstruction of the pre-Hellenic subduction stage (15-13 Ma)

The next test is a schematic reconstruction of the Pre-Hellenic subduction stage based on the kinematics we just discussed. Although the tectonic evolution involved a considerable amount of deformation, we use three successive rigid rotations. This quite crude geometric reconstruction (Figs. 6 and 7) allows us to illuminate the three basic elements of the geologic history, the HS/NU anticlockwise bulk rotation of the Aegean buttress, the additional simultaneous clockwise rotation of the buttress, and finally the clockwise rotation of the Albania-Greece coastline about a pole near the Scutari-Pec fault. The justification for this reconstruction is the observation that the Hellenic Scarp margin has not been significantly modified and consequently has kept its rectilinearity in a first approximation, as discussed earlier.

First, we apply to the Hellenic Scarp south of fault F4 (line CD') the clockwise rotation with respect to pole HS/NU that puts the outer backstop limit C'D' in contact with the extremities of faults F5 to F7. The necessary angle is 18° which is equivalent to 10 Myr at the present rate of rotation, or about 350 km along fault F6.

Second, we apply a 40° anticlockwise rotation to the southern Albania-Northern Greece coastline (GD in Figure 6) about a pole situated on the Scutari Pec fault (GR/EU, G for short in Figure 6) to resorb the 40° clockwise rotation detected by paleomagnetism. This second rotation is indeed the one necessary to maintain continuity between GD and DC.

684 Third, to simulate the clockwise rotation of the Hellenic margin, we choose a pole of rotation
685 at point D (37.4°N, 21.7°E) on the coastline at the northern limit of the zone of clockwise
686 rotation and we assume a single finite rotation with respect to this pole. The location of D is
687 close to the 40 km isodepth curve of the slab and thus corresponds approximately to the
688 southwestern limit of the deep Moho area of Greece. Strictly speaking, this rotation should
689 be applied simultaneously with the AN/NU rotation as the position of its pole of rotation,
690 linked to the Hellenic margin, is also displaced by the AN/NU rotation. For simplicity, we
691 choose instead to apply these rotations successively. We apply the anticlockwise rotation
692 (37°) that puts the Hellenic margin CD' in line with the Albania-Greece coastline after the
693 18° AN/NU clockwise rotation has been applied. The final result is the same (see discussion
694 below). Note however that the slip lines shown in Figure 7 are not those that would be
695 predicted by the summed finite rotation. The 37° necessary for the fit are in fair agreement
696 with the paleomagnetic results. Finally, we restore Anatolia in its pre-NAF position (6.8°
697 clockwise with respect to AN/NU) pole.

698
699 To recover the actual slip lines, consider what happens when effecting simultaneously the
700 HS/NU and the D rotations. D being the center of rotation, the trajectory of point D is
701 predicted by the HS/NU anticlockwise rotation. Moving away from point D to the southeast,
702 the actual trajectories deviate more and more from those predicted by the HS/NU pole.
703 However, this deviation is small within the area covered by the Sachpazi faults, as
704 demonstrated in Figure 7. We emphasize that even this highly simplified reconstruction
705 implies a very large amount of deformation by extension in the trail of the Albania-Northern
706 Greece rotation and of the Hellenic Scarp rotation, as originally suggested by Dewey and
707 Şengör (1979).

708

709 We propose that this reconstruction applies to the pre-Hellenic subduction state. We
710 mentioned above that the present subduction rate is larger than it was in the past. Thus, the
711 age to which this reconstruction applies is at least 10 Ma but probably more. This is first
712 because the slip rate along the NAF must have increased with time because the total offset
713 along the fault is much smaller than the offset predicted with the present rate of slip. Şengör
714 et al. (2005) have proposed that the rate of slip increased approximately linearly with time
715 since 11-10 Ma. Second, the geological evidence indicates that the subduction zone was
716 initiated a few millions of years before the NAF (15 instead of 11-10). We use a simple
717 scheme to estimate the increase of subduction rate with time. We assume initially a
718 subduction rate equal to the present relative motion of the Hellenic Scarp with respect to
719 Anatolia, which is 13 mm/yr of extension, and we assume further a linear increase with time.
720 This gives an average subduction rate of 24 mm/yr and a total subduction time of 15 Myr, to
721 produce 360 km, the approximate length of fault F6. This age of duration of the subduction
722 fits the geological constraints.

723

724 A first important result of this reconstruction is to confirm that the northern margin of Aegea
725 about 15 Ma had a smooth curving shape. We pointed out in part 1 and in the introduction
726 that recent reconstructions, obtained in a completely different way, led to the same result
727 (see section *The Apulia Block and the northern EMNT*). The common conclusion since
728 McKenzie (1972) is that the arc running from the coast of Albania to Turkey had a “simple
729 shape that suggests that it formed the boundary of a single plate before this period”. We
730 agree and attribute this simple shape to the transform (EMNT) which limited the

731 Mediterranean to the north. The transform fault had maintained its integrity between Lower
732 Cretaceous (120 Ma) and Middle Miocene (15 Ma). We have to note here that Garfunkel
733 (2004) had proposed the existence of a transform fault acting along the northern margin of
734 the Eastern Mediterranean during its Lower Jurassic to earlier formation exactly along what
735 we now call the EMNT east of Apulia. He also assumed that the transform fault was destroyed
736 by the Hellenic subduction after 15 Ma. However, to the west, he assumed that this transform
737 connected with what we called in part 1 the Apulian Transform, instead of prolonging it
738 northwestward as we do.

739
740 This is indeed the second major result of this reconstruction: the trace of the EMNT that we
741 obtained in the first part of this paper coincides with the reconstructed boundary (see Fig.
742 6). This corroborates our hypothesis for the location of the EMNT and confirms the
743 conclusion we reached earlier that the Hellenic scarp is a preserved remnant of this
744 transform.

745
746 A third striking character of the reconstruction is the asymmetry of this geologic
747 transformation. This is because a single break occurred along the EMNT between the
748 southern extremity of the Hellenic Scarp (C in Figure 6) and Antalya (B). As a result, the 1000
749 km length of EMNT from the Scutari-Pec fault (G in Figure 7) to Antalya (B in Figure 7)
750 rotated southwestward in a first approximation about a pivot situated near the Scutari Pec
751 fault. But there was an important difference between the northwestern part facing the
752 Apulia block and the southeastern Hellenic Scarp facing the deep Mediterranean. The
753 Hellenic scarp part advanced through subduction of oceanic Mediterranean lithosphere

while the northeastern part advanced through subduction of the thinned Apulian continental lithosphere. The oceanic subduction was going at an average velocity of about 30 mm/yr while the continental subduction rate varied between 20 mm/yr to the southeast and 0 mm/yr near point G, averaging 10 mm/yr. The oceanic subduction maintained its steady progression southwestward, approximately along the limit between the Herodotus-Levant very thickly sedimented thinned continental crust basins and the Central Mediterranean oceanic basins (see part 1). The continental subduction instead decreased rapidly northeastward to zero. The two systems were completely different and a major structural discontinuity separated them. The Hellenic scarp motion was powered in great part by the sharp gradient of potential energy from the continental lithosphere to the Mediterranean one. No such large gradient existed to the northwest to help in forcing the subduction of the Apulian thinned continental crust.

Finally, we emphasize a remarkable consequence of this asymmetry: the margin between C (the southern extremity of the Hellenic Scarp) and B (Antalya) was created during the last 15 Myr. It is the product of extreme extension and anticlockwise rotation. These extreme extension rates are a common result of all recent reconstructions (e.g. van Hinsbergen and Schmid 2012). And paleomagnetism has indeed detected a phase of rapid anticlockwise rotation in southwestern Anatolia, west and northwest of Antalya, about 20° between 16 and 5 Ma (van Hinsbergen et al. 2010). This anticlockwise rotation is contemporaneous with the 15-13 to 8 Ma western clockwise rotation that has been related to the motion along the detachment systems of the Cyclades and the Rhodope (van Hinsbergen et al. 2005; Jolivet et al. 2015) as was pointed out earlier. In addition, Jolivet et al. (2015) emphasized that during the same period granitoids migrated within Aegea from east to west indicating that a lot of

778 crustal material was remelted. All of this is compatible with extreme thinning propagating
779 diachronously from NE to SW to the west and NNE/SSW to N/S to the east.

780

781 ***The evolution of Anatolia-Aegea***

782

783 Anatolia-Aegea is probably the best studied example of a newly formed continental block
784 (less than 15 Ma) that is still in rapid evolution. Following Ketin's initial usage in 1948, we
785 use the term block instead of plate because plate implies internal rigidity, whereas Anatolia-
786 Aegea suffered significant deformation within its Aegea-Western Anatolia portion.
787 Alternatively, Dewey and Şengör 1979, used the German term *Scholle*, long in use in Central
788 Europe for the same purpose, to emphasize this. The absence of internal rigidity is
789 corroborated by the fact that England et al. (2016) have been able to account for the present
790 dynamics of this Anatolia-Aegea system by considering it as a thin fluid sheet.

791

792 An important question is whether the block appeared simultaneously with its boundaries or
793 did it precede them? We show next that the Anatolia-Aegea block acquired a kinematic
794 identity before it had acquired permanent plate boundaries. By kinematic identity we mean
795 that the motion of the block with respect to adjacent lithospheric blocks can be described
796 fairly accurately by Euler vectors. By boundaries, we mean relatively narrow (a few tens of
797 kilometers at most) plate-boundary-scale zones of shear that absorb the relative motions
798 with respect to the adjacent blocks. The oblique Mercator maps of Figures 5 and 8 beautifully
799 illustrate what we mean by kinematic identity as they show the well-defined pattern of

increasing southeastward extension within Aegea along the general direction imposed by the rotation of the Hellenic buttress. In this projection, the western limit of the Anatolian shelf is parallel with the Hellenic Scarp. The amount of extension between the two limits increases from the top to fault F10. This results in a component of dextral shear strain which is absorbed along a few localized faults, such as those proposed by Brun et al. (2016) and by distributed clockwise rotation. Note on the bathymetry of Figure 8 that the extensional basins behind Crete appear to be limited by what appears to be E-W segments in this projection. Note also the evidence for numerous structural trends parallel with this direction along the easternmost portion of the Hellenic subduction zone. These latter trends correspond there to sinistral and not dextral faults because they absorb differential motion between Aegea and the deep Mediterranean basin.

The present kinematics

We know from GPS studies that the motion of the Hellenic Arc with respect to Nubia is close to 35 mm/yr whereas the motion of the Cyprus arc is only 5 mm/yr opposite Cyprus and increases rapidly westward to more than double at its western end. This difference of motion is accompanied by important internal deformation within Aegea. We show here that our interpretation in terms of a rotating rigid Hellenic Slab buttress is compatible with the description of the present kinematics within Aegea based on extensive GPS surveying. A remarkable result established by the geodetic surveys of Aegea is the existence of an Aegea block which is not affected at present by significant deformation. This block is limited by the two shear zones (GSZ and WASZ) mentioned earlier (Reilinger et al. 2006; Floyd et al. 2010; Le Pichon and Kreemer 2010; Royden and Papanikolaou 2011). Le Pichon et al. (2016)

823 discussed the tectonic significance of this undeforming Aegea block. Although rapid
824 extension within the present Aegea block was very important between Serravallian (14-12
825 Ma) and Messinian (7 to 5 Ma), it ceased to exist there after Messinian time (5 Ma) while
826 extension continued to the east within the WASZ, to the west within N/S grabens cutting
827 through Peloponnesus and was initiated to the north within the GSZ between 2 and 1 Ma ago
828 (Armijo et al. 1996; Sorel 2000; Flotté 2003). Le Pichon et al. (2016) attributed this cessation
829 of extension about 5 Ma within the present Aegea undeforming block to the shielding effect
830 of the cold oceanic Ionian slab progressing below the Aegean lithosphere. In addition, the
831 propagation of the NAF within the northern Aegean Sea after 2.5 Ma must have definitely
832 stabilized the cessation of extension, as the NAF localized the shear strain. England et al.
833 (2016) propose that the presence of the undeformed Aegea block is simply the consequence
834 of the inverse relationship between the viscosity of the lithosphere and the deviatoric stress.
835 However, this is a problem similar to the discussion of the anteriority between the hen and
836 the egg. Why did the deviatoric stress reach a very low value below most of southern Aegea
837 5 Ma? Whatever the cause (s) for the formation of the undeformed Aegea block, its presence
838 after 5 Ma led to a redistribution of the deformation within Aegea that we now discuss.

839
840 A difficulty in discussing the present-day kinematics comes from the fact that many of the
841 GPS measurements have been made within this Aegea block. Most of the discussions of the
842 kinematics of Aegea ignore the deformation presently going on west of this block and they
843 identify the subduction motion with the rigid Aegea motion with respect to Nubia, which is
844 not correct. A notable exception is the study of Vernant et al. (2014) who identify smaller
845 blocks west of the Aegea block. We adopt their description of the motion of Aegea. To
846 describe the present motion between the Hellenic Scarp Buttress and Aegea, we use the

847 simple kinematics obtained in the paragraph *The permanency of the kinematics of Anatolia-*
848 *Aegea with respect to Nubia and Eurasia*. We then compute the Aegea/Anatolia motion and
849 the Hellenic Scarp Buttress/Aegea motion. Since about 5 Ma, the Hellenic Scarp
850 Buttress/Anatolia motion that we described earlier adapted itself to the presence of the
851 undeformed Aegea block by distributing the motion to the west and east. We determine
852 these two motions below.

853

854 We first use the motion of the Aegea block determined by the GPS study of Vernant et al.
855 (2014) to compute the Aegea/Anatolia differential motion using the kinematics of Anatolia
856 of Le Pichon and Kreemer (2010). The euler pole is at 38.37°N, 18.95°E, with a rate of
857 rotation of 1.42°/Myr. This produces left-lateral N/S motion moving the Aegea block
858 southward of Anatolia at a rate of about 27 mm/yr. As a first approximation, the WASZ now
859 acts as a left-lateral shear zone that accommodates the relative southward component of
860 motion of the Aegea block. The left-lateral motion is produced by distributed E/W
861 extensional grabens opening to the west with anticlockwise rotations (e.g. Floyd et al. 2010).

862

863 We then determine the present instantaneous motion between this Aegea block and the rigid
864 Hellenic buttress using the euler pole obtained above (43.02°N, 11.05°E, -1.98°/Myr) that
865 governs the instantaneous motion of the Hellenic Scarp buttress with respect to Anatolia.
866 The differential motion distributed between the Aegea block and the Hellenic Scarp buttress
867 corresponds to E/W extension over Peloponnesus, about 7 mm/yr to the north and 10
868 mm/yr to the south. This extension is manifest within the Argos, Lakonia and Messinia gulfs
869 which correspond to grabens opening southward. Figure 9 after Le Pichon et al. (1995) uses

78 triangulation displacement vectors between 1895 and 1975 to show that there is indeed
a westward relative displacement of about 10 mm/yr of southwestern Peloponnesus with
respect to southeastern Peloponnesus presumably absorbed along the N/S system of normal
faults extending from Central Peloponnesus to western Crete along these Argos, Lakonia and
Messinia extensional systems. Although these triangulation displacements are less precise
than the permanent GPS network, they provide a denser pattern of measurements within
the region of interest. They do confirm that the distributed extension there is indeed the one
expected within our model, that is an approximately E/W extension increasing southward
within Peloponnesus to about 10 mm/yr. Note that Vernant et al. (2014) identified a
relatively poorly defined West Peloponnesus block that also predicts E/W extension within
Peloponnesus.

881

882

883 **The permanency of the kinematics of Anatolia-Aegea with respect to Nubia and** 884 **Eurasia**

885 The Sachpazi faults are a key element of our discussion. This is because the Hellenic slab,
886 that is part of the Nubia plate, is rigid and recorded the lines of slip left by the Aegean motion.
887 They demonstrate that about 400 to 500 kilometers of East Mediterranean lithosphere have
888 been subducted during the last 15 Myr with essentially the same southwestward direction
889 of motion. The lines of slip, governed by the HS/NU pole of rotation (Figs. 6 and 7), show that
890 the oceanic portion of the Mediterranean is subducted at a fast rate, whereas the Herodotus
891 extended continental crust covered by a huge sedimentary section (see part 1) is subducted
892 at a much slower rate.

893

894 The second information we have on the finite kinematics of this block comes from the NAF.
895 The present kinematics based on GPS show that the NAF corresponds in a first
896 approximation to a flow line about the present AN/EU pole, slightly different from and
897 compatible with the AN/NU pole (32°N , 32°E , $1.31^{\circ}/\text{Ma}$). Thus, Eastern and Central Anatolia
898 have formed a single block with a motion governed by the same kinematics since the
899 initiation of motion. These kinematics are in direct continuity of those of Aegea-Western
900 Anatolia as indicated by the Sachpazi faults. However, the Hellenic Scarp has been moving
901 faster than the Anatolia inducing extension within Aegea as explained in the discussion of
902 the reconstruction.

903

904 Here we discuss the kinematics of extension predicted by our reconstruction, using simple
905 finite kinematics. To determine the motion of the Hellenic rigid buttress with respect to
906 Anatolia, we combine the HS/NU euler pole (30.3°N , 31.4°E , $1.78^{\circ}/\text{Myr}$) with the clockwise
907 rotation adopted in the reconstruction about pole D of Figure 3 (37.4°N , 21.7°E). As the finite
908 rotation about pole D is 37° , we use a rotation rate of $3.7^{\circ}/\text{Myr}$, one tenth of the total finite
909 rotation. This is because the NS/NU rotation rate of $1.78^{\circ}/\text{Myr}$ is one tenth of the finite
910 rotation of 18° . The resulting euler pole (43.02°N , 11.05°E , $-1.98^{\circ}/\text{Myr}$) governs the present
911 instantaneous motion of the Hellenic Scarp buttress with respect to Nubia in our
912 reconstruction. We can now combine the resulting pole with the AN/NU pole to obtain the
913 instantaneous pole describing the differential motion between the Hellenic buttress and
914 Anatolia (39.4°N , 20.9°E , $2.03^{\circ}/\text{Myr}$). This pole predicts extension velocities of 10 to 18
915 mm/yr from northwest to southeast along a general southwest-northeast direction close to

916 the direction given by the Sachpazi faults. It is of course a crude approximation, as discussed
917 above, but it enables us to estimate the predicted differential motion distributed between
918 the Aegea block and the Hellenic Scarp buttress. The main point is that the extension within
919 Aegea is accompanied by a strong dextral component.

920

921 It is easy to identify on the free air gravity map of Figure 2 the part of Aegea that has been
922 extended according to these kinematics. It coincides with the zone of strong lineated positive
923 anomalies limited to the northeast from Evia to Mykonos by the linear North Cycladic
924 Detachment that can be followed northwest to the Vardar (see Fig. 1 of Jolivet et al. 2010).
925 This limit then turns 90° to the east toward Samos. A dextral fault, active during Middle to
926 Upper Miocene, identified by Philippon et al. (2012), follows this Mykonos-Samos branch of
927 the detachment and extends further west toward Peloponnesus. Brun et al. (2016) call this
928 fault the Myrthes-Ikaria fault. They suggest the existence of other similar dextral faults to
929 the south within the Cretan Basin. All these dextral faults are along the same direction as the
930 Sachpazi faults. The extension direction was parallel to these faults, increasing
931 southeastward and reaching a maximum at the level of Fault F10. Beyond this point, there
932 is no additional clockwise rotation, as we know that Crete did not rotate (see above). The
933 40° angle between Crete and the Hellenic Scarp reflects this absence of clockwise rotation to
934 the south of F10 and accounts for the decrease of extension further south. To the south of
935 F10, the excess velocity with respect to the HS/NU rotation could only be accommodated by
936 dextral faults. The gravity gradient map of Figure 4 also neatly displays the pattern of
937 extension.

938

The total amount of extension since 15 Myr has to be very large since the 400-500 km of subduction are much larger than the geologically measured offset on the NAF (less than 100 km, Şengör et al. 2005). We conclude that since 15 Ma the pattern of motion with respect to the adjacent Mediterranean basins has been quite constant even though the velocities have increased through time. However, although the Aegea-Anatolia block has followed the same kinematic pattern since the beginning of its motion, it has been affected by significant deformation, mostly extension within Aegea-West Anatolia, as discussed above. Figure 7 in particular demonstrates that the systematic southwestward motion of the Hellenic Scarp buttress with respect to the adjacent Mediterranean basin has been the major controlling factor of the dynamics of this block. This is what one would expect if the gradient of potential gravity across the Hellenic margin played the principal role in the dynamics of this system.

The weak lithosphere of Anatolia-Aegea

Şengör et al. (2005) and Le Pichon et al. (2016) pointed out that the lower crust and mantle of Anatolia and Aegea are highly ductile and that the NAF follows closely the limit of thick relatively cold lithosphere that borders the Anatolia block to the north. The same thing is true for the San Andreas Fault, now located at the limit of thick lithosphere (a lithospheric lid with a thickness larger than 30 km) next to very thin lithosphere (Melbourne and Helmberger 2001). Le Pichon et al. (2016) have proposed that whenever the NAF attempts to enter the relatively weaker lithosphere of Anatolia to the south, it has been pushed back against the thick northern cold lithosphere as the low viscosity lithospheric mantle is being dragged into the motion of Anatolia. As a consequence, the whole lithosphere of the Anatolia-Aegea block is quite hot and ductile (cf. Şengör et al. 2008; Göğüş et al. 2017). This fact is

962 supported by the relatively low viscosity obtained in their modelling by England et al. (2016)
963 for this lithosphere. In addition, as we have seen, within Western Anatolia-Aegea, this ductile
964 lithosphere is submitted to large extensional stresses. In the absence of mature localized
965 boundaries, the strain is widely distributed over its whole width. Brittle distension roots in
966 a shallow decollement above the ductile lower crust and the normal mode of shearing within
967 the upper brittle crust is through rotation of en-echelon normal faults over fairly wide shear
968 zones (Dewey and Şengör 1979; Şengör 1979; McKenzie and Jackson 1983).

969
970 We then have a paradox, discussed by Le Pichon et al. (2016). If the Aegean-Anatolian
971 lithosphere is mechanically so weak, how is it possible that traction at its southwestern
972 extremity can be transmitted over 1500 km to its eastern extremity? McKenzie (1972) and
973 then Le Pichon (1982) had proposed that the westward movement of the Anatolian
974 lithosphere responded to a gradient in gravitational energy, as the elevation decreases
975 regularly from 2 km on the Anatolian Plateau to the east to below sea level in the Aegean
976 Sea. The more detailed information we now have on the correlation between decrease in
977 crustal thickness and topography from east to west validates this proposition (Le Pichon et
978 al. 2016; England et al. 2016). However, the present situation is in complete contrast with
979 the situation that existed 11 Ma when the East Anatolian Plateau was below sea level (Şengör
980 et al. 2003, 2008) whereas western Turkey topography was very high (Şengör 1991) and
981 most of what is now the Central and Southern Aegean Sea were just beginning to subside
982 below sea level (Drooger and Meulenkamp 1973; see discussion in Le Pichon et al. 2016).
983 The last marine sediments deposited above the present East Anatolian Plateau date from
984 Serravallian, 13-11 Ma (Şengör et al. 2003). The uplift thus started 11 Ma and was
985 accompanied by volcanic activity that became plateau-wide at 8 Ma while the Aegean Sea

continued to subside (Şengör et al. 1985, 2008). Şengör et al. (2003) attributed part of the uplift of the East Anatolian Plateau to slab break-off, arguing that the asthenosphere appeared to lie immediately below the crust, as evidenced by its anomalously low seismic velocity. The existence of a slab break-off was also proposed by Faccenna et al. (2006) and by many others since, as tomographic studies have demonstrated that the plunging African slab below Turkey is detached from the African plate. The present linear decrease in topographic gradient and continental crust thickness between the Anatolia Plateau and the Aegean Sea was consequently progressively established after 8 Ma. Its establishment was only possible because of the ductility of the lower crust and uppermost mantle as discussed by Le Pichon et al. (2016). This topographic gradient, once formed, was essential to enable Anatolia to follow the motion of Aegea pulled along its Hellenic Scarp margin by the strong potential gravity gradient across it.

But Anatolia began its westward motion at a time when this potential gravity gradient was not yet established, and possibly opposite to the actual motion. The delay between the beginning of the subduction along the Hellenic Trench at 15 Ma and the beginning of uplift of the East Anatolian Plateau at 11-8 Ma is reflected in the 4-5 Myr delay between the motion of Aegea and the motion of Anatolia. During the Middle and Upper Miocene, the pull of the Hellenic margin scarp was the main element of the dynamics of the Aegea-Anatolia system and, as a result, distension was very important within Aegea and mainly localized in the wake of the Hellenic Scarp, before the gradient of gravitational potential energy across Anatolia-Aegea was fully reversed. This establishes the existence of a pull force acting on Aegea from the west during Middle and Upper Miocene. On the other hand, the existence of east-west thrusting within Central Anatolia during Pliocene indicates at this time the

existence of a push force acting within Anatolia from the east. Was the effect of the pull across the Hellenic slab and the uplift of the Anatolian Plateau sufficient to trigger the motion of Anatolia, explaining the progressive increase of velocity of Anatolia with time? Or rather, should we, following Le Pichon and Kreemer (2000) invoke an additional effect: an underlying counterclockwise asthenospheric flow along the eastern edge of the African slab, owing to subduction rollback, after the slab detachment below the Anatolian Plateau? Only, a detailed analysis of the evolution from Middle Miocene to Present of the strain pattern within Anatolia-Aegea would allow to fully answer these questions. It appears to us that the dynamics of this area are the result of a complex pattern of forces to which the gradient of potential gravity adjusts, smoothing out the transition from push to pull and possibly underlying traction.

The formation of the Anatolia-Aegea boundaries

We have solid evidence that the present subduction system was initiated 15 Ma. We have direct evidence by drilling (see *The backstop*) that 15 Ma marks the beginning of the formation of the accretionary wedge (Kastens 1991). We also have indirect evidence because 15 Ma is the date of simultaneous initiation of the clockwise rotation to the west and anticlockwise rotation to the east within southern Aegean (van Hinsbergen et al. 2005, 2010), the clockwise rotation having been related to the contemporaneous activity of the Aegean detachment system (Jolivet et al. 2015) (see *The kinematic significance of the Sachpazi faults*). Finally, we have additional indirect evidence related to the formation of the outer rises of the Hellenic Trough, which already existed by Tortonian time (see *The Hellenic Subduction Trough*).

1033

1034 In contrast, the northern boundary is not yet completely established as the NAF has not yet
1035 reached the northern extremity of the Hellenic trench. Its existence as a single fault does not
1036 extend beyond the western extremity of the North Aegean Trough. This is important as the
1037 NAF, when established, has an extremely low friction coefficient that Provost et al. (2003)
1038 estimated from modelling to be 0.05. It becomes close to a free slip boundary. This is possible
1039 because friction has developed a vertical tabular zone with extremely low friction coefficient
1040 rooting within the mantle. In the brittle upper part of the crust, this is done through
1041 production of soft material (succession of ultracataclasite, cataclasite, brecciae, dense
1042 fracturing, Chester et al. 1993), which requires a long length of slip. Şengör et al. (2005) used
1043 the Tchalenko (1970) model to explain the formation of what Tchalenko called the Principal
1044 Displacement Zone where strain localization has occurred within the very low friction core.
1045 Tchalenko had shown that the formation of the Principal Displacement Zone requires a
1046 displacement comparable to the width of the initially sheared zone, hence the delay in the
1047 formation of the fault and its westward propagation. Şengör et al. (2005) showed that the
1048 earliest evidence that the fault existed to the east dates from 11 Ma. They could then show
1049 that the eastern half east of Marmara propagated westward at an average rate of 110
1050 km/Myr.

1051

1052 Le Pichon et al. (2016) showed that within Western Anatolia-Aegea the NAF propagated
1053 westward, having reached the eastern extremity of the Marmara Sea 4.5 Ma, the Gulf of Saros
1054 2.5 Ma and its present western extremity within the last million years. This evolution is
1055 compatible with an overall westward propagation rate of about 140-150 km/Myr along its

present 1400 km length. As discussed by Le Pichon et al. (2016), within the Western Anatolia-Aegea distensional environment, the shear is initially distributed over the whole width of the block, then it tends to concentrate within a shear zone about 200 km wide with clockwise rotating rifts (Şengör 1982; Taymaz et al. 1991), such as the present Gulf of Corinth within the Grecian Shear Zone, before the fault finally cuts through the shear zone.

The failure of the EMNT

The initiation of the Hellenic subduction zone was due to the failure of the EMNT at the western extremity of the Antalya obduction front about 15 Ma. The transform faults that framed the Eastern Mediterranean had maintained their identity for about 100 Myr. Here, we attempt to understand more fully the context of this failure, in light of the evolution of the AF/EU relative motion based on the Atlantic magnetic anomalies (Rosenbaum et al. 2002). However, the EMNT along the eastern part of the Eastern Mediterranean area was affected, between the Turonian and the Maestrichtian, by a several thousand kilometers long obduction driven orogen that Şengör and Stock (2014) call the Ayyubid Orogen (see part 1 The kinematic context of the late Cretaceous obduction). This Ayyubid Orogen represents a partial failure of the former transform faults framework, failure more advanced along the Croissant Ophiolitique than over the Syrian Arc. But the transform framework appears to have maintained its integrity as discussed earlier.

The western limit of this partial failure along the EMNT (B in figure 6) coincides with the western limit of what are now the Antalya nappes. At the time the eastern extremity of what is now Crete was much closer to this limit where 75 Myr later the EMNT failed, failure that

would lead to the initiation of the Hellenic subduction. Garfunkel (2004) pointed out the existence of a peculiar late assemblage in Eastern Crete of Late Cretaceous intrusive and metamorphic rocks in Crete. The Cretaceous intrusive rocks and the metamorphic rocks there belong to two separate settings: the Cretaceous metamorphic rocks are mostly in an allochthonous position. Some of the Cretaceous igneous rocks are clearly related to the nappes. Some may be of more southerly origin. In fact, these 'southerly' Cretaceous intrusions begin showing up in Crete, where the evidence for Cretaceous ophiolitic obduction ends. This may have been a tectonic 'hick-up', an attempt to begin a subduction, that failed, but managed to destroy the northern transform fault boundary of the Eastern Mediterranean. The Antalya nappes system is situated at the northern end of the NNE/SSW Herodotus/central oceanic basins boundaries (see part 1 section *The extended continental Levant and Herodotus basins*). We propose that this boundary coincides with the transfer fault proposed by Şengör and Stock (2014, see their Figure 15). This reinforces the proposition we made that the Herodotus/oceanic basin boundary is a major structural boundary within the Mediterranean that has conditioned its evolution. We propose that the late Cretaceous orogenic episode is the cause of the localization of the failure of the EMNT there.

According to Rosenbaum et al. (2002), the shortening between Africa and Eurasia stopped for 9 Myr during Palaeocene between anomalies 30 and 25 (Trümpy's, 1960, "Palaeocene restoration"). From then on the pattern of AF/EU motion changed drastically as there were about 700 kilometers of pure shortening between anomaly 25 (56 Ma) and Present. The shortening velocity first increased progressively to about 20 mm/yr during Oligocene-Aquitania (30-20 Ma) and then decreased to about 7 mm/yr afterwards. However, Arabia

that was linked to Africa before Miocene (Anomaly 6, 20 Ma), continued at a slightly reduced velocity (25 mm/yr at 38°N, 48°E) while Africa was trailing behind resulting in the opening of the Red Sea-Gulf of Aden (McQuarrie et al. 2003). In addition, as the relative velocity of Arabia/Eurasia changed from NE to N, this resulted in partitioning of the motion between shortening and an equivalent component of dextral motion of about 15 mm/yr between Arabia and Eurasia. The formation of the backstop and then of the accretionary prism occurred during this Lower-Middle Miocene period of rearrangement of the kinematics of the Africa/Arabia/Eurasia new plate system.

Conclusion

The central tenet of part 1 is that both the Alpine Tethys and the Eastern Mediterranean Sea are the result of left-lateral shearing of the Adria block as Africa was separated from Eurasia during the initial post-Triassic opening of the Central Atlantic Ocean. The Eastern Mediterranean Sea was thus created as a very long narrow pull-apart that we compare to the Cayman Shear zone the Caribbean (Rosencrantz 1995). We propose that this Eastern Mediterranean shear zone continued to be active during early Cretaceous time, although the Africa-Eurasia motion deviated somewhat from the previous Jurassic trend. The transform faults that framed the Eastern Mediterranean Sea during its formation in Jurassic played an important role during the following geological history. The Eastern Mediterranean North Transform fault (EMNT) formed the northern boundary of this Eastern Mediterranean shear

zone. This fault has been destroyed over a large portion by the Hellenic subduction. In this paper, we have reconstructed the northern margin of the Eastern Mediterranean Sea prior to the Hellenic subduction using the faults identified within the Hellenic slab by Sachpazi et al. (2016a and 2016b) and we have shown that it coincided with the EMNT. We achieved that by investigating the Hellenic subduction zone since its origin. The Sachpazi faults within the Hellenic slab were a key element of our discussion. This is because the slab, which is part of the Nubia plate, is rigid and the faults within it coincide with lines of slip congruent with the relative motion of the Aegean block over it. These faults demonstrate that about 400 to 500 kilometers of East Mediterranean lithosphere have been subducted with essentially the same southwestward direction of motion during the last 15 Myr. This has important implications for the Neogene evolution of Anatolia-Aegea that we have discussed.

Finally, although we were able to bring considerable new light on the formation of both the Alpine Tethys and the Eastern Mediterranean Sea, demonstrating that it was governed by the Africa/Eurasia post-Triassic motion, we have not made a detailed reconstruction of the area in Triassic prior to the opening of the Atlantic Ocean. Such a reconstruction would require major work as it implies in particular discussing the pre-drift Africa-Eurasia-America fit. We emphasize however that our approach imposes significant new constraints on such a fit.

Acknowledgments

1146 The first author is grateful for the invitation to contribute to this special issue. We thank
1147 Nicolas Chamot-Rooke for pointing out to us the relevance of the gravity gradient maps on
1148 the Hellenic backstop and for discussion. We thank Jean Mascle for contributing bathymetric
1149 maps of the Eastern Mediterranean and David Sandwell for sending us his latest gravity data.
1150 We thank Jean-Pierre Brun for a review that helped us clarify our presentation. We thank
1151 George Davis for a detailed review that was most helpful and an anonymous reviewer for
1152 important suggestions. We thank Dimitrios Papanikolaou for pointing out that we should
1153 discuss the nature of the limit between the Hellenic Trench and the continental slope. Finally,
1154 we thank Zvi Garfunkel who encouraged us to publish this paper that he considered to be a
1155 major step forward in understanding the Mediterranean.

1156

Draft

References

- Armijo, R., Meyer, B., King, G.C.P., Rigo, A., and Papanastassiou, D. 1996. Quaternary evolution of the Corinth Rift and its implications for the Late Cainozoic evolution of the Aegean. *Geophys. J. Int.*, **126**: 11–53.
- Aubouin, J., Bonneau, M., Davidson, J., Leboulenger, P., Matesco, S., and Zambetakis, A. 1976. Esquisse structurale de l'arc Egéen externe: des Dinarides aux Taurides. *Bull. Soc. Geol. France*, **18**: 327-336.
- Biryol, C.B., Beck, S.L., Zandt, G., and Özacar, A.A. 2011. Segmented African lithosphere beneath the Anatolian region inferred from teleseismic P-wave tomography. *Geophys. J. Int.*, **184**: 1037-1057. doi:10.1111/j.1365-246X.2010.04910.x
- Bijwaard, H., Spakman, W. and Engdahl, E.R. 1998. Closing the gap between global and regional mantle tomography. *J. Geophys. Res.*, **103**: 30055-30078. doi:10.1029/98JB02467.
- Bocchini, G.M., Brüstle, A., Becker, D., Meier, T., van Keken, P.E., Ruscic, M., Papadopoulos, G.A., Rische, M., and Friederich, W. 2018. Tearing, segmentation and backstepping of subduction in the Aegean, new insights from seismicity. *Tectonophysics*, **734**: doi: 10.1016/j.tecto. 2018.04.002.
- Brun, J.P., Faccenna, C., Gueydan, F., Sokoutis, D., Philippon, M., Kydonakis, K., and Gorini, C. 2016. Effects of slab rollback acceleration on Aegean extension. *Bul. Geol. Soc. Greece*, **50**: 5-14. doi: org/10.12681/bgsg.11697.
- Campistron-Brosolo, L., and Mascle, J. 2015. Morpho-Bathymetry of the Eastern Mediterranean Sea. CGMW-CIESM.

- 1178 Chamot-Rooke, N., Rangin, C., and Le Pichon, X. 2005. DOTMED, Deep offshore tectonics of
1179 the Eastern Mediterranean, A synthesis of deep marine data in the Eastern Mediterranean.
1180 Mémoires Soc. Géol. France, n.s., 177.
- 1181 Chester, F.M., Evans, J.P., and Biegel, R.L. 1993. Internal structure and weakening
1182 mechanisms of the San Andreas fault. *J. Geophys. Res.*, **74**: 3821-3827.
- 1183 Chousianitis, K., Ganas, A., and Evangelidis, C.P. 2015. Strain and rotation rates patterns of
1184 mainland Greece from continuous GPS data and comparison between seismic and geodetic
1185 moment release. *J. Geophys. Res. (Solid Earth)*, **120**: 3909-3931.
1186 doi:10.1002/2014JB011762.
- 1187 Dewey, J. F., and Şengör, A.M.C. 1979. Aegean and surrounding regions: complex multiplate
1188 and continuum tectonics in a convergent zone. *Geol. Soc. America Bull.*, **1(90)**: 84-92.
- 1189 Drooger, C.W., and Meulen Kamp, J.E. 1973. Stratigraphic contribution to geodynamics in the
1190 Mediterranean Area: Crete as a case history. *Bull. Soc. Geol. Greece*, **10**: 193-200.
- 1191 England, P., Houseman, G., and Nocquet, J.M. 2016. Constraints from GPS measurements on
1192 the dynamics of deformation in Anatolia and the Aegean. *J. Geophys. Res. Solid Earth*, **121**:
1193 8888-8916. doi:10.1002/2016JB013382.
- 1194 Faccenna, C, Bellier O, Martinod J, Piromallo C, and Regard V. 2006. Slab detachment
1195 beneath eastern Anatolia: a possible cause for the formation of the North Anatolian Fault.
1196 *Earth Planet. Sci. Lett.*, **242**: 85–79.
- 1197 Faccenna, C., Becker, T.W., Auer, L., Billi, A., Boschi, L., Brun, J.P., Capitanio, F.A., Funiciello, F.,
1198 Horváth, F., Jolivet, L., Piromallo, C., Royden, L., Rossetti, F., and Serpelloni, E. 2014. Mantle
1199 dynamics in the Mediterranean. *Rev. Geoph.*, **52**: 283-332. doi: 10.1002/2013RG000444.

- 1200 Flotté, N. 2003. Caractérisation Structurale et Cinématique d'un Rift sur Détachement: Le
1201 Rift de Corinthe-Patras, Grèce. Thèse de Doctorat, Orsay: Univ. Paris-Sud.
- 1202 Floyd, M.A., Billiris, H., Paradissis, D., Veis, G., Avallone, A., Briole, P., McClusky, S., Nocquet, J.-
1203 M., Palamartchouk, K., Parsons, B., and England, P.C. 2010. A new velocity field for Greece:
1204 Implications for the kinematics and dynamics of the Aegean. *J. Geophys. Res. Solid Earth*, **115**:
1205 B10403. doi:10.1029/2009JB007040.
- 1206 Gallais, F., Gutscher, M.A., Graindorge, D., Chamot-Rooke, N., and Klaeschen, A.D. 2011.
1207 Miocene tectonic inversion in the Ionian Sea (central Mediterranean): Evidence from
1208 multichannel seismic data. *J. Geophys. Res.*, **116**: B12108. doi:10.1029/2011JB008505.
- 1209 Garfunkel, Z. 2004. Origin of the Eastern Mediterranean basin: a reevaluation.
1210 *Tectonophysics*, **391**: 11-34.
- 1211 Göğüş, O. H., Pysklywec, R.N., Şengör, A.M.C., and Gün, E. 2017. Drip tectonics and the
1212 enigmatic uplift of the Central Anatolian Plateau. *Nature Communications*, **8**:
1213 doi:10.1038/s41467-017-01611-3.
- 1214 Govers, R., and Wortel, M.J.R. 2005. Lithosphere tearing at STEP faults: Response to edges of
1215 subduction zones. *Earth Planet. Sci. Lett.*, **236**: 505–523.
- 1216 Hansen, S., Evangelidis, C.P., and Papadopoulos, G.A. 2019. Imaging slab detachment within
1217 the Western Hellenic Subduction zone. *Geoscience Commun.*, 10.1029/GC007810.
- 1218 Howell, A., Palamartchouk, K., Papanikolaou, X., Paradissis, D., Raptakis, C., Copley, A.,
1219 England, P., and James Jackson, J. 2017a. The 2008 Methoni earthquake sequence: the
1220 relationship between the earthquake cycle on the subduction interface and coastal uplift in
1221 SW Greece. *Geophys. J. Int.*, **208**: 1592–1610.

- 1222 Howell, A., Jackson, J., Copley, A., McKenzie, D., and Nissen, E. 2017b. Subduction and vertical
1223 coastal motions in the eastern Mediterranean. *Geophys. J. Int.*, **211**: 593-620.
1224 doi:10.1093/gji/ggx307.
- 1225 Huguen, C., Chamot-Rooke, N., Loubrieu, B., and Mascle, J. 2006. Morphology of a pre-
1226 collisional, salt-bearing, accretionary complex: The Mediterranean Ridge (Eastern
1227 Mediterranean). *Marine Geophysical Researches*, **27**: 61-75. doi:10.1007/s11001-005-
1228 5026-5.
- 1229 Jolivet, L., and Brun, J.P. 2010. Cainozoic geodynamic evolution of the Aegean. *Int. J. Earth Sci.*
1230 (*Geol. Rundsch.*), **99**: 109 -138. doi : 10.1007/s00531-008-0366-4.
- 1231 Jolivet, L., Lecomte, E., Huet, B., Denèle, Y., Lacombe, O., Labrousse, L., Le Pourhiet, L., and
1232 Mehl, C. 2010. The North Cycladic Detachment System. *Earth Planet. Sci. Let.*, **289**: 87-104.
1233 doi:10.1016/j.epsl.2009.10.032.
- 1234 Jolivet, L., Menant, A., Sternai, P., Rabillard, A., Arbaret, L., Augier, R., Laurent, V., Beaudoin,
1235 A., Graseman, A., Huet, B., Labrousse, L., and Le Pourhiet, L. 2015. The geological signature of
1236 a slab tear below the Aegean. *Tectonophysics*, **659**: 166-182.
1237 doi.org/10.1016/J.tecto.2015.08.004.
- 1238 Kastens, K.A. 1991. Rate of outward growth of the Mediterranean Ridge accretionary
1239 complex. *Tectonophysics*, **199**: 25-50.
- 1240 Ketin, İ. 1948. Über die tektonisch-mechanischen Folgerungen aus den großen anatolischen
1241 Erdbeben des letzten Dezenniums. *Geologische Rundschau*, **36**: 77-83.

- 1242 Kissel, C. and Laj, C. 1988. A tertiary geodynamical evolution of the Aegean Arc: a
1243 paleomagnetic reconstruction. *Tectonophysics*, **146**: 183-201. doi:10.1016/0040-
1244 1951(88)90090-X.
- 1245 Kissel, C., Laj, C., Poisson, A., and Görür, N. 2003. Palaeomagnetic reconstruction of the
1246 Cainozoic evolution of the eastern Mediterranean. *Tectonophysics*, **362**: 199-217.
- 1247 Le Pichon, X. 1982. Land-locked oceanic basins and continental collision: the eastern
1248 Mediterranean as a case example. In *Mountain Building Processes*. Edited by K.J. Hsü,
1249 Academic Press, London. pp. 201–211.
- 1250 Le Pichon, X., and Angelier, J. 1979. The Hellenic arc and trench system: a key to the
1251 neotectonic evolution of the Eastern Mediterranean area. *Tectonophysics*, **60**: 1-42.
- 1252 Le Pichon, X., and Angelier, J. 1981. The Aegean Sea. *Phil. Trans. R. Soc. Lond. A*, **300**: 357-
1253 372. doi: 10.1098/rsta.1981.0069.
- 1254 Le Pichon, X., and Kreemer, C. 2010. The Miocene-to-present kinematic evolution of the
1255 Eastern Mediterranean and Middle East and its implications for dynamics. *Ann Rev Earth*
1256 *Planet. Sci.*, **38**: 323–351.
- 1257 Le Pichon, X., Francheteau, J., and Bonnin, J. 1973. *Plate Tectonics*. Elsevier, 300 pp.
- 1258 Le Pichon, X., Chamot-Rooke, N., Lallemant, S., Noomen, R., and Veis, G. 1995. Geodetic
1259 determination of the Kinematics of Central Greece with respect to Europe. *J. Geophys.*
1260 *Res.*, **100**: 12675-12690.
- 1261 Le Pichon, X., Lallemant, S.J., Chamot-Rooke, N., Lemeur, D., and Pascal, G. 2002. The
1262 Mediterranean Ridge backstop and the Hellenic nappes. *Marine Geology*, **186**: 111-125.

- 1263 Le Pichon, X., İmren, C., Rangin, C., Şengör, A.M.C., and Siyako, M. 2014. The South Marmara
1264 Fault. *International Journal of Earth Sciences (Geol Rundsch)*, **103**: 219–231. doi:
1265 10.1007/s00531-013-0950-0.
- 1266 Le Pichon, X., Şengör, A.M.C., Kende, J., İmren, C., Henry, P., Grall, C., and Karabulut, H. 2016.
1267 Propagation of a strike-slip plate boundary within an extensional environment: the
1268 westward propagation of the North Anatolian Fault. *Can. J. Earth Sci.*, **53(11)**: 1416–1439.
1269 doi: 10.1139/cjes-2015-0129.
- 1270 McKenzie, D.P. 1970. Plate tectonics of the Mediterranean region. *Nature*, **226(5242)**: 239–
1271 243.
- 1272 McKenzie, D.P. 1972. Active tectonics of the Mediterranean region. *Geophys. J. R. Astron. Soc.*,
1273 **30**: 109–185.
- 1274 McKenzie, D., and Jackson, J. 1983. The relationship between strain rates, crustal thickening,
1275 palaeomagnetism, finite strain and fault movement within a deforming zone. *Earth Planet.*
1276 *Sci. Lett.*, **65**: 182–202.
- 1277 McQuarrie, N., Stock, J.M., Verdel, C., and Wernicke, B.P. 2003. Cainozoic evolution of
1278 Neotethys and implications for the causes of plate motions. *Geophys. Res. Lett.*, **30**: 2036–
1279 2040.
- 1280 Melbourne, T., and Helmberger, D. 2001. Mantle control of plate boundary deformation.
1281 *Geophys. Res. Lett.*, **28**: 4003–4006.
- 1282 Özbakır, A.D., Govers, R., and Wortel, R. 2017. Active faults in the Anatolian-Aegean plate
1283 boundary region with Nubia. *Turkish Journal of Earth Sciences*, **26**: 30–56. doi: 10.3906/yer-
1284 1603-4.

- 1285 Özeren, M.S. and Holt, W.E. 2010. The dynamics of the eastern Mediterranean and eastern
1286 Turkey. *Geophys. J. Int.*, **183**: 1165-1184.
- 1287 Pearce, F.D. 2015. Seismic imaging of the western Hellenic subduction zone: The relationship
1288 between slab composition, retreat rate, and overriding lithosphere genesis. PhD thesis,
1289 Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of
1290 Technology, Cambridge, MA, USA.
- 1291 Pérouse, E., Chamot-Rooke, N., Rabaute, A., Briole, P., Jouanne, F., Georgiev, I., and Dimitrov,
1292 D. 2012. Bridging onshore and offshore present-day kinematics of central and eastern
1293 Mediterranean: Implications for crustal dynamics and mantle flow. *Geochem. Geophys.*
1294 *Geosyst.*, **13(9)**: Q09013. doi:10.1029/2012GC004289.
- 1295 Philippon, M., Brun, J.P., and Gueydan, F. 2012. Deciphering subduction from exhumation in
1296 the segmented Cycladic Blueschist Unit (Central Aegean, Greece). *Tectonophysics*, **524**: 116-
1297 134, doi:10.1016/j.tecto.2011.12.025.
- 1298 Provost, A.S., Chéry, J., and Hassani, R. 2003. 3-D mechanical modeling of the GPS velocity
1299 field along the North Anatolian Fault. *Earth Planet. Sci. Lett.*, **209**: 361–377.
- 1300 Reilinger, R., McClusky, S., Vernant, P., Lawrence, S., Ergintav, S., Cakmak, R., Ozener, H.,
1301 Kadirov, F., Guliev, I., Stepanyan, R., Nadariya, M., Habubia, G., Mahmoud, S., Sakr, K.,
1302 ArRajehi, A., Paradissis, D., Al-Aydrus, A., Prilepin, M., Guseva, T., Evren, E., Dmitrotsa, A.,
1303 Filikov, S.V., Gomez, F., Al-Ghazzi, R., and Karam, G. 2006. GPS constraints on continental
1304 deformation in the Africa-Arabia-Eurasia continental collision zone and implications for the
1305 dynamics of plate interactions. *J. Geophys. Res.*, **111(B5)**: doi:10.1029/2005JB004051.

- 1306 Rosenbaum, G., Lister, G.S., and Duboz, C. 2002. Relative motions of Africa, Iberia and Europe
1307 during Alpine orogeny. *Tectonophysics*, **359**: 117–129.
- 1308 Rosencrantz, E. 1995. Opening of the Cayman Trough and the evolution of the northern
1309 Caribbean Plate boundary. Geological Society of America Annual Meeting. New Orleans, 27,
1310 153.
- 1311 Royden, L.H., and Papanikolaou, D.J. 2011. Slab segmentation and late Cainozoic disruption
1312 of the Hellenic arc. *Geochem. Geophys. Geosyst.*, **12(3)**: Q03010.
1313 doi:10.1029/2010GC003280.
- 1314 Ryan, W.B.F. 1970. The stratigraphy of the Eastern Mediterranean, part II, Thesis, Columbia
1315 University, New York.
- 1316 Sachpazi, M., Laigle, M., Charalampakis, M., Diaz, J., Kissling, E., Gesret, A., Becel, A., Flueh, E.,
1317 Miles, P., and Hirn, A. 2016a. Segmented Hellenic slab rollback driving Aegean deformation
1318 and seismicity. *Geophys. Res. Lett.*, **43**: 651-658. doi:10.1002/2015GL066818.
- 1319 Sachpazi, M., Laigle, M., Charalampakis, M., Sakallariou, D., Flueh, E., Sokos, E., Deskalaki, E.,
1320 Galvé, A., Petrou, P., and Hirn, A. 2016b. Slab segmentation controls the interplate slip motion
1321 in the SW Hellenic subduction: new insight from the 2008 Mw 6.8 Methoni interplate
1322 earthquake: Slab segmentation and interplate slip. *Geophys. Res. Lett.*, **43**: 9619-9626.
1323 doi:10.1002/2016GL070447.
- 1324 Sandwell, D.T., Müller, R.D., Smith, W.H., Garcia, E., and Francis, R. 2014. New global marine
1325 gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure. *Science*,
1326 **346(6205)**: 65-67.

- 1327 Şengör, A.M.C. 1979. The North Anatolian Transform Fault: its age, offset and tectonic
1328 significance. *Jour. Geol. Soc. London*, **136**: 269-282.
- 1329 Şengör, A.M.C. 1982. Ege'nin neotektonik evrimini yöneten etkenler (Factors governing the
1330 neotectonic evolution of the Aegean). In O. Erol and V. Oygür, eds., *Batı Anadolu'nun Genç*
1331 *Tektoniği ve Volkanizması Paneli*, Türk. Jeol. Kur., Ankara pp. 59-71.
- 1332 Şengör, A.M.C. 1987. Cross-faults and differential stretching of hangingwalls in regions of
1333 low-angle normal faulting: examples from western Turkey. *Geol. Soc. London Spec. Pub.*, **28**:
1334 575-589.
- 1335 Şengör, A.M.C. 1991. Timing of orogenic events: a persistent geological controversy. in
1336 *Modern Controversies. In Geology (Proceedings of the Hsü Symposium edited by D.W.*
1337 *Müller, J.A. McKenzie, and H. Weissert: Academic Press, London, pp. 405-473.*
- 1338 Şengör, A.M.C., and Stock, J. 2014. The Ayyubid orogen: an ophiolite obduction-driven orogen
1339 in the Late Cretaceous of the Neo-Tethyan south margin. *Geoscience Canada*, **41**: 224-254.
1340 doi.org/10.12789/geocanj.2014.41.042.
- 1341 Şengör, A. M. C., and Zabcı, C. in press. The North Anatolian Fault and the North Anatolian
1342 Shear Zone. In Kuzucuoğlu, C., Ciner, A., and Kazancı, N., editors, *Landforms and Landscapes*
1343 *of Turkey*, Springer, Heidelberg.
- 1344 Şengör, A.M.C., Görür, N., and Şaroğlu, F. 1985. Strike-slip faulting and related basin
1345 formation in zones of tectonic escape: Turkey as a case study. In Biddle, K.T. and Christie-
1346 Blick, N., eds, *Strike-slip Deformation, Basin Formation, and Sedimentation*, Soc. Econ.
1347 *Palaeont. Min. Spec. Pub. 37* (in honor of J.C. Crowell), p. 227-264.

- 1348 Şengör, A.M.C., Özeren, S., Genç, T., and Zor, E. 2003. East Anatolian high plateau as mantle-
1349 supported, north-south shortened domal structure? *Geophys. Res. Lett.*, **30(24)**:
1350 doi:10.1029/2003GL017858.
- 1351 Şengör, A.M.C., Tüysüz, O., İmren, C., Sakıncı, M., Eyidoğan, H., Görür, N., Le Pichon, X., and
1352 Rangin, C. 2005. The North Anatolian Fault: A new look. *Annu. Rev. Earth Planet. Sci.*, **33**: 37–
1353 112.
- 1354 Şengör, A.M.C., Özeren, M.S., Sakıncı, M., Özbakır, A.D., and Kayan, İ. 2008. Eastern Turkish
1355 high plateau as a small Turkic-type orogen: implications for post-collisional crust-forming
1356 processes. *Earth Science Reviews*, **90**: 1-48.
- 1357 Serpetsidaki, A., Elias, P., Ilieva, M.P., Bernard, P., Briole, P., Deschamps, A., Lambotte, S., Lyon-
1358 Caen, H., Sokos, E., and Tselentis, G-A. 2014. New constraints from seismology and geodesy
1359 on the Mw = 6.4 2008 Movri (Greece) earthquake: evidence for a growing strike-slip fault
1360 system. *Geophys. J. Int.*, **198**: 1373-1386.
- 1361 Shaw, B., and Jackson, J. 2010. Earthquake mechanisms and active tectonics of the Hellenic
1362 subduction zone. *Geophys. J. Int.*, **181(2)**: 966-984. doi:10.1111/j.1365-246X.2010.04551.x.
- 1363 Shaw, B., Ambraseys, N., England, P.C., Floyd, M.A., Gorman, G.J., Higham, T.F.G., Jackson, J.A.,
1364 Nocquet, J.-M., Pain, C.C., and Piggott, M.D. 2008. Eastern Mediterranean Tectonics and
1365 tsunami hazard inferred from A.D.365 earthquake. *Nature Geoscience*, **1(4)**: 268-276.
- 1366 Sorel, D. 2000. A Pleistocene and still-active detachment fault and the origin of the Corinth-
1367 Patras rift, Greece. *Geology*, **28(1)**: 83–86.

- 1368 Spakman, W., van der Lee, S., and van der Hilst, R. 1993. Travel time tomography of
1369 the European-Mediterranean mantle down to 1400 km. *Phys. Earth Planet. Int.*, **79(1-2)**: 3-
1370 74.
- 1371 Stampfli, G.M., and Borel, G.D. 2002. A plate tectonic model for the Palaeozoic and Mesozoic
1372 constrained by dynamic plate boundaries and restored synthetic oceanic isochrons. *Earth*
1373 *and Planetary Science Letters*, **196**: 17–33.
- 1374 Taymaz, T., Jackson, J., and McKenzie, D. 1990. Earthquake mechanisms in the Hellenic
1375 Trench near Crete. *Geophys. J. Int.*, **102**: 695-731.
- 1376 Taymaz, T., Jackson, J., and McKenzie, D. 1991. Active tectonics of the north and central
1377 Aegean Sea. *Geophys. J. Int.*, **106**: 433-490.
- 1378 Tchalenko, J.S. 1970. Similarities between shear zones of different magnitudes. *Geol. Soc. Am.*
1379 *Bull.*, **81**: 1625-1640.
- 1380 Trümpy, R. 1960. Palaeotectonic evolution of the Central and Western Alps. *Bull Geol Soc*
1381 *Am*, **71**: 843–908.
- 1382 Van Hinsbergen, D.J.J., and Schmid, S.M. 2012. Map view restoration of Aegean-West
1383 Anatolian accretion and extension since Eocene. *Tectonics*, **31(5)**:
1384 doi:10.1029/2012TC003132.
- 1385 Van Hinsbergen, D.J.J., Langereis, C.G., and Meulenkamp, J.E. 2005. Revision of the timing,
1386 magnitude and distribution of Neogene rotations in the western Aegean region.
1387 *Tectonophysics*, **396(1-2)**: 1-34. doi:10.1016/j.tecto.2004.10.001.

1388 Van Hinsbergen, D.J.J., Dekkers, M.J., Bozkurt, E., and Koopman, M. 2010. Exhumation with a
1389 twist: Palaeomagnetic constraints on the evolution of the Menderes metamorphic core
1390 complex, western Turkey. *Tectonics*, **29(3)**: TC3009. doi:10.1029/2009TC002596.

1391 Vernant, P., Reilinger, R., and McClusky, S. 2014. Geodetic evidence for low coupling on the
1392 Hellenic subduction plate interface. *Earth Planet. Sci. Lett.*, **385**: 122-129.

1393

Draft

Captions

Figure 1

Bathymetry of the Eastern Mediterranean after Campistron-Brosolo and Mascle (2015). Fractures F1 to F9 within the Hellenic slab (heavy white lines and heavy dashed white lines) are after Sachpazi et al. (2016a and 2016b). Proposed additional fractures F10 to F12 (white dots). Black lines, small circles about the HS/NU rotation pole (red small circle). Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Thick dashed black lines: probable extension of backstop below Mediterranean Ridge. Continuous coloured contours on top of the slab are after Sachpazi et al. (2016a and 2016b). Dashed 20 to 60 km contours are after Vernant et al. (2014). Dashed 100 and 140 km black contours are after Shaw and Jackson (2010). Violet arrow is slip vector of the 2008 Methoni earthquake after Howell et al. (2017a). Black arrows are slip vectors of subduction earthquakes after Shaw and Jackson (2010). KF, Kefalonia fault; NAF, North Anatolian fault. Fault plane solutions of earthquakes within slab (red, P axis, black, T axis) are after Shaw and Jackson (2010). Thick white lines, approximate direction of P axis within slab. Black circle is the AN/NU pole (32.71°N, 34.2°E, 1.29°/Ma, Le Pichon and Kreemer 2010).

Figure 2

Free air gravity map of Eastern Mediterranean Sea from Sandwell et al. (2014). Fractures F1 to F9 within the Hellenic slab (white lines and heavy dashed white lines) are after Sachpazi et al. (2016a and 2016b). Proposed additional fractures F10 to F12 (white dots). Dashed red small circles about HS/NU rotation pole shown by red dot. Backstop outer

1416 limit in green and inner limit in dashed blue after Huguen et al. (2006). Dashed black:
1417 probable extension of backstop below Mediterranean Ridge. "+" identify three different
1418 gravity highs related to the flexure of the subducting Eastern Mediterranean lithosphere.
1419 Red faults on Medina axial gravity high are a set of reverse faults active during Tortonian
1420 (Gallais et al. 2011). Heavy black line is small circle following the Lybian Egyptian
1421 Transform margin centered on 52.0°N, 39.0°E. Heavy black dashed lines are small circles
1422 about same center identifying the Apulian Transform and the EMNT.

1423 Figure 3

1424 Aegean extensional areas and directions of Aegean basin development after map 6 of
1425 Chamot-Rooke et al. (2005). Hellenic nappes front at sea from Figure 5 of Aubouin et al.
1426 (1976). Backstop outer limit in green and inner limit in dashed blue after Huguen et al.
1427 (2006). Note the remarkable agreement of expected Hellenic nappe front with the outer
1428 limit of the backstop. Dashed black: probable extension of backstop below the
1429 Mediterranean Ridge.

1430 Figure 4

1431 Gravity Gradient Map from Sandwell et al. (2014) outlining the passage of the Sachpazi
1432 fractures F1 to F9 below the backstop. This leads to the proposal of additional fractures F10
1433 to F12. Backstop outer limit in green and inner limit in dashed blue after Huguen et al.
1434 (2006). Dashed black: probable extension of the backstop below Mediterranean Ridge.
1435 Dashed red small circles about the Hellenic Scarp rotation pole.

1436 Figure 5

1437 Oblique Mercator projection of Figure 4 with pole of projection at location of Hellenic Scarp
1438 rotation pole. In this projection, faults should be along lines of latitude if they are small
1439 circles about the pole. F1 to F9 in continuous red lines are the Sachpazi faults. Dashed lines
1440 are projected extensions of the faults on both sides of the backstop to indicate the locations
1441 of the faults below the backstop. F10 to F12 are possible additional faults suggested by
1442 trends in backstop. Backstop outer limit in green and inner limit in dashed blue after
1443 Huguen et al. (2006). Dashed yellow: probable extension of backstop below Mediterranean
1444 Ridge.

1445 Figure 6

1446 Schematic reconstruction at the beginning of the Hellenic subduction. Black: positions of
1447 present coast lines and of backstop and of the Sachpazi faults F1 to F9. Red, reconstructed
1448 positions at the beginning of the Hellenic subduction. Violet: intermediate stage of
1449 reconstruction obtained by rotating Greece south of D and limits of backstop by 18° about
1450 HS/NU pole. This puts the outer limit of backstop in contact with extremities of Sachpazi
1451 faults. To obtain the final position an additional rotation of 37° with respect to point D is
1452 added. Note the good agreement between the positions of the reconstructed margin and
1453 the one of the EMNT obtained independently earlier.

1454 Figure 7

1455 Five successive reconstructions at equal angle intervals showing the flow lines from
1456 present to pre-subduction stage.

1457 Figure 8

1458 Oblique Mercator projection of bathymetry similar to Figure 5 with pole of projection at
1459 location of Hellenic Scarp rotation pole. In this projection, faults should be along lines of
1460 latitude if they are small circles about the pole. F1 to F9 in continuous red lines are the
1461 Sachpazi faults. Dashed lines are projected extensions of the faults on both sides of the
1462 backstop to indicate the locations of the faults below the backstop. F10 to F12 are possible
1463 additional faults suggested by trends in backstop. Backstop outer limit in green and inner
1464 limit in dashed blue after Huguen et al. (2006). Dashed yellow: probable extension of
1465 backstop below the Mediterranean Ridge. See discussion in text.

1466 Figure 9

1467 Triangulation displacement vectors with respect to Attica from Le Pichon et al. (1995,
1468 Figures 4 and 6). Note how western Peloponnesus moves westward with respect to Attica
1469 due to a distributed N/S extensional system: Argos (Ar), Lakonia (Lak) and Messinia (Me)
1470 are identified within this system. Co (Gulf of Corinth), Pat (Gulf of Patras), Tri (Lake of
1471 Trikhonis), Am (Gulf of Amvrakia), KF (Kefalonia dextral fault)

Fig. 1. Bathymetry of the Eastern Mediterranean after Campistron-Brosolo and Mascle (2015). Fractures F1 to F9 within the Hellenic slab (heavy white lines and heavy dashed white lines) are after Sachpazi et al. (2016a and 2016b). Proposed additional fractures F10 to F12 (white dots). Black lines, small circles about the HS/NU rotation pole (red small circle). Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Thick dashed black lines: probable extension of backstop below Mediterranean Ridge. Continuous coloured contours on top of the slab are after Sachpazi et al. (2016a and 2016b). Dashed 20 to 60 km contours are after Vernant et al. (2014). Dashed 100 and 140 km black contours are after Shaw and Jackson (2010). Violet arrow is slip vector of the 2008 Methoni earthquake after Howell et al. (2017a). Black arrows are slip vectors of subduction earthquakes after Shaw and Jackson (2010). KF, Kefalonia fault; NAF, North Anatolian fault. Fault plane solutions of earthquakes within slab (red, P axis, black, T axis) are after Shaw and Jackson (2010). Thick white lines, approximate direction of P axis within slab. Black circle is the AN/NU pole (32.71°N, 34.2°E, 1.29°/Ma, Le Pichon and Kreemer 2010).

307x217mm (300 x 300 DPI)

Fig. 2. Free air gravity map of Eastern Mediterranean Sea from Sandwell et al. (2014). Fractures F1 to F9 within the Hellenic slab (white lines and heavy dashed white lines) are after Sachpazi et al. (2016a and 2016b). Proposed additional fractures F10 to F12 (white dots). Dashed red small circles about HS/NU rotation pole shown by red dot. Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Dashed black: probable extension of backstop below Mediterranean Ridge. "+" identify three different gravity highs related to the flexure of the subducting Eastern Mediterranean lithosphere. Red faults on Medina axial gravity high are a set or reverse faults active during Tortonian (Gallais et al. 2011). Heavy black line is small circle following the Lybian Egyptian Transform margin centered on 52.0°N, 39.0°E. Heavy black dashed lines are small circles about same center identifying the Apulian Transform and the EMNT.

307x217mm (300 x 300 DPI)

Fig. 3. Aegean extensional areas and directions of Aegean basin development after map 6 of Chamot-Rooke et al. (2005). Hellenic nappes front at sea from Figure 5 of Aubouin et al. (1976). Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Note the remarkable agreement of expected Hellenic nappe front with the outer limit of the backstop. Dashed black: probable extension of backstop below the Mediterranean Ridge.

307x217mm (300 x 300 DPI)

Fig. 4. Gravity Gradient Map from Sandwell et al. (2014) outlining the passage of the Sachpazi fractures F1 to F9 below the backstop. This leads to the proposal of additional fractures F10 to F12. Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Dashed black: probable extension of the backstop below Mediterranean Ridge. Dashed red small circles about the Hellenic Scarp rotation pole.

211x175mm (300 x 300 DPI)

Fig. 5. Oblique Mercator projection of Figure 4 with pole of projection at location of Hellenic Scarp rotation pole. In this projection, faults should be along lines of latitude if they are small circles about the pole. F1 to F9 in continuous red lines are the Sachpazi faults. Dashed lines are projected extensions of the faults on both sides of the backstop to indicate the locations of the faults below the backstop. F10 to F12 are possible additional faults suggested by trends in backstop. Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Dashed yellow: probable extension of backstop below Mediterranean Ridge.

198x146mm (300 x 300 DPI)

Fig. 6. Schematic reconstruction at the beginning of the Hellenic subduction. Black: positions of present coast lines and of backstop and of the Sachpazi faults F1 to F9. Red, reconstructed positions at the beginning of the Hellenic subduction. Violet: intermediate stage of reconstruction obtained by rotating Greece south of D and limits of backstop by 18° about HS/NU pole. This puts the outer limit of backstop in contact with extremities of Sachpazi faults. To obtain the final position an additional rotation of 37° with respect to point D is added. Note the good agreement between the positions of the reconstructed margin and the one of the EMNT obtained independently earlier.

306x203mm (300 x 300 DPI)

Fig. 7. Five successive reconstructions at equal angle intervals showing the flow lines from present to pre-subduction stage.

306x201mm (300 x 300 DPI)

Fig. 8. Oblique Mercator projection of bathymetry similar to Figure 5 with pole of projection at location of Hellenic Scarp rotation pole. In this projection, faults should be along lines of latitude if they are small circles about the pole. F1 to F9 in continuous red lines are the Sachpazi faults. Dashed lines are projected extensions of the faults on both sides of the backstop to indicate the locations of the faults below the backstop. F10 to F12 are possible additional faults suggested by trends in backstop. Backstop outer limit in green and inner limit in dashed blue after Huguen et al. (2006). Dashed yellow: probable extension of backstop below the Mediterranean Ridge. See discussion in text.

198x146mm (300 x 300 DPI)

Fig. 9. Triangulation displacement vectors with respect to Attica from Le Pichon et al. (1995, Figures 4 and 6). Note how western Peloponnesus moves westward with respect to Attica due to a distributed N/S extensional system: Argos (Ar), Lakonia (Lak) and Messinia (Me) are identified within this system. Co (Gulf of Corinth), Pat (Gulf of Patras), Tri (Lake of Trikhonis), Am (Gulf of Amvrakia), KF (Kefalonia dextral fault)

249x231mm (300 x 300 DPI)